




Agriculture
Canada

BUDD'S FLORA

OF THE
CANADIAN PRAIRIE
PROVINCES



LIBRARY - BIBLIOTHÈQUE



**Agriculture
Canada**

**DEPARTMENTAL LIBRARY
BIBLIOTHÈQUE DU MINISTÈRE
ÉDIFICE SIR JOHN CARLING BLDG.
OTTAWA ONTARIO
K1A 0C5**

LIBRARY - BIBLIOTHÈQUE

Digitized by the Internet Archive
in 2011 with funding from
Agriculture and Agri-Food Canada – Agriculture et Agroalimentaire Canada

BUDD'S FLORA

OF THE CANADIAN PRAIRIE PROVINCES

Revised and enlarged by
J. Looman
Research Station
Swift Current, Saskatchewan
and
K.F. Best
Research Station
Regina, Saskatchewan

Research Branch
Agriculture Canada
Publication 1662 1987

© Minister of Supply and Services Canada 1987

Available in Canada through

Authorized bookstore agents
and other bookstores

or by mail from

Canadian Government Publishing Centre
Supply and Services Canada
Hull, Quebec, Canada K1A 0S9

Catalogue No. A53-1662/1979

Canada: \$38.50

ISBN 0-660-10233-1

Other Countries: \$46.20

Price subject to change without notice

First printed 1979

Reprinted 1981

Revised 1987



A tribute to A. C. Budd

Archibald Charles Budd was born in London, England, on 28 April 1889. He attended Bellenden Road Higher Grade School, where as top boy he earned a merit certificate. He took commercial training at Camberwell Grammar School, and attended Choumert Road Evening School and Kings College. From 1905 until 1909 he clerked in the Civil Service at the Customs House and the General Post Office.

A prize of 250 pounds for winning a limerick contest made it possible for Mr. Budd to emigrate to Canada in 1910 as a land seeker. He first worked as a farmhand at Waldeck, Saskatchewan, and later homesteaded and farmed at Burnham until 1922.

A lifelong interest in nature study helped him when he began working at the Dominion Experimental Station at Swift Current, Saskatchewan, in 1926. As an assistant to the late Dr. Sidney Barnes of the Soils Research Laboratory, Mr. Budd assisted in studies on the taxonomy of weeds and the physiology of weed seeds until 1944. Although he had no formal training in botany, Mr. Budd was a keen student of taxonomy. An avid reader of botanical works and an astute observer, he soon became an authority on the prairie flora.

From 1944 until his retirement in 1957, Mr. Budd was Range Botanist and he made an extensive study of plant life in the prairie region. As curator of the herbarium, he built up one of the finest plant collections in Western Canada. Mr. Budd also helped train many professional agrologists, who are managing and appraising rangeland and are carrying out ecological studies requiring a knowledge of applied botany.

For many years Mr. Budd was a member and director of the Saskatchewan Natural History Society. He often contributed articles to *The Blue Jay*, the magazine published by the Society. He assisted in the establishment of a herbarium in the Museum of Natural History in Regina, Saskatchewan, and he built up a large personal collection of plants, which he later donated to the Swift Current Museum.

In 1949 he prepared a preliminary draft of a key to prairie plants, which he called *Flora of the Farming and Ranching Areas of the Canadian Prairies*. Mimeographed copies were enthusiastically received by amateurs and professional botanists throughout Western Canada. In 1952 a revised and expanded version entitled *Plants of the Farming and Ranching Areas of the Canadian Prairies* was published by the Experimental Farms Service of the Canada Department of Agriculture. After many revisions and the addition of more drawings, *Wild Plants of the Canadian Prairies* was published in 1957, the year of Mr. Budd's retirement. Until his death on 30 December 1960, he cooperated on revisions that were incorporated in the editions printed in 1964 and 1969.

As a tribute to Archie and his faithful and devoted work, which was the basis for this edition, the title of this first major revision in which he did not have a direct input has been changed to:

Budd's Flora of the Canadian Prairie Provinces

J. Looman and K. F. Best

Contents

A tribute to A. C. Budd	3
Preface	7
Zones of vegetation	9
How to use the plant key	12
Keys to main groups	20
Genera and species	32
Tabular classification	790
Glossary	793
The spelling of common names	803
Additions and corrections to the 1979 edition	804
Index to common names	805
Index to scientific names	831



Preface

This publication, like its predecessor, *Wild Plants of the Canadian Prairies*, is intended for amateur botanists, agricultural representatives, farmers, and ranchers. For this reason, emphasis has been placed on keys for identification of species. However, the book will also be useful to professional botanists.

This publication is not just a revised edition of *Wild Plants of the Canadian Prairies*; it has also been very much expanded. The scope of the work has been greatly increased to include the areas of the Boreal forests and the Rocky Mountains. These additions were considered necessary because, since the last revision of *Wild Plants*, many parts of the Prairie Provinces have become accessible to travelers. The sections on grasses, sedges, and willows have been completely rewritten and expanded.

An attempt has been made to include all native species presently known to occur in the Prairie Provinces, as well as species, native or introduced, which are likely to be found along roads, railroads, rivers, and lakeshores; but undoubtedly some species have been missed. Introduced species, which occur occasionally as escapes from cultivation but do not persist outside the garden environment, are mentioned but not described.

Further changes include measurements given in metric units (SI, International System of Units) and photographs for illustrative purposes. In addition, rare plants have been identified as such.

The basis for rarity is the number of occurrences in the Prairie Provinces, as well as the number of plants per occurrence. Thus, a plant indicated as rare is known either from only a few locations in an area, or from several locations, but each with few plants. Very rare plants are known from very few places—perhaps only one or two—and are not abundant even there.

Please do not collect very rare and rare species. If identification of a species is uncertain, send a set of good photographs to a professional botanist for positive identification.

The 1987 revision was undertaken by Dr. J. Waddington of the Swift Current Research Station. It includes changes to the key for the chloripetalous dicotyledons (p. 26), the sympetalous dicotyledons (p. 29), the Leguminosae (p. 463), and group 3 of the Compositae (p. 704). A list of additions and corrections to the 1979 edition is provided on p. 804, including cross-references to the increasingly common European–Russian classification of perennial Triticeae. Finally, the index has been modified to reflect these changes.

Metric conversion:

0.9 m = 1 yard

15 cm = 6 inches

2.5 cm = 1 inch

25 mm = 1 inch




Fig. 1. Vegetation zones.

Zones of vegetation

Manitoba, Saskatchewan, and Alberta occupy an area of almost 2 000 000 km². As a result of differences in climate and soil type, several vegetation types can be distinguished, each with some species occurring mainly there. Therefore, these vegetation types can be used to designate species distribution. The major types outlined in Fig. 1 follow.

The Prairies

The Prairies include the almost treeless grasslands in the southern parts of Saskatchewan and Alberta. Rainfall in this area is low, summers are warm, and the vegetation is composed mainly of drought-tolerant species. Most of the grassland is short-grass prairie, with spear grass, blue grama, June grass, and other low-growing plants predominating. On eroded areas several uncommon, specially adapted cushion plants can be found, and the three native cacti belong to this vegetation type. Transitions to the midgrass prairie occur at the boundaries of the Prairies and on slopes with north to east exposures. Shrubs and trees occur almost entirely in deep coulees and ravines.

The Parklands

The Parklands form a broad belt around the Prairies. They consist of open grassland alternating with tree groves. This belt can be divided into three sections: eastern, central, and western Parklands. In the eastern Parklands, the grassland is tall-grass prairie, and big bluestem, porcupine grass, prairie cord grass, and switch grass predominate. Tree groves contain bur oak, ash, Manitoba maple, and balsam poplar. The central Parklands have midgrasses in the grassland openings, which consist of rough fescue, western porcupine grass, and Hooker's oat grass, and on the light high-lime soils, little bluestem. Tree groves consist predominantly of aspen poplar and often include some willows. In the western Parklands, grassland is midgrass prairie, containing fescues, spear grass, and poverty oat grass. Tree groves are predominantly aspen poplar, often interspersed with willows and balsam poplar. Drought is not as prevalent in the Parklands as in the Prairies; soils are Dark Brown in the southern parts and Black in the more northern parts.

The Boreal forest

The Boreal forest is the vast area covered with mostly coniferous forest situated north of the Parklands. Its southern margin intergrades with the Parklands, but farther north the Boreal forest is dominated by white spruce on

uplands, black spruce on lowlands, and jack pine on light sandy soils. Areas that have been logged in recent years are covered with deciduous forest, in which aspen poplar, balsam poplar, and birch are the main species. Wetlands are sedge–reed swamps, grading into muskeg, where large tussocks of peat moss, Labrador tea, willows, swamp birch, larch, and black spruce occur. An area in southeastern Manitoba varies from the general area in that it supports white pine, white cedar, ground hemlock, and other eastern or southeastern species. In Manitoba and northeastern Saskatchewan the Boreal forest includes outcrops of Precambrian rock, on which several ferns and saxifrages with numerous species of lichens can be found.

Rocky Mountains

The western boundary of the Parklands and part of the Boreal forest grade into the Rocky Mountains. In the mountains the climate varies sharply with differences in elevation. In the lower montane zone to about 1500 m, fescue prairie occupies exposed slopes. Coniferous forest of spruce, lodgepole pine, Douglas-fir, and western red cedar, aspen poplar woods, or mixed forest are found on less exposed slopes and in ravines. At higher altitudes the open grasslands gradually disappear, to make place for vegetation dominated by shrubs, mainly willows and ground birch, and for the coniferous forest. At altitudes of about 2200–2400 m, the upper tree limit is reached, and alpine meadows, with many low sedges and grasses, mosses, and lichens predominating. The Rocky Mountains can be divided into the southern section, reaching north to approximately the Calgary–Banff line, and the northern section above this line. Several plant species, such as common camas, bear-grass, Pursh's silky lupine, and balsamroot, occur only in the southern section.

Exceptional areas

Besides the major vegetation types, there are several types that occur in areas where the climate or soil differs from that prevailing in the area dominated by a major vegetation type. The most important ones follow.

Cypress Hills – Wood Mountains

This ridge of hills rises above the surrounding Prairies with elevations ranging from about 870 m to 1350 m. The ridge is interrupted repeatedly by gaps of lowland, but in the lower elevations the vegetation is similar throughout. Midgrass prairie intergrades with fescue prairie and aspen poplar groves in coulees and ravines. At elevations above 1200 m the grasslands become pure fescue prairie, and above 1300 m, coniferous forest, similar to that of the southern Rocky Mountains, occurs.

Riding Mountain – Duck Mountain

These areas of hills and plateaus rise 200–300 m above the surrounding plains in western Manitoba and eastern Saskatchewan. The vegetation is mainly coniferous forest, similar to that of the Boreal zone to the north. However, exposed areas and slopes support Parkland vegetation interspersed with small areas of fescue prairie vegetation.

Peace River district

With a climate warmer and drier than what is customary for this latitude, the Peace River district offers a Parkland landscape, with southern exposed slopes and other exposed areas covered with midgrass prairie, dominated by spear grass. Somewhat eroded areas contain brittle prickly-pear, as well as several other species characteristic of the southern prairies. Tree groves are dominated by aspen poplar, with willows, chokecherries, pin cherries, and particularly Saskatoon berries often plentiful.

Sand dune areas

Several areas of sandy soils and mobile sand dunes are characterized by a vegetation type that is quite different from that of the heavier soils surrounding them. The vegetation of sand dune areas depends on the zone in which they occur. The Middle, Great, and Little Sandhills in southeastern Alberta and southwestern Saskatchewan are in the Prairies, and are characterized by short-grass prairie on stabilized dry sand; sand grass, sand dropseed, Indian rice grass, and sand dock in open sand; little bluestem on moist calcareous sand; and large areas of western snowberry, silverberry, choke cherries, and aspen poplar in moist dune valleys. Several other species of shrubs occur in small groups.

In the Parklands, sand areas are mostly in midgrass prairie, with rough fescue and western porcupine grass dominating the dry locations and little bluestem the moister locations. Aspen poplar mixed with some balsam poplar and willows occupy the sites that have good moisture conditions. In the eastern Parklands, tall grasses dominate the grasslands in the sand dunes. These include big bluestem, porcupine grass, and switch grass. Willows, aspen poplar, and bur oak form the tree groves, with white spruce gradually taking over if the tree cover remains undisturbed.

Saline areas

Many large and small areas with poor drainage and areas around lakes and sloughs have a high to very high salt content, often visible by a white salt

incrustation when they are dry. The vegetation on these areas is made up of species with special adaptations for the extreme saline conditions. Most of the species may also be found along the seacoasts, or they have affinities with species found there: sea-blite, red samphire, seaside arrow-grass, and various salt grasses, which are indigenous in these areas.

How to use the plant key

Plants are classified into various groups according to their structure and method of reproduction. The first of these are divisions, which are separated into subdivisions. The subdivisions are divided into classes, and the classes into families. Families are split into genera, and each genus contains one or more species. The species is sometimes further divided into forms, subspecies, and varieties.

The keys that follow contain paired contrasting statements of particular characteristics, and a plant must agree with one of them. Find out which of the pair it agrees with and then continue to the name or number following that statement until you find the family name. Go through the key to the family, then the genus, and finally the species. Continue to follow the statements until you find the species.

Imagine that the plant in the sketch (Fig. 2) is actually in front of you and that you have no idea what it is and you want to identify it.

Turn to p. 20. Begin at the first key, and do not worry about the scientific names. If there are any terms or expressions that you do not understand, turn to the Glossary and the diagrams.

- Clause 1. Because it obviously has flowers and will undoubtedly have seeds, it is in the SPERMATOPHYTA division. Go to the next entry, numbered 2.
- Clause 2. It is certainly not an evergreen tree bearing cones, therefore it must be an ANGIOSPERM. Go to Clause 3.
- Clause 3. Because the leaves are not parallel-veined, the plant must belong to the DICOTYLEDONEAE.

Now look through the section "KEY TO THE FAMILIES" and find the portion headed "Class: DICOTYLEDONEAE" (p. 23). If you look carefully at the flower, you will find that the petals are not entirely separate but are joined near the base, therefore the plant is a SYMPETALOUS dicotyledon, section C. Turn to the heading of that section on p. 29.

- Clause 1. It is obviously a herb and not a shrub or tree; go to Clause 8.
- Clause 8. It is a green plant; turn to Clause 11.
- Clause 11. The flowers are neither in heads nor in spikes; go on to Clause 19.
- Clause 19. The ovary is superior, as you will find by referring to the Glossary and the diagram of flower parts; go to Clause 25.
- Clause 25. The petal portion (corolla) is regular (*see* Glossary); go to Clause 28.



Fig. 2. A specimen (*Lysimachia ciliata* L.) illustrating the use of the key.

- Clause 28. It does not have a milky juice; go to Clause 30.
Clause 30. It is certainly not a twining plant; go to Clause 31.
Clause 31. The stamens are directly in front of the petals, therefore it should belong to the PRIMULACEAE, or primrose family, which is described on p.580.

Turning to p.580, under PRIMULACEAE, start again at Clause 1. Note that the leaves are not all basal, thus go to Clause 5.

- Clause 5. The leaves are opposite; go to Clause 6.
Clause 6. Petals are present; go to Clause 7.
Clause 7. The stem leaves are normal; go to Clause 8.
Clause 8. The flowers are yellow, therefore it appears to belong to the genus *Lysimachia*, or loosestrife.

The genera are in alphabetical order; you will find *Lysimachia* on p.583. Again begin with Clause 1 of the key to the species, and note that the plant is flowering, and does not have bulblets in the upper axils of the leaves, so go to Clause 2.

- Clause 2. The flowers are borne on separate stalks in the leaf axils; go to Clause 3.
Clause 3. The leaves are lanceolate (*see* diagram of leaf shapes, p. 15) and rounded at the base; go to Clause 4.
Clause 4. The petioles of the lower leaves are about 1 cm long and ciliate, therefore it is apparently a specimen of *Lysimachia ciliata* L., or fringed loosestrife. Now read the description of that species and you will find that it agrees in every way with the specimen. Apparently the keying has been correct.

The *L.* after the scientific name means that the Swedish botanist Linnaeus gave that name to this plant. The other name is a synonym, or another name, given to it by another botanist. In this case, because Rafinesque thought that the plant should be in a separate genus, *Steironema*, he took the specific name from Linnaeus and placed it after *Steironema*, but divided the credit for the name with Linnaeus, and therefore the authority is (L.) Raf. Because later botanists agreed with Linnaeus, Rafinesque's name for this plant became a synonym.




Fig. 3. Shapes of simple leaves.





Fig. 4. Types of divided leaves.



A.C. Budd

Fig. 5. Leaf variations.




Fig. 6. Types of inflorescences.




Fig. 7. Flower parts.




Fig. 8. Types of flowers.




Fig. 9. Types of fruits.

Keys to main groups

Key to Divisions, Subdivisions, and Classes

1. Plants without true flowers, reproducing by spores. Division: PTERIDOPHYTA, p. 32
Plants with flowers, reproducing by seeds. Division: SPERMATOPHYTA, p. 56 2
2. Plants bearing flowers having neither styles nor stigmas, the naked ovules (seeds) on the upper side of a scale. Mostly evergreen trees and shrubs bearing cones or cone-like fruit. Subdivision: GYMNOSPERMAE, p. 56
Plants bearing flowers with the ovules enclosed in an ovary with styles or stigmas, and bearing seeds in a closed ovary. Subdivision: ANGIOSPERMAE, p. 61 3
3. Stems, when present, without central pith or annular layers; leaves usually parallel-veined; flower parts generally in threes or multiples of three; seeds with only one cotyledon or seed leaf. Class: MONOCOTYLEDONEAE, p. 61
Stems with a central pith, or if woody, the wood generally arranged in annular layers; leaves net-veined; seeds with two cotyledons or seed leaves. Class: DICOTYLEDONEAE, p. 279

Key to the Families

Division: PTERIDOPHYTA—ferns and fern allies

1. Leaves reduced to scales, whorled at the internodes of strongly jointed stems, forming a sheath. EQUISETACEAE, p. 47
Leaves reduced to green scales, awl-shaped, or larger, not forming a sheath; stems not strongly jointed. 2
2. Leaves linear, grass-like, sheathing at base. ISOETACEAE, p. 50
Leaves not linear or grass-like. 3
3. Leaves scale-like or awl-shaped. 4
Leaves larger and broader. 5
4. Sporangia borne in the upper axils of ordinary leaves, or in terminal terete strobiles; spores uniform in size. LYCOPODIACEAE, p. 51
Sporangia borne in terminal four-sided strobiles; spores of two different sizes. SELAGINELLACEAE, p. 55

- 5. Fronds (leaves) long-petioled, palmately divided into 4 leaflets. MARSILEACEAE, p. 47
Fronds not divided into 4 leaflets. 6
- 6. Plants with a single stalk, bearing a sterile frond below, a fertile one above; sporangia arranged in two rows. OPHIOGLOSSACEAE, p. 32
Plants not as above. 7
- 7. Sporangia naked, borne on much modified divisions at the tip or middle of fronds. OSMUNDACEAE, p. 35
Sporangia covered, borne on fertile fronds similar to the sterile ones or on modified fronds. POLYPODIACEAE, p. 35

Division: SPERMATOPHYTA—seed-bearing plants

Subdivision: GYMNOSPERMAE—plants with naked seeds

- 1. Cones reduced to a single ovule, fruit a one-seeded berry. TAXACEAE, p. 56
Cones many-seeded, fleshy or woody. 2
- 2. Leaves scale-like or awl-shaped, often overlapping; opposite or whorled. CUPRESSACEAE, p. 60
Leaves linear or needle-shaped, not overlapping; alternate or clustered. PINACEAE, p. 56

Subdivision: ANGIOSPERMAE—plants with covered seeds

Class: MONOCOTYLEDONEAE—monocotyledons

Stems, when present, without central pith or annual layers; leaves usually parallel-veined; flower parts mostly in threes or multiples of three; seeds with only one cotyledon (seed leaf).

- 1. Plants aquatic, floating or submerged, with floating leaves or emersed inflorescence. 2
Plants not aquatic, or if growing in water, most of the plant emersed. 4
- 2. Plants small, floating, leaf-like; without differentiation between stems and leaves. LEMNACEAE, p. 245
Plants larger, generally rooting; clear differentiation between stem and leaves. 3
- 3. Perianth absent or single and inconspicuous; plants monoecious. ZOSTERACEAE, p. 65

Perianth present, in two whorls; plants dioecious.	HYDROCHARITACEAE, p. 73	
4. Perianth absent or inconspicuous.		5
Perianth present, conspicuous, in two series of 3 segments, often brightly colored.		11
5. Inflorescence a dense, single, cylindrical spike 8–15 cm long, 1–2.5 cm thick.		6
Inflorescence not as above; if a single spike, less than 1 cm thick.		7
6. Spike fleshy, subtended by a large bract or spathe.	ARACEAE, p. 246	
Spike not fleshy, bractless.	TYPHACEAE, p. 61	
7. Flowers in globular heads, the upper ones staminate, the lower ones pistillate.	SPARGANIACEAE, p. 63	
Flowers not in globular heads.		8
8. Perianth in two whorls, each of three segments.		9
Perianth not in two whorls.		10
9. Flowers in spike-like racemes; perianth of greenish sepals and petals.	JUNCAGINACEAE, p. 68	
Flowers in branched inflorescences; perianth segments scale-like.	JUNCACEAE, p. 246	
10. Flowers enclosed in two-ranked bracts with the lowest (glumes) empty; perianth reduced to minute lodicules or lacking; culms usually hollow; leaves two-ranked.	GRAMINEAE, p. 73	
Flowers solitary in the axil of a single bract (scale); perianth reduced to bristles, or scales, or lacking; culms usually solid; leaves three-ranked.	CYPERACEAE, p. 181	
11. Perianth small, consisting of 3 green sepals and 3 deciduous whitish petals.	ALISMACEAE, p. 69	
Perianth usually large, consisting of sepals and petals usually alike, but petals not deciduous, variously colored.		12
12. Flowers irregular, stamens 1 or 2.	ORCHIDACEAE, p. 270	
Flowers regular, stamens 3 or 6.		13
13. Stamens 3, flowers borne in a spathe.	IRIDACEAE, p. 268	
Stamens 6, flowers not borne in a spathe.		14
14. Ovary inferior.	AMARYLLIDACEAE, p. 268	
Ovary superior.		15
15. Leaves sheathing the stem; inflorescence a cyme, subtended by strongly sheathing bracts.	COMMELINACEAE, p. 246	
Leaves not sheathing the stem; inflorescence not subtended by strongly sheathing bracts.	LILIACEAE, p. 256	

Class: DICOTYLEDONEAE—dicotyledons

Stems with a central pith, or, if woody, the wood generally arranged in annual layers; leaves net-veined; seeds with two cotyledons (seed leaves).

- A. APETALOUS Flowers with only one floral ring, with sepals but not petals (p. 23).
- B. CHORIPETALOUS Flowers with two floral rings, and with each petal distinct from the others (p. 25).
- C. SYMPETALOUS Flowers with two floral rings, but with the petals wholly or partly united forming a tube or bell (p. 29).

A. Apetalous dicotyledons
(flowers with only one floral ring, with sepals but not petals)

- 1. Herbs. 2
 Shrubs or trees. 23
- 2. Plants parasitic on branches of trees;
 leaves reduced to scales. LORANTHACEAE, p. 308

 Plants not parasitic on trees; rooted in
 soil or water. 3
- 3. Water plants. 4
 Land plants. 7
- 4. Leaves all simple. 5
 Some or all leaves divided into filiform
 segments. 6
- 5. Leaves opposite. CALLITRICHACEAE, p. 514
 Leaves whorled. HALORAGACEAE, p. 543
 (*Hippuris*)
- 6. Plants free-floating, rootless. CERATOPHYLLACEAE, p. 361
 Plants rooted at the bottom. HALORAGACEAE, p. 544
 (*Myriophyllum*)
- 7. Stamens or pistils in separate flowers on
 the same plant. 8
 Stamens and pistils in each flower. 11
- 8. Plants with milky sap; fruit 3-seeded,
 splitting open at maturity. EUPHORBIACEAE, p. 511
 Plants without milky sap. 9
- 9. Plants twining, climbing; female flowers
 and fruit in catkin-like spikes. CANNABINACEAE, p. 304
 Plants not climbing; fruit not catkin-like. 10
- 10. Stigmas 2; flowers without calyx; fruit
 enclosed in 2 bracts. CHENOPODIACEAE, p. 326
 (*Atriplex*)

Stigmas 1; flowers with calyx; fruit enclosed by 2–4 perianth segments; leaves often bearing stinging hairs.	URTICACEAE, p. 304	
11. Plants stemless with a single pair of large reniform leaves; calyx 3-lobed.	ARISTOLOCHIACEAE, p. 308	
Plants with stems or stemless, but with more than one pair of leaves.		12
12. Leaves with stipules free or sheathing the stem.		13
Leaves without stipules.		14
13. Leaves opposite; stipules membranous; plant cushion-like.	CARYOPHYLLACEAE, p. 354 (<i>Paronychia</i>)	
Leaves alternate; stipules forming a sheath above nodes.	POLYGONACEAE, p. 308	
14. Leaves opposite.		15
Leaves alternate; lower ones sometimes opposite.		18
15. Leaves scale-like; stem fleshy; flowers imbedded in stem; plants of saline areas.	CHENOPODIACEAE, p. 338 (<i>Salicornia</i>)	
Leaves not scale-like; calyx colored.		16
16. Plants climbing.	RANUNCULACEAE, p. 369 (<i>Clematis</i>)	
Plants not climbing.		17
17. Erect herbs; inflorescence in panicles or clusters.	NYCTAGINACEAE, p. 344	
Low plants; flowers solitary in leaf axils.	PRIMULACEAE, p. 583 (<i>Glaux</i>)	
18. Fruit fleshy when ripe.	SANTALACEAE, p. 305	
Fruit dry when ripe.		19
19. Leaves orbicular or reniform; sepals 4.	SAXIFRAGACEAE, p. 421 (<i>Chrysplenium</i>)	
Leaves not orbicular or reniform.		20
20. Leaves compound or simple, but very deeply divided.	RANUNCULACEAE, p. 364 (<i>Anemone</i>)	
Leaves not compound and not very deeply divided.		21
21. Perianth segments 6, in two series; flowers colored.	POLYGONACEAE, p. 309 (<i>Eriogonum</i>)	
Perianth segments 2–5; flowers small, numerous, greenish.		22
22. Bracts and perianth segments dry and membranous.	AMARANTHACEAE, p. 342	

Membranous bracts absent; perianth segments greenish.	CHENOPODIACEAE, p. 324	
23. Male flowers, at least, in catkins or aments.		24
Flowers not in catkins or aments.		27
24. Seeds each with a tuft of hairs.	SALICACEAE, p. 279	
Seeds without a tuft of hairs.		25
25. Styles 3 or more; fruit an acorn.	FAGACEAE, p. 303	
Styles 2; fruit not an acorn.		26
26. Three flowers in axil of each bract of male catkin.	BETULACEAE, p. 299	
One flower in axil of each bract of male catkin; fruit a nut.	BETULACEAE, p. 301 (<i>Corylus</i>)	
27. Climbing plants; flowers with colored sepals; fruit with a persistent feathery style.	RANUNCULACEAE, p. 369 (<i>Clematis</i>)	
Plants not climbing; fruit without a feathery style.		28
28. Leaves pinnately compound.		29
Leaves simple.		30
29. Fruit a double samara; leaflets mostly 3–5.	ACERACEAE, p. 515	
Fruit a single samara; leaflets mostly 5–11.	OLEACEAE, p. 587	
30. Leaves linear, evergreen, 2.5–7 mm long; plants with decumbent stems.	EMPETRACEAE, p. 514	
Leaves wider, deciduous; plants with erect stems.		31
31. Fruit fleshy when ripe.		32
Fruit dry when ripe.		33
32. Leaves silvery or brownish scurfy.	ELAEAGNACEAE, p. 533	
Leaves aromatic, not silvery or brownish scurfy.	MYRICACEAE, p. 299	
33. Shrubs without stipules; branches spiny.	CHENOPODIACEAE, p. 339 (<i>Sarcobatus</i>)	
Trees with stipules; flowers appearing before leaves.	ULMACEAE, p. 303	

B. Choripetalous dicotyledons

(flowers with two floral rings, and with each petal distinct from the others)

1. Succulent, spiny plants with leaves absent or scale-like, inconspicuous.	CACTACEAE, p. 531	
Plants with leaves.		2

2. Leaves all basal, and tubular or with tentacles for catching insects.	3
Leaves normal.	4
3. Plants with solitary flowers; leaves tubular with lid-like lobe on top.	SARRACENIACEAE, p. 418
Plants with flowers in racemes; leaves flat with glandular tentacles.	DROSERACEAE, p. 418
4. Herbs.	5
Shrubs or trees.	41
5. Plants aquatic; leaves submerged or floating.	6
Plants terrestrial or semiaquatic.	8
6. Leaves mostly floating, large, suborbicular or reniform.	NYMPHAEACEAE, p. 361
Leaves mostly submerged, small, coarsely to finely dissected.	7
7. Flowers pedicellate, mostly white or yellow, perfect.	RANUNCULACEAE, p. 362
Flowers sessile, axillary, monoecious.	HALORAGACEAE, p. 543
8. Plants with a single ternately compound stem leaf, appearing as 3 long-petioled leaves.	BERBERIDACEAE, p. 384 (<i>Caulophyllum</i>)
Plants with more than one leaf.	9
9. Plants with colored milky juice.	PAPAVERACEAE, p. 384
Plants not with colored milky juice.	10
10. Leaves opposite, whorled, or basal.	11
Some or all leaves alternate.	18
11. Low plants with an involucre of 4 petal-like bracts; fruit a red drupe.	CORNACEAE, p. 563
Plants without large petal-like bracts.	12
12. Mud plants with small axillary flowers.	ELATINACEAE, p. 525
Terrestrial plants.	13
13. Styles single.	14
Styles 2 or more.	16
14. Ovary inferior, flowers 2- or 4-merous (parts).	ONAGRACEAE, p. 534
Ovary superior.	15
15. Petals 5, stamens 10.	PYROLACEAE, p. 567
Petals 4 or 6, stamens 12.	LYTHRACEAE, p. 534
16. Sepals 2.	PORTULACACEAE, p. 345
Sepals more than 2.	17
17. Leaves glandular-dotted, stamens united at base into 3–5 bundles.	HYPERICACEAE, p. 522
Leaves not glandular-dotted, stamens not united at base.	CARYOPHYLLACEAE, p. 347

18. Calyx irregular, some sepals smaller than others.	19
Calyx regular.	20
19. Leaves lobed or divided.	RANUNCULACEAE, p. 362
Leaves entire.	CISTACEAE, p. 525
20. Leaves with stipules.	21
Leaves without stipules, or having glands.	26
21. Stamens numerous, united into a column; leaves palmately veined.	MALVACEAE, p. 520
Stamens usually separate or partly so, not in a column.	22
22. Flowers irregular in shape.	23
Flowers regular in shape.	24
23. Corolla pea-like; fruit a legume.	LEGUMINOSAE, p. 462
Corolla with one petal spurred or sac-like; fruit a 3-valved capsule.	VIOLACEAE, p. 526
24. Stamens usually numerous; ovary of one or more carpels, either separate or enclosed by a fleshy receptacle.	ROSACEAE, p. 436
Stamens 5 or 10; ovary of 5 united carpels.	25
25. Leaves palmately divided; fruit with a long beak.	GERANIACEAE, p. 504
Leaves of 3 leaflets; fruit not beaked.	OXALIDACEAE, p. 505
26. Stamens usually more than 10.	27
Stamens 10 or fewer.	30
27. Carpels separate.	RANUNCULACEAE, p. 362
Carpels united.	28
28. Sepals 4; fruit a pod; annuals.	CAPPARIDACEAE, p. 386
Sepals 4–8; fruit a capsule opening at top.	29
29. Flowers few, large, solitary, and terminal; petals over 2.5 cm long.	LOASACEAE, p. 531
Flowers many, small, in terminal spikes.	RESEDACEAE, p. 417
30. Ovary inferior.	31
Ovary superior.	33
31. Parts of flowers in twos or fours.	ONAGRACEAE, p. 534
Flowers in umbels; parts of flowers mostly in fives.	32
32. Styles 5; fruit fleshy.	ARALIACEAE, p. 544
Styles 2; stems hollow.	UMBELLIFERAE, p. 545
33. Carpels 3–5.	34
Carpels 1 or 2.	37
34. Sepals 3; one petal-like and spurred; fruit an explosive capsule.	BALSAMINACEAE, p. 508
None of petals or sepals spurred; fruit not explosive.	35

35. Sepals 2.	PORTULACACEAE, p. 345	
Sepals 3–5.		36
36. Stamens 5, united at base; fruit a capsule.	LINACEAE, p. 507	
Stamens 8–10; fleshy-leaved plants.	CRASSULACEAE, p. 418	
37. Corolla irregular in shape.		38
Corolla regular in shape.		39
38. Petals 3; sepals 5, two large, colored, and petal-like.	POLYGALACEAE, p. 510	
Petals 4; sepals 2.	FUMARIACEAE, p. 385	
39. Petals 5; receptacle cup-shaped.	SAXIFRAGACEAE, p. 420	
Petals 4; sepals 4.		40
40. Stamens 6, of equal length; leaves of 3 leaflets.	CAPPARIDACEAE, p. 386	
Stamens usually 6, four long and two short.	CRUCIFERAE, p. 389	
41. Plants climbing.		42
Plants not climbing.		45
42. Leaves opposite; styles plumose in fruit.	RANUNCULACEAE, p. 369 (<i>Clematis</i>)	
Leaves alternate; styles not plumose in fruit.		43
43. Plants with tendrils; leaves simple or digi- tately compound.	VITACEAE, p. 519	
Plants with twining stems; leaves simple.		44
44. Leaves orbicular-cordate, peltate near the margin, 3- to 7-lobed.	MENISPERMACEAE, p. 384	
Leaves ovate-oblong, finely serrate.	CELASTRACEAE, p. 517	
45. Leaves opposite.		46
Leaves alternate.		49
46. Plants low shrubs.		47
Trees.		48
47. Leaves 15–35 mm long; flowers pink to red.	ERICACEAE, p. 575 (<i>Kalmia</i>)	
Leaves about 3 mm long; flowers yellow.	CISTACEAE, p. 525	
48. Trees with palmately lobed leaves; fruit a 2-seeded samara.	ACERACEAE, p. 515	
Shrubs or low trees; petals 4; fruit fleshy.	CORNACEAE, p. 563	
49. Leaves compound.		50
Leaves simple.		53
50. Leaflets leathery, spinose-tipped.	BERBERIDACEAE, p. 382	
Leaflets not leathery or spinose-tipped.		51
51. Petal (standard) solitary; fruit a legume.	LEGUMINOSAE, p. 463 (<i>Amorpha</i>)	
Petals 5; fruit not a legume.		52
52. Flowers perfect; stamens numerous.	ROSACEAE, p. 436	
Flowers dioecious or perfect; stamens 5.	ANACARDIACEAE, p. 514	

53. Stems armed with prickles; leaves several at the nodes.	BERBERIDACEAE, p. 382	
Stems not armed with prickles, or if prickled, only one leaf per node.		54
54. Stamens numerous.		55
Stamens less than 10.		56
55. Fruit with a tongue-shaped bract adhering to the peduncle.	TILIACEAE, p. 520	
Fruit without a tongue-shaped bract.	ROSACEAE, p. 436	
56. Inflorescence terminating the branches, half-round or ovoid; fruit a 3-lobed drupe.	RHAMNACEAE, p. 518 (<i>Ceanothus</i>)	
Inflorescence not as above; fruit not a 3-lobed drupe.		57
57. Fruit a berry; stamens 3.	SAXIFRAGACEAE, p. 427 (<i>Ribes</i>)	
Fruit a dry capsule.		58
58. Plants with leathery green leaves; mostly prostrate.	PYROLACEAE, p. 567 (<i>Chimaphila</i>)	
Plants not with leathery green leaves; mostly upright.	ERICACEAE, p. 571	

C. Sympetalous dicotyledons

(flowers with two floral rings, but with the petals wholly or partly united forming a tube or bell)

1. Shrubs or trees.		2
Herbs.		8
2. Leaves opposite or whorled.		3
Leaves alternate.		6
3. Trees with pinnately compound leaves; stamens 2; fruit winged.	OLEACEAE, p. 587	
Shrubs; stamens 5–10.		4
4. Stems bearing a solitary flower; dwarf shrubs.	DIAPENSIACEAE, p. 580	
Stems bearing few to several flowers; large shrubs.		5
5. Stamens 5; ovary inferior; fruit fleshy.	CAPRIFOLIACEAE, p. 670	
Stamens 5–10; ovary superior; fruit a capsule.	ERICACEAE, p. 571	
6. Flowers in heads; fruit 1-seeded.	COMPOSITAE, p. 683	
Flowers not in heads; fruit many-seeded.		7
7. Ovary inferior; fruit fleshy.	VACCINIACEAE, p. 577	
Ovary superior; fruit usually a capsule, sometimes a mealy-fleshy drupe.	ERICACEAE, p. 571	
8. Plants parasitic or saprophytic, without chlorophyll (green coloring matter).		9
Plants green or with some chlorophyll.		11

9. Plants twining and attached to stems of other plants.	CONVOLVULACEAE, p. 599 (<i>Cuscuta</i>)	
Plants growing out of soil or attached to roots of other plants.		10
10. Corolla regular; stamens 6–10.	MONOTROPACEAE, p. 570	
Corolla 2-lipped; stamens 4.	OROBANCHACEAE, p. 661	
11. Flowers in heads or in form resembling a head.		12
Flowers in long or short spikes.		17
Flowers not in heads or spikes.		20
12. Flowers in true heads with an involucre (bracts).		13
Flowers in form resembling a head, opening in irregular order.		15
13. Male and female flowers in separate heads.	COMPOSITAE, p. 683 (Group 1)	
Flowers perfect, or male and female flowers in same head.		14
14. Stamens united by their anthers.	COMPOSITAE, p. 683 (Groups 2 and 3)	
Stamens separate.	DIPSACACEAE, p. 681	
15. Leaves mostly basal, 1–3 times divided into 3 leaflets; flowers very small, greenish.	ADOXACEAE, p. 677	
Leaves not with 3 leaflets; flowers usually colored.		16
16. Corolla 2-lipped; leaves opposite, stems square.	LABIATAE, p. 620	
Corolla regular; stems round.	POLEMONIACEAE, p. 600	
17. Flowers inconspicuous, on long spikes; leaves all basal.	PLANTAGINACEAE, p. 663	
Flowers brightly colored, irregular.		18
18. Fruit a many-seeded capsule.	SCROPHULARIACEAE, p. 638	
Fruit not a many-seeded capsule.		19
19. Flowers alternate or very crowded; fruit of 4 nutlets.	VERBENACEAE, p. 619	
Flowers opposite, not crowded; fruit an achene, sharply reflexed.	PHRYMACEAE, p. 663	
20. Ovary inferior.		21
Ovary superior.		26
21. Climbing plants with tendrils.	CUCURBITACEAE, p. 677	
Not climbing plants.		22
22. Leaves basal or alternate; plants often with milky juice.		23
Leaves opposite; plants without milky juice.		24

23. Corolla regular, bell-shaped; anthers free.	CAMPANULACEAE, p. 677	
Corolla irregular, split down one side; anthers united into a tube.	LOBELIACEAE, p. 680	
24. Stamens 3.	VALERIANACEAE, p. 681	
Stamens 4 or 5.		25
25. Leaves with stipules, appearing whorled.	RUBIACEAE, p. 667	
Leaves without stipules.	CAPRIFOLIACEAE, p. 670	
26. Corolla irregular.		27
Corolla regular.		29
27. Plants aquatic; leaves bearing bladders; corolla spurred.	LENTIBULARIACEAE, p. 659	
Plants not aquatic; leaves not bearing bladders.		28
28. Stems square; leaves opposite; fruit with 4 nutlets.	LABIATAE, p. 620	
Stems not square; fruit a many-seeded capsule.	SCROPHULARIACEAE, p. 638	
29. Plants with milky juice.		30
Plants without milky juice.		31
30. Stamens united; styles distinct; pollen grains united into a mass.	ASCLEPIADACEAE, p. 593	
Stamens distinct; styles united; pollen grains separate.	APOCYNACEAE, p. 592	
31. Twining plants; corolla large, funnel- form.	CONVOLVULACEAE, p. 597	
Plants not twining.		32
32. Stamens directly in front of corolla lobes.	PRIMULACEAE, p. 580	
Stamens alternating with corolla lobes, or twice their number.		33
33. Stamens 8 or 10.	PYROLACEAE, p. 567	
Stamens 4 or 5.		34
34. Fruit with 4 nutlets; style from center of lobes of ovary; inflorescence usually scorpioid.	BORAGINACEAE, p. 609	
Fruit a berry or capsule; style terminal.		35
35. Ovary 1-celled with seeds borne on wall of cell.		36
Ovary 2- or 3-celled with seeds borne on axis in center of cell.		37
36. Flowers in cymose clusters, leaves opposite.	GENTIANACEAE, p. 589	
Flowers scorpioid or solitary in leaf axils, leaves alternate or basal.	HYDROPHYLLACEAE, p. 605	
37. Ovary 3-celled; fruit a capsule.	POLEMONIACEAE, p. 600	
Ovary 2-celled; fruit a berry or a capsule.	SOLANACEAE, p. 631	

Genera and species

Division: PTERIDOPHYTA—ferns and fern allies

OPHIOGLOSSACEAE—adder's-tongue family

Botrychium grape fern

Plants from fleshy roots with sheath at base of frond (leaf) stalk, containing the following year's bud. Fertile frond bearing the spore-bearing bodies spike-like or raceme-like. Sterile frond leaf-like and lobed.

1. Sterile frond oblong, ovate, or somewhat triangular-ovate, longer than wide, usually glabrous 2
Sterile frond triangular, wider than long, pubescent, at least when young. 6
2. Sterile fronds simple, trilobed, or pinnate; pinnae without midrib. 3
Sterile fronds bipinnate; pinnae with conspicuous midrib. 4
3. Sterile frond petiolate, simple, trilobed, or pinnate with 2 or 3 pairs of pinnae. *B. simplex*
Sterile frond sessile, pinnate with 3–9 pairs of pinnae. *B. lunaria*
4. Pinnae triangular to rhomboid-ovate, about as long as wide. *B. boreale*
Pinnae lanceolate to oblong, longer than wide. 5
5. Pinnae lanceolate to linear-lanceolate, acute at apex; midrib pronounced. *B. lanceolatum*
Pinnae ovate to oblong, obtuse at apex; midrib not pronounced. *B. matricariifolium*
6. Sterile frond petiolate, 2- or 3-pinnate, fleshy; pinnae obtuse at apex. *B. multifidum*
Sterile frond sessile, 3- or 4-pinnate, not fleshy; pinnae acute at apex. *B. virginianum*

Botrychium boreale Milde var. *obtusilobum* (Rupr.) Brown

Plants to 30 cm high. Leaf inserted near middle of stem, 10–20 cm long, subsessile, triangular-ovate, bipinnate; pinnae ovate, acute, pinnatifid. Rare; in grassland and open slopes near tree line; southern Rocky Mountains.

Botrychium lanceolatum (Gmel.) Angst.

Plants to 30 cm high. Leaf inserted at base of fertile frond, to 25 cm long, subsessile, triangular-ovate, bipinnate; pinnae lanceolate to linear-lanceolate, serrate or pinnatifid, with midrib pronounced. Rare; in moist grassland; southern Rocky Mountains, Cypress Hills.

Botrychium lunaria (L.) Swartz.

MOON FERN

Plants to 30 cm high. Leaf inserted near middle of stem, 5–20 cm long, sessile, oblong, pinnate, with 3–9 pairs of fan-shaped pinnae without midrib. Rare; in grassland and open areas; throughout the Prairie Provinces.

Botrychium matricariifolium Braun.

CHAMOMILE-LEAVED GRAPE FERN

Plants to 30 cm high. Leaf inserted above middle of stem, 5–20 cm long, subsessile, oblong to ovate, bipinnate; pinnae oblong to ovate, with midrib. Rare; in moist grassland, muskeg, wooded shores; throughout the Prairie Provinces.

Botrychium multifidum (Gmel.) Rupr.

THICK-LEAVED GRAPE FERN

Plants to 30 cm high. Leaf inserted near base of stipe, 5–20 cm long, triangular, wider than long, bi- or tri-pinnate; pinnae lanceolate, with midrib pronounced. Rare; in moist meadows, margins of woods; Boreal forest.

Botrychium simplex E. Hitchc.

GRAPE FERN

Plants 10–15 cm high. Leaf inserted near middle of stem, 5–8 cm long; stipe 2–4 cm long; pinnae subovate, cuneate at base. Rare; in meadows, margins of woods; Parklands and Boreal forest.

Botrychium virginianum (L.) Swartz. (Fig. 10)

VIRGINIA GRAPE FERN

Plants to 80 cm high. Leaf inserted near middle of stem, 20–50 cm long, triangular, wider than long, sessile, 3- or 4-pinnate; pinnae and segments oblong to ovate, dentate, with midrib pronounced. Rare; in moist woods; Boreal forest, Cypress Hills, Riding Mountain.

(Fig. 10 overleaf)



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Botrychium virginianum* (L.) Sw.

LOCATION: Waskegou Narrows

COLL: J. Nusche

DET: A. LOGMAN

COLL. NO: 15584

DATE: 20 July 1970

ELEVATION: 600 m

HABITAT: Coniferous
woods



Fig. 10. Virginia grape fern, *Botrychium virginianum* (L.) Swartz.

OSMUNDACEAE—royal fern family

Osmunda royal fern

Osmunda claytoniana L.

INTERRUPTED FERN

Plants with stout rhizomes. Fronds 50–100 cm high; stipes shorter than blade, 2–5 mm thick, tomentose when young; blades pinnate-pinnatifid; pinnae in 15–20 pairs, alternate or subopposite. Fertile pinnae 1–5 pairs, 3–6 cm long, 2 cm wide, in middle of blade, bipinnate, with segments oblong, soon withering; sterile pinnae oblong-lanceolate, to 12 cm long, 3 cm wide, acute, deeply pinnatifid, with segments in 10–15 pairs. Sporangia about 0.5 mm wide, dark brown. Southeastern boreal forest.

POLYPODIACEAE—fern family

Ferns with fronds spirally coiled in the bud, and usually bearing green leaf-like fronds. Spores usually borne in little clusters (sori) and usually under a membranous cover (indusium).

1. Sori enclosed in modified globular segments of fertile fronds; fronds becoming blackish in age. 2
 Sori exposed or covered by the reflexed margin of the fertile fronds. 3
2. Fronds scattered along rhizome; sterile fronds once pinnatifid; fertile fronds bipinnate. *Onoclea*
 Fronds tufted on a caudex; sterile fronds surrounding fertile ones. *Matteuccia*
3. Sori marginal or appearing marginal, covered by margin or marginal lobes. 4
 Sori not marginal. 8
4. Sterile and fertile fronds dissimilar; fertile ones longer than sterile ones. *Cryptogramma*
 Sterile and fertile fronds not dissimilar. 5
5. Sori appearing marginal, but borne on veins of reflexed marginal lobes. *Adiantum*
 Sori marginal or submarginal, but not borne on reflexed lobes. 6
6. Rhizome without scales, hairy only; fronds and stipes (stalks) to 60 cm high. *Pteridium*
 Rhizomes scaly; fronds and stipes to 25 cm high. 7
7. Stipes wiry, purplish brown; scales of rhizome rusty brown; blades with 6–12 pairs of pinnae. *Pellaea*

- Stipes soft and hairy or glabrate; scales of rhizome orange; blades 3-pinnate. *Cheilanthes*
8. Indusium absent; blades pinnatisect, with segments linear-oblong, denticulate. *Polypodium*
 Indusium present or absent; blades bipinnatifid to tripinnate. 9
9. Indusium attached by its base under sporangia; veins of fronds not reaching margins. *Woodsia*
 Indusium laterally attached or absent; veins reaching margins. 10
10. Indusium linearly attached. 11
 Indusium attached at a point or absent. 12
11. Indusia not crossing veins; scales of rhizomes with short, broad cells; plants small, up to 20 cm high. *Asplenium*
 Indusia usually crossing veins; scales of rhizomes with long, narrow cells; plants large, up to 100 cm high. *Athyrium*
12. Indusium ovate or oblong; fronds delicate, light green. *Cystopteris*
 Indusium orbicular, reniform, or absent. 13
13. Indusium attached at its center; fronds leathery, with pinnae spinulose-toothed. *Polystichum*
 Indusium attached laterally by a deep notch or absent; fronds not leathery. *Dryopteris*

Adiantum maidenhair

Adiantum pedatum L.

MAIDENHAIR FERN

Plants with rhizomes 2–5 mm thick. Stipes reddish brown, forked above into two branches, each bearing 3–5 pinnae on upper side. Pinnae 15–25 cm long, 3–4 cm wide; pinnules 15–20 pairs, thin, glaucous, oblong, with outer margins incised. Sori oblong-lunate, covered by the reflexed margin of pinnule lobe. Rare; southern Rocky Mountains.

Asplenium spleenwort

Asplenium viride Huds.

GREEN SPLEENWORT

Plants with a short, thin rhizome 1 mm thick. Stipes short, brownish at base. Fronds delicate, 5–15 cm long; pinnae 9–16 pairs, opposite to subopposite at base, alternate above, 3–8 mm long, 2–4 mm wide, with apex obtuse, base cuneate, and margins toothed. Sori elongate; indusia hyaline, attached at one side. Rare; on limestone cliffs; southern Rocky Mountains.

Athyrium spleenwort

Indusium absent; blades tripinnate-pinnatifid. *A. alpestre*

Indusium present; blades mostly subbipinnate. *A. filix-femina*

Athyrium alpestre (Hoppe) Rylands ALPINE SPLEENWORT

Plants with a stout short-creeping rhizome, covered with old stipe bases. Fronds 30–50 cm long, tufted, with stipe erect, shorter than blade; blade lanceolate, bi- or tri-pinnate; pinnae 20–25 pairs, ascending, oblong-lanceolate; pinnules in 15 or more pairs, deeply toothed or lobed. Sori round; indusium lacking. Rare; in alpine meadows; southern Rocky Mountains.

Athyrium filix-femina (L.) Roth LADY FERN

Plants with thick creeping to suberect rhizomes, covered with scales. Fronds 40–100 cm long, tufted, with stipe stout, scaly at base, shorter than, to as long as, blade; blade lanceolate to ovate-lanceolate in outline, sparingly scaly, glabrous or somewhat glandular, bipinnate or tripinnate; pinnae 20–30 pairs, ascending, linear-lanceolate; pinnules 1–2 cm long, serrate to deeply pinnate-lobed. Sori elongate; indusia ciliate. Two varieties distinguished: var. *filix-femina*, with stipe about as long as blade; basal scales to 6 mm long, blackish; indusium hooked at apex, short ciliate; southeastern Boreal forest; and var. *cyclosorum* (Ledeb.) Moore, with stipe about one-third as long as blade; basal scales 10–12 mm long, pale brown; indusium often horseshoe-shaped, long ciliate; Rocky Mountains.

Cheilanthes lip fern

Cheilanthes fei Moore LIP FERN

Plants from short branching rhizomes with abundant scales. Fronds 4–15 cm long; stipe as long as, or shorter than, blade, purplish, long pilose above; blade linear-oblong, 1–3 cm wide, tripinnate; pinnae 7–15 pairs, subopposite, distant below, ascending, ovate to oblong; pinnules 3–6 pairs, diminishing in size toward tip, opposite to subopposite; segments 2 or 3 pairs, ovate to suborbicular, sessile, lobate or incised, sparsely pubescent above, densely brownish pubescent below. Sori marginal, covered by the unmodified margin. Rare; on limestone rocks and cliffs; southern Rocky Mountains.

Cryptogramma rock brake

Rhizome short; fronds densely tufted. *C. crispa*

Rhizome slender, long-creeping; fronds solitary or few together, distant. *C. stelleri*

Cryptogramma crispa (L.) R. Br. PARSLEY FERN

Plants with short rhizomes and numerous fibrous roots and scales. Fronds densely tufted, glabrous. Sterile fronds erect to spreading; stipe straw-colored, to 7 cm long; blade 5–7 cm long, ovate; pinnae 4–6 pairs, alternate or the lowest ones subopposite; pinnules 2–4 pairs; segments elliptic, with bases cuneate and margins dentate. Fertile frond erect, to 20 cm high; stipe 1.5–2.0 times as long as blade; pinnation similar to that of sterile fronds, but segments linear, with margins revolute. Sori on vein tips. On outcrops of Precambrian rocks in

Boreal forest; Rocky Mountains. In Canada these plants are usually distinguishable from the Eurasian var. *crispa* as var. *acrostichoides* (R. Br.) Clarke (Fig. 11) with thicker fronds and paler scales.

Cryptogramma stelleri (Gmel.) Prantl

ROCK BRAKE

Plants with scaly, pilose, creeping rhizomes. Glabrous fronds scattered along the rhizome. Sterile fronds erect; stipe brownish at base, 6–9 cm long; blade ovate, 3–6 cm long; pinnae thin, 5–6 pairs, alternate or sometimes opposite, with 1–3 segments. Fertile fronds to 20 cm high; stipes 12–15 cm; blades bipinnate or sometimes tripinnate; segments linear-lanceolate, entire, margins revolute. Sori on vein tips. Rare; on shaded, usually limestone rock; southern Rocky Mountains.

Cystopteris bladder fern

- Stipes much shorter than blade; blades much narrower at base than long. *C. fragilis*
 Stipes often more than twice as long as blade; blades as wide at base as long. *C. montana*

Cystopteris fragilis (L.) Bernh. (Fig. 12)

FRAGILE FERN

Plants with creeping unbranched rhizomes. Fronds solitary or in small tufts, 10–30 cm long; stipes 3–12 cm long; blade 7–18 cm long, light green, ovate or ovate-lanceolate, bipinnate to tripinnate, glabrous, or with a few hairs at base of pinnae; pinnae in 9–15 pairs, ascending, subopposite to alternate; segments ovate, margins dentate. Sori on veins; indusium hood-like. Fairly common; in deep, shaded, wooded ravines, on moist slopes, and rock ledges; throughout the Prairie Provinces.

Cystopteris montana (Lam.) Bernh.

MOUNTAIN BLADDER FERN

Plants with slender, black, creeping and branching rhizomes; scales conspicuous, pale brown. Fronds scattered, few, 20–35 cm high; stipes 12–20 cm; blades 8–15 cm long, tripinnate; pinnae in 7–13 pairs, the lower ones large, opposite or subopposite, bi- or tri-pinnatifid, the higher ones alternate, pinnales 12–15 mm long; segments dentate. Sori on veins; indusium inconspicuous. Rare; on wet calcareous rock or slopes; southern Rocky Mountains.

Dryopteris shield fern

1. Blades short-pubescent above; segments ciliate; stipe bundles 2, united at base of blade. 2
 Blades glabrous above; segments not ciliate; stipe bundles free above base of blade. 3
2. Blades triangular in outline; stipes scaly and pilose. *D. phegopteris*
 Blades lanceolate in outline; stipes glabrous. *D. thelypteris*
3. Rhizomes thin, blackish; stipe bundles 2; indusium absent. *D. disjuncta*



HEPATICUM IN THE MOUNTAINS OF SWITZERLAND, 1847.
 H. B. K. Bot. Lin. Tab. 100. f. 1. 1825.
 H. B. K. Bot. Lin. Tab. 100. f. 1. 1825.
 H. B. K. Bot. Lin. Tab. 100. f. 1. 1825.
 H. B. K. Bot. Lin. Tab. 100. f. 1. 1825.
 H. B. K. Bot. Lin. Tab. 100. f. 1. 1825.
 H. B. K. Bot. Lin. Tab. 100. f. 1. 1825.

Fig. 11. Parsley fern, *Cryptogramma crispa* (L.) R. Br. var. *acrostichoides* (R. Br.) Clarke.



Fig. 12. Fragile fern, *Cystopteris fragilis* (L.) Bernh.

- Rhizomes thick; stipe bundles more than
2. 4
4. Blades bipinnate-pinnatifid to tripinnate;
segments spinulose toothed. *D. austriaca*
Blades pinnate-pinnatifid to subbipin-
nate; segments not clearly spinulose
toothed. 5
5. Blades glandular on rachis and veins. *D. fragrans*
Blades not glandular. 6
6. Rhizomes short, erect; indusia large, kid-
ney-shaped; basal segments lobed. *D. filix-mas*
Rhizomes thick, short-creeping; indusia
small; basal segments deeply cleft. *D. cristata*

Dryopteris austriaca (Jacq.) Woynar

SPINULOSE SHIELD FERN

Plants with thick rhizomes. Fronds tufted, to 1 m high; stipes shorter than blades; blades 40–60 cm long, 20–30 cm wide at base, bipinnate-pinnatifid or tripinnate-pinnatifid; pinnae in 10–15 pairs; pinnules in 12–20 pairs; segments oblong, spinulose-toothed. Sori on veins; indusium kidney-shaped. A rather variable species, of which many varieties have been described, or treated as distinct species. In Canada the plant is usually named var. *spinulosa* (Muell.) Fiori (*D. spinulosa* (Muell.) Watt.). In moist woods and on slopes; throughout Boreal forest, Rocky Mountains. Rare in Parklands.

Dryopteris cristata (L.) Gray

CRESTED SHIELD FERN

Plants with short-creeping thick rhizomes. Fronds 35–80 cm high; stipes shorter than blades, 15–30 cm; sterile fronds spreading, shorter and wider than erect fertile fronds; blades lanceolate, pinnate-pinnatifid; pinnae in 10–20 pairs, to 8 cm long, with the lower ones spaced; segments ovate oblong, serrate or biserrate, with teeth acute. Sori on the veins; indusia smaller than sori. In damp woods and marshes; Boreal forests, Rocky Mountains.

Dryopteris disjuncta (Rupr.) Morton

OAKFERN

Plants with dark brown to blackish, scaly, slender rhizomes. Fronds to 50 cm high; stipes slender, longer than blades, shiny; blades deltoid, bipinnate-pinnatifid to tripinnate-pinnatifid, delicate; pinnae in 4–7 pairs, with basal ones up to half as long as blades; pinnules oblong, obtuse. Sori on the veins; indusium lacking. Rather rare; moist woods, talus slopes, damp; Precambrian rock outcrop, Boreal forests.

Dryopteris filix-mas (L.) Schott

MALE FERN

Plants with thick, erect rhizomes. Fronds 40–100 cm high, tufted; stipes shorter than blades, with brown, denticulate scales; blades 30–90 cm long, lanceolate, pinnate-pinnatifid to subbipinnate; pinnae in 20–25 pairs, to 15 cm long, 4 cm wide, lanceolate, acuminate; segments in 15–20 pairs, to 2.5 cm long, 1 cm wide, serrate. Sori on veins near midrib; indusia large, horseshoe-shaped. On slopes and cliffs; southern Rocky Mountains.

Plants with short, erect rhizomes. Fronds 25–40 cm high; stipes 6–10 cm long, densely scaly, glandular; blades to 35 cm long, linear-lanceolate, pinnate-pinnatifid; pinnae in 20–25 pairs, to 2.5–3.0 cm long, 1 cm wide, linear oblong; segments in 6–10 pairs, oblong, crenate; veins scaly and glandular. Sori on veins; indusia large, rounded. Rare; on dry cliffs and slopes; Boreal forest.

Dryopteris phegopteris (L.) Christ

BEECH-FERN

Plants with slender, creeping, densely hairy and scaly rhizomes. Fronds 40–50 cm long, solitary; stipes longer than blades, pilose and scaly throughout; blades to 25 cm long, almost as wide at base, pinnate-pinnatifid; pinnae in 8–12 pairs, with the lower ones subopposite, often reflexed; segments oblong, obtuse, with the margins entire or somewhat crenate, pubescent. Sori on veins, submarginal; indusia lacking. Rare; on moist slopes; Boreal forest.

Dryopteris thelypteris (L.) Gray

MARSH FERN

Plants with extensively creeping black, somewhat scaly rhizomes. Fronds solitary; sterile fronds to 40 cm long, with stipes about as long as blades; fertile fronds to 75 cm long, with stipes longer than blades. Stipes black below, glabrous; blades lanceolate, pinnate-pinnatifid; pinnae in 10–15 pairs; sterile pinnae to 10 cm long, 2 cm wide, with segments oval, and margin entire; fertile pinnae with the segments oblong, to 4 mm wide, and margins revolute. Sori on veins; indusia small, ciliate. Rare; in the margins of marshes and wet woods; Manitoba.

Matteuccia ostrich fern*Matteuccia struthiopteris* (L.) Tod.


OSTRICH FERN, FIDDLE HEADS

Plants large, with fronds typically densely tufted around crown of erect, scaly rhizome. Fronds of two kinds. Stipe of sterile fronds shorter than blade, to 40 cm long; blade up to 100 cm long, oblong-lanceolate, pinnate-pinnatifid; pinnae in 20–30 pairs; lower pinnae short, reflexed; middle pinnae long, ascending, with segments oblong and margins revolute. Fertile fronds 40–60 cm long; stipe thick, rigid, dark brown; pinnae to 5 cm long, linear, with margins strongly revolute. Sori on the veins, covered by margin. In Canada the plants are distinguishable as var. *pensylvanica* (Willd.) Mort. (Fig. 13) by having light brown scales on rhizome and lower stipe; in the Eurasian type these scales have a black central band. Syn.: *Onoclea struthiopteris* (L.) Hoffm., *Pteretis nodulosa* (Michx.) Nieuwl., *P. pensylvanica* (Willd.) Fern. Often plentiful; moist woods; Boreal forest, Rocky Mountains. Rare in Parklands.

Onoclea sensitive fern*Onoclea sensibilis* L.

SENSITIVE FERN

Plants with thick, scaly rhizomes 5–7 mm in diam. Fronds to 100 cm long, solitary or few together; stipes longer than blades, glabrous. Sterile fronds pinnatifid; blades 20–40 cm long and as wide at base; rachis winged above; lower segments opposite, lobate to somewhat pinnatifid. Fertile fronds shorter than



HERBARIUM OF THE UNIVERSITY OF PENNSYLVANIA
 JOHN H. RAVENHILL, CURATOR
 THE UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, PA.
 1887
 No. 1000
 MATTEUCCIA STRUTHIOPTERIS (L.) TOD. var. PENNSYLVANICA (WILLD.) MORT.
 COLLECTED BY J. H. RAVENHILL
 PENNSYLVANIA, 1887




Fig. 13. Ostrich fern, *Matteuccia struthiopteris* (L.) Tod. var. *pennsylvanica* (Willd.) Mort.

sterile ones; blade 12–15 cm long, bipinnate; pinnae in bead-like segments with revolute margins. Sori globose; indusium hood-like and covering sori. Southeastern Boreal forest.

Pellaea cliff brake

Pellaea glabella Mett.

PURPLE CLIFF BRAKE

Plants with short, erect rhizomes with yellow brown scales. Fronds to 25 cm long; stipes shorter than blades, subdimorphic; blades pinnate or bipinnate at base; pinnae in 5–10 pairs, usually ascending; sterile segments ovate to ovate-oblong, to 10 mm wide at base; fertile segments linear-oblong, to 5 mm wide at base, with margins revolute. Sori submarginal, at tip of veins; indusia absent. Two varieties are recognized: var. *simplex* Butters, having cells of scales long-linear, 10–15 times as long as wide, occurring in southern Rocky Mountains; var. *nana* (Rich.) Cody, having plants usually smaller than in var. *simplex*, with cells of scales oblong-lanceolate, 3–5 times as long as wide. Both varieties occur on rock cliffs, usually limestone; Boreal forests.

Polypodium rock tripe

Polypodium vulgare L.

COMMON ROCK TRIPE

Plants with stout rhizomes, 1.5–3 mm thick, covered with fibrous scales. Fronds mostly distant, 10–30 cm long; stipes shorter than blades, scaly at base. Blades pinnatisect, with segments in 12–20 pairs, alternate to subopposite, spreading. Sori round, submarginal; indusia absent. In Canada plants are distinguishable from the European var. *vulgare* by having paraphyses (thin sterile filaments) mixed in the sporangia, and smaller rhizome scales, and have been named var. *virginianum* (L.) D.C. Eaton (Fig. 14); occur in Precambrian rock outcrops in Boreal forest, northern Rocky Mountains. A var. *columbianum* Gilbert, usually smaller, with fewer segments in fronds, occurs in southern Rocky Mountains.

Polystichum rock tripe

Polystichum lonchitis (L.) Roth

HOLLY FERN

Plants with erect or ascending, thick rhizomes, to 1 cm in diam. Fronds 25–50 cm long; stipes short or almost none, persistently scaly; blades linear-lanceolate in outline, pinnate; pinnae in 25–45 pairs, spreading at right angles, with margin spinulose-toothed, scaly below. Sori round; indusia arising from center of sori, lacerate. Southern Rocky Mountains.

Pteridium brake fern

Pteridium aquilinum (L.) Kuhn

BRACKEN FERN

Plants with pubescent, creeping rhizomes. Fronds large; stipes about as long as blades; vascular bundles in cross section resembling “an eagle with spreading wings.” Blades bipinnate- or tripinnate-pinnatifid; pinnae in 6–9 pairs, opposite or nearly so. Pinnulae alternate, with segments oblong or linear-oblong, and margins revolute. Sori linear, marginal, continuous, covered



HERBARIUM OF THE RESEARCH STATION
SAINT JOHN'S, Nfld.

PLANTS OF: SACKETT-HESMAN, Lake St. John, Labrador

SPECIES: *Polypodium vulgare* L.
var. *virginianum* (L.) Eaton

LOCATION: McKay Lake

COLL: J. Lesman DET:

COLL. NO: 10479 DATE: 29 July 1971 ELEV: 300 m

HABITAT: In crevices of Precambrian rock

Fig. 14. Common rock tripe, *Polypodium vulgare* L. var. *virginianum* (L.)
D. C. Eaton.

by the revolute margin; indusium, if present, delicate, continuous along the margin. A variable cosmopolite, in which several species have been recognized, but now generally considered a single species with two subspecies and several more or less well-defined varieties. In Canada these plants are usually described as ssp. *aquilinum* var. *pubescens* Underw., with the fronds commonly pubescent below, especially along the midribs and margins. Boreal forest, Rocky Mountains.

Woodsia

1. Blades stipitate glandular; stipes not articulate. 2
Blades not glandular; stipes articulate below middle. 3
2. Fronds with long hairs on veins and midribs of pinnulae and segments. *W. scopulina*
Fronds lacking hairs. *W. oregana*
3. Stipe brown, scaly, and pubescent. 4
Stipe straw-colored, glabrous. *W. glabella*
4. Stipe slender, less than 1 mm thick; midribs of pinnae not or scarcely scaly. *W. alpina*
Stipe stout; midribs of pinnae scaly. *W. ilvensis*

Woodsia alpina (Bolton) S. F. Gray

Plants with scaly rhizomes; scales 4–6 mm long, 1–2 mm wide, brown, toothed. Fronds 6–15 cm long; stipes shorter than blades, brown, somewhat pilose and scaly, articulate below middle; blades 1–2 cm wide, linear, pinnate-pinnatifid; rachis somewhat hairy and scaly; pinnae in 8–15 pairs, deeply pinnatifid, hairy below, usually without scales; segments in 2 or 3 pairs, oblong to suborbicular. Sori at the apex of secondary veins; indusium with few long septate hair-like lobes. Shaded cliffs; southeastern Boreal forest.

Woodsia glabella R. Br.

Plants with slender scaly rhizomes; scales 3–4 mm long, 1–1.5 mm wide, brown, toothed. Fronds 5–15 cm long; stipes shorter than blades, straw-colored, glabrous, scaly at base only, articulate below middle; blades 7–15 mm wide, linear, glabrous, without scales, pinnate; pinnae in 8–15 pairs, deltoid or rounded in outline, trilobate or pinnatifid, with 2 or 3 pairs of segments. Sori at apex of veins; indusia with numerous septate hairs. Shaded moist cliffs of calcareous or dolomitic rock; Boreal forest, Rocky Mountains.

Woodsia ilvensis (L.) R. Br.

Plants with stout scaly rhizomes; scales 4–6 mm long, to 1 mm wide, brown and toothed. Fronds to 20 cm long; stipes shorter than blades, shiny brown, stout, to 1 mm thick, scaly and pubescent, articulate below middle; blades 2–3.5 cm wide, pinnate-pinnatifid; rachis scaly and hirsute; pinnae in 10–15 pairs, the lower ones subopposite, the higher ones alternate, deeply pinnatifid; segments in 2–7 pairs, oblong, with the apex rounded, scaly, and hairy

below. Sori submarginal, confluent; indusia with a fringe of long hairs. Crevices in Precambrian rock outcrops; Boreal forest.

Woodsia oregana D. C. Eaton

Plants with stout scaly rhizomes; scales 3–5 mm long, to 0.5 mm wide, subentire. Fronds to 25 cm long; stipes shorter than blades, dark brown, stout to 1.5 mm thick, scaly at base, not articulate; blades lanceolate, to 5 cm wide, pinnate-pinnatifid to pinnate-bipinnatifid, glabrous or glandular pubescent; pinnae in 10–15 pairs, subopposite, deltoid to lanceolate, mostly bipinnatifid; segments in 5–7 pairs, with margins often revolute. Sori marginal, in part confluent; indusia divided into few narrow segments. Occur in rock crevices. The typical var. *oregana* has blades sparingly glandular to glabrous; rare in southern Rocky Mountains, Lake Athabasca. The var. *cathcartiana* (Robins.) Morton has blades copiously, finely glandular; rare in Cypress Hills.

Woodsia scopulina D. C. Eaton

Plants with stout scaly rhizomes; scales 4–5 mm long, 1.0–1.5 mm wide. Fronds 15–35 cm long; stipes shorter than blades, stout, brown, pubescent, scaly at base, not articulate; blades linear lanceolate, 3–7 cm wide, bipinnate; rachis somewhat pubescent, not scaly; pinnae in 9–17 pairs, subopposite, lanceolate, ovate or deltoid-ovate; pinnulae in 7–10 pairs, oblong, blunt, crenate to subpinnatifid, pubescent and somewhat glandular below. Sori submarginal at tips of veins; indusia with 3–6 filamentous scales. Occur in rock crevices; Boreal forest, Rocky Mountains.

MARSILEACEAE—marsilea family

Marsilea pepperwort

Marsilea mucronata A. Br.

HAIRY PEPPERWORT

Low growing from slender creeping rootstocks. Leaves borne singly on thin stalks 5–15 cm long, divided into 4 triangular leaflets, each 3–10 mm long. Spores borne in ovoid bean-like containers (sporocarps) on short stalks near base of plant and covered with hair-like scales. Not common; in slough bottoms at several locations in the south central prairies. Syn.: *Marsilea vestita* Hook & Grev.

EQUISETACEAE—horsetail family

Equisetum horsetail

Perennial rush-like plants, with stems fluted or grooved, and joints or nodes solid but surrounded by a toothed sheath. Fertile stems with a terminal cone in which spores are borne.

1. Branches compound; sheaths bright red-dish brown. *E. sylvaticum*

- Branches not compound, sometimes absent; sheaths not bright reddish brown. 2
2. Stems usually much-branched with whorls of branches. 3
 Stems usually not branched above the ground. 6
3. Plants with hollow branches. 4
 Plants with solid branches. 5
4. Center cavity of stem very small. *E. palustre*
 Center cavity of stem large. *E. fluviatile*
5. Fertile stems not branched, soon withering. *E. arvense*
 Fertile stems branching toward top, only the tip withering. *E. pratense*
6. Stems low, slender, tufted, 5- to 10-grooved. 7
 Stems tall, stouter, many-grooved. 8
7. Stems solid; sheaths 3-toothed. *E. scirpoides*
 Stems with small central cavity; sheaths 5- to 10-toothed. *E. variegatum*
8. Sheaths almost as broad as high, cylindrical, turning black or gray with black bands above and below. *E. hyemale*
 Sheaths higher than broad, somewhat funnel-shaped, usually with a narrow black band. *E. laevigatum*

Equisetum arvense L. (Fig. 15)

COMMON HORSETAIL

A plant with annual stems withering at the end of the season; stems of two kinds: fertile, unbranched stems bearing the spore-containing cone at the summit; and sterile, much-branched stems. Plants 8–25 cm high; sheaths having 8–12 brownish teeth. Sterile stems with whorls of branches. Common; in wet places; throughout the Prairie Provinces, especially on sandy soils.

Equisetum fluviatile L.

SWAMP HORSETAIL

An annual-stemmed species 10–100 cm high with whorls of hollow branches. Sheaths flaring and bearing about 18 dark brown teeth. Common; in marshes and shallow water; Boreal forest. Syn.: *E. limosum* L.

Equisetum hyemale L. var. *affine* (Engelm.) A. A. Eaton

COMMON SCOURING-RUSH

A perennial-stemmed, unbranched species 30–100 cm high, with broad, conspicuous sheaths, usually grayish with black bands above and below. Common; where subsoil is moist, in sandhill areas, creek flats, and lake margins; throughout the Prairie Provinces. Syn.: *E. prealtum* Raf.




Fig. 15. Common horsetail, *Equisetum arvense* L.

Equisetum laevigatum A. Br.

SMOOTH SCOURING-RUSH

A species 30–100 cm high, usually with annual stems, but with some stems perennial. Sheaths green and flaring at the summit, often with a black band, or a gray or black band at the base of the sheath. Cone rounded or with a tip. On light soils, and in moist areas; throughout Prairies and Parklands. Includes *E. kansanum* Schaffn.

Equisetum palustre L.

MARSH HORSETAIL

An annual-stemmed plant 15–50 cm high, with a very small cavity in the center of stem. Usually whorls of hollow branches. Sheaths usually somewhat flaring at the top and bearing 8 brownish but white-margined teeth. Fairly common; in wet soil; throughout Parklands and Boreal forest.

Equisetum pratense Ehrh.

MEADOW HORSETAIL

An annual-stemmed species 15–50 cm high, with fertile stems appearing before sterile ones. Sterile stems much-branched, but fertile ones only branched when old. Fairly common; in moist, sandy soils; Parklands, Boreal forest, Cypress Hills, and other favorable locations.

Equisetum scirpoides Michx.

DWARF SCOURING-RUSH

A tufted species with thread-like, unbranched, solid stems 8–15 cm high. Sheaths 3-toothed; cones very small. In swamps, wet spruce woods; Boreal forest, Cypress Hills, Riding Mountain – Duck Mountain.

Equisetum sylvaticum L.

WOODLAND HORSETAIL

A pretty species with conspicuous sheaths bearing large, loose, reddish brown teeth. Both fertile and sterile stems, annual, and branched with solid, compound (divided) branches. Fairly common; in moist woodlands; Boreal forest, Cypress Hills, Riding Mountain – Duck Mountain.

Equisetum variegatum Schleich.

VARIEGATED HORSETAIL

A low, tufted perennial-stemmed species 15–50 cm high, with 5–10 teeth on each sheath; teeth black with a white border and a bristle-like tip. Occur occasionally; in wet places; Boreal forest.

ISOETACEAE—quillwort family

Isoetes quillwort

Submersed aquatic plants with a 2-lobed corm. All leaves with a basal sporangium, and a triangular ligule above the sporangium. Spores of two kinds: megaspores, 0.25 mm in diam or larger; and microspores, visible only through a microscope.

Megaspores 0.3–0.5 mm in diam, white or bluish; surface with small tubercles or wrinkled. *I. bolanderi*

Megaspores 0.4–0.6 mm in diam; surface distinctly spinulose. *I. echinospora*

Isoetes bolanderi Engelm.

Plants submersed. Leaves 6–25, bright green, soft, 6–15 or 20 cm long; stomata few. Sporangia 3–4 mm long, covered for one-quarter to one-third of their length by the ligule. Megaspores 0.3–0.5 mm, microspores 25–30 μm . Rare; in lakes and ponds; southern Rocky Mountains.

Isoetes echinospora Dur. var. *braunii* (Dur.) Engelm.

Plants submersed or emersed. Leaves 10–45, coarse, pale green, 6–15 cm long; stomata few toward the leaf tip. Sporangia 4–5 mm long, covered for one-half to three-quarters of their length by the ligule. Megaspores 0.4–0.6 mm, microspores 25–30 μm . Rare; in lakes and ponds; Boreal forests.

LYCOPODIACEAE—club-moss family

Lycopodium club-moss

Perennial, low, usually trailing plants with short, stiff, single-nerved, overlapping leaves. Spores sulfur-colored, borne in spore cases (sporangia) on the upper surfaces or on the axils of the leaves, on ascending or aerial branches, often aggregated in strobili. Spores containing much oil, inflammable.

1. Sporangia borne in zones along stem, not in strobili. *L. selago*
Sporangia borne in strobili. 2
2. Strobili green, not much different from stems; plants without rhizomes. *L. inundatum*
Strobili yellow, much different from stems; plants with rhizomes. 3
3. Strobili borne on a long peduncle. 4
Strobili sessile on stem. 5
4. Leaves to 7 mm long, awl-shaped, with a hair-like tip; branches not flattened. *L. clavatum*
Leaves to 3 mm long, scale-like, with apex acuminate; branches strongly flattened. *L. complanatum*
5. Erect stems much-branched, resembling a small shrub. *L. obscurum*
Erect stems not much-branched. 6
6. Leaves in 6–10 ranks; plants coarse, stems often very long, prickly. *L. annotinum*
Leaves in 4–5 ranks; plants not coarse. 7
7. Branchlets much-flattened; leaves strongly appressed; trailing stems about 2 mm thick. *L. alpinum*

Branchlets not much-flattened; leaves spreading; trailing stems about 1–2 mm thick. *L. sabinifolium*

Lycopodium alpinum L.

The horizontal stems 30–60 cm long, rooting throughout, 2–2.5 mm thick, with few bract-like yellow leaves. Erect stems repeatedly forked, 4–10 cm long, with branchlets flattened. Strobili sessile, solitary at the end of a leafy peduncle, 1–2 cm long. Rocky Mountains.

Lycopodium annotinum L.

STIFF CLUB-MOSS

A trailing, prostrate plant often 100–200 cm long, with stiff, linear-lanceolate, sharp-tipped leaves about 3–7 mm long, crowded along stems and branches. Aerial or upright branches 10–20 cm high, tipped with yellowish fruiting spike 15–25 mm long. Often found in moist woodlands; Boreal forest, Riding Mountain – Duck Mountain, Cypress Hills.

Lycopodium clavatum L.

RUNNING-PINE

A prostrate, trailing plant, with the main stem up to 100–200 cm long. Leaves very similar to the previous species but tipped with a fine bristle. Fruiting spikes borne on a bract-covered stalk 3.5–12 cm long. Not common; Boreal forest.

Lycopodium complanatum L. (Fig. 16)

TRAILING CLUB-MOSS

A plant with the main stem trailing on or slightly under the ground surface. Leaves small and tightly clasping the stem, giving the stem a very narrow smooth appearance, quite different from other club-mosses. Fruiting spikes 15–25 mm long, cylindrical, and borne on a chaffy stalk 5–15 cm high. Fairly common in pine woods and damp forests; Boreal forest.

Lycopodium inundatum L.

A prostrate plant with arching stems frequently rooting. Strobili few, 1.5–4 cm long, to 1 cm thick, at the tip of a leafy erect stem. Rare; in bogs; Boreal forest. Known only from Saskatchewan.

Lycopodium obscurum L. (Fig. 17)

GROUND-PINE

A species with the main stem creeping horizontally below the surface of the ground. Aerial upright branches appearing like miniature evergreen trees growing to 10–25 cm high, with tightly packed, spreading leaves about 3–4 mm long, linear-lanceolate, and sharp-tipped. Fruiting spikes almost stalkless, 1.5–3 cm long. Occasionally found in moist woods; Boreal forest, Cypress Hills.

Lycopodium sabinifolium Willd.

GROUND-FIR

A species with the creeping stem bearing few scaly leaves. Erect stems forking repeatedly; branchlets somewhat flattened at the tips, 5–20 cm long. Strobili sessile, solitary or in pairs on leafy stems, 1.5–2.5 cm long. Rare in open woods; Boreal forest. In Canada the plants are distinguishable as var.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

PLANTS OF: SASKATCHEWAN, Lac La Poudre District

SPECIES: *Lycopodium complanatum* L.


LOCATION: Lussier Lake

COLL: J. Loewen DET:

COLL. NO: 16612 DATE: 5 August 1971 ELEV: 4200

HABITAT: Coniferous forest

Fig. 16. Trailing club-moss, *Lycopodium complanatum* L.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

PLANTS OF: SASKATCHEWAN, MANITOBA PROVINCE
SPECIES: *Lycopodium obscurum* L.

LOCATION: Eckey Lake

COLL: J. Lozman DATE: July 1971

COLL. NO: 10,25 ELEV: 320m

HABITAT: Coniferous forest

Fig. 17. Ground-pine, *Lycopodium obscurum* L.

sitchense (Rupr.) Fern., having the fertile branches much longer than the sterile ones; in the typical form, fertile and sterile branches are about equal in length, with only the strobili elevated.

Lycopodium selago L.

Plants with short horizontal stems and tufted erect stems, forking repeatedly, to 30 cm high. Leaves 8-ranked, about 5 mm long, erect, and appressed. Sporangia borne in axils of leaves in alternating zones; each season's growth having a sterile basal zone, with a fertile zone at the summit. Peat bogs, muskeg; Boreal forest, Rocky Mountains.

SELAGINELLACEAE—spike-moss family

Selaginella little club-moss

1. Leaves acute, not bristle-tipped, rootlets few. *S. selaginoides*
Leaves bristle-tipped, rootlets many. 2
2. Plants blue green, glaucous, loosely tufted. *S. wallacei*
Plants green, densely tufted. 3
3. Bristle 1.0–2.0 mm, conspicuous. *S. densa*
Bristle 0.3–0.5 mm, inconspicuous. *S. rupestris*

Selaginella densa Rydb.

PRAIRIE SELAGINELLA

A low, densely matted plant, with stems rooting almost their whole length and densely branched. Plants covered thickly with tiny leaves up to 3 mm long, each tipped by a minute bristle, and varying from green to yellowish according to age and condition. Strobili 10–25 mm long, covered with somewhat triangular, green, much overlapping bracts; spore containers occurring singly in the axils of these bracts. Although not usually noticed, this inconspicuous plant is probably one of the commonest and most dominant plants of the drier and more open prairies. It helps stop erosion and perhaps builds up soil by decaying organic matter, but it has no forage value and it increases with overgrazing and abuse of prairie pastures. Very plentiful on drier light soils, and eroded spots in grassland; throughout Prairies and Parklands.

Selaginella rupestris (L.) Spring

Very similar to the preceding species, but with bristles short and plants less conspicuously bristly. On rock outcrops, and in open pine forests; Boreal forest.

Selaginella selaginoides (L.) Link

Small, mostly inconspicuous plants. Prostrate stems 2–5 cm long, with few rootlets. Fertile stems erect, 6–10 cm high, or rarely to 20 cm; strobili 1.5–3 cm long, almost cylindrical, open. Not common; in boggy areas, wet meadows; Boreal forest, Rocky Mountains.

Selaginella wallacei Hieron.

Very similar to *S. densa* and *S. rupestris*, but much more loosely spreading, and glaucous blue green. Apical leaves usually with a short bristle; stem leaves often not bristle-tipped. Over rocks and on dry slopes; southern Rocky Mountains.

Division: SPERMATOPHYTA—seed-bearing plants

Subdivision: GYMNOSPERMAE—naked-seeded plants

TAXACEAE—yew family

Taxus yew

- Shrub, usually straggling, with ascending or rarely erect stems up to 2 m high. *T. canadensis*
- Shrub or small tree with straight trunk, up to 10 m high. *T. brevifolia*

Taxus brevifolia Nutt. WESTERN YEWE

A tree 10–15 m high, erect, with a straight trunk. Leaves 2-ranked along branches, giving branches a flat appearance. Tip of leaves reflexed. Rare; southern Rocky Mountains.

Taxus canadensis Marsh. GROUND HEMLOCK

A shrub, usually with straggling to ascending branches, rarely with erect stems. Branches appearing flat. Tip of leaves not reflexed. Rare; in moist woods and muskeg; southeastern Boreal forest.

PINACEAE—pine family

- 1. Leaves borne in clusters of 2 to many. 2
 - Leaves borne singly. 3
- 2. Leaves in clusters of 2–5, evergreen; base of clusters enclosed in a chaffy sheath. *Pinus*
 - Leaves in clusters of 10–40, deciduous; base of clusters without sheath. *Larix*
- 3. Leaves 4-sided, not appearing 2-ranked; leaf scars prominent on twigs. *Picea*
 - Leaves flat, appearing 2-ranked; leaf scars not prominent on twigs. 4
- 4. Cones drooping, with scales persistent; bracts longer than scales, 3-lobed; leaf scars oval. *Pseudotsuga*
 - Cones erect, with scales deciduous; bracts shorter than scales, rounded; leaf scars round. *Abies*

Abies fir

Abies balsamea (L.) Mill.

BALSAM FIR

A tall tree, with gray fairly smooth bark having numerous resinous blisters. Leaves needle-like, 2–3 cm long, flat and stalkless, shiny dark green above and whitish below, with 8–10 lines of stomata. Male and female flowers borne on the same tree, with males yellowish to red and females purple. Cones dark purple, somewhat oblong, 5–10 cm long. Common; in Boreal forest, Rocky Mountains.

The var. *fallax* (Eng.) Boiv. (= *A. lasiocarpa* (Hook.) Endl.) has needles more glaucous, narrower, and with 10–12 lines of stomata below. Southern Rocky Mountains.

Larix larch

- 1. Twigs tomentose pubescent, even when old. *L. lyallii*
- Twigs glabrous, or pubescent only when young. 2
- 2. Needles 1–2 cm long; scales of cones glabrous. *L. laricina*
- Needles 3 cm long or more; scales of cones puberulent. *L. occidentalis*

Larix laricina (DuRoi) K. Koch

TAMARACK

A rather slender tree 6–15 m high; bark reddish brown, with small, flaky scales. Leaves needle-like, 1–2 cm long, very pale green, in clusters of 10–20 along the twigs, turning yellow in autumn and dropping off. Fruits small cones, developing the first year and soon shed. Common; in swamps and marshy woods; Boreal forest, Rocky Mountains, Riding Mountain – Duck Mountain.

Larix lyallii Parl.

ALPINE LARCH

A tree, rarely over 10 m high; bark thin, furrowed, with reddish brown loose scales. Leaves 30–40 in a cluster, bluish green, 4-sided. Seed cones ellipsoid-oblong, with scales hairy below, fringed. At high altitudes, at timberline in Rocky Mountains.

Larix occidentalis Nutt.

WESTERN TAMARACK

A large tree reaching to 50 m high; bark reddish brown, deeply furrowed at base, forming large flutes. Leaves 15–30 in a cluster, triangular. Seed cones oblong, with scales tomentose below when young. Rare; southern Rocky Mountains.

Picea spruce

Shapely evergreen trees with 4-sided needle-like leaves scattered around the twigs. Cones pendulous, maturing the first year.

- Branchlets not hairy; cones falling the first winter; cones oblong-cylindric, often over 3 cm long. *P. glauca*
- Branchlets somewhat hairy; cones remaining on tree for several years; cones oval or ovoid, not over 3 cm long. *P. mariana*

Picea glauca (Moench) Voss

WHITE SPRUCE

A shapely tree 7–20 m high, with scaly, brown bark. Leaves bluish green 1–2.5 cm long. Female inflorescence crimson; cones cylindric 2.5–5 cm long, with smooth-margined scales. Very plentiful throughout the Prairie Provinces, except Prairies and Parklands. A variety, the western white spruce, *Picea glauca* (Moench) Voss var. *albertiana* (S. Brown) Sarg., in which the cone scales are erose or somewhat toothed at the margins, is the common variety of southern Rocky Mountains.

Picea mariana (Mill.) BSP.

BLACK SPRUCE

A less shapely tree than the previous species 7–15 m high. Bark grayish to reddish brown, scaly. Needles 1–2 cm long, bluish green. Male inflorescence dark red, and female purplish; cones 1–4 cm long, ovoid, remaining on trees for several seasons. Fairly common; in wet or swampy woodlands; throughout Boreal forest.

Pinus pine

Tall evergreen trees with leaves borne in clusters; cones maturing in second season.

1. Leaves 2 in a cluster. 2
 Leaves 3–5 in a cluster. 4
2. Leaves 7–17 cm long; cones ovoid-conical, terminal or almost so. *P. resinosa*
 Leaves less than 7 cm long; cones conical, lateral at maturity. 3
3. Leaves 2–4 cm long, thick and rigid, twisted and spreading; cones curved toward tips of branches; scales without prickles. *P. banksiana*
 Leaves 3–6 cm long, not usually twisted and spreading; cones spreading at right angles to branches; scales with a prickle. *P. contorta*
4. Leaves 3 in a cluster, 8–20 cm long; cones to 18 cm long, scales with a recurved prickle. *P. ponderosa*
 Leaves 5 in a cluster. 5
5. Cones ovoid, with thick scales. 6
 Cones cylindric or subcylindric, with thin scales. 7

6. Cones to 7 cm long. *P. albicaulis*
 Cones to 20 cm long. *P. flexilis*
7. Leaves 8–13 cm long, pale green and
 glaucous; cones to 15 cm long; eastern
 species. *P. strobus*
 Leaves 5–10 cm long, bluish green and
 glaucous; white bands of stomata;
 cones 10–20 cm long; western species. *P. monticola*

Pinus albicaulis Engelm.

WHITEBARK PINE

Small alpine tree, with crooked and twisted trunk, or reduced to a shrub. Bark smooth, whitish; twigs yellowish, hairy. Leaves 4–8 cm long, dark green, stiff. Cones oval to subglobose, 3–7 cm long, purplish, with scales forming a stout protuberance, not prickly. Rare; at timberline; Rocky Mountains.

Pinus banksiana Lamb.

JACK PINE

A tree up to 20 m high, with thin, reddish brown bark. Needle-like leaves generally somewhat twisted, 2–5 cm long, borne in twos, yellowish green. Male inflorescence yellow, female dark purple; cones 2–5 cm long, curved, generally in pairs, with unarmed scales and usually pointing toward apex of branches. Abundant; on sandy soils; Boreal forest.

Pinus contorta Dougl. var. *latifolia* Engelm.

LOGEPOLE PINE

A tree similar to *P. banksiana*. Difficult to distinguish between the two species, but lodgepole pine usually having darker and less twisted needles, darker and thinner bark, and less curved cones often bending backward and pointing downward. Cone scales bearing a small, recurved prickle at tip. Very common; in southern Rocky Mountains and Cypress Hills, but apparently intergrading with jack pine in northwestern Boreal forest. Syn.: *P. murrayana* Balf. The species *P. contorta* and *P. banksiana* are sometimes considered as varieties of a single species: *P. divaricata* (Ait.) Dumont, var. *divaricata* (= *P. banksiana*) and var. *latifolia* (Engelm.) Boiv. (= *P. contorta* var. *latifolia*).

Pinus flexilis James

LIMBER PINE

A small tree with a short, stout trunk, and thick branches; bark light gray becoming dark in age. Leaves 3–7 cm long, rigid with 1–4 rows of stomata on all sides. Seed cones 8–20 cm long, light brown to purplish, scales thickened and curved inward at apex. Open rocky slopes and hilltops; at altitudes between 1300 and 2000 m in southern Rocky Mountains.

Pinus monticola Dougl.

WESTERN WHITE PINE

Trees to 50 m high, with a short-branched symmetrical crown. Leaves 5–10 cm long, in 5-leaved clusters, bluish green. Cones cylindrical, 10–20 cm long, thin-scaled. Only found as young trees in southern Rocky Mountains.

Pinus ponderosa Dougl.

WESTERN YELLOW PINE

Trees to 75 m high, with a spire-like crown, or flat-topped in poor sites. Branchlets orange when young. Leaves in clusters of 3, to 25 cm long. Cones

7–15 cm long, with thin scales, thickened at apex, and armed with a slender prickle. Only found as young trees in southern Rocky Mountains.

Pinus resinosa Ait.

NORWAY PINE

Trees to 40 m high, with thick, brown, furrowed bark. Leaves in pairs, 10–15 cm long, dark green. Cones ovoid, 4–8 cm long, spreading, with apex thickened and a smooth protuberance. Southeastern Boreal forest.

Pinus strobus L.

EASTERN WHITE PINE

Trees to 50 m high, with thick, furrowed bark. Leaves slender, in clusters of five, 8–13 cm long, pale green, glaucous. Cones cylindrical, 10–15 cm long, with scales having a protuberance at tip. Rare; southeastern Boreal forest.

Pseudotsuga Douglas-fir

Pseudotsuga menziesii (Mirb.) Franco

DOUGLAS-FIR

Trees to 50 m high, with dark brown, thick, furrowed bark. Lower branches often drooping. Leaves 2–3 cm long, flat, narrowed to a short stalk. Cones pendent, 5–10 cm long; bracts with 3 teeth, projecting beyond scales. Not common; Rocky Mountains.

CUPRESSACEAE—cypress family

Seeds in a small dry cone; branchlets flattened; medium-sized trees. *Thuja*

Seeds in a small, bluish, berry-like cone; branchlets not flattened; low or trailing shrubs. *Juniperus*

Juniperus juniper

Low shrub with short, awl-shaped or scale-like leaves, opposite or in whorls. Fruit composed of 3–6 fleshy scales, each containing a seed and joined to form a fleshy berry-like cone.

1. Leaves in whorls of 3, linear, awl-shaped, sharp-pointed. *J. communis*

Leaves opposite, scale-like. 2

2. Shrubs, prostrate or ascending. *J. horizontalis*

Shrubs or small trees, mostly with a well-developed trunk. *J. scopulorum*

Juniperus communis L.

LOW JUNIPER

A shrub about 1–1.5 m high in some places but usually very low. Leaves 5–12 mm long, narrowly awl-shaped, pointed and dark green below, whitish and grooved above. Cones berry-like in leaf axils, bluish with a bloom, 6–10 mm in diam. The var. *depressa* Pursh is fairly common on light, rocky soil throughout most of the Prairie Provinces. Syn.: *J. sibirica* Burgsd.

Juniperus horizontalis Moench (Fig. 18)

CREeping JUNIPER

A prostrate shrub, with long, gnarled, woody stems often 3–5 m long. Leaves scale-like, overlapping, each about 1.5 mm long and forming narrow branchlets up to 12 mm long. Cones bluish and berry-like about 6 mm long, at the ends of branchlets. Very common throughout drier parts of area, forming large mats on dry banks and sandy hillsides. Syn.: *Sabina horizontalis* (Moench) Rydb.

Juniperus scopulorum Sarg.

ROCKY MOUNTAIN JUNIPER

Usually a small tree 2–3 m high, or a large upright shrub with several stems; otherwise hardly distinguishable from *J. horizontalis*. Southern Rocky Mountains.

Thuja arbor-vitae

Leaves of branchlets and twigs with a conspicuous resin gland. *T. occidentalis*

Leaves of branchlets and twigs not with a conspicuous resin gland. *T. plicata*

Thuja occidentalis L.

WHITE CEDAR

A tree to 20 m high, with widely spreading branches. Branchlets forming a flattened spray, with soft twigs. Leaves strongly appressed, keeled, with a light green or yellow resin gland below apex. Cones about 10 mm long, oblong-ovoid. Moist woods; southeastern Boreal forest.

Thuja plicata D. Don

WESTERN RED CEDAR

Similar to the preceding species, but with leaves not distinctly keeled, and the resin gland inconspicuous. Moist woods; southern Rocky Mountains.

Subdivision: ANGIOSPERMAE

Class: MONOCOTYLEDONEAE

TYPHACEAE—cattail family

Typha cattail

Staminate part of spike contiguous with pistillate part. *T. latifolia*

Staminate part of spike separated from pistillate part by 3–5 cm. *T. angustifolia*

Typha angustifolia L.

NARROW-LEAVED CATTAIL

Marsh or aquatic plants to 3 m high, with extensive creeping roots. Leaves 4–10 mm wide, to 30 cm long. Spikes 25–40 cm long, with staminate and pistillate parts about equal, separated by 3–5 cm of bare stem. Rare; southeastern Boreal forest.




Fig. 18. Creeping juniper, *Juniperus horizontalis* Moench.

Similar to the preceding species, but with leaves to 30 mm wide. Spikes with staminate and pistillate parts contiguous. Very common; in slough margins, marshes, lakeshores, and riverbanks; throughout the Prairie Provinces.

SPARGANIACEAE—bur-reed family

Sparganium bur-reed

- 1. Stigmas 2; achenes broadly obpyramidal, truncate at summit. *S. eurycarpum*
- Stigma 1; achenes tapering at both ends, often stipitate. 2
- 2. Beak of achene less than 1.5 mm long or lacking; stipe of achene less than 1 mm long or lacking; staminate head solitary. 3
- Beak of achene 1.5–6 mm long; stipe of achene 1–5 mm long; staminate heads 2 or more. 4
- 3. Achenes short-beaked; pistillate heads all borne in leaf axils; staminate head separated from the uppermost pistillate one. *S. minimum*
- Achenes beakless; some pistillate heads borne above leaf axils; staminate head contiguous with the uppermost pistillate one. *S. hyperboreum*
- 4. Beak of achene flattened, strongly curved; leaves all floating, not keeled. *S. fluctuans*
- Beak of achene not flattened or strongly curved; leaves erect or floating, keeled. 5
- 5. Beak of achene about as long as body. *S. chlorocarpum*
- Beak of achene about half as long as body. 6
- 6. Leaves 2–5 mm wide, rounded on back; fruiting heads 1–2 cm in diam. *S. angustifolium*
- Leaves 5–10 mm wide, flat on back; fruiting heads 2–2.5 cm in diam. *S. multipedunculatum*

Sparganium angustifolium Michx.

NARROW-LEAVED BUR-REED

Stems floating and elongated, or erect to 30–50 cm high; leaves 4–10 mm wide, flat. Inflorescence with staminate heads 2–4, distant; pistillate heads 2 or 3; fruiting heads 7–10 mm in diam. Not common; in slow running water.

Sparganium chlorocarpum Rydb. var *acaule* (Beeby) Fern. STEMLESS BUR-REED

Stems slender, erect, to 75 cm high; leaves 2–10 mm wide, overtopping inflorescence. Staminate heads 4–9, mostly distant; pistillate heads 1–4; fruiting heads 1.5–2.5 cm in diam.




Fig. 19. Common cattail, *Typha latifolia* L.

Sparganium eurycarpum Engelm.

BROAD-FRUITED BUR-REED

Stems stout, 50–150 cm high; leaves 7–10 mm wide, flat. Inflorescence branched; staminate heads 2–20; pistillate heads 1–3; fruiting heads 20–25 mm in diam.

Sparganium fluctuans (Morong) Robins.

Stems floating, to 1.5 m long; leaves flat, 3–8 mm wide, thin. Inflorescence branched; staminate heads 4–6; pistillate heads 3–6, mostly on the branches; fruiting heads 10–20 mm in diam.

Sparganium hyperboreum Laest.

NORTHERN BUR-REED

Stems slender, 10–30 cm long; leaves 1–5 mm wide, thick. Staminate head one; pistillate heads 1–3, with at least 1 head above the axils; fruiting heads 5–12 mm in diam. Rare; in shallow water; Boreal forests.

Sparganium minimum (Hartm.) Fries

SMALL BUR-REED

Stems slender, floating, to 50 cm long; leaves 3–7 mm wide. Staminate head one; pistillate heads 1–3, all axillary; fruiting heads 5–7 mm in diam. Rare; in shallow water; Boreal forest.

Sparganium multipedunculatum (Morong) Rydb.

MANY STALKED BUR-REED

Stems stout, to 80 cm high; leaves 5–12 mm wide, flat. Staminate heads 1–4, crowded; pistillate heads 1–5, above the axils, with lower ones often on peduncles; fruiting heads 20–25 mm in diam. Not common; in shallow water; throughout the Prairie Provinces.

ZOSTERACEAE—pondweed family

Annual or perennial aquatic or marsh plants growing entirely in water. Roots usually fibrous, often growing from the lower nodes of the stem. Leaves varying in shape from thread-like to broad, and all either floating or submersed. Flowers inconspicuous, with neither sepals nor petals.

- 1. Flowers bisexual, appearing above the surface of the water; stamens 2–4; leaves alternate. 2
- Flowers unisexual, usually developed below the surface of the water; stamens 1; leaves opposite or all basal from the crown of the root. 3
- 2. Stamens 4; flowers in spikes; fruit without stem. *Potamogeton*
- Stamens 2; flowers not in spikes; fruit long-stemmed. *Ruppia*
- 3. Annual plants; stemless; inflorescence either solitary in leaf axils or in spikes on summit of scape. *Lilaea*
- Plants with stems; leaves opposite. 4

4. Inflorescence of solitary flowers in axils of leaves. *Najas*
 Inflorescence clusters of flowers in axils of leaves. *Zannichellia*

Lilaea flowering quillwort

Lilaea scillioides (Poir.) Hauman

FLOWERING QUILLWORT

An annual marsh or mud plant with narrow leaves, circular in cross section, and clustered. Plants 8–15 cm high, bearing spikes up to 1 cm long of mixed male and female flowers and also solitary female flowers enclosed in a sheath at the base of the leaves. Fruits small achenes, with those of the flowers of spikes winged and ridged, and those of the basal flowers larger and not winged. A plant of the Pacific Coast, with very few records known from the Canadian Prairie Provinces, these being reported in saline sloughs in the vicinity of Cypress Hills and the southeastern part of Alberta. Some authorities place this species in a separate family, the Lilaeaceae, or quillwort, family. Syn.: *L. subulata* Humb. & Bonpl.

Najas naiad

Najas flexilis (Willd.) Rostk. & Schmidt

SLENDER NAIAD

An annual aquatic plant. Leaves opposite, 1.0–2.5 cm long, with widened bases and conspicuous sheaths. Flowers inconspicuous and borne in leaf axils, with male and female inflorescence borne on the same plant. Rare; in shallow lakes and slow-moving water; near Edmonton and north of Winnipeg.

Potamogeton pondweed

Perennial aquatic plants with fibrous roots from the lower nodes of the stems. Leaves generally submersed but in one species some floating; varying from thread-like to broad. Flowers with neither sepals nor petals, sometimes borne on spikes projecting from water and sometimes in axils of leaves.

1. Plants with all leaves similar and submersed. 2
 Plants with broader leaves floating and narrower leaves submersed. *P. gramineus*
2. Leaves broad, their bases clasping stem. *P. richardsonii*
 Leaves linear and thread-like. 3
3. Stipules free from base of leaf. *P. foliosus*
 Stipules fused with base of leaf, forming a sheath at least 1 cm long. 4
4. Stigmas raised on evident style; nutlets with 2 keels. *P. pectinatus*
 Stigmas inconspicuous; nutlets not keeled. 5
5. Plant slender; sheaths close around stem. *P. interior*
 Plant coarse; sheaths 2–5 times diameter of stem. *P. vaginatus*

Potamogeton foliosus Raf. LEAFY PONDWEED

More likely to be found toward the eastern part of the Prairie Provinces, but not common.

Potamogeton gramineus L. VARIOUS-LEAVED PONDWEED

Easily distinguishable by broad floating leaves and narrower submersed ones. Not common; but may be expected in lakes; Boreal forest.

Potamogeton interior Rydb. INLAND PONDWEED

Fairly common in alkaline ponds in eastern portion of the Prairie Provinces.

Potamogeton pectinatus L. SAGO PONDWEED, FENNEL-LEAVED PONDWEED

Our commonest thread-leaved pondweed; found in ponds, lakes, and streams; throughout the Prairie Provinces.

Potamogeton richardsonii (Benn.) Rydb. RICHARDSON'S PONDWEED

Easily recognizable by broad clasping leaves. One of the commonest species; found in ponds and streams; throughout the Prairie Provinces.

Potamogeton vaginatus Turcz. SHEATHED PONDWEED

Distinguishable from sago pondweed by the broadened stipular sheath at the base of the narrow leaves. Fairly common; in larger sloughs and lakes; throughout the Prairie Provinces.

Ruppia ditch-grass

Submersed perennial plants having hair-like stems and thread-like single-nerved leaves with a membranous sheath at base. Flowers perfect and clustered on a slender stem, the stem elongating into a spiral coil and curling up after fertilization of the flowers. Not common; in brackish and saline sloughs; throughout the entire Prairie Provinces.

Leaf sheaths 6–15 mm long; fruit 2 mm long
or less. *R. maritima*

Leaf sheaths 18–30 mm long; fruit 3 mm or
longer. *R. occidentalis*

Ruppia maritima L. DITCH-GRASS

Widespread, but uncommon in the Prairie Provinces.

Ruppia occidentalis S. Wats. WESTERN DITCH-GRASS

Commoner than *R. maritima*; in sloughs over the entire Prairie Provinces.

Zannichellia horned-pondweed

Zannichellia palustris L. HORNED-PONDWEED

A submersed, branching aquatic plant having thread-like, opposite, single-nerved leaves with membranous sheaths at the base. Flowers of both sexes

borne in axils of leaves; fruits curved, nut-like, 3–5 mm long, with a short beak from which this plant derives its common name. Very common; in brackish ponds; throughout Prairies and Parklands.

JUNCAGINACEAE—arrow-grass family

Flowers in a bracted, few-flowered raceme;

stems leafy. *Scheuchzeria*

Flowers in a bractless, many-flowered spike-

like raceme; stems leafless. *Triglochin*

Scheuchzeria

Scheuchzeria palustris L.

A rush-like bog plant with stems 10–40 cm high. Leaves 10–40 cm long, with the upper ones reduced to bracts; sheaths of basal leaves often 10 cm long, with ligule to 12 mm long. Inflorescence a few-flowered raceme; flowers white; perianth segments 3 mm long, 1-nerved, membranous. Follicles 4–8 mm long, with lower ones on pedicels to 25 mm long. Rare; in peat bogs and lake-shores; Boreal forest.

Triglochin arrow-grass

Perennial marsh or semiaquatic herbs with short rootstocks. Leaves linear or rush-like, semicylindric, all basal and clustered, and bearing membranous sheaths. Flowers perfect, borne in tall, slender, spike-like racemes. Fruit capsules splitting open at maturity. **Poisonous** to cattle and sheep.

Plant stout; carpels 6; fruit oblong or ovoid,

obtuse at base. *T. maritima*

Plant slender; carpels 3; fruit linear or club-

shaped, tapering at base. *T. palustris*

Triglochin maritima L. (Fig. 20)

SEASIDE ARROW-GRASS

A stout plant, with rootstock but no stolons. Leaves long, narrow, and half cylindric, up to 30 cm long and 3 mm wide, with bases usually covered with old leaf sheaths. Flowering stem 40–80 cm high, with flowers in a raceme at summit; raceme up to 50 cm long. Fruit about twice as long as thick, 6 mm long and 2–3 mm in diam, on short stalks. Common; found over the whole area in marshy and wet places. **Poisonous** to cattle and sheep.

Triglochin palustris L.

MARSH ARROW-GRASS

A slender plant, with rootstock and slender stolons. Leaves similar to the preceding species, but usually about 10–20 cm long. Flowers on slender stalks. Fruit about 3–5 times as long as thick, usually 6–8 mm long and 1 mm in diam, on fine stems paralleling the stalk. Found in marshy places throughout the Prairie Provinces, but not nearly so common as *T. maritima*. **Poisonous** to cattle and sheep.



Fig. 20. Seaside arrow-grass,
Triglochin maritima L.

ALISMACEAE—water-plantain family

- Flowers all perfect; fruit a single ring of
carpels. *Alisma*
- Flowers of one sex, with lower ones female
and upper ones male; fruit in dense globular
heads. *Sagittaria*

Alisma water-plantain

Perennial aquatic or marsh plants growing from stout corm-like root-stocks producing offshoots. Leaves generally ovate or oblong, with several parallel veins; borne on long stems from the crown of the root. Flowers perfect, with 3 green sepals and 3 white petals. Fruits flat-sided short-beaked achenes.

Flowering stems usually not extending above the leaves, its branches generally curved downward; achenes as wide as long; leaves generally ovate-lanceolate. *A. gramineum*

Flowering stems extending above the leaves, its branches ascending; achenes longer than wide; leaves usually ovate. *A. plantago-aquatica*

Alisma gramineum K. C. Gmel. NARROW-LEAVED WATER-PLANTAIN

Leaves usually long and narrow or ovate-lanceolate, 30–80 mm long. Flowering stems usually shorter than the leafage. Common in Prairies. Syn.: *A. geyeri* Torr.

Alisma plantago-aquatica L. (Fig. 21) COMMON WATER-PLANTAIN

Leaves oblong to ovate, 5–18 cm long. Inflorescence diffuse, to 30 cm or higher, much overtopping the leafage. Common throughout the Prairie Provinces.

Sagittaria arrowhead

Perennial water or marsh plants from fleshy rootstocks. Leaves all basal, usually broadly arrow-shaped, borne on long stems; occasionally reduced to mere thickened stems (called phyllodia) replacing blades. Flowers having 3 sepals and 3 waxy white petals; female flowers, borne lowest on the stem, developing before male flowers. Fruits achenes, crowded into globular heads.

Beak of achenes erect and very minute. *S. cuneata*

Beak of achene horizontal and long. *S. latifolia*

Sagittaria cuneata Sheld. (Fig. 22) ARUM-LEAVED ARROWHEAD

Readily identified by its broad arrow-shaped leaves and waxy white flowers, borne in whorls of 3 on the long stem. Leaves up to 10–15 cm long, but submersed stems often lacking blade, the thickened stem (phyllodia) replacing the blade. Flowers 6–12 mm in diam, later forming globular seed heads up to 15 mm in diam, green at first turning black later. Very common; in water and marshy habitats; in all parts of the Prairie Provinces.

Sagittaria latifolia Willd. BROAD-LEAVED ARROWHEAD

Resembling *S. cuneata*, but differing in having the beak of achene horizontal and fairly long. Leaves very variable, being either broadly or quite narrowly arrow-shaped. Not common over most of the Prairie Provinces, but found in central and eastern Parklands.



Fig. 21. Common water-plantain, *Alisma plantago-aquatica* L.



Fig. 22. Arum-leaved arrowhead, *Sagittaria cuneata* Sheld.

HYDROCHARITACEAE—frog's-bit family

Elodea waterweed

Elodea canadensis Michx.

CANADA WATERWEED

Submersed perennial aquatic plants with fibrous roots springing from the nodes of the stems. Leaves not stalked, borne in whorls of 2–4, oblong-ovate, to 5 mm long. Flowers rarely found, but borne on the end of an apparent stalk, with male flowers on one plant and female on another. Not common but probably widespread; in still or slow-moving water; throughout the Prairies and Parklands.

GRAMINEAE—grass family

Annual or perennial herbs (Fig. 23), with stems usually hollow except at the nodes. Leaves borne on two sides of stem, one growing from each node, and consisting of a sheath, usually split, enveloping the main stem and a blade, which is a continuation of the sheath, growing at an angle to the stem. Blades flat or rolled, narrow, and without stalks. Inflorescence (Fig. 24) in spikes, racemes, or panicles, each composed of spikelets borne on a rachis. Each spikelet consisting of a series of bracts alternating on either side of a rachilla. The spikelet breaking off below or above the empty glumes; this point of articulation is important in identifying grasses. These bracts are called glumes. The lowermost glumes empty or sometimes one or both missing or replaced by bristles; other glumes containing the floret, each having an enveloping palea and lemma; flowering glumes often having bristles called awns. Flower usually perfect, but sometimes unisexual. In the perfect flower, feather-like stigmas arising from the style, usually having three stamens, and the ovary developing into a caryopsis (or grain). A magnifying glass may be needed for identifying the various parts of a grass plant.

Gramineae is such a large family that subdivisions have been made. The family has been divided into two subfamilies, Panicoideae and Festucoideae. These subfamilies are further divided into tribes. Species in 10 of these, Paniceae, Andropogoneae, Phalarideae, Hordeae, Chlorideae, Agrostideae, Avenae, Festuceae, Zizanieae, and Oryzeae (Fig. 25) may be found in the Prairie Provinces.

The following characters distinguish grasses from sedges and rushes:

Character	Gramineae—grasses (p. 73)	Cyperaceae—sedges (p. 181)	Juncaceae—rushes (p. 246)
Culm	Usually hollow; cylindrical or flattened	Filled with pith, rarely hollow; usually 3-sided	Filled with sponge-like pith, cylindrical
Nodes	Conspicuous	Indistinct	Indistinct
Leaf arrangement	Two-ranked	Three-ranked	Principal leaves basal or nearly so
Leaf blade	Usually flat, often folded, involute or bristle-like; glabrous or pubescent	Flat, plicate, or bristle-like; rarely pubescent	Channeled or round; usually glabrous
Leaf margins	Smooth, scabrous, or ciliate	Usually scabrous	Smooth




Fig. 23. Structure of a typical grass.




Fig. 24. Types of inflorescences of grasses.




Fig. 25. Floral characteristics of grass tribes.

Keys to the Grasses

1. Spikelets one- to many-flowered; sterile florets, if present, above the fertile ones (except in Phalarideae, having two sterile florets below the fertile one); articulation usually above the glumes; spikelets usually laterally compressed. Subfamily 1. FESTUCOIDEAE 2
 Spikelets with one fertile floret, and a sterile floret, if present, below the fertile one; articulation below the glumes; spikelets compressed dorsally. Subfamily 2. PANICOIDEAE 9
2. Spikelets with one fertile floret above and two (or rarely one) sterile florets below; no sterile florets above the fertile florets. Tribe 6. PHALARIDEAE
 Spikelets with one to many fertile florets; sterile florets above the fertile florets. 3
3. Spikelets unisexual, one-flowered; the rachis disarticulating below the glumes. Tribe 8. ZIZANIEAE
 Spikelets perfect or if unisexual not as above; the rachis disarticulating above the glumes. 4
4. Spikelets disarticulating below the glumes, one-flowered; strongly compressed laterally; glumes small or absent. Tribe 7. ORYZEAE
 Spikelets disarticulating above or below the glumes, at least one of these well-developed. 5
5. Spikelets sessile on the rachis in spikes. 6
 Spikelets pedicellate in open or contracted, sometimes spike-like, panicles. 7
6. Spikes terminal, solitary; spikelets arranged singly or in twos or threes on opposite sides of the rachis; the rachis disarticulating in some species. Tribe 2. HORDEAE
 Spikes usually digitate or racemose; spikelets arranged on one side of the rachis in two rows. Tribe 5. CHLORIDEAE
7. Spikelets one-flowered, with only occasionally two florets, in some spikelets in *Muhlenbergia*. Tribe 4. AGROSTIDEAE
 Spikelets two- to many-flowered. 8
8. Glumes shorter than the first floret; lemmas awnless or awned from the tip or a bifid apex. Tribe 1. FESTUCEAE
 Glumes as long as the lowest floret, sometimes enclosing the whole spikelet; lemmas awnless or awned from the back. Tribe 3. AVENEAE

9. Glumes membranous; fertile lemma and palea indurate, sterile lemma like the glumes. Tribe 9. PANICEAE
 Glumes indurate; fertile and sterile lemmas equal in texture. Tribe 10. ANDROPOGONEAE

FESTUCOIDEAE subfamily

Tribe 1. FESTUCEAE

1. Rachilla with long silky hairs; inflorescence a large, plume-like panicle; tall reeds. *Phragmites*
 Rachilla without silky hairs; inflorescence not plume-like; not tall reeds. 2
2. Glumes exceeding the uppermost floret in the spikelet, these shiny. *Dupontia*
 Glumes shorter than the lowest floret. 3
3. Plants dioecious with long rhizomes or stolons; lemmas without a tuft of hairs at the base. *Distichlis*
 Plants monoecious, or if dioecious lemmas with a tuft of hairs at the base, or plants annual. 4
4. Lemmas prominently 3-nerved. 5
 Lemmas 5- to many-nerved, often obscurely so. 7
5. Inflorescence hidden among sharp-pointed leaves; plants annual (Chlorideae). *Munroa*
 Inflorescence not as above. 6
6. Spikelets many-flowered; glumes and lemmas keeled; plants annual. *Eragrostis*
 Spikelets 2-flowered; perennial. *Catabrosa*
7. Callus of florets bearded. 8
 Callus of florets not bearded. 9
8. Plants with rhizomes; lemmas erose at tip, awnless. *Scolochloa*
 Plants with fibrous roots; lemmas bifid at tip, awned. *Schizachne*
9. Spikelets strongly compressed, arranged in one-sided clusters on stiff panicle branches. *Dactylis*
 Spikelets and inflorescence not as above. 10
10. Lemmas obscurely nerved; panicle loose, with drooping branches; plants large. *Arctophila*
 Lemmas distinctly nerved, or if not, plants not as above. 11

11. Lemmas keeled on back.	12
Lemmas rounded on back, or keeled only slightly toward tip.	13
12. Spikelets large; lemmas 2-lobed at tip, awned.	<i>Bromus</i>
Spikelets small; lemmas not 2-lobed, awnless.	<i>Poa</i>
13. Glumes papery; lemmas firm, strongly nerved; sterile florets at tip often reduced to rudiments.	<i>Melica</i>
Glumes not papery; sterile florets not differing from the others.	14
14. Lemmas parallel-nerved, with nerves not converging at the tip of the lemma.	15
Lemmas narrowing at the tip, with nerves converging.	16
15. Plants of freshwater habitats; lemmas prominently nerved; leaf sheaths closed, flattened.	<i>Glyceria</i>
Plants of saline habitats; lemmas not prominently nerved; leaf sheaths open, terete.	<i>Puccinellia</i>
16. Lemmas 2-lobed at the tip, awned or awn-tipped from between the lobes.	<i>Bromus</i>
Lemmas not 2-lobed, awned from the tip or awnless.	17
17. Lemmas awnless; leaves with boat- shaped tip.	<i>Poa</i>
Lemmas awned; leaves not with boat- shaped tip.	<i>Festuca</i>

Tribe 2. HORDEAE

1. Spikelets one at each node of the rachis, or at least never more than one at all nodes.	2
Spikelets 2 or 3 at each node, or at least at most of the nodes.	5
2. Spikelets placed with the narrow side toward the rachis; the glume on that side absent, the terminal spikelet with both glumes developed.	<i>Lolium</i>
Spikelets placed with the broad side toward the rachis; both glumes devel- oped in all spikelets.	3
3. Plants perennial or weedy annuals.	<i>Agropyron</i>
Plants annual, cultivated.	4
4. Glumes 3-nerved, ovate.	<i>Triticum</i>
Glumes 1-nerved, subulate.	<i>Secale</i>

- 5. Spikelets 1-flowered, 3 at each node; the lateral pair pedicellate, usually reduced to awns. *Hordeum*
- Spikelets 2- to 6-flowered, usually 2 at each node; florets all alike. 6
- 6. Rachis continuous. *Elymus*
- Rachis disarticulating. 7
- 7. Glumes short-awned, or awnless; lemmas with awns to 2 cm long. *Agrohordeum*
- Glumes and lemmas both with long divergent awns. *Sitanion*

Tribe 3. AVENEAE

- 1. Lemmas awnless or rarely very short-awned. 2
- Lemmas distinctly awned. 3
- 2. Articulation of rachis above the glumes; glumes about equal in length and width. *Koeleria*
- Articulation of rachis below the glumes; the first glume linear, the second one much wider. *Sphenopholis*
- 3. Articulation of rachis below the glumes; lemmas bearing a bent awn; panicle spike-like. *Trisetum*
- Articulation of rachis above the glumes. 4
- 4. Awn flattened, twisted, borne from a notch between apical teeth of lemma; collar with tuft of hairs. *Danthonia*
- Awn not flattened, borne well below the apex of the lemma; collar glabrous. 5
- 5. Glumes 20–35 mm long, 7- to 9-nerved; plants annual; spikelets pendulous. *Avena*
- Glumes less than 20 mm long, 1- to 5-nerved; spikelets not pendulous. 6
- 6. Spikelets 3- to 7-flowered, 10–15 mm long; blades flat or folded, obscurely nerved, glaucous. *Helictotrichon*
- Spikelets 2-flowered, usually less than 10 mm long. 7
- 7. Lemmas keeled, awned from above the middle. *Trisetum*
- Lemmas rounded on the back, awned from below the middle. *Deschampsia*

Tribe 4. AGROSTIDEAE

- 1. Lemmas thicker and much harder than the glumes. 2

Lemmas not thicker and harder than the glumes.	5
2. Spikelets dorsally compressed; glumes shiny; lemmas awnless; rachis disarticulating below the glumes.	<i>Milium</i>
Spikelets not dorsally compressed; glumes dull; lemmas awned; rachis disarticulating above the glumes.	3
3. Awn weak, short, and deciduous; fruit plump.	<i>Oryzopsis</i>
Awn firm, persistent; fruit slender.	4
4. Awn of lemma simple.	<i>Stipa</i>
Awn of lemma trifid.	<i>Aristida</i>
5. Rachis disarticulating below the glumes.	6
Rachis disarticulating above the glumes.	8
6. Glumes long-awned; panicle dense, spike-like, appearing silky.	<i>Polypogon</i>
Glumes awnless.	7
7. Panicle open, with spreading branches; florets stipitate; glumes not united.	<i>Cinna</i>
Panicle dense, spike-like; florets not stipitate; glumes united at base.	<i>Alopecurus</i>
8. Lemmas awned at the apex or short-pointed.	<i>Muhlenbergia</i>
Lemmas awnless, or the awn inserted below the apex.	9
9. Inflorescence dense and spike-like.	<i>Phleum</i>
Inflorescence an open, branched, panicle.	10
10. Florets bearing a tuft of hairs at the base.	11
Florets without hairs at the base.	12
11. Lemmas awned from the middle or near the base; 3- to 5-nerved; the glumes longer than the lemma.	<i>Calamagrostis</i>
Lemmas awnless, 1-nerved; the first glume shorter than the lemma.	<i>Calamovilfa</i>
12. Glumes longer than the lemma; palea small or obsolete; callus often somewhat hairy.	<i>Agrostis</i>
Glumes shorter than the lemma; palea well-developed; callus smooth.	13
13. Lemma 1-nerved; mature grain plump, free from the lemma and palea.	<i>Sporobolus</i>
Lemma 3- to 5-nerved; mature grain not plump, not readily freed from the lemma and palea.	<i>Arctagrostis</i>

Tribe 5. CHLORIDEAE

1. Plants with imperfect flowers; monoecious or dioecious. *Buchloe*
Plants with perfect flowers. 2
2. Spikelets with more than one perfect floret; inflorescence a few-flowered head hidden among sharp-pointed leaves. *Munroa*
Spikelets with only one perfect floret; inflorescence not as above. 3
3. Spikelets with one or more sterile florets above the fertile ones. *Bouteloua*
Spikelets without sterile florets. 4
4. Spikes very slender; leaves short and narrow. *Schedonnardus*
Spikes not very slender; leaves long and wide. 5
5. Glumes unequal, narrow; spikelets 6–15 mm long; plants with scaly rhizomes. *Spartina*
Glumes equal, boat-shaped; spikelets 3 mm long; plants with fibrous roots, annual. *Beckmannia*

Tribe 6. PHALARIDEAE

- Lower florets staminate; spikelets brownish, shiny; glumes rounded on the back. *Hierochloe*
Lower florets neutral; spikelets greenish, not shiny; glumes boat-shaped, keeled. *Phalaris*

Tribe 7. ORYZEAE

- One genus; glumes lacking; lemmas awnless. *Leersia*

Tribe 8. ZIZANIEAE

- One genus; spikelets unisexual. *Zizania*

PANICOIDEAE subfamily

Tribe 9. PANICEAE

1. Spikelets with one to many bristles forming an involucre. *Setaria*
Spikelets without bristles. 2
2. Glumes or sterile lemmas awned. *Echinochloa*
Glumes and sterile lemmas awnless. 3

- 3. Spikelets in digitate slender racemes at tip of culms. *Digitaria*
- Spikelets in open panicles with slender branches. *Panicum*

Tribe 10. ANDROPOGONEAE

Racemes reduced to one or a few joints; these racemes numerous in large open panicles. *Sorghastrum*

Racemes several- to many-jointed; these racemes solitary or digitately clustered. *Andropogon*

Agrohordeum wild rye

Agrohordeum macounii (Vasey) Lepage

MACOUN'S WILD RYE

Plants densely tufted, erect, 50–100 cm high. Sheath glabrous or sometimes pubescent; blades to 5 mm wide, scabrous. Spike slender, erect to somewhat nodding, 4–12 cm long; rachis disarticulating at maturity; spikelets overlapping, 8–10 mm long, mostly 2-flowered; glumes 5–8 mm long, very narrow; lemmas 7–10 mm long; awns 1–2 cm long. Moist, often alkali flats; Prairies and Parklands.

Agropyron wheatgrass

- 1. Plants with rhizomes. 2
- Plants with fibrous roots. 7
- 2. Stomata forming fine white lines on underside of blades; rhizomes long, yellow white. *A. repens*
- Stomata not forming white lines. 3
- 3. Glumes rigid, 10–12 mm long; plants glaucous, bluish green; auricles often purplish. *A. smithii*
- Glumes not rigid, mostly 6–9 mm long; plants green or gray green; auricles yellowish green. 4
- 4. Awns of lemmas recurved or divergent. 5
- Awns of lemmas straight. 6
- 5. Lemmas pubescent. *A. albicans*
- Lemmas glabrous or scabrous. *A. albicans*
- 6. Lemmas pubescent. *A. dasystachyum*
- Lemmas glabrous or scabrous. *A. riparium*
- 7. Plants annual, introduced. *A. triticeum*
- Plants perennial 8
- 8. Spikes short, with spikelets very closely spaced on the rachis; introduced species. *A. cristatum*
- Spikes elongated, with spikelets not closely spaced on the rachis; native species. 9

9. Spikelets awnless or awn-tipped. 10
 Spikelets long-awned, with awns 1–4 cm
 long. 12
10. Glumes narrow; rachilla scaberulous;
 blades involute. *A. spicatum*
 Glumes wide, 2–2.5 mm; rachilla villous;
 blades flat or nearly so. 11
11. Glumes with thin margins, awn-tipped;
 lemmas usually pubescent; spike 3–8
 cm long. *A. latiglume*
 Glumes not with thin margins, awnless;
 lemmas glabrous; spike 10–25 cm long. *A. trachycaulum*
12. Awn straight or nearly so; blades glab-
 rous, lax; spikes 5–20 cm long. *A. subsecundum*
 Awns divergent or bent. 13
13. Spikelets not closely spaced on the rachis,
 barely overlapping. *A. spicatum*
 Spikelets closely spaced on the rachis,
 overlapping. 14
14. Culms tufted, decumbent to ascending,
 often flexuous; spikes often nodding. *A. scribneri*
 Culms erect; spikes erect. *A. bakeri*

Agropyron albicans Scribn. & Smith

AWNED NORTHERN WHEATGRASS

Plants with tufted culms 40–70 cm high arising from slender rhizomes. Blades flat to involute, 1–3 mm wide, glabrous. Spike 6–15 cm long; spikelets loosely overlapping, 4- to 8-flowered, 1–1.5 cm long; glumes 6–9 mm long, sparsely pubescent; lemmas 7–10 mm long, densely or sparsely pubescent; awn 1–1.5 cm long, divergent at maturity. Prairies.

A variety with glabrous lemmas has been named var. *griffithsii* (Scribn. & Smith) Beetle.

Agropyron bakeri E. Nels.

BAKER'S WHEATGRASS

Plants loosely tufted, with erect culms 50–100 cm high. Sheaths and blades glabrous, scabrous, or sparsely pubescent. Blades flat, to 8 mm wide. Spikes 5–10 cm long; spikelets 10–15 mm long, 3- to 5-flowered, loosely imbricate; glumes 8–10 mm long, awn-tipped; lemmas 10–12 mm long, with the awn to 4 cm long, divergent and recurved when dry. Prairies, southern Rocky Mountains.

Agropyron cristatum (L.) Gaertn. (Fig. 26)

CRESTED WHEATGRASS

Plants densely tufted, with culms 30–50 cm high. Sheaths scabrous or the lowest ones pubescent; blades to 8 mm wide, scabrous to pubescent above. Spikes 2–7 cm long, flat; spikelets 8–15 mm long, 3- to 5-flowered, densely crowded, spreading to ascending; glumes 4–6 mm long, awn-tipped; lemmas 6–8 mm long, awnless or awn-tipped. Introduced forage grass; widely sown for pasture and hay, and escaped from cultivation in many areas. Usually, crested wheatgrass includes several taxa besides *A. cristatum*: *A. desertorum* (Fisch.)




Fig. 26. Crested wheatgrass, *Agropyron cristatum* (L.) Gaertn.

Schult., with more rounded spikes and smaller, less spreading spikelets; *A. cristatiforme* Sarkar, with the spikelets very densely crowded and the culm villose below the spike; and *A. pectiniforme* R. & S., with the spikelets spaced in the spike.

Agropyron dasystachyum (Hook.) Scribn.

NORTHERN WHEATGRASS

Plants with tufted culms 40–70 cm high arising from slender creeping rhizomes. Sheath glabrous to slightly scabrous; blades to 6 mm wide, often involute, strongly veined and scabrous above. Spike 6–15 cm long, often involute, strongly veined, and scabrous above; spikelets 10–15 mm long, 4- to 8-flowered, loosely to closely imbricate; glumes 6–9 mm long, acute or awn-tipped; lemmas 8–10 mm long, more or less densely pubescent, awnless or with a short awn. Grasslands; throughout Prairies and Parklands.

Agropyron latiglume (Scribn. & Sm.) Rydb.

BROAD-GLUMED WHEATGRASS

Plants loosely tufted, with culms 20–50 cm high, ascending to geniculate. Blades 3–5 mm wide, flat, short-pubescent on both sides. Spikes 3–7 cm long; spikelets 10–15 mm long, 3- to 5-flowered, closely imbricated; glumes 7–10 mm long, very broad, thin-margined, awn-tipped; lemmas 10–12 mm long, pubescent, awn-tipped or awnless. Alpine meadows; Rocky Mountains, southwest Alberta.

Agropyron repens (L.) Beauv. (Fig. 27)

QUACK GRASS, COUCH GRASS

Plants with culms 50–100 cm high, arising in tufts from long, creeping, thick yellowish white rhizomes. Sheaths at first often soft pubescent, later glabrous; blades 6–10 mm wide, flat, usually sparsely pubescent above, and the underside glabrous, with stomata visible as fine white lines. Spikes 5–15 cm long; spikelets 10–15 mm long, 4- to 7-flowered; glumes 6–8 mm long, awn-tipped; lemmas 8–10 mm long, with the awn 2–8 mm long. An introduced species, often becoming weedy in waste places and gardens, rarely in cultivated fields.

Agropyron riparium Scribn. & Sm.

STREAMBANK WHEATGRASS

Plants with culms 30–80 cm high, arising in tufts from creeping slender rhizomes. Sheaths glabrous; blades 1–3 mm wide, often involute. Spikes 5–10 cm long; spikelets 10–15 mm long, 5- to 7-flowered, closely imbricated. Glumes 6–10 mm long, awnless; lemmas 8–10 mm long, glabrous or sparsely pubescent along margins, awnless. Ravines; Prairies.

Agropyron scribneri Vasey

Plants densely tufted, with culms 10–30 cm high, ascending-spreading. Sheaths usually more or less pubescent; blades 1–3 mm wide, flat, more or less pubescent, mostly basal; culm leaves very short. Spikes 3–7 cm long, often nodding or flexuous; spikelets 8–12 mm long, 3- to 5-flowered, densely crowded; glumes 6–8 mm long, tapering into an awn; lemmas 8–10 mm long, tapering into an awn; awn 15–20 mm long, strongly divergent. Alpine slopes; Rocky Mountains, southwest Alberta.



Fig. 27. Quack grass, *Agropyron repens* (L.) Beauv.

Agropyron smithii Rydb.

WESTERN WHEATGRASS

Plants with culms 30–60 cm high, arising singly or in small tufts from long creeping rhizomes. Sheaths glabrous, with auricles often purplish; blades 3–6 mm wide, stiff, prominently veined, usually very glaucous. Spikes 7–15 mm long, erect; spikelets 10–20 mm long, 6- to 10-flowered; glumes 10–12 mm long, rigid, tapering to a short sharp awn; lemmas 10–12 mm long, glabrous to somewhat pubescent. Moist areas, moderately alkaline river flats; Prairies and Parklands. A form with both glumes and lemmas pubescent has been described as var. *molle* (Scribn. & Sm.) Jones, and has the same distribution as the species.

Agropyron spicatum (Pursh) Scribn. & Smith

BLUEBUNCH WHEATGRASS

Plants densely tufted, often forming large bunches, with erect culms 60–100 cm high. Sheaths glabrous; blades 3–5 mm wide, flat to loosely involute, glabrous to minutely pubescent below, finely pubescent above, usually glaucous. Spikes 7–17 cm long, sometimes longer; spikelets 10–15 mm long, 6- to 8-flowered, mostly shorter than the internodes; glumes 6–8 mm long, acute or awn-tipped; lemmas 8–10 mm long, with the awn 10–20 mm long, strongly divergent. Prairies, southwest Alberta. A form with awnless lemmas has been described as var. *inerme* Heller.

Agropyron subsecundum (Link) Hitchc. (Fig. 28)

AWNED WHEATGRASS

Plants loosely tufted, with erect culms 50–100 cm high. Sheaths densely pubescent in young plants, becoming glabrous; blades 6–10 mm wide, flat, pubescent when young. Spikes 6–15 or 20 cm long, erect to slightly nodding; spikelets 12–20 mm long, imbricated, 5- to 7-flowered, often twisted to one side of the rachis; glumes 12–15 mm long, broad, tapering into a short awn; lemmas 10–12 mm long, with the awn 10–30 mm long. Moist areas, fescue prairie, woodlands; Parklands and Boreal forest. Often considered to be a variety of *A. trachycaulum* (Link) Malte var. *unilaterale* (Cass.) Malte.

Agropyron trachycaulum (Link) Malte (Fig. 29)

SLENDER WHEATGRASS

Plants tufted, with culms 50–100 cm high, erect or somewhat decumbent at base. Sheaths glabrous, sometimes purplish at base; blades 4–6 mm wide, flat, somewhat scabrous on both sides. Spikes 10–25 cm long, usually erect to slightly nodding; spikelets 15–20 mm long, 5- to 8-flowered, somewhat imbricated or remote; glumes 10–12 mm long, awnless; lemmas 12–15 mm long, awnless or awn-tipped. Moist areas, around sloughs and lakes, open woods and meadows; throughout the Prairie Provinces.

Agropyron triticeum Gaertn.

Plants loosely tufted, annual, with culms 10–30 cm high, decumbent at base or erect. Blades 2–3 mm wide, flat. Spikes 10–15 mm long, oval to ovate, thick; spikelets 5–7 mm long, densely crowded; glumes 4–6 mm long, awnless; lemmas 6–8 mm long, awnless. Introduced; southeast Alberta.




Fig. 28. Awned wheatgrass, *Agropyron subsecundum* (Link) Hitchc.



Fig. 29. Slender wheatgrass, *Agropyron trachycaulum* (Link) Malte.

Agrostis bent grass

- 1. Palea present, at least half as long as the lemma. 2
 Palea absent or minute. 4
- 2. Plants tufted; rachilla prolonged behind the palea as a bristle. *A. thurberiana*
 Plants rhizomatous or stoloniferous; rachilla not prolonged behind the palea. 3
- 3. Plants often with long stolons; panicle contracted even when in flower; blades to 3 mm wide. *A. stolonifera* var. *major*
 Plants often rhizomatous, sometimes rooting at the nodes of decumbent stems; panicle open when in flower; blades usually more than 3 mm wide. *A. stolonifera*
- 4. Panicle narrow, with the branches appressed or ascending. 5
 Panicle open, with the branches spreading or reflexed. 6
- 5. Culms usually less than 3 dm high; panicle to 6 cm long, purple. *A. variabilis*
 Culms usually more than 3 dm high; panicle to 30 cm long, greenish. *A. exarata*
- 6. Panicle very diffuse; spikelets about 2 mm long; lemmas awnless. *A. scabra*
 Panicle not very diffuse; spikelets about 3 mm long; lemmas awned. *A. borealis*

Agrostis borealis Hartm.

NORTHERN BENT GRASS

Plants tufted, with erect culms 5–40 cm high. Sheaths glabrous; leaves 1–3 mm wide, mostly basal. Panicle 1–10 cm long, with lower branches whorled, spreading; glumes 2–3 mm long; lemma awned, slightly shorter than glumes, with the awn usually twisted, bent, and exserted. Slopes and moist areas; Rocky Mountains, Boreal forest.

Agrostis exarata Trin.

SPIKE REDTOP

Plants tufted, with culms 20–100 cm high, slender to somewhat stout. Sheaths glabrous to somewhat scabrous; blades 2–10 mm wide, scabrous. Panicle 5–30 cm long, narrow, dense or slightly open; glumes 2.5–4 mm long, acuminate to awn-tipped, scabrous in the keel; lemma 1.5–2 mm long, with the midrib excurrent as a short awn. Moist areas; Prairies and Parklands.

Agrostis scabra Willd. (Fig. 30)

ROUGH HAIR GRASS

Plants tufted, sometimes densely so, with erect culms 30–70 cm high. Sheaths glabrous; blades 1–3 mm wide, scabrous. Panicle 15–25 cm long, diffuse, with the branches spreading to ascending, branching above the middle; glumes 2–2.5 mm long, acuminate; lemma 1.5–1.7 mm long, sparsely pubescent at base. Meadows, open woods, waste places; throughout the Prairie Provinces.




Fig. 30. Rough hair grass, *Agrostis scabra* Willd.

Plants tufted, often with rhizomes, with culms erect or decumbent, or forming stolons, which root at the internodes. Sheaths glabrous; blades 4–8 mm wide, flat or somewhat involute to folded, scabrous on both sides. Panicle 5–30 cm high, usually pyramidal; glumes 2–3.5 mm long, acuminate, somewhat pubescent on the keel; lemma 1.5–2.3 mm long, sometimes short-awned; palea about half as long as lemma. Moist areas in Boreal forest, or as escape from cultivation. Often listed as *A. alba* L., but this name appears to be based on a rather depauperate specimen of *Poa nemoralis* L. and is therefore invalid. Our plants are in part var. *genuina* (Schur) A. & G. (Fig. 31), with rather numerous and creeping rhizomes, and var. *major* Farw. forma *palustris* (Huds.) J. & W., lacking or with few short rhizomes.

Agrostis thurberiana Hitchc.

Plants in small tufts, with erect culms 20–40 cm high. Sheaths glabrous; blades 1–2 mm wide, mostly basal, crowded. Panicle 5–7 cm long, somewhat drooping; glumes 1.5–2 mm long, often purplish; lemmas about as long as glumes; palea about 1.5 mm. Bogs and meadows; Rocky Mountains, southwest Alberta.

Agrostis variabilis Rydb.

Plants densely tufted, with erect culms 10–25 cm high. Sheaths glabrous; blades about 1 mm wide, flat. Panicle 2–5 cm long, narrow, with branches ascending; glumes 2–2.5 mm long, purple; lemma 1.5 mm long. Rocky slopes and creek banks; Rocky Mountains, southwest Alberta.

Alopecurus water foxtail

- 1. Panicles thick, about 10 mm wide. 2
 Panicles slender, about 5 mm wide. 3
- 2. Panicles very woolly, ovoid, to 4 cm long;
 plants with rhizomes. *A. alpinus*
 Panicles not very woolly, cylindric, to 10
 cm long; plants without rhizomes. *A. pratensis*
- 3. Culms erect or spreading; awn of lemma
 barely exerted; anthers orange yellow. *A. aequalis*
 Culms usually decumbent, geniculate;
 awn of lemma much exerted; anthers
 pale yellow or purplish. *A. geniculatus*

Alopecurus aequalis Sobol.

Plants forming large tufts or sods, with culms 15–60 cm high, erect or spreading. Blades 1–4 mm wide, gray green. Panicle slender, cylindric, 2–7 cm long, 4 mm wide; glumes 2.5–3 mm long; lemmas 2.5 mm long, with the awn inserted at the middle, hardly exerted. Anthers orange. Moist to wet soils, around sloughs, lakeshores, river flats; common throughout the Prairie Provinces.

Alopecurus alpinus J. E. Smith

Plants from creeping rhizomes, with stiff culms 10–60 cm high, erect or decumbent at base. Sheaths glabrous, often inflated. Blades 3–5 mm wide.




Fig. 31. Redtop, *Agrostis stolonifera* L. var. *genuina* (Schur.) A. & G.

Panicle thick, ovoid or oblong, 1–4 cm long, 1 cm wide, woolly; glumes 3–4 mm long, woolly; lemmas 3–4 mm long, with the awn inserted near the base, exserted. Rare; southwest Saskatchewan, southern Rocky Mountains.

Alopecurus geniculatus L.

WATER FOXTAIL

Plants forming tufts or sods, with culms to 60 cm high, at first spreading, later geniculate ascending, often rooting at the lower nodes. Blades to 6 mm wide, gray green, strongly nerved. Panicle cylindrical, to 5 cm long, 7 mm wide; glumes 2.5–3 mm long, short pubescent toward the apex; lemmas 2.5 mm long, short pubescent, with the awn inserted at the base, well-exserted. Anthers light yellow. Rare; moist and wet soils; throughout the Prairie Provinces.

Alopecurus pratensis L.

MEADOW FOXTAIL

Plants from short rhizomes, with culms to 100 cm high, erect or decumbent at base. Sheaths glabrous; blades to 6 mm wide or wider, flat. Panicle cylindrical, 3–10 cm long, often wider than 1 cm; glumes about 5 mm long, whitish with green veins; keel and margins ciliate; lemmas 4 mm long, with the awn inserted at the base, 1 cm long, exserted. Introduced forage grass, occasionally seeded in moist meadows; throughout the Prairie Provinces.

Andropogon bluestem

1. Racemes solitary on peduncles. *A. scoparius*
Racemes 2–5 on peduncles, digitate. 2
2. Plants with long rhizomes; spikes villous. *A. hallii*
Plants with short rhizomes; spikes not
villous. *A. gerardi*

Andropogon gerardi Vitman (Fig. 32)

BIG BLUESTEM

Plants in large tufts, with offshoot on short rhizomes; culms 100–150 cm high. Sheaths pubescent; blades flat, 6–10 mm wide, somewhat pubescent at base, otherwise glabrous, blue green to glaucous. Racemes 3–6, 5–10 cm long, usually purplish; sessile spikelet 7–10 mm long, with first glume having awn 10–20 mm long, geniculate, twisted below; pedicellate spikelet 7–10 mm long, awnless, staminate. Prairies, southeastern Parklands.

Andropogon hallii Hack.

SAND BLUESTEM

Plants in tufts with extensive creeping rhizomes; culms 100–150 cm high; sheaths somewhat pubescent to glabrous; blades 6–10 mm wide, glabrous, blue green to purplish. Racemes 3–6, 5–10 cm long, purplish, densely villous with gray or yellowish hairs; sessile spikelet 7–10 mm long, with first glume having awn 5 mm long; pedicellate spikelet 7–10 mm long, awnless, staminate. Sand hills; southeastern Parklands.

Andropogon scoparius Michx. (Fig. 33)

LITTLE BLUESTEM

Plants densely tufted, with short scaly rhizomes; erect culms 30–70 cm high. Sheaths keeled, glabrous; blades 5–8 mm wide, flat to folded, light green to blue green, glaucous. Racemes 3–6 cm long, several to many on long ascending peduncles, with rachis pilose; sessile spikelet 6–8 mm long, with the



Fig. 32. Big bluestem, *Andropogon gerardi* Vitman.




Fig. 33. Little bluestem, *Andropogon scoparius* Michx.

awn 8–15 mm long; pedicellate spikelet reduced, short-awned. Usually calcareous soils; Prairies, southern Manitoba, southern Saskatchewan, southwest Alberta.

Arctagrostis

Arctagrostis latifolia (R. Br.) Griseb.

Plants with creeping rhizomes; tufted culms 30–100 cm high, or occasionally to 150 cm. Sheath smooth, glabrous; blades 4–10 mm wide, flat, rather lax. Panicle 3–25 or to 30 cm long, somewhat open to narrow, with the branches 1–12 cm long, erect and appressed to ascending, usually 3 or 4 at one side of the stem, often interrupted or lobed; spikelets 2.5–5 mm long, with first glume 2–3.5 mm long, 1- to 3-nerved, and second glume 3–5 mm long, 3-nerved; lemmas 3.5–5 mm long, 5-nerved, pubescent, with the palea as long as the lemma, obscurely to strongly 3-nerved, pubescent. A species of variable size, appearance, and color; panicle ranging from yellowish green through purple-tinged to deep purple; anthers yellow or purple. Rare; in marshes and tundra; Boreal forest.

Arctophila

Arctophila fulva (Trin.) Anderson

Plants with creeping rhizomes; culms 30–90 cm high, erect or decumbent. Sheaths strongly ribbed; blades 4–10 mm wide, flat. Panicle 6–20 cm long, usually open, with branches 5–8 cm long, ascending to spreading, and the lower ones often reflexed; spikelets 3–5 mm long, usually purplish, 2- to 6-flowered, with the first glume 2–3 mm long, and the second glume 3–4 mm long; lemmas 2.5–4 mm long, glabrous, obtuse. Rare; wet shores and marshes; Boreal forest.

Aristida three-awn

Plants easily distinguished by the 3 lemmas.

Aristida longiseta Steud.

RED THREE-AWN

Plants tufted, with erect culms 20–30 cm high. Sheaths scabrous with minute prickles. Blades 1–2 mm wide, often involute or bristle-like, very scabrous on both sides. Panicle 3–4 cm long without the awns, narrow, erect, few-flowered; branches ascending or appressed; first glume 8–10 mm long; second glume 15–20 mm long; lemmas 12–15 mm long, terete, with awns 6–8 cm long, divergent. Rare; in grassland, on slopes; Prairies.

Avena oat

Spikelets mostly 2-flowered; awn usually straight or absent; lemmas glabrous. *A. sativa*

Spikelets mostly 3-flowered; awn stout, twisted, geniculate; lemmas pubescent. *A. fatua*

Avena fatua L. (Fig. 34)

WILD OAT

Plants tufted, with culms 30–75 cm high or sometimes to 100 cm, erect, stout. Sheaths smooth; blades 4–8 mm wide, flat. Panicle open, with branches ascending, spreading; spikelets mostly 3-flowered; glumes 2–2.5 cm long; the lower lemmas 15–20 mm long, the upper ones shorter, pubescent with stiff brown hairs, with the awn to 4 cm long, twisted and geniculate. A noxious weed in cultivated fields. Has become very common in many areas; throughout Prairies and Parklands.

Avena sativa L.

OAT

The cultivated form, derived from *A. fatua*, differing from it in having mostly 2-flowered spikelets, glabrous lemmas, and the awn small and straight or lacking. Widely cultivated throughout the Prairie Provinces.

Beckmannia slough grass

Beckmannia syzigachne (Steud.) Fern. (Fig. 35)

SLOUGH GRASS

Plants tufted, with culms 30–70 cm high, erect or decumbent, and sometimes spreading, geniculate ascending. Sheaths glabrous; blades to 12 mm wide, light green. Panicle 10–25 cm long; branches 1–5 cm long, appressed to ascending; spikes crowded, 1–2 cm long; spikelets 1-flowered, 3 mm long, with glumes 3 mm long, wrinkled, deeply keeled; lemmas 7.5 mm long. Wet areas, slough margins, lakeshores, valuable as hay, and readily eaten by cattle; throughout the Prairie Provinces.

Bouteloua grama

Spikes 10–50, pendulous. *B. curtipendula*

Spikes 1–3, spreading. *B. gracilis*

Bouteloua curtipendula (Michx.) Torr.

SIDE-OATS GRAMA

Plants tufted, from short scaly rhizomes, with erect culms 50–75 cm high. Sheaths usually pubescent; blades to 7 mm wide, flat to convolute, with few scattered hairs. Spikes 35–50, usually pendulous, secund; spikelets 5–8 in a spike, 6–10 mm long; fertile lemma 6 mm, mucronate; sterile lemma with 3 awns. Rare; in grassland; southeastern Parklands.

Bouteloua gracilis (HBK.) Lag. (Fig. 36)

BLUE GRAMA

Plants densely tufted, from very short scaly rhizomes, with erect culms 20–50 cm high. Sheaths glabrous or sparsely pubescent; blades to 3 mm wide, light green to dark or reddish green, twisted. Spikes 1–3, usually 2, 2.5–5 cm long, falcate, spreading at maturity; spikelets numerous, 70–80, about 5 mm long; first glume 2–3 mm, second glume 3–4.5 mm; fertile lemma 3–4 mm long, pubescent; sterile lemma densely bearded. Common; in grassland; Prairie and Parkland.



Fig. 34. Wild oat, *Avena fatua* L.



Fig. 35. Slough grass, *Beckmannia syzigachne* (Steud.) Fern.




Fig. 36. Blue grama, *Bouteloua gracilis* (HBK.) Lag.

Bromus brome

1. Plants with rhizomes. 2
Plants with fibrous roots. 3
2. Lemmas glabrous; blades and culms glabrous or somewhat scabrous. *B. inermis*
Lemmas pubescent, at least along the margins; blades pilose above; culms pubescent at nodes. *B. pumpehianus*
3. First glume 1-nerved. 4
First glume 3- to 5-nerved. 7
4. Plants annual; sheath and blade softly pubescent; spikelets drooping; culms to 5 dm high. *B. tectorum*
Plants perennial; sheath and blades pubescent or glabrous; culms to 1 m high. 5
5. Lemmas rather evenly pubescent on the back, but more densely so on lower margins. *B. purgans*
Lemmas pubescent along margins and lower part only, with the upper part glabrous. 6
6. Sheath and blades soft pubescent; blades 5–8 mm wide; lemmas narrow; ligules prominent; awns 6–8 mm long. *B. vulgaris*
Sheath and blades glabrous or short pubescent; blades 4–6 mm wide; lemmas broad; ligules inconspicuous. *B. ciliatus*
7. Plants annual; sheath and blade softly villose. 8
Plants perennial; sheath and blade glabrous or sparingly pubescent. 9
8. Spikelets 5–8 mm wide, inflated; first glume 5-nerved; awns strongly divergent. *B. squarrosus*
Spikelets less than 5 mm wide; first glume 3-nerved; awns flexuous. *B. japonicus*
9. Spikelets strongly flattened, with lemmas keeled; panicle with ascending branches. *B. marginatus*
Spikelets not strongly flattened, with lemmas rounded on the back; panicle drooping. 10
10. Second glume 3-nerved. *B. porteri*
Second glume 5-nerved. *B. kalmii*

Plants loosely caespitose, with 2–4 culms together 60–100 cm high. Culms and sheaths glabrous or short pubescent at the nodes; blades 4–10 mm wide, glabrous to sparsely villose. Panicle 10–20 cm long, with branches slender, often flexuous, spreading or drooping, up to 15 cm long. Spikelets drooping, 1–2 cm long, 4- to 10-flowered, with first glume lance-subulate, 5–7 mm long, 1-nerved, and second glume lanceolate, 7–10 mm long, 3-nerved; lemmas 8–11 mm long, pubescent along the lower half to three-quarters margin, glabrous or nearly so on the back, with the awn to 5 mm long. Woods, and forest openings; throughout the Prairie Provinces.

Bromus inermis Leyss.

SMOOTH BROME

Erect culms 50–100 cm high, from long creeping rhizomes. Culms, sheaths, and blades usually glabrous or nearly so; blades 5–12 mm wide. Panicle 6–20 cm long, usually contracted and secund at maturity, with 1–4 branches at each node. Spikelets 1.5–2.5 cm long, 7- to 10-flowered, with first glume 6–9 mm long, 1-nerved, and second glume 8–10 mm long, 3-nerved; lemmas 10–12 mm long, 5- to 7-nerved, somewhat pubescent at base, awnless or short-awned. Introduced for forage, and now acclimatized in many areas, particularly in low-lying areas; throughout Prairies and Parklands.

Bromus japonicus Thunb.

JAPANESE CHESS

Plants caespitose, with culms erect or ascending 40–70 cm high. Culms long pubescent on the nodes, with sheaths densely villous; blades 1–3 mm wide, long villous. Panicle 10–20 cm long, open, with branches slender, spreading or drooping, somewhat flexuous. Spikelets 1–2 cm long, 5- to 9-flowered, with first glume 4–6 mm long, broad, 3-nerved, and second glume 6–8 mm long, 5-nerved; lemmas 7–9 mm long, 9-nerved, with awn 8–10 mm long, twisted and flexuous at maturity. Weed along roadsides, waste places, and fields; southwest Alberta.

Bromus kalmii Gray

Plants loosely caespitose, with few culms together 50–100 cm high. Culms usually pubescent at the nodes, with lower sheaths pubescent, and upper ones glabrous or nearly so; blades 5–10 mm wide, pubescent on both sides. Panicle 5–10 cm long, with branches slender, flexuous, drooping. Spikelets few, 1.5–2.5 cm long, 5- to 10-flowered, with first glume 6–7 mm long, 3-nerved, and second glume 7–9 mm long, 5-nerved; lemmas 8–10 mm long, 7-nerved, pubescent on the back, densely so along margins, with the awn 2–3 mm long. Rare; margins of woods, openings; southeastern Parklands.

Bromus marginatus Nees

Plants caespitose, with few culms together 50–100 cm high. Sheaths usually retrorsely pubescent; blades to 12 mm wide, scabrous to sparsely pubescent. Panicles 10–20 cm long, with branches ascending to spreading; spikelets 15–25 mm long, 6- to 10-flowered, with first glume 6–9 mm long, 3-nerved, and second glume 10–15 mm long, 5-nerved; lemmas 10–15 mm long, keeled, minutely pubescent, with the awn 5–7 mm long. Open woods; southern Rocky Mountains.

Bromus porteri (Coult.) Nash

Plants loosely cespitose, with culms 30–60 cm high, slender, pubescent on the nodes. Sheaths pilose to glabrous; blades 2–4 mm wide, scabrous. Panicle nodding, 5–10 cm long, with branches slender flexuous, spreading or drooping. Spikelets 2–2.5 cm long, 5- to 7-flowered, with first glume 5–7 mm long, 3-nerved, and second glume 7–9 mm long, 3-nerved; lemmas 9–12 mm long, densely and evenly pubescent on the back, with the awn 2–4 mm long. Margins of woods and openings; Alberta.

Bromus pumpellianus Scribn.

NORTHERN AWNLESS BROME

Culms erect from long creeping rhizomes. Sheaths usually pubescent, purplish at base; blades to 12 mm wide, scabrous below, often pubescent above. Panicle 6–20 cm long, with branches ascending or the lower ones divergent, usually contracted at maturity, 1–4 branches at each node; spikelets 1.5–3.5 cm long, 7- to 10-flowered, with first glume 6–9 mm long, 1-nerved, and second glume 8–11 mm long, 3-nerved; lemmas 9–12 mm long, 5- to 7-nerved, pubescent at base and along margins, awnless or the awn 1–2 mm long. Open woods, margins, and shrubbery; Cypress Hills, Rocky Mountains.

Bromus purgans L.

CANADA BROME

Plants loosely cespitose, with few culms together 60–100 cm high. Culms pubescent at nodes; sheaths usually retrorsely pubescent; blades 5–10 mm wide, sparsely pubescent to glabrous on both surfaces. Panicle 5–20 cm long, nodding, with branches elongate, spreading or drooping; spikelets 2–3 cm long, 7- to 10-flowered, with first glume 5–8 mm long, 1-nerved, and second glume 6–10 mm long, 3-nerved; lemmas 9–12 mm long, 7-nerved, thinly to densely and evenly pubescent on whole back, with the awn 2–8 mm long. Woods and forest margins; Parklands and Boreal forest.

Bromus squarrosus L.

FIELD BROME

Plants cespitose, with culms 20–30 cm high. Sheaths and blades densely pubescent. Panicles 10–20 cm long, nodding, with branches slender, spreading or drooping; spikelets 1.5–2.0 cm long, 7- to 10-flowered, with first glume 4–6 mm long, 5-nerved, and second glume 5–8 mm long, 7-nerved; lemmas 6–9 mm long, 7-nerved, glabrous, with the awn about 10 mm long, spreading or recurved. Waste places and roadsides.

Bromus tectorum L. (Fig. 37)

DOWNY CHESS

Plants cespitose, with erect culms 30–60 cm high. Sheaths and blades densely long pubescent. Panicle drooping, 10–20 cm long, with many rather short flexuous branches drooping; spikelets 2–3 cm long, 6- to 10-flowered, pubescent throughout, with first glume 5–7 mm long, 1-nerved, and second glume 8–10 mm long, 3-nerved; lemmas 10–12 mm long, 5- to 7-nerved, with the awn 12–17 mm long, straight. Waste places and roadsides.

Bromus vulgaris (Hook.) Shear

Plants cespitose, with culms 70–100 cm high, erect, slender, pubescent on the nodes. Sheaths and blades more or less pubescent; blades 7–12 mm wide.



Fig. 37. Downy chess, *Bromus tectorum* L.

Panicle 10–15 cm long, with branches slender, ascending to divergent, somewhat drooping; spikelets 1.5–2.5 cm long, 5- to 9-flowered, with first glume 5–8 mm long, 1-nerved, and second glume 7–9 mm long, 3-nerved; lemmas 8–10 mm long, sparsely pubescent on the back, with the awn 6–8 mm long. Woods and openings; southwest Alberta.

Buchloe buffalo grass

Buchloe dactyloides (Nutt.) Engelm.

BUFFALO GRASS

Plants stoloniferous, forming a dense sod, with culms of staminate plants to 20 cm high and culms of pistillate plants shorter than the leaves. Sheaths somewhat inflated; blades 1–2 mm wide, sparsely pubescent. Staminate spikes 5–15 mm long, 2 or 3 on a peduncle; spikelets 2-flowered, sessile, closely imbricated, with first glume 1–1.5 mm long, and second glume 1.5–2 mm long; both glumes 1-nerved; lemmas to 4 mm long, 3-nerved. Pistillate spikes with 4 or 5 spikelets, usually 2 spikes on a peduncle; the peduncle short, included in the sheaths of the upper leaves; first glume of pistillate floret 4 mm long, and second glume 5 mm long, firm, thick, rigid, rounded and expanded in the middle, enveloping the floret, and having 3 awn-like lobes at the summit; lemma membranous, 3-nerved, with the summit 3-lobed. Very rare; Prairies.

Calamagrostis reed grass

Medium- to tall-growing creeping-rooted grasses, with open or narrow panicles of small-flowered spikelets. Few or no basal leaves. The tall species usually occur in low, moist localities; found in all types of vegetation across the Prairie Provinces.

1. Awn geniculate, longer than or about as long as the glumes, usually protruding from glumes. 2
 - Awn straight, as long as or shorter than the glumes, usually not protruding from glumes. 5
2. Awn longer than the glumes. 3
 - Awn not longer than the glumes. 4
3. Panicle open, with the branches spreading; glumes acuminate, glabrous, 4–6 mm long; callus hairs half as long as lemma; culms to 40 cm high. *C. deschampsioides*
 - Panicle narrow, with the branches appressed to erect; glumes acute, minutely scabrous, 6–8 mm long; callus hairs one-third as long as lemma; culms 60–100 cm high. *C. purpurascens*
4. Plants strongly rhizomatous, with culms often solitary; blades about 2 mm wide, stiff, involute; collar glabrous. *C. montanensis*
 - Plants rhizomatous, with culms tufted; blades about 4 mm wide, rather lax, flat; collar pubescent. *C. rubescens*

- 5. Panicle open, with the branches spreading and often drooping. *C. canadensis*
 Panicle narrow and more or less contracted, with the branches appressed or ascending. 6
- 6. Blades and upper part of culm scabrous; ligules of upper leaves 4–8 mm long, lacerate; panicle dense, to 25 mm wide. *C. inexpansa*
 Blades smooth or scabrous only at tip; culms smooth except under the panicle; ligules of upper leaves 1–3 mm long, entire; panicle dense or loose, about 10 mm wide. 7
- 7. Panicle lax, purplish; spikelets 4.5–5.5 mm long; lemmas 3.5–4 mm long; leaves short, about 2 mm wide. *C. lapponica*
 Panicle stiff, usually brownish; spikelets 2.5–4.5 mm long; lemmas 2–3.5 mm long; leaves long, about 4 mm wide. *C. neglecta*

Calamagrostis canadensis (Michx.) Beauv.

MARSH REED GRASS

Plants tufted, with creeping rhizomes; culms 60–120 to 150 cm high. Sheaths glabrous; blades 6–10 mm wide, flat, rather lax. Panicle 10–20 cm long, mostly nodding, open to dense and somewhat contracted; glumes 3–4 mm long, acute, somewhat scabrous; lemma 2–3 mm long, thin, smooth, with callus hairs abundant, as long as the lemma; awn delicate, straight, inserted just below the middle. In marshes, lakeshores, meadows, and moist woods; throughout the Prairie Provinces.

In the northern part of the boreal zone var. *scabra* (Presl) Hitchc., with larger spikelets, glumes 4.5–6 mm long, and lemmas 3.5–4 mm, and var. *robusta* Vasey, with glumes 3–5.6 mm and lemmas 3–3.5 mm, have been found.

Calamagrostis deschampsii Trin.

Plants small, tufted; culms 15–40 cm high. Sheaths smooth; blades 2–4 mm wide, glabrous. Panicle 4–8 cm long, open, pyramidal, with branches spreading, bearing spikelets toward the tips; spikelets purplish; glumes 4–6 mm long, acuminate; lemma 3–5 mm long, with callus hairs half as long as the lemma; awn 8–10 mm long, bent, exerted. Boreal forests, Hudson Bay, Manitoba.

Calamagrostis inexpansa A. Gray

NORTHERN REED GRASS

Plants tufted, with slender creeping rhizomes; culms 40–100 cm high. Sheaths glabrous, often purplish at base; blades 2–4 mm wide, flat or somewhat involute, firm. Panicle 5–15 cm long, contracted, dense, with branches ascending-appressed at maturity; glumes 3–4 mm long, acuminate; lemmas 2.5–3.5 mm long, with callus hairs half to three-quarters as long as the lemma; awn inserted above the middle, a little longer than the lemma, straight. One of the common slough grasses; around sloughs, lakeshores, marshes; throughout the Prairie Provinces.

Calamagrostis lapponica (Wahl.) Hartm.

Plants with creeping rhizomes; culms 30–60 cm high, mostly solitary. Sheaths scabrous; leaves 1–3 mm wide, scabrous toward the tip. Panicle to 10 cm, narrow, contracted to somewhat open, with branches ascending; glumes 4.5–5.5 mm long; lemmas 3.5–4 mm long, with callus hairs as long as the lemma; awn delicate, curved or weakly bent. Boreal forest. Generally considered to be var. *nearctica* Porsild.

Calamagrostis montanensis Scribn. (Fig. 38)

PLAINS REED GRASS

Plants with extensively creeping rhizomes; culms 20–40 cm high, solitary or in small tufts, stiffly erect. Sheaths rather chartaceous; leaves 2–3 mm wide, bluish green, stiff, flat to involute. Panicle 5–10 cm long, erect, interrupted, dense, usually whitish; spikelets 4–5 mm long; glumes acuminate; lemmas 4–4.5 mm long, with callus hairs about half as long as the lemma; awn inserted above the base, 5 mm long, geniculate, exserted. In moist or moderately dry grassland; throughout Prairies and Parklands.

Calamagrostis neglecta (Ehrh.) Gaertn., Mey. & Schreb. NARROW REED GRASS

Plants with long slender rhizomes; culms to 100 cm high, solitary or few in a small tuft. Sheaths glabrous; blades 2–5 mm wide, flat to convolute, usually scabrous on both sides, gray green. Panicle 5–10 cm long, very narrow, usually contracted, greenish to brownish; spikelets 3–3.5 mm; glumes acuminate; lemmas about as long as glumes, with callus hairs about three-quarters as long as the lemma; awn as long as the lemma, straight. Mostly associated with *C. canadensis* and *C. inexpansa* in somewhat acid wet soils; throughout the Prairie Provinces.

Calamagrostis purpurascens R. Br.

PURPLE REED GRASS

Plants tufted; rhizomes, if present, very short; erect culms 40–60 cm high. Sheaths scabrous; blades 2–4 mm wide, flat or somewhat involute, scabrous. Panicle 5–15 cm long, dense, reddish to purplish; spikelets 6–8 mm long; glumes acuminate, scabrous; lemmas as long as the glumes, with callus hairs less than half as long as the lemma; awn exserted near the base, about 1 cm long, geniculate, exserted. Rocky Mountains, Cypress Hills.

Calamagrostis rubescens Buckl.

PINE REED GRASS

Plants with creeping rhizomes; culms closely spaced, often sod-forming or solitary. Sheaths smooth, pubescent at the collar; blades 2–5 mm wide, flat, somewhat lax, scabrous. Panicle 6–15 cm long, often somewhat loose, with ascending branches, mostly contracted at maturity, usually rather dense, often purplish; spikelets 4–5 mm long; glumes narrow, acuminate; lemmas 4.5 mm long; callus hairs less than half as long as the lemma; awn inserted at the base, geniculate, exserted. Open pine woods; Boreal forest, Cypress Hills, southern Rocky Mountains.




Fig. 38. Plains reed grass, *Calamagrostis montanensis* Scribn.

Calamovilfa sand grass

Calamovilfa longifolia (Hook.) Scribn. (Fig. 39)

SAND GRASS

Plants with long, yellowish, thick and tough, scaly rhizomes; culms 50–150 cm high, usually solitary. Sheaths usually smooth, rarely somewhat pubescent; blades to 12 mm wide, flat to involute, firm, dark green. Panicle 15–35 cm long, with branches ascending, and often appressed, contracted at maturity; spikelets 6–7 mm long; first glume 4 mm; lemma 6–6.5 mm, with callus hairs copious, 4–5 mm long. In sandy areas of Prairies and Parklands; throughout the Prairie Provinces.

Catabrosa brook grass

Catabrosa aquatica (L.) Beauv.

BROOK GRASS

Plants with long rhizomes; culms 10–40 cm long, decumbent, rooting at the nodes. Sheaths often somewhat compressed, glabrous, often purplish at base; blades 4–8 mm wide, glabrous, light green, flat, lax. Panicle 10–30 cm long, pyramidal, with branches slender, spreading or sometimes reflexed; spikelets about 3 mm long, usually purplish, with first glume 1 mm long, and second glume 1.5 mm long; lemma 2.5–3 mm long, broad, strongly 3-nerved, with the apex scarious, irregular; palea as long as the lemma. In streams, ponds, swampy areas; throughout the Prairie Provinces in Prairies and Parklands.

Cinna wood grass

Cinna latifolia (Trev.) Griseb.

SLENDER WOOD GRASS

Plants loosely tufted, with culms to 150 cm high. Sheaths smooth; blades 8–15 mm wide, flat, lax. Panicle 15–30 cm long, greenish or yellowish, with branches slender, spreading or drooping; spikelets 3.5–4.0 mm long; glumes 3–3.5 mm long; lemma 3 mm long, 3-nerved, with awn to 1 mm long. Throughout Boreal forests.

Dactylis orchard grass

Dactylis glomerata L. (Fig. 40)

ORCHARD GRASS

Plants forming large tussocks; culms to 100 cm high. Sheaths strongly compressed, keeled; blades 4–10 mm wide, flat or folded, pale green or somewhat glaucous. Panicles 5–20 cm long, with branches solitary or 2 together, secund, ascending, appressed at maturity; spikelets 5–7 mm long, mostly 3- to 4-flowered, clustered at the tips of the branches, with first glume 5–6 mm long, and second glume 6–7 mm long, often ciliate on the keel; lemmas 6–8 mm long, mucronate, ciliate on the keel and margins. Introduced; seeded in orchards and irrigation projects; occasionally escaped.



Fig. 39. Sand grass, *Calamovilfa longifolia* (Hook.) Scribn.



Fig. 40. Orchard grass, *Dactylis glomerata* L.

Danthonia oat grass

- 1. Lemmas glabrous except on the margins. 2
 Lemmas pubescent on whole back. 4
- 2. Panicle usually with a single spikelet;
 culms seldom more than 25 cm high. *D. unispicata*
 Panicle usually with several spikelets;
 culms often more than 25 cm high. 3
- 3. Panicle narrow, with the branches
 appressed; glumes 10–15 mm long;
 blades pilose. *D. intermedia*
 Panicle open, with the branches spread-
 ing; glumes 15–20 mm long; blades
 glabrous. *D. californica*
- 4. Glumes about 10 mm long; lemmas about
 4 mm long; blades often curled. *D. spicata*
 Glumes about 20 mm long; lemmas about
 10 mm long; blades not curled. *D. parryi*

Danthonia californica Boland. CALIFORNIA OAT GRASS

Plants tufted, with culms 30–80 cm high, glabrous, tending to break at the nodes. Sheaths glabrous; blades 1–3 mm wide, glabrous, pilose at the collar. Panicle 5–8 cm long, usually with 2–5 spikelets on slender, flexuous branches; spikelets 20–25 mm long, with glumes 15–20 mm long; lemma 8–10 mm long, with apical teeth long aristate, pilose at base and callus; awn twisted, geniculate, 15–20 mm long. Rare; Prairies, southern Rocky Mountains. Generally considered to be var. *americana* (Scribn.) Hitchc., usually having foliage pilose.

Danthonia intermedia Vasey (Fig. 41) TIMBER OAT GRASS

Plants tufted, with culms to 50 cm high. Sheaths pilose-pubescent; blades to 3 mm wide, flat to involute, short pubescent above, long scattered pubescent below. Panicle 2–5 cm long, with branches appressed ascending, each bearing a single spikelet; spikelets 12–15 mm long, with glumes 10–15 mm long; lemmas 7–8 mm long, with apical teeth acuminate-aristate, appressed pilose along lower margins and callus; awn twisted, geniculate, 10–15 mm long. In fescue grasslands; in Parklands, Boreal zone, Rocky Mountains, Cypress Hills, Wood Mountain.

Danthonia parryi Scribn. PARRY OAT GRASS

Plants forming large tough tussocks, with stout culms 30–60 cm high. Sheaths glabrous; blades to 3 mm wide, flat to involute, pilose at the collar, otherwise glabrous. Panicle 3–7 cm long, with branches appressed-ascending, 1–2 cm long, each bearing a single spikelet; spikelets 20–25 mm long, with glumes about 20 mm long; lemma about 10 mm long, with apical teeth acuminate, pilose on the back and margins; awn twisted and geniculate, 15–20 mm long. Prairies, southern Rocky Mountains.

Danthonia spicata (L.) Beauv. POVERTY OAT GRASS

Plants usually in rather small tufts, with culms 20–50 cm high. Sheaths usually pilose-pubescent; blades 1–3 mm wide, flat to involute, often flexuous,



Fig. 41. Timber oat grass, *Danthonia intermedia* Vasey.

usually pilose-pubescent, especially at the collar. Panicle 2–5 cm long, with stiff branches 2–3 cm long, the lower ones with 2 or 3 spikelets, the upper ones bearing a single spikelet; spikelets to 15 mm long, with glumes 10–12 mm long; lemmas 3–5 mm long, with apical teeth acuminate, sparsely pubescent across the back. Rock outcrops, dry prairie; Parklands, Boreal forest, Rocky Mountains.

Danthonia unispicata (Thurb.) Munro

ONE-SPIKE OAT GRASS

Plants tufted, sod-forming when abundant, with culms 15–25 cm high. Sheaths pilose, or the lower ones glabrous; blades to 3 mm wide, flat, light green, pilose. Panicle a single spikelet, or occasionally 2 or 3 spikelets, 10–15 mm long, with glumes 10–12 mm long; lemmas 3–5 mm long, glabrous, with the callus pubescent. Cypress Hills, southern Rocky Mountains.

Deschampsia hair grass

1. Plants annual; very few, short leaves. *D. danthonioides*
Plants perennial; several to many leaves. 2
2. Panicle spike-like, with the branches
appressed; leaves filiform. *D. elongata*
Panicle open, with the branches spreading;
leaves not filiform. 3
3. Glumes longer than the florets; blades
flat, rather soft. *D. atropurpurea*
Glumes shorter than the florets; blades
rigid, translucent between the ribs. *D. caespitosa*

Deschampsia atropurpurea (Wahl.) Scheele

MOUNTAIN HAIR GRASS

Plants loosely tufted, with culms 40–80 cm high, purplish at base. Sheaths glabrous; blades 4–6 mm wide, flat, thin. Panicle 5–10 cm long, loose, open, with few slender branches; spikelets purplish, with glumes about 5 mm long, the second glume 3-nerved; lemmas 2.5 mm long, with callus hairs 0.8–1.2 mm long; awn of the first floret straight, that of the second one geniculate, exerted. Southern Rocky Mountains.

Deschampsia caespitosa (L.) Beauv. (Fig. 42)

TUFTED HAIR GRASS

Plants densely tufted, with culms 60–120 cm high. Sheaths compressed, keeled, glabrous; blades to 5 mm wide, flat to folded, prominently veined, translucent between the veins. Panicle 10–25 cm long, loose, open, often nodding, with branches slender, bearing spikelets toward the tips, spreading, the lower ones to 10 cm long; spikelets 4–5 mm long, mostly 2-flowered, with the first glume 2 mm long, 1-nerved, and the second glume 3 mm long, 3-nerved; florets distant, with the lemma of the first floret about 2 mm long, and that of the second floret 1.5 mm long; awns about 1 mm long, fragile. On poorly drained, rather fertile soils; throughout the Prairie Provinces.

Deschampsia danthonioides (Trin.) Munro

ANNUAL HAIR GRASS

Plants tufted, with culms 15–60 cm high, slender, erect. Sheaths glabrous; blades 2 mm wide, glabrous, very few. Panicle 7–25 cm long, open, with branches very slender, mostly in twos, ascending; spikelets borne toward the




Fig. 42. Tufted hair grass, *Deschampsia caespitosa* (L.) Beauv.

tips, with glumes 4–8 mm long, 3-nerved; lemmas 2–3 mm long, pilose at base; awn 4–6 mm long, geniculate. Not yet reported, but probably in southern Rocky Mountains.

Deschampsia elongata (Hook.) Munro

SLENDER HAIR GRASS

Plants densely tufted, with culms 30–100 cm high, slender, erect. Sheaths glabrous to somewhat scabrous; blades 1–1.5 mm wide, flat or folded, soft. Panicle to 30 cm long, narrow, with slender branches appressed-ascending; spikelets short appressed, pedicellate, with glumes 4–6 mm long, 3-nerved; lemmas 2–3 mm long; awns about 4 mm long, straight. Meadows and slopes; southern Rocky Mountains.

Digitaria crabgrass

First glume evident, with sheaths pubescent. *D. sanguinalis*

First glume rudimentary or lacking, with sheaths glabrous. *D. ischaemum*

Digitaria ischaemum (Schreb.) Schreb.

SMOOTH CRABGRASS

Plants sod-forming, with culms to 60 cm high, at first erect, but soon decumbent and spreading, and branching. Sheaths glabrous, smooth; blades to 10 mm wide, narrowed at the base. Racemes 4–10 cm long, usually 2–6, reddish purple; spikelets 2–2.5 mm long, with the first glume rudimentary or lacking, and the second glume 1.5 mm long, 3-nerved; sterile lemma about 1.5 mm, 5-nerved; fertile lemma 1.5 mm, pubescent. Rare; in waste areas; Manitoba.

Digitaria sanguinalis (L.) Scop.

CRABGRASS

Plants sod-forming, with culms to 60 cm high, decumbent and spreading, often rooting at the nodes. Sheaths, at least the lower ones, papillose-pilose; blades 5–10 mm wide, pubescent. Racemes to 15 cm long, usually having 3–6 spikelets 3 mm long, with the first glume scale-like, and the second glume 1.5–2 mm long, 3-nerved; sterile lemma to 2 mm long, 7-nerved; fertile lemma to 2 mm long, dark reddish purple. Rare; in waste areas; Manitoba, Saskatchewan, Alberta.

Distichlis alkali grass

Distichlis stricta (Torr.) Rydb. (Fig. 43)

ALKALI GRASS, SALT GRASS

Plants sod-forming, from scaly creeping rhizomes, with tufted culms 10–40 cm high. Sheaths glabrous; blades 2–4 mm wide, flat to involute, yellowish green. Panicles 2–6 cm long, with branches erect to ascending; staminate spikelets 10–15 mm long, 8- to 15-flowered; pistillate spikelets about 10 mm long, 7- or 9-flowered; glumes 3–6 mm long; lemma 2.5–4 mm long, firm. Salt marshes, around saline sloughs, and dry saline areas; throughout Prairies and Parklands.



Fig. 43. Alkali grass, *Distichlis stricta* (Torr.) Rydb.

Dupontia

Dupontia fisheri R. Br.

Plants from creeping rhizomes, with culms 15–60 cm high, erect to decumbent at the base. Sheaths smooth, glabrous; blades 2–4 mm, flat or involute, with pronounced midrib, yellowish green. Panicle 3–10 cm long, narrow, with short branches ascending to appressed; spikelets 4–8 mm long, shiny bronze to purplish, having 3–5 florets, with the first glume 1.5–5 mm long, and the second glume 2–6 mm long, enclosing the spikelet; lemmas 3–5 mm long, pubescent, especially on the midrib toward the base, occasionally at the base only. Rare; saline marshes; along Hudson Bay.

Echinochloa barnyard grass

Echinochloa crusgalli (L.) Beauv. (Fig. 44)

BARNYARD GRASS

Plants tufted, with culms to 100 cm high, erect to decumbent, stout. Sheaths flat, compressed, keeled, glabrous; leaves 6–15 mm wide, flat or V-shaped, keeled below. Panicle 10–20 cm long; racemes erect to spreading; spikelets crowded, ovate, about 3 mm long, green- or reddish-tinged, with the first glume 1 mm long, broadly ovate, acute, 3-nerved, and the second glume 3 mm long, 5-nerved, acuminate; sterile lemma about 3 mm long, with awn 1–5 cm long; glumes and sterile lemma pubescent; fertile lemma 2.5 mm long, glabrous. Considered a weed in gardens, fields, and waste places; Prairies and Parklands. *Echinochloa pungens* (Poir.) Rydb. has been distinguished by the pubescence of glumes and sterile lemma, having hairs with a papillate base. However, *E. crusgalli* in Europe has, usually, as much papillosity as *E. pungens*. The var. *frumentacea* (Roxb.) Wight, with thick racemes and inflated spikelets, has been grown for forage, and is occasionally found. In Europe, plants with erect culms have been differentiated as var. *erecta* Soest, and those with the culms all prostrate and only the panicle ascending, as var. *depressa* Soest. Further differentiation can be made on the basis of length of awn: those with long awns, 2–5 cm, are f. *longiseta* (Doll) J. & W.; those with most of the awns shorter than 1 cm, or reduced to a mere tip, are f. *breviseta* (Doll) J. & W.

(Fig. 44 overleaf)



Fig. 44. Barnyard grass, *Echinochloa crusgalli* (L.) Beauv.

Elymus wild rye

Tufted, usually fairly tall grasses with spike-like awned heads. Closely related to the wheat grasses, but the double floret at each node distinguishing the rye grasses from the wheat grasses, having only one floret at the node of the rachis. Fairly palatable to livestock.

1. Plants with rhizomes. 2
Plants without rhizomes. 6
2. Lemmas awnless. 3
Lemmas awned. 4
3. Glumes lanceolate; rhizomes long-creeping. *E. arenarius*
Glumes subulate; rhizomes, when present, short and stout. *E. cinereus*
4. Awn of lemma 2–4 cm long, strongly divergent at maturity; spikes typically curved. *E. canadensis*
Awn of lemma about 1 cm long, straight; spikes typically erect. 5
5. Lemmas densely purplish pubescent; glumes awn-tipped; awn of lemmas about 5 mm long. *E. innovatus*
Lemmas with hyaline pubescence, awn of glumes about as long as glume; awn of lemmas 5–10 mm long. *E. hirtiflorus*
6. Lemmas awnless. 7
Lemmas awned or awn-tipped. 8
7. Glumes long ciliate; lemmas evenly pubescent across the back; plants to 0.7 m high. *E. junceus*
Glumes not ciliate; lemmas sparsely pubescent along margins, to subglabrous; plants to 1.5 m high. *E. cinereus*
8. Awns 2–4 cm long, strongly divergent at maturity. 9
Awns less than 2 cm long, straight or nearly so. 10
9. Glumes about 1 mm wide, 3- to 5-nerved; spikes usually strongly curved. *E. canadensis*
Glumes about 0.5 mm wide, 1-nerved or nerveless; spikes semierect, flexuous. *E. interruptus*
10. Awns of the lemmas about twice as long as the body of the lemma. *E. glaucus*
Awns of the lemmas usually much shorter than the body of the lemma. 11
11. Glumes indurated at the base and strongly outward bent; lemmas glabrous; blades lax. *E. virginicus*

Glumes not indurated or bent; lemmas pubescent, at least on the margins; blades stiff. *E. angustus*

Elymus angustus Trin. ALTAI WILD RYE

Plants densely tufted, with culms 60–100 cm high. Sheaths glabrous; blades to 15 mm wide, flat, stiff, prominently veined. Spike 10–20 cm long; spikelets 20–25 mm long; glumes 10–18 mm long; lemmas 10–20 mm long. Introduced; forage grass; Saskatchewan.

Elymus arenarius L. SEA LIME GRASS

Plants from creeping rhizomes, with culms to 100 cm high, stout, erect. Sheaths glabrous; blades to 15 mm wide, hard and stiff, involute in drying. Spike to 30 cm long, 2.5 cm wide, stiffly erect; spikelets to 2.5 cm long, usually 2 at the upper and lower nodes, 3 at the middle nodes, mostly 3-flowered; glumes 15–25 mm long, lanceolate, pubescent; lemmas 20–25 mm long, acuminate, short pubescent. Boreal zone, along coast of Hudson Bay and Lake Athabasca. Plants with the culm pubescent below the spike and the glumes and lemmas more densely pubescent are *E. arenarius* ssp. *mollis* (Trin.) Hult.

Elymus canadensis L. (Fig. 45) CANADA WILD RYE

Plants tufted, with short rhizomes at least when young; culms 100–150 cm high. Sheaths glaucous; blades 10–20 mm wide, flat, sometimes convolute, dark green to glaucous. Spike 10–25 cm long, nodding; spikelets 20–30 mm long without awns, mostly 3 or 4 at a node; glumes 15–20 mm long, pubescent, with the awn as long as the body; lemmas 10–15 mm long, pubescent, with the awns 2–4 cm long, divergent. Beaches, riverbanks, sandy areas, and sand dunes; throughout the Prairies and Parklands.

Elymus cinereus Scribn. & Merr. GIANT WILD RYE

Plants densely tufted, forming large tussocks, with very short rhizomes; erect culms 60–120 cm high. Sheaths glabrous; blades 10–15 mm wide, flat to convolute, thick and stiff. Spike 10–25 cm long, thick, dense; spikelets 15–20 mm long, 3–5 at a node; glumes awl-shaped, with the first one 8–10 mm long, and the second one 10–15 mm long; lemmas 10–15 mm long, somewhat pubescent. Rare; riverbanks, slopes, and ravines; southern Saskatchewan, southern Alberta.

Elymus glaucus Buckl. SMOOTH WILD RYE

Plants tufted, with culms 60–120 cm high, erect or ascending. Sheaths glabrous or scabrous; blades 10–15 mm wide, flat, becoming involute on drying. Spike 5–20 cm long, erect to somewhat nodding; spikelets 20–25 mm long; glumes 8–15 mm long, lanceolate, acuminate or awn-tipped; lemmas about 10 mm long, with the awn 10–20 mm long, erect to spreading. Not common; open woods and meadows; Parklands.

Elymus hirtiflorus Hitchc. BLUE WILD RYE

Plants tufted, with slender creeping rhizomes; erect culms 40–90 cm high. Sheaths glabrous; blades 1–4 mm long, flat or involute, firm. Spike 5–15 cm



Fig. 45. Canada wild rye, *Elymus canadensis* L.

long, erect; spikelets about 10 mm long; glumes 8–10 mm long, hirsute, tapering into an awn; lemmas 8–9 mm long, hirsute, with an awn 5–10 mm long. Woods, riverbanks; southern Rocky Mountains.

Elymus innovatus Beal

HAIRY WILD RYE

Plants in small tufts from long creeping rhizomes; erect culms 50–80 cm high. Sheaths glabrous, or the lower ones scabrous; blades 6–12 mm wide, flat to convolute, scabrous on both sides. Spike 5–12 cm long, usually dense, villous; spikelets 10–15 mm long; glumes 10–12 mm long, subulate, purplish or grayish villous; lemmas about 10 mm long, awn-tipped, villous. Common throughout Boreal forest; in woods, clearings, and openings; rare in Parklands and Prairies.

Elymus interruptus Buckl.

VARIABLE-GLUMED WILD RYE

Plants tufted, with erect culms 70–100 cm high. Sheaths glabrous; blades 5–12 mm wide. Spike 8–15 cm long, flexuous or somewhat nodding; spikelets 20–30 mm long; glumes about 10 mm long, awl-shaped; lemmas 10–12 mm long, scabrous or hirsute, with the awn 1–3 cm long, flexuous or divergent. Rare; woods and openings; Boreal forest, Saskatchewan, Riding Mountain.

Elymus junceus Fisch.

RUSSIAN WILD RYE

Plants densely tufted, with erect culms 30–60 cm high. Sheaths glabrous, smooth; blades 3–5 mm wide, flat to convolute, scabrous on both sides, grayish green. Spike 5–12 cm long, erect; spikelets 12–15 mm long, 2 or 3 at a node; glumes 4–5 mm long; lemmas 5–7 mm long. Introduced forage grass; becoming established in coulees and roadsides in various locations in the Prairies, rarely in Parklands.

Elymus virginicus L.

VIRGINIA WILD RYE

Plants in loose, small tufts; with erect culms 60–120 cm high. Sheaths glabrous; blades 5–12 mm wide, flat, scabrous on both sides. Spike 5–15 cm long, usually erect or somewhat flexuous; spikelets 10–15 mm long; glumes 13–17 mm long, 5-nerved, much widened at base; lemmas 10–12 mm long, tapering into an awn, about 1 cm long. In woods and openings, along rivers in Parklands; rare in Prairies.

Eragrostis love grass

1. Plants stoloniferous, rooting at the nodes of prostrate culms. *E. hypnoides*
Plants not stoloniferous, with culms erect or decumbent. 2
2. Palea long ciliate on the keels; spikelets mostly 1.5–2 cm long, 15- to 40-flowered. *E. cilianensis*
Palea scabrous; spikelets shorter, 8- to 15-flowered. *E. poaeoides*

Plants mat-forming, with culms 10–50 cm high, ascending to spreading. Sheaths pilose at the collar; blades 2–7 mm wide, flat; the nerves and margins with depressed glands, causing the unpleasant odor of the plant. Panicles 5–20 cm long, erect; the branches ascending with glands; glumes about 2 mm long, deciduous; lemmas 2–2.5 mm long. Rare in southern Parklands of Manitoba and southeast Saskatchewan; a weed in waste areas and gardens.

Eragrostis hypnoides (Lam.) BSP.

Plants mat-forming, with culms stoloniferous, creeping and flowering culms 10–20 cm high. Sheaths glabrous; blades 1–2 mm wide, scabrous to pubescent above. Panicle 1–6 cm long, few-flowered; spikelets 5–15 mm long, 8- to 12-flowered; glumes 1.5–2 mm long; lemmas 1.5–2.0 mm long. Rare; on sandy shores in southern Parklands of Manitoba, southeast Saskatchewan.

Eragrostis poaeoides Beauv.

Plants mat-forming, with culms 10–50 cm high, ascending to spreading, branching. Sheaths long pubescent toward the collar; blades glabrous. Panicle 5–10 cm long; branches with depressed glands; glumes 1.5–2 mm; lemmas 1.5–2 mm, with the keel glandular. Very rare weedy species; at Saskatoon, Sask.

Festuca fescue

Small- to medium-growing grasses with paniculate heads. Useful forage grasses and very palatable to stock.

- 1. Leaf blades flat, up to 10 mm wide; plants not densely tufted. 2
 - Leaf blades narrow, to 4 mm wide, usually involute; plants mostly densely tufted. 4
- 2. Lemmas tipped with an awn 5–20 mm long; panicle open, drooping. *F. subulata*
 - Lemmas awnless. 3
- 3. Panicle erect; spikelets 8–12 mm long; lemmas thin, with scarious margins. *F. elatior*
 - Panicle diffuse, with the branches spreading to reflexed; lemmas firm. *F. obtusa*
- 4. Plants annual; culms solitary or in small tufts. *F. octoflora*
 - Plants perennial. 5
- 5. Glumes firm, much shorter than the spikelet. 6
 - Glumes thin, dry, not much shorter than the spikelet. 10
- 6. Panicle open, with the branches mostly in pairs, ascending or spreading; lemmas awnless or nearly so. *F. viridula*

- Panicle narrow, with the branches mostly appressed; lemmas awned. 7
7. Plants loosely tufted, often with short matted rhizomes; culms often decumbent, red at the base; underside of leaf glossy. *F. rubra*
Plants densely tufted; roots fibrous. 8
8. Awns as long as, or longer than, the lemmas; blades filiform, soft; panicle secund. *F. occidentalis*
Awns shorter than the lemmas; blades firm, involute, rigid. 9
9. Culms usually more than 40 cm high; panicle 10–20 cm long, with branches somewhat spreading; blades glaucous blue green. *F. idahoensis*
Culms usually less than 40 cm high; panicle to 10 cm long, narrow; blades gray green. *F. ovina*
10. Spikelets with 2 or 3 florets, the third floret sterile; glumes subequal, as long as the first lemma. *F. hallii*
Spikelets with 3–6 florets, mostly all fertile; glumes unequal, shorter than the first lemma. 11
11. Plants grayish or bluish green; shoots coarse, with the culms 40–100 cm high. *F. campestris*
Plants yellowish or dark green; shoots small, with the culms 30–60 cm high. *F. altaica*

Festuca altaica Trin. ex Ledeb.

NORTHERN ROUGH FESCUE

Plants tufted, with fibrous root system; culms 30–60 cm high. Sheaths glabrous to scabrous; blades 1–2.5 mm wide, 7-nerved, somewhat pubescent, flat to involute. Panicle 8–15 cm long, open to somewhat contracted; nodes with 1–3 flexuous branches, the lower ones spreading to ascending; spikelets 10–13 mm long, 3- to 5-flowered, light green often diffused with purple; first glume 4–5 mm long, 1-nerved; second glume 6–7 mm long, faintly 3-nerved; lemmas 6–8 mm long, 5-nerved, scabrous to more or less pubescent; palea equaling the lemma. In the typical form the culm and sheath are glabrous and usually lustrous. The form with scabrous to softly pubescent culms and scabrous sheaths is f. *scabrella* (Torr.) Loom. (*F. scabrella* Torr. ex Hook.). Grassland and open woods; northern Rocky mountains.

Festuca campestris Rydb.

ROUGH FESCUE

Plants tufted, with fibrous root system, often forming large tussocks; shoots coarse, with culms 40–120 cm high, erect, stout. Sheaths and culms glabrous, often lustrous; blades 2–4 mm wide, 7-nerved, rough, usually sparsely short pubescent, gray green or bluish green, flat or involute, often breaking off at the collar. Panicle 10–20 cm long, open to somewhat con-

tracted at anthesis; nodes with 1–3 flexuous branches, the lower ones spreading to reflexed; spikelets 11–16 mm long, 3- to 6-flowered; first glume 3–6 mm long, 1-nerved; second glume 4–8 mm long, 3-nerved; lemmas 7–9 mm long, 5-nerved, glabrous to scabrous or slightly pubescent on the margins, occasionally diffused with purple. One of the important grasses in the prairies in the southern Rocky Mountains and foothills. Syn.: *F. scabrella* Torr. var. *major* Vasey, *F. doreana* Loom.

Festuca elatior L.

MEADOW FESCUE

Plants sod-forming; culms 50–100 cm high, the decumbent culms sometimes rooting at the nodes. Sheaths glabrous, reddish or purplish at the base; blades 5–8 mm wide, prominently veined, dull green above, and slightly keeled, glossy green below. Panicle 10–20 cm long, erect or nodding at the top, often somewhat secund, contracted before and after flowering; lower branches mostly in twos; spikelets linear-cylindric, 8–12 mm long, 6- to 10-flowered, yellow green or violet-tinged; first glume 3 mm long, 1-nerved; second glume 4 mm long, 3-nerved; lemmas 6–7 mm long, obscurely 5-nerved, with the apex scarious. Introduced forage grass, for seeding in moist meadows; Parklands. Tall fescue, *F. elatior* var. *arundinacea* (Schreb.) Wimm., is used for seeding in wet areas.

Festuca hallii (Vasey) Piper

PLAINS ROUGH FESCUE

Plants tufted, with fibrous roots and often more or less well-developed rhizomes; culms 20–60 cm high, glabrous, often lustrous. Sheaths glabrous, lustrous, often diffused with purple; blades 1–1.5 mm wide, sparsely short pubescent, mostly gray green, 5-nerved, always involute. Panicle 6–15 cm long, open to contracted at flowering; the lowest node with 1 or 2 branches, these ascending to contracted; spikelets 7–8 mm long, with 2 or 3 florets, the third floret usually infertile; glumes membranous, often diffused with purple, lustrous; first glume 6–7 mm long, 1-nerved; second glume 6–8 mm long, 1- to 3-nerved; lemmas 7–8 mm long, 5-nerved, scabrous to short pubescent, especially on the margins, often diffused with purple. An important grass in the Parkland prairie, Cypress Hills, and Wood Mountain; sparse on sheltered slopes in Prairies.

Festuca idahoensis Elmer. (Fig. 46)

IDAHO FESCUE

Plants densely tufted, with culms to 100 cm high; whole plant blue green and glaucous. Sheaths flattened, keeled, smooth; blades to 2 mm wide, folded, filiform, erect. Panicle 10–20 cm long; branches ascending to appressed; spikelets 4–7 mm long, 5- to 7-flowered; glumes 2.5 mm and 3 mm long; lemmas almost terete, 6–7 mm, with the awn 2–4 mm long. Fescue prairie, Cypress Hills, southern Rocky Mountains.

Festuca obtusa Biehler

Plants loosely tufted, with culms 50–100 cm high, solitary or few together. Sheaths glabrous; blades 4–7 mm wide, flat, lax, glossy. Panicle 10–20 cm long, loose, somewhat nodding; spikelet 4–8 mm long, 3- to 5-flowered; glumes about 3 and 4 mm long; lemmas about 4 mm long, coriaceous. Open woods; southeast Boreal forest and Parklands.




Fig. 46. Idaho fescue, *Festuca idahoensis* Elmer.

Festuca occidentalis Hook.

Plants densely tufted, with erect culms 40–100 cm high. Sheaths glabrous; blades to 2 mm wide, involute, filiform, soft and smooth. Panicle 7–20 cm long, loose, often drooping at the top; branches solitary or in twos; spikelets 6–10 mm long, 3- to 5-flowered, mostly on slender pedicels; glumes 3 and 4 mm long; lemmas 5–6 mm long, thin, with awn 6–10 mm long, straight. Woods and shrubbery, southern Rocky Mountains.

Festuca octoflora Walt.

SIX-WEEKS FESCUE

Plants tufted, with erect culms 15–30 cm high. Sheaths flattened, keeled, scabrous or minutely pubescent; blades to 2 mm wide, filiform, twisted, strongly veined, dark green. Panicle 4–8 cm long; spikelets 6–8 mm long, 5- to 10-flowered, crowded; first glume 3–3.5 mm long, 1-nerved; second glume 3.5–4.5 mm long, 3-nerved; lemmas 4–5 mm long, with the awn 3–7 mm long. Not common; moist open ground in Prairies.

Festuca ovina L.

SHEEP FESCUE

Plants densely tufted, with culms 10–30 cm high; whole plant gray green. Sheaths glabrous, smooth; blades about 1 mm wide, tightly rolled, filiform. Panicle 5–7 cm long, narrow, contracted before and after flowering; spikelets 4–7 mm long, green, sometimes violet-tinged, 3- to 8-flowered; first glume 1.5 mm long; second glume 2 mm long; lemmas 3–4 mm long, obscurely 5-nerved. In open prairie and forest margins throughout Parklands; in openings and open woods in Boreal forest; rare in moist areas in Prairies. A variable species, which has been divided into several varieties and forms. Plants from the Prairie Provinces are usually distinguished as *F. ovina* var. *saximontana* (Rydb.) G1. (Syn.: *F. saximontana* Rydb.). However, these plants conform quite well to *F. ovina* ssp. *eu-ovina* Hack. var. *vulgaris* Koch, with very narrow filiform leaves, and the lemma with an awn more than 1 mm long.

Festuca rubra L.

RED FESCUE

Plants loosely tufted, with short rhizomes; culms mostly 30–80 cm high, erect to somewhat decumbent at the base. Sheaths finely pubescent, reddish to purplish at the base; blades to 3 mm wide, thick, V-shaped to tightly folded, dark green. Panicle 5–20 cm long, erect or somewhat nodding; branches spreading during and after flowering; spikelets 7–10 mm long, 4- to 6-flowered; first glume 2.5–3 mm long; second glume 3–3.5 mm long; lemmas 5–7 mm long, often somewhat pubescent, awn-tipped or with a short awn, to about 3 mm long. Rocky Mountains, Boreal forest in Alberta. Introduced and cultivated for pasture, hay, and lawns throughout the Prairie Provinces. Very variable; most of our plants belong to ssp. *eu-rubra* Hack. var. *genuina* Hack., the loosely tufted rhizomatous form. Plants with a small panicle and densely pubescent, rather small spikelets are subvar. *arenaria* (Osb.) Hack.; Lake Athabasca, York Factory.

Festuca subulata Trin.

Plants tufted, with erect culms 50–100 cm high. Sheaths glabrous; blades 3–10 mm wide, flat, thin, lax. Panicle 15–40 cm long, open, drooping; branches in twos and threes, spreading to reflexed, the lower ones to 15 cm long; spike-

lets 7–10 mm long, 3- to 5-flowered, very open; first glume 3 mm long; second glume 5 mm long; lemmas about 4 mm long, scabrous toward the apex, tapering into an awn 5–20 mm long. Rare; moist woods; Rocky Mountains.

Festuca viridula Vasey

Plants loosely tufted, with erect culms 50–100 cm high. Sheaths glabrous or scabrous; blades 4–6 mm wide, flat or involute, soft, erect. Panicle 10–15 cm long, open; branches mostly in pairs, ascending to spreading; spikelets 6–12 mm long, 3- to 6-flowered; first glume 4–6 mm long; second glume 5–7 mm long; lemma 6–8 mm long, membranous. Montane meadows; southern Rocky Mountains.

Glyceria manna grass

Mostly medium to tall semiaquatic grasses, found in sloughs, shallow water, and marshes. Quite palatable to livestock.

1. Spikelets linear, almost terete; sheaths strongly flattened. 2
 Spikelets ovate or oblong; sheaths not strongly flattened. 3
2. Spikelets less than 2 cm long, slender pediceled; the upper glume 2–3 mm long; lemmas 3–4 mm long; blades 2–5 mm wide. *G. borealis*
 Spikelets 2–4 cm long, sessile; the upper glume 3–4 mm long; lemmas 6–8 mm long; blades 4–8 mm wide. *G. fluitans*
3. Upper glume 3-nerved; lemmas with 5 prominent nerves; ligules 5 mm or longer. *G. pauciflora*
 Upper glume 1-nerved; lemmas with 7 prominent nerves; ligules less than 5 mm long. 4
4. Upper and lower glumes of about equal length, both 2–2.5 mm long. 5
 Upper glumes about 1 mm long, lower glumes about 0.5 mm long. 6
5. Glumes acute, greenish white; lemmas purple with very narrow scarious margins; blades to 15 mm wide, with culms to 1.5 m high. *G. grandis*
 Glumes obtuse, bronze purple; lemmas purple with broad scarious margins; blades 3–6 mm wide, with culms to 0.6 m high. *G. pulchella*
6. Plants strongly rhizomatous, with culms to 0.8 m high; blades usually less than 5 mm wide; panicle to 2 dm long. *G. striata*

Plants not strongly rhizomatous tufted,
with culms to 1.5 m high; blades 5–10
mm wide; panicle 2–3 dm long. *G. elata*

Glyceria borealis (Nash) Batch.

NORTHERN MANNA GRASS

Plants with creeping rhizomes; culms to 100 cm high, solitary or in tufts. Sheaths compressed, keeled; blades 2–4 mm wide, flat or folded. Panicle 20–40 cm long, with branches to 10 cm long, mostly drooping; spikelets 10–15 mm long, 6- to 12-flowered, often appressed, linear; glumes about 1.5 and 3 mm long; lemmas 3–4 mm long, thin, strongly 7-nerved. Wet areas, slough margins, lakeshores; common in Boreal forest; rare in Parkland and Prairie.

Glyceria elata (Nash) Hitchc.

TALL MANNA GRASS

Plants with creeping rhizomes; culms 100–150 cm high, often in large tufts or tussocks, dark green. Sheaths compressed, keeled; blades 6–12 mm wide, flat, lax. Panicle 15–30 cm long, oblong, with branches spreading, drooping, the lower ones often reflexed; spikelets 4–6 mm long, ovate, 6- to 8-flowered; glumes about 0.6 and 1.3 mm long; lemmas 3 mm long, strongly 7-nerved. Moist woods, riverbanks, wet meadows; southern Rocky Mountains.

Glyceria fluitans (L.) R. Br.

MANNA GRASS

Plants with creeping rhizomes, loosely sod-forming; culms to 100 cm high, lax, ascending. Sheaths compressed, keeled; blades 4–7 mm wide, dull green, flat, with margins and midrib scabrous. Panicle 10–50 cm long, narrow; spikelets 20–25 mm long, loosely 5- to 12-flowered; first glume about 3 mm long; second glume 4 mm long; lemmas 6–7 mm long, strongly 7-nerved, the 5 middle ones almost parallel. Probably found in Manitoba.

Glyceria grandis S. Wats. (Fig. 47)

TALL MANNA GRASS

Plants with creeping rhizomes, loosely sod-forming; culms to 200 cm high, solitary or in small tufts. Sheaths to 1 cm thick, round or compressed above, somewhat scabrous; blades 10–15 mm wide, flat, scabrous on the margins. Panicle 20–40 cm long, usually nodding, rather dense; spikelets 6–8 mm long, 4- to 7-flowered; first glume 1.5–2 mm long; second glume 2–2.5 mm long; lemmas about 2.5 mm long, mostly purplish, strongly 7-nerved. Slough margins, lakeshores, riverbanks; throughout the Prairie Provinces. Often considered a variety or subspecies of *G. maxima* (Hartm.) Holmb., namely var. *americana* (Torr.) Boiv., or ssp. *grandis* (Wats.) Hult.

Glyceria pauciflora Presl

SMALL-FLOWERED MANNA GRASS

Plants with long creeping rhizomes; culms 50–100 cm high, in small tufts or solitary. Sheaths compressed, keeled, smooth; blades 5–15 mm wide, thin, flat, lax. Panicle 10–20 cm long, open to somewhat dense, flexuous; spikelets 4–5 mm long, 5- or 6-flowered, mostly purplish; glumes 1 and 1.5 mm long; lemmas 2–2.5 mm long, prominently 5-nerved. Wet areas; Rocky Mountains.

Glyceria pulchella (Nash) K. Schum.

GRACEFUL MANNA GRASS

Plants with rhizomes; culms 40–60 cm high, loosely tufted. Sheaths compressed, keeled; blades 2–6 mm wide, flat, yellow green. Panicle 10–15 cm



Fig. 47. Tall manna grass, *Glyceria grandis* S. Wats.

long, with branches spreading to drooping; spikelets 5–6 mm long, mostly purplish; glumes 1 and 1.5 mm long; lemmas 2–2.5 mm long, purplish with a broad scarious margin. Uncommon; wet areas, meadows; Boreal forest, Peace River district.

Glyceria striata (Lam.) Hitchc.

FOWL MANNA GRASS

Plants with long creeping rhizomes; culms 30–80 cm high, often in large clumps. Sheaths compressed, keeled; blades 2–5 mm wide, flat or folded, light green. Panicle 10–20 cm long, erect or nodding at the tip; spikelets 3–4 mm long; first glume 0.5–0.8 mm long; second glume 1–1.5 mm long; lemmas 1.5–2.2 mm long, strongly 7-nerved, somewhat scarious at the apex. Common in shallow water; throughout the Prairie Provinces.

Helictotrichon oat grass

Helictotrichon hookeri (Scribn.) Henr.

HOOKER'S OAT GRASS

Plants densely tufted, with erect culms 20–40 cm high. Sheaths compressed, keeled; blades 2–5 mm wide, flat or folded, blue green, glaucous. Panicle 5–10 cm long, narrow, with branches erect or ascending, mostly bearing a single spikelet; spikelets 12–18 mm long, 3- to 6-flowered; glumes 12–15 mm long; lemmas 10–12 mm long, firm, brown; rachilla white villous; awn 10–15 mm long, twisted and geniculate. Fairly common; in moist to moderately dry prairie; in Prairies and Parklands, Rocky Mountains, Cypress Hills, Riding Mountain, Duck Mountain.

Hierochloe sweet grass

Staminate lemmas bearing awns 5–8 mm long. *H. alpina*

Staminate lemmas, awnless. *H. odorata*

Hierochloe alpina (Swartz) R. & S.

HOLY GRASS

Plants with short rhizomes; culms 10–40 cm high, erect, tufted, leafy shoots at base. Sheath glabrous; blades 1–2 mm wide, with the basal ones elongate; culm leaves short. Panicle 3–4 cm long, narrow, with branches short, ascending; spikelets 6–8 mm long, broad; glumes 4–5 mm long, ovate; staminate lemmas about 5 mm long, ciliate on the margins, with awns 5–8 mm long, inserted below the tip; fertile lemma 5 mm long, appressed pubescent. Alpine and Arctic meadows, Boreal forest, Hudson Bay, possibly Rocky Mountains.

Hierochloe odorata (L.) Beauv. (Fig. 48)

SWEET GRASS

Plants with long rhizomes; culms 30–60 cm high, erect, often solitary or with a few leafy shoots. Sheaths smooth; blades 2–6 mm wide, those of the shoots elongate; culm leaves seldom more than 5 cm long. Panicle 10–15 cm long, pyramidal, to 7 cm wide at the base; branches spreading, somewhat flexuous; spikelets about 6 mm long, lustrous golden yellow; glumes 5–6 mm long, ovate; staminate lemmas about 4 mm long, awnless; fertile lemma 4 mm long, brown, appressed pubescent. Wet areas; throughout the Prairie Provinces.



Fig. 48. Sweet grass, *Hierochloa odorata* (L.) Beauv.

Hordeum barley

- 1. Plants perennial; awns slender; auricles missing. *H. jubatum*
- Plants annual; awns stout; auricles present or missing. 3
- 2. Blades with prominent auricles. *H. vulgare*
- Blades without auricles. *H. pusillum*

Hordeum jubatum L. (Fig. 49)

WILD BARLEY

Plants densely tufted; culms 30–60 cm high, erect or decumbent at the base. Sheaths pubescent, glaucous, often purplish; blades 2–6 mm wide, flat, often twisted, usually villose above, densely puberulent below, bluish green. Spike 5–10 cm long, often nodding; lateral spikelets reduced to 1–3 spreading awns; glumes of perfect floret awn-like, spreading; lemma 6–8 mm long, with the awn as long as the glumes. Several varieties are distinguishable by the length of the awns. In var. *jubatum* the awns 2.5–6 cm long, the spike about as wide as long; var. *caespitosum* (Scribn.) Hitchc. having awns 1.5–3 cm long, spikes about twice as long as wide; var. *boreale* (Hitchc.) Boiv. (= *H. brachyantherum* Nevski) having awns less than 1.5 cm long, and the spike several times longer than wide. Varieties *jubatum* and *caespitosum* very common in wetlands, brackish marshes, roadsides; throughout the Prairie Provinces. Variety *boreale* not common; occurring occasionally in saline areas; in Prairies.

Hordeum pusillum Nutt.

LITTLE BARLEY

Plants annual, with tufted culms 10–30 cm high. Sheaths soft pubescent; blades 1–3 mm wide, flat, gray green, pubescent. Spike 2–7 cm long, erect; spikelets 8–15 mm long without the awns; glumes of fertile spikelets 8–10 mm long, much widened above the base, narrowing into an awn 8–15 mm long; lemmas 6–8 mm long, that of the central spikelet awned, those of the lateral spikelets awn-tipped. Open ground, alkali flats; southern Rocky Mountains.

Hordeum vulgare L.

BARLEY

Plants annual, with erect tufted culms 30–100 cm high. Sheaths somewhat pubescent; blades 5–15 mm wide, strongly auricled, flat, usually pubescent. Spike 2–10 cm long without the awns; spikelets 8–15 mm long, sessile; glumes narrow, divergent at base, with a stout awn; lemmas with an awn 10–15 cm long. Cultivated in two main forms: 2-rowed barley (*H. distichon* L.), having sterile lateral spikelets; and 6- or 4-rowed barleys (*H. hexastichon* L.), having all fertile florets. In 4-rowed barley the lateral spikelets overlapping. Both types consisting of several cultivated varieties (cultivars) occurring along roadsides and railways, and in waste areas, but lacking persistence.

Koeleria june grass

Koeleria gracilis Pers. (Fig. 50)

JUNE GRASS

Plants densely tufted, with erect culms 10–50 cm high. Sheaths glabrous to more or less densely pubescent; blades 1–4 mm wide, flat or involute, bluish




Fig. 49. Wild barley, *Hordeum jubatum* L.



Fig. 50. June grass, *Koeleria gracilis* Pers.

green, more or less pubescent on one or both sides or glabrous. Panicle 3–10 cm long, spike-like, often interrupted below, contracted at maturity; spikelets 4–4.5 mm long, 2- or 3-flowered; glumes 3–4 mm long; lemmas 3–4 mm long, lustrous. Occurring in four forms: ssp. *nitida* (Nutt.) Domin is glabrous throughout, or has at most the lower sheaths somewhat pubescent; in ssp. *eugracilis* Domin at least the sheaths are pubescent; in var. *typica* Domin the blades are pubescent with long spreading hairs; in var. *glabra* Domin the blades are short pubescent or glabrate. Very common throughout grasslands and occasionally found in forest openings or open forest, especially on light calcareous soils throughout the Prairie Provinces.

Leersia cut grass

Leersia oryzoides (L.) Sw.

RICE CUT GRASS

Plants with slender rhizomes; culms 70–100 cm high, erect or decumbent at the base. Sheaths retrorse scabrous; blades 6–12 mm wide, with the margins very rough, spinulose. Panicle 10–20 cm long, with branches slender, spreading or ascending, bearing spikelets along the upper half or two-thirds, the lower branches often included in the sheath; spikelets 4–6 mm long, 3–8 forming a spike-like raceme; lemmas about 4 mm long, stiffly ciliate on the keel and nerves; glumes absent. Wet meadows, riverbanks, and lakeshores; southeastern Parklands, Boreal forest.

Lolium rye grass

1. Glumes as long as, or longer than, the spikelets. 2
 Glumes shorter than the spikelets. 3
2. Florets compressed, 9–10 mm long. *L. persicum*
 Florets rounded, 6–8 mm long. *L. temulentum*
3. Lemmas awnless or nearly so. *L. perenne*
 Lemmas, at least the upper ones, awned;
 awns 3–5 mm long. *L. multiflorum*

Lolium multiflorum Lam.

ITALIAN RYE GRASS

Plants annual or short-lived perennial, with culms to 100 cm high, erect or ascending. Sheaths glabrous, or somewhat scabrous; blades 3–5 mm wide, prominently nerved, scabrous above, with auricles usually well-developed. Spike 10–30 cm long; spikelets 1–4 cm long, numerous, mostly 10- to 15-flowered; glumes 8–10 mm long, mostly 7-nerved; lemmas 7–8 mm long, usually awned. Introduced; often in mixtures for lawn grass and occasionally persisting.

Lolium perenne L.

PERENNIAL RYE GRASS

Plants sod-forming; culms 30–60 cm high, erect or geniculate ascending. Sheaths smooth and glabrous, usually compressed, not keeled; blades to 6 mm wide, keeled, prominently veined above, glossy bright green below, with small auricles. Spike to 25 cm long, often somewhat nodding; spikelets 15–20 mm long, acute, awnless. Introduced; seeded in short-term pasture and hayland, and in mixtures for lawn grass.

Plants annual, tufted; culms 30–60 cm high, branching at the lower nodes. Sheaths glabrous, round or slightly compressed; blades 2–6 mm wide, flat to convolute, twisted. Spike 8–12 cm long; spikelets 15–20 mm long, distant to somewhat overlapping; glumes 10–15 mm long; lemmas about 10 mm long, with the awn 5–12 mm long. Introduced; a troublesome weed in grainfields, gardens, and waste areas; has become widespread throughout the Prairies and Parklands.

Lolium temulentum L.

Plants annual, loosely tufted; culms 30–80 cm high, mostly unbranched or with little branching. Sheaths somewhat scabrous; blades 4–8 mm wide, flat, scabrous above. Spike to 25 cm; spikelets 10–25 mm long, numerous, barely imbricate; 6- to 15-flowered; glumes 12–20 mm long; lemmas 6–8 mm long, with the awn 3–5 mm long, or rarely awnless. Introduced; weedy in fields and gardens; Prairies and Parklands. The seed of this species contains a **poisonous** narcotic.

Melica melic grass

- 1. Lemmas awned; plants not bulbous at base. *M. smithii*
- Lemmas awnless; plants mostly bulbous at base. 2
- 2. Glumes narrow; lemmas narrow, long acuminate, pubescent. *M. subulata*
- Glumes broad; lemmas broad, acute or obtuse, glabrous. 3
- 3. Spikelets ascending on stout pedicels; first glume more than half as long as the spikelet. *M. bulbosa*
- Spikelets spreading on slender flexuous pedicels; first glume less than half as long as the spikelet. *M. spectabilis*

Melica bulbosa Geyer

Plants with short rhizomes; culms 30–60 cm high, bulbous at base. Sheaths flat; blades 2–4 mm wide, flat to involute, glabrous to somewhat pubescent. Panicle 10–15 cm long, narrow, with the short branches appressed; spikelets 7–15 mm long, papery; glumes 6–8 mm long; lemmas 5–8 mm long. Open woods and meadows; southern Rocky Mountains.

Melica smithii (Porter) Vasey

Plants with elongated rhizomes; culms 50–100 cm high, not bulbous at the base. Sheaths retrorsely scabrous; blades 6–12 mm wide, flat, lax. Panicle 10–25 cm long, erect to nodding at the tip; branches solitary, distant, spreading to reflexed; spikelets 15–20 mm long, 3- to 6-flowered, often purplish; first glume 3–6 mm long; second glume 4–8 mm long; lemmas 8–10 mm long, with the awn 3–5 mm long, inserted at the bifid apex. Moist woods, meadows; southern Rocky Mountains.



Fig. 51. Persian darnel, *Lolium persicum* Boiss. & Hohen.

Melica spectabilis Scribn.

PURPLE ONION GRASS

Plants with short rhizomes; culms 30–60 cm high, bulbous at base. Sheaths pubescent; blades 2–4 mm wide, flat to involute; panicle 10–15 cm long, with the flexuous branches spreading-ascending; spikelets 10–15 mm long, purplish; first glume 4–6 mm long; second glume 5–7 mm long; lemmas 6–10 mm long, strongly 7-nerved. Open woods and meadows; southern Rocky Mountains.

Melica subulata (Griseb.) Scribn.

ALASKA ONION GRASS

Plants with short rhizomes; culms 60–100 cm high, mostly bulbous at base. Sheaths retrorsely scabrous to long pubescent; blades 2–5 mm wide, flat, thin, scabrous above. Panicle 10–20 cm long, narrow, with the branches appressed-ascending; spikelets 15–20 mm long, narrow; first glume 5–7 mm long; second glume 6–10 mm long; lemmas about 12 mm long, 7-nerved, pubescent. Moist woods; Rocky Mountains.

Milium millet grass

Milium effusum L.

MILLET GRASS

Plants with short, rather stout rhizomes; culms 30–70 cm high, erect from a bent base. Sheaths smooth; blades 7–12 mm wide, flat, lax. Panicles 10–20 cm long, open, pyramidal, with the slender branches spreading, and the lower ones often reflexed; spikelets 3–3.5 mm long; glumes 3–3.5 mm long, rounded, scaberulous; lemmas 2.5–3 mm long. Very rare; moist woods and clearings; Boreal forest.

Muhlenbergia muhly

A very variable genus with panicles usually narrow.

- 1. Panicle open with divergent capillary branches; spikelets long pedicellate. *M. asperifolia*
 Panicle contracted, with the branches appressed or ascending; spikelets short pedicellate or sessile. 2
- 2. Panicle very narrowly linear, usually not more than 2 mm wide; blades 1–2 mm wide. 3
 Panicle not narrowly linear, usually about 5 mm wide; blades 2–8 mm wide. 4
- 3. Plant with rhizomes; glumes ovate, 1–1.5 mm long, less than half as long as the spikelet. *M. richardsonis*
 Plant with fibrous roots and hard, scaly, bulb-like base; glumes acuminate-cuspidate, 2–2.5 mm long, more than half as long as the spikelet. *M. cuspidata*

4. Lemmas awned, the awn to 10 mm long; hairs at base of lemma copious, as long as the lemma; glumes awnless or awn-tipped. *M. andina*
 Lemmas not awned; hairs at base of lemma not conspicuous, usually less than half as long as lemma; glumes awnless, awn-tipped, or awned. 5
5. Glumes awnless or awn-tipped, about as long as the lemma. *M. mexicana*
 Glumes awned, much longer than the lemmas. 6
6. Sheath keeled; ligule 1–1.5 mm long; culms usually branching from the middle nodes; internodes smooth. *M. racemosa*
 Sheath not keeled; ligule minute; culms simple or branching from the base; internodes puberulent. *M. glomerata*

Muhlenbergia andina (Nutt.) Hitchc. (Fig. 52)

FOXTAIL MUHLY

Plants with elongated, wiry, and scaly rhizomes; culms 20–60 cm high, erect, puberulent below the nodes. Sheaths somewhat scabrous; blades 2–6 mm wide, scabrous. Panicles 7–15 cm long, spike-like; spikelets 3–4 mm long; glumes 3–4 mm long, scabrous on the keel; lemmas 2.5–3.5 mm long, tapering into an awn 4–8 mm long, with copious hairs at base. Very rare; wet mud soils; Duck Mountain, Boreal forest.

Muhlenbergia asperifolia (Nees & Mey.) Parodi

SCRATCH GRASS

Plants with thin, scaly rhizomes; culms 10–30 cm high, compressed, branching at the base. Sheaths flattened, keeled; blades to 2 mm wide, flat, very scabrous above, smooth below. Panicles 5–15 cm long, diffuse, with branches very slender, at first erect or ascending, later spreading; spikelets 1.5–2 mm long; glumes 1–2 mm long; lemmas 2 mm long. Not common; damp or marshy calcareous or moderately alkaline soils; Prairies.

Muhlenbergia cuspidata (Torr.) Rydb.

PRAIRIE MUHLY

Plants with hard, bulb-like scaly bases; culms 10–30 cm high, slender, wiry, densely tufted. Sheaths somewhat flattened, glabrous; blades 3 mm wide, flat to folded, prominently veined, hard. Panicles 5–10 cm long, very narrow, with branches short, appressed; spikelets 2–3 mm long; glumes 1–1.5 mm long; lemmas about 2.5–3 mm long. Slopes and crests of moderately to strongly eroded calcareous slopes; Prairies.

Muhlenbergia glomerata (Willd.) Trin.

BOG MUHLY

Plants with long, branching scaly rhizomes; culms 20–50 cm high, occasionally branching at the base. Sheaths scabrous; blades 2–5 mm wide, ascending, flat. Panicles 3–7 cm long, usually interrupted, narrow; spikelets 5–6 mm long; glumes about 2 mm long, with a stiff awn 3–5 mm long; lemmas about 3 mm long, awnless, long pilose at the base. Bogs and swamps; Boreal forest, where var. *cinnoides* (Link) Hermann is found.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Muhlenbergia andina* (Nutt.) Hitchc.

LOCATION: Laurie Lake

COLL: J. Looman

DET:

COLL. NO. 15015

DATE: 10 August 1970

ELEVATION: 720 m.

HABITAT: Pooey bearn




Fig. 52. Foxtail muhly, *Muhlenbergia andina* (Nutt.) Hitchc.

Plants with creeping scaly rhizomes; culms 30–60 cm high, erect or ascending, somewhat branching below. Sheaths scabrous; blades 2–4 mm wide, flat, lax. Panicles 10–15 cm long, with short densely flowered ascending branches; spikelets 2–3 mm long; glumes 1.5–2 mm long, awn-tipped; lemmas about 2 mm long, awn-tipped, long pilose at the base. Not common; margins of woods, moist grassland; eastern Parklands.

Muhlenbergia racemosa (Michx.) BSP.

Plants with creeping scaly rhizomes; stout culms 20–50 cm high. Sheaths flattened, keeled, scabrous; blades 3–6 mm wide, flat to folded, scabrous on both sides. Panicles 5–15 cm long, narrow, often interrupted; spikelets 5–7 mm long; glumes 4–4.5 mm long, awn-tipped; lemmas 2.5–3.5 mm long, occasionally awn-tipped, long pilose at base. Not common; meadows, margins of woods, coulees and ravines; Prairies and Parklands.

Muhlenbergia richardsonis (Trin.) Rydb. (Fig. 53)

Perennial with numerous thin, hard, scaly rhizomes; culms 5–40 cm high, densely tufted, wiry, erect or decumbent. Sheaths round, smooth; blades 1–2 mm wide, flat or involute, scabrous above, smooth below. Panicles 3–10 cm long, very narrow; spikelets 2–3 mm long; first glume 1 mm long; second glume 1.5 mm long; lemmas 2.5–3 mm long. Common; in moist, often alkaline, grasslands; Prairies, Parklands, parts of Boreal forest, Peace River district.

Munroa false buffalo grass

Munroa squarrosa (Nutt.) Torr.

Plants annual, forming mats to 50 cm across; culms prostrate, with internodes to 10 cm long. Sheaths round, pilose or ciliate at the throat, inflated; blades to 3 mm wide, stiff, in bundles at nodes and tips of branches. Spikelets 8–12 mm long, 2- to 4-flowered, in groups of two or three at the tips of culms; first glume 4–6 mm long; second glume 6–8 mm long; both glumes 1-nerved; lemmas 3–5 mm long, 3-nerved, with a conspicuous tuft of hairs halfway along the margin. Very rare; in dry grassland; Prairies.

Oryzopsis rice grass

Long-leaved, tufted grasses of medium height with rather large, rice-like seeds, found in various locations.

- 1. Panicles diffuse, with regularly dichotomous branches; glumes with a long firm tip, much exceeding the long silky lemma. *O. hymenoides*

Panicles not diffuse, with branches erect or somewhat spreading; glumes not sharp-pointed; lemmas short pubescent or glabrous. 2



Fig. 53. Mat muhly, *Muhlenbergia richardsonis* (Trin.) Rydb.

2. Lemmas glabrous, with awns 5–10 mm long; blades flat or involute, 0.5–2 mm wide. *O. micrantha*
 Lemmas pubescent. 3
3. Blades flat, evergreen, 4–10 mm wide; spikelets 6–8 mm long not including the awns; awns 5–10 mm long. *O. asperifolia*
 Blades mostly involute or filiform; spikelets 3–5 mm long. 4
4. Blades flat to involute; panicle open, lax, with flexuous, ascending or spreading branches. *O. canadensis*
 Blades filiform; panicle narrow, with ascending or appressed branches. 5
5. Panicle branches erect or appressed; awns 4–6 mm long, geniculate; glumes acute. *O. exigua*
 Panicle branches loosely ascending; awns 0.5–2 mm long, straight; glumes obtuse. *O. pungens*

Oryzopsis asperifolia Michx.

WHITE-GRAINED MOUNTAIN RICE GRASS

Plants tufted, with culms 20–70 cm high, at first erect, later spreading to prostrate. Sheaths smooth or somewhat scabrous, dark purple at base; blades of two types: some 1–5 cm long, others 20–40 cm long, all 3–10 mm wide, flat to convolute, dark green. Panicles 5–10 cm long, few-flowered, narrow, with branches ascending-appressed; spikelets 6–9 mm long; glumes 6–8 mm long, mostly 5- to 7-nerved; lemmas 7–9 mm long, sparsely pubescent on the back, densely so at the base, with the awn 5–10 mm long. Rather common in woods, Boreal forest; less common in Parklands.

Oryzopsis canadensis (Poir.) Torr.

CANADIAN RICE GRASS

Plants tufted, with erect culms 30–60 cm high. Sheaths scabrous; blades 2–4 mm wide, flat to involute, scabrous. Panicles 5–10 cm long, open, with branches flexuous, ascending to spreading; spikelets about 5 mm long, on long slender pedicels; glumes 4–5 mm long; lemmas about 3 mm long, appressed-pubescent, with the awn 1–2 cm long, weakly twice geniculate. Not common; in woods; Boreal forest, Parklands.

Oryzopsis exigua Thurb.

LITTLE RICE GRASS

Plants densely tufted, with culms 15–30 cm high, stiffly erect. Sheaths smooth or somewhat scabrous; blades 2–4 mm wide, filiform to involute, stiffly erect. Panicles 3–6 cm long, narrow, with branches ascending-appressed; spikelets about 4 mm long, short-pedicel; glumes 3–4 mm long; lemmas 3–4 mm long, appressed pubescent, with the awn 4–8 mm long, geniculate. Not common; woods and clearings; southern Rocky Mountains.

Oryzopsis hymenoides (Roem. & Schult.) Ricker (Fig. 54)

INDIAN RICE GRASS

Plants densely tufted, with culms 30–60 cm high. Sheaths smooth or somewhat scabrous, prominently veined; blades 2–5 mm wide, to 50 cm long,



Fig. 54. Indian rice grass, *Oryzopsis hymenoides* (Roem. & Schult.) Ricker.

mostly involute, coarsely veined. Panicles 10–20 cm long, diffuse, with branches slender, in pairs, branching forked, and flexuous pedicels; spikelets 6–7 mm long, solitary; glumes 6–7 mm long, papery; lemmas about 3 mm long, almost black at maturity, densely pilose with white hairs as long as the lemma; the awn about 4 mm long, straight. On sandy soils and slopes, sand dunes; Prairies and Parklands. An important grass in the sand hills, very resistant to wind action, and a good sand-binder. Palatable to livestock, and fairly resistant to grazing.

Oryzopsis micrantha (Trin. & Rupr.) Thurb. LITTLE-SEED RICE GRASS

Plants rather densely tufted, with culms 30–70 cm high, erect, slender. Sheaths glabrous; blades 0.5–2 mm wide, flat or involute, scabrous. Panicles 10–15 cm long, open, with branches spreading to reflexed, single or in pairs; spikelets about 4 mm long; glumes 3–4 mm long, thin; lemmas 2–2.5 mm long, glabrous or appressed pubescent, with the awn 5–10 mm long, straight. Rare; in shrubbery on sandy soils; Prairies, eastern Parklands.

Oryzopsis pungens (Torr.) Hitchc. NORTHERN RICE GRASS

Plants densely tufted, with culms 20–40 cm high, erect, slender. Sheaths smooth or somewhat scabrous; blades 1–2 mm wide, flat or involute, strongly nerved, erect. Panicles 3–6 cm long, narrow, with branches ascending-appressed or spreading; spikelets 3–4 mm long, few; glumes 3–4 mm long, often bronze-colored; lemmas 3–4 mm long, densely pubescent, with the awn 1–3 mm long. Rather common; open woods, clearings, mostly on light soils; Boreal forest.

Panicum millet

Annual or perennial grasses of various habits and habitats, with glumes unequal, the first often being very minute. Mostly found in the moister, eastern part of the Prairie Provinces.

- 1. Plants annual. 2
 Plants perennial. 3
- 2. Panicles erect, diffuse; spikelets 2–4 mm long. *P. capillare*
 Panicles arching, not diffuse; spikelets 4.5–5.5 mm long. *P. miliaceum*
- 3. Plants with hard, scaly creeping rhizomes. *P. virgatum*
 Plants without rhizomes, often with specialized form in autumn. 4
- 4. Spikelets less than 2 mm long. *P. lanuginosum*
 Spikelets more than 2 mm long. 5
- 5. Spikelets 3.5–4 mm long, soft villous; sheaths papillose-hispid. *P. leibergii*
 Spikelets glabrous or pilose, not long villous. 6
- 6. Spikelets of autumn plants often hidden in lower sheaths. 7

- Spikelets of autumn plants on branches of the culms. 9
7. Spikelets 3–4.5 mm long, beaked, exceeding the fruit. *P. depauperatum*
- Spikelets 2–3.5 mm long, beakless, as long as the fruit. 8
8. Spikelets 2.5–3.5 mm long, sheaths densely pilose. *P. perlongum*
- Spikelets 2.2–2.7 mm long, sheaths glabrous or nearly so. *P. linearifolium*
9. Spikelets 2.5–3 mm long, pilose; culms of autumn plants branching at base, forming bushy tufts. *P. wilcoxianum*
- Spikelets 3–4 mm long, glabrous or minutely pubescent; autumn plants branching from the nodes, not forming bushy tufts. 10
10. Spikelets 3–3.4 mm long; panicle branches spreading; culms of autumn plants branching at upper nodes. *P. oligosanthos*
- Spikelets 3.5–4 mm long; panicle branches stiffly erect; culms of autumn plants branching at lower nodes. *P. xanthophysum*

Panicum capillare L.

WITCH GRASS

Plants annual, tufted, with culms 20–80 cm high, erect or spreading, papillose-hispid. Sheaths dull green, conspicuously papillose-hispid; leaves 5–15 mm wide, hispid on both sides, papillose-ciliate at the base. Panicles to more than half the height of plants; the plant very diffuse, densely flowered; spikelets 2–2.5 mm long; the entire panicle breaking off at maturity. Not common; waste places, sandy prairie. The var. *occidentale* Rydb. having the panicle to two-thirds the height of the plants, long exserted; rare in west; Parklands.

Panicum depauperatum Muhl.

PANIC GRASS

Plants tufted; the summer plants with several to many erect culms 20–30 cm high. Sheaths glabrous or papillose-pilose; blades 2–5 mm wide, flat to involute when drying. Panicles 5–10 cm long, exserted; spikelets 3–4 mm long, pointed, glabrous or sparsely pubescent. Autumn plants similar, but with panicles reduced, partly hidden in the basal leaves. Rare; sandy areas, open woods; southeastern Boreal forest, Parklands.

Panicum lanuginosum Ell.

SOFT MILLET

Summer plants in large clumps; culms 40–70 cm high, lax, spreading, villos throughout, except under the nodes. Sheaths densely velvety pubescent; blades 5–10 mm wide, flat, pubescent on both sides. Panicles 5–15 cm long; spikelets about 1–2 mm long, pubescent. Autumn plants with culms decumbent or spreading, repeatedly branching from the middle nodes, with branches again repeatedly branching and forming leafy inflorescences. Rare; open pine woods, sandy areas; southeastern Boreal forests, Parklands.

Panicum leibergii (Vasey) Scribn.

Summer plants tufted; culms 30–70 cm high, erect or geniculate at the base, pilose to scabrous. Sheaths papillose-hispid, with hairs spreading; blades 6–15 mm wide, erect or ascending, thin, papillose-hispid on both sides. Panicles 5–15 cm long; spikelets 3.5–4 mm, strongly papillose-hispid. Autumn plants spreading, branching from the middle and lower nodes. Rare; dry prairie and clearings; southeastern Boreal forest, Parklands.

Panicum linearifolium Scribn.

Summer plants densely tufted; culms 20–40 cm high, slender, erect. Sheaths papillose-pilose; blades 2–4 mm wide, erect, usually exceeding the panicles. Panicles 5–10 cm long, with branches flexuous, ascending; spikelets 2.2–2.7 mm long, sparsely pilose. Autumn plants with reduced panicles hidden among the basal leaves. Rare; rock outcrops in southeastern Boreal forest.

Panicum miliaceum L.

BROOMCORN MILLET

Plants annual, tufted; culms 20–80 cm high, stout, erect or decumbent at the base. Sheaths pilose; blades 10–20 mm wide, to 30 cm long, pilose to glabrate on both sides. Panicles 10–30 cm long, included at the base, nodding, branches ascending, scabrous; spikelets 4.5–5.5 mm long; fruit 3 mm long, straw-colored to reddish brown. Introduced; escaped in various places.

Panicum oligosanthos Schult. var. *scribnerianum* (Nash) Fern.

Summer plants tufted; culms 40–80 cm high, appressed-pubescent. Sheaths papillose-pubescent; blades 5–8 mm wide, to 15 cm long, glabrous or nearly so above, coarsely puberulent below. Panicles 5–15 cm long; spikelets 3–3.5 mm long, sparsely pubescent. Autumn plants freely branching from the upper nodes, erect to spreading. Rare; sandy prairie, southeastern Parklands.

Panicum perlongum Nash

LONG-STALKED PANIC GRASS

Summer plants in small tufts, with the whole plant pilose; culms 20–40 cm high. Sheaths pilose; blades 2–4 mm wide, pubescent on both sides. Panicles 5–10 cm long, with branches appressed-ascending; spikelets 2.5–3.5 mm long, sparingly pubescent. Autumn plants with numerous reduced panicles. Rare; open woods on sandy soil; southeastern Parklands.

Panicum virgatum L. (Fig. 55)

SWITCH GRASS

Plants with long, scaly rhizomes; culms 80–150 cm high, tufted, erect, tough. Sheaths round, glabrous, white to purplish-tinged below; blades 6–12 mm wide, distinctly veined, long pubescent above at base, otherwise glabrous. Panicles 15–50 cm long, open to diffuse; spikelets 3.5–5 mm long; fruit narrowly ovate. Very rare; prairies, open woods; southeastern Parklands.

Panicum wilcoxianum Vasey

SAND MILLET

Summer plants tufted, papillose-hirsute throughout; culms 10–25 cm high; blades 3–6 mm wide, to 10 cm long, involute-acuminate. Panicles 2–5 cm long; spikelets 2.5–3 mm long, papillose-pubescent. Autumn plants branching from the nodes, forming bushy tufts with rigid, erect blades. Rare; sand hill prairie, clearings; Parklands.



Fig. 55. Switch grass, *Panicum virgatum* L.

Panicum xanthophysum A. Gray

Summer plants tufted, yellowish green; culms 20–60 cm high, scabrous. Sheaths somewhat papillose-pilose; blades 10–20 mm wide, to 15 cm long, erect, with base ciliate, otherwise glabrous. Panicles 5–10 cm long, very narrow, with branches stiffly erect, few-flowered; spikelets 3.5–4 mm long, pubescent. Autumn plants branching from the lower nodes, erect or ascending. Open pine woods; southeastern Boreal forest.

Phalaris canary grass

Perennial with rhizomes; panicles narrow; outer glumes not winged. *P. arundinacea*

Annual; panicles spike-like, ovate; outer glumes winged. *P. canariensis*

Phalaris arundinacea L. (Fig. 56) REED CANARY GRASS

Plants with thick rhizomes, to 4 mm in diam, scaly, dark brown; culms to 2 m high, stiffly erect, smooth, with up to 10 leaves. Sheaths glabrous, distinctly veined; blades to 20 mm wide, to 20 cm long, flat, somewhat scabrous. Panicles 10–20 cm long, with the lower branches 3–5 cm long, spreading during flowering, later appressed; glumes 3–5 mm long, narrowly winged; lemmas 3–4 mm long, appressed-pubescent; sterile lemmas 1 mm long, villous. Native; in wet areas, marshes, riverbanks, and lakeshores; Boreal forest. Introduced for forage in irrigated haylands.

Phalaris canariensis L. CANARY GRASS

A medium-growing annual with rather pretty ovoid head, 20–35 mm long, pale with green stripe on glumes. Found where seed for a caged bird has been scattered.

Phleum timothy

Panicles elongate-cylindric; awns less than half the length of empty glumes. *P. pratense*

Panicles short, ovoid or oblong; awns about half the length of empty glumes. *P. alpinum*

Phleum alpinum L. (Fig. 57) ALPINE TIMOTHY

Plants with short rhizomes; culms 20–50 cm high, densely tufted, decumbent at the base. Sheath round, glabrous, inflated near the middle; blades 4–8 mm wide, 2–15 cm long, scabrous above and on the margins. Panicles 1–4 cm long, to 1 cm wide, bristly; glumes about 5 mm long, ciliate on the keel, with the awn 2–3 mm long. Mountain meadows, bogs, and wet areas; Rocky Mountains, Cypress Hills; occasionally in Boreal forest.

Phleum pratense L. (Fig. 58) TIMOTHY

Plants with a swollen, bulb-like base; culms 50–80 cm high, often forming large clumps. Sheaths glabrous, green, often purplish at base; blades 6–12 mm wide, to 30 cm long, flat, often twisted, light green or grayish green. Panicles, usually 5–10, occasionally to 20 cm long, to 1 cm thick; glumes 3–5 mm long,



Fig. 56. Reed canary grass, *Phalaris arundinacea* L.



Fig. 57. Alpine timothy, *Phleum alpinum* L.



Fig. 58. Timothy, *Phleum pratense* L.

with a green midrib, ciliate on the keel, with the awn 1–2 mm long, stout. Introduced forage grass; escaped and established in many parts of Parklands, Boreal forest, and in moist areas in Rocky Mountains.

Phragmites giant reed grass

Phragmites communis Trin.

COMMON REED GRASS

Plants with long, extensively creeping rhizomes, to 3 cm thick; culms 1–4 m high, erect, occasionally decumbent and stoloniferous. Sheaths glabrous, purplish at the base; blades 20–40 mm wide, to 40 cm long, acuminate, glabrous. Panicles 15–40 cm long, with branches spreading in flower, often drooping, densely flowered; spikelets 6–15 mm long; first glume 5–7 mm long; second glume 10–12 mm long; lemmas about 12 mm long; hairs of the rachilla exceeding the spikelets. Wet places in Parklands, Boreal forest; rarely in springy places in the Prairies.

Poa blue grass

A very large and difficult genus, mainly of low- to medium-growing species and found in all types of habitat. The leaves have boat-shaped tips and usually have a double line down the midrib.

Valuable forage grasses. Many species grow very early in the season.

1. Plants with rhizomes. 2
 Plants with fibrous roots. 10
2. Culms strongly compressed, flat; panicle narrow, with the branchlets bearing spikelets to near the base. *P. compressa*
 Culms terete or somewhat flattened. 3
3. Lemmas pubescent, at least on the nerves. 5
 Lemmas not pubescent. 4
4. Panicle open, with branches spreading; lemmas keeled; rhizomes long. *P. nervosa*
 Panicle narrow, dense; lemmas rounded on the back; rhizomes short. *P. ampla*
5. Lemmas less than 4 mm long. 6
 Lemmas more than 4 mm long. 7
6. Lemmas cobwebby at base; keel and marginal nerves sparsely pubescent toward the base. *P. pratensis*
 Lemmas not cobwebby at base; densely pubescent on lower keel and marginal nerves. *P. arida*
7. Lemmas cobwebby at base. *P. arctica*
 Lemmas not cobwebby at base. 8
8. Panicle narrow, with branches ascending; spikelets 7–10 mm long; leaves folded or involute, stiff. *P. fendleriana*

- Panicle open, with branches spreading;
leaves flat or folded, not stiff. 9
9. Panicle usually small, less than 10 cm
long; lemmas purplish, suffused with
orange at the tip. *P. arctica*
Panicle usually large, 15–20 cm long; lem-
mas not purplish; foliage blue green. *P. glaucifolia*
10. Plants annual; blades soft and flat, often
rugose; ligule white. *P. annua*
Plants perennial. 11
11. Plants densely tufted; blades narrow. 12
Plants not densely tufted; blades flat. 20
12. Lemmas cobwebby at base. 13
Lemmas not cobwebby at base. 14
13. Spikelets less than 4.5 mm long; lemmas
less than 3 mm long; blades 2–3 mm
wide, flat. *P. nemoralis*
Spikelets 5–6 mm long; lemmas about 4
mm long; blades 1 mm wide, folded,
lax. *P. pattersonii*
14. Lemmas not pubescent. 15
Lemmas crisp pubescent, silky or villose,
at least at base. 16
15. Blades of culm leaves filiform; spikelets
7–9 mm long. *P. cusickii*
Blades of culm leaves flat; spikelets 5–6
mm long. *P. epilis*
16. Lemmas rounded on the back; spikelets
not flattened. 17
Lemmas keeled; spikelets flattened. 18
17. Blades short, often curled; culms usually
less than 30 cm high; panicles less than
10 cm long. *P. secunda*
Blades 10 cm or longer, not curled; culms
30–60 cm high; panicles to 15 cm long. *P. canbyi*
18. Plants usually blue green; spikelets 2- and
3-flowered; panicle rather lax. *P. glauca*
Plants not blue green; spikelets 4- and 5-
flowered; panicle stiff, erect. 19
19. Blades 5–10 cm long; culms 5–20 cm
high; blades about 1 mm wide, lax. *P. pattersonii*
Blades 1–5 cm long; culms 10–20 cm
high; blades 1–1.5 mm wide, stiff. *P. rupicola*
20. Lemmas cobwebby at base. 21
Lemmas not cobwebby at base. 23
21. Spikelets 5–6 mm long; panicle usually
5–10 cm long; culms solitary or few
together. *P. leptocoma*

- Spikelets 3–4 mm long; panicle usually 10–30 cm long; culms loosely tufted. 22
22. Culms erect from a decumbent base; ligule to 3 mm long; blades long, grayish green beneath. *P. palustris*
- Culms decumbent to spreading, often rooting at nodes; ligule to 7 mm long; blades glossy green below. *P. trivialis*
23. Panicle narrow, with branches ascending. 24
- Panicle with spreading or drooping branches. 25
24. Blades involute, rather stiff; lemmas 3.5–4 mm long. *P. juncifolia*
- Blades flat or folded, 1–3 mm wide; lemmas 4–6 mm long. *P. ampla*
25. Blades flat and short, 3–6 cm long, 2–4 mm wide; spikelets purplish or bronze. *P. alpina*
- Blades folded or involute; spikelets greenish. 26
26. Basal leaves filiform; panicles with lower branches spreading or reflexed; spikelets rounded. *P. gracillima*
- Basal leaves flat; panicles with drooping branches; spikelets flattened. *P. stenantha*

Poa alpina L.

ALPINE BLUE GRASS

Plants tufted, sod-forming, with erect culms 10–30 cm high. Sheaths glabrous, keeled; blades 2–5 mm wide, flat, short. Panicles 1–8 cm long, compact short pyramidal or ovoid, with branches spreading, or the lower ones reflexed; spikelets 4–6 mm long, purplish; glumes 1.5–2 mm long; lemmas 3–4 mm long, pubescent to villous on keel and marginal nerves. Rocky Mountains, Boreal forest.

Poa ampla Merr.

BIG BLUE GRASS

Plants densely tufted, with short rhizomes; erect culms 60–80 cm high. Sheaths smooth or somewhat scabrous; blades 1–3 mm wide, green or glaucous. Panicles 10–15 cm long, narrow, mostly dense; spikelets 8–10 mm long, 4- to 7-flowered; glumes 2.5–5 mm long; lemmas 4–6 mm long. Meadows and slopes; Rocky Mountains.

Poa annua L.

ANNUAL BLUE GRASS

Plants annual or biennial, sod-forming; culms 5–20 cm high, ascending or spreading, sometimes rooting at the internodes. Sheaths somewhat compressed, glabrous; blades 1–4 mm wide, flat or somewhat folded, thin, light green. Panicles 3–10 cm long, pyramidal, often secund, with branches spreading, the lower ones often reflexed; spikelets about 3 mm long, 4- and 5-flowered, green- or purple-tinged; first glume 1.5–2 mm long, 1-nerved; second glume 2–2.5 mm long, 3-nerved; glumes 2.5–3 mm long, 5-nerved, subglabrous

to short pubescent. Introduced; weedy in gardens in the Prairies; increasingly common in Parklands and Boreal forest.

Poa arctica R. Br.

ARCTIC BLUE GRASS

Plants with creeping rhizomes; culms 10–30 cm high, erect, with the base decumbent, tufted. Sheaths glabrous; blades 2–3 mm wide, flat to folded. Panicles 5–10 cm long, open, pyramidal, with the lower branches spreading to reflexed; spikelets 5–8 mm long, 3- and 4-flowered; first glume 1.5–2 mm long; second glume 2–2.5 mm long; lemmas 3.5–4 mm long, densely villous-pubescent on the keel and marginal nerves. Alpine and high boreal meadows; Rocky Mountains, Boreal forests.

Poa arida Vasey

PLAINS BLUE GRASS

Plants with creeping rhizomes; culms 20–50 cm high, erect, solitary or few together. Sheaths glabrous; blades 2–3 mm wide, folded, stiff. Panicles 2–10 cm long, narrow, with branches appressed-ascending; spikelets 5–7 mm long, 4- to 8-flowered; first glume 2–4 mm long; second glume 2.5–4.5 mm long; lemmas 3–4 mm long, densely pubescent on the back. Dry to moist, often somewhat alkali prairies; Prairies, Parklands.

Poa canbyi (Scribn.) Piper (Fig. 59)

CANBY BLUE GRASS

Plants densely tufted, with erect culms 30–80 cm high. Sheaths somewhat compressed, scabrous; blades to 4 mm wide, flat to folded, green and glaucous. Panicles 10–15 cm long, narrow, compact or somewhat loose, with branches short, appressed or ascending; spikelets 5–6 mm long, 3- to 5-flowered; first glume 2.5–3.5 mm long; second glume 3–4.5 mm long; lemmas 3–4.5 mm long, obscurely 5-nerved, more or less pubescent on the nerves, at least toward the base. Moist, often alkali meadows; Prairies, Parklands.

Poa compressa L.

CANADA BLUE GRASS

Plants with creeping rhizomes; culms 15–50 cm, solitary or few together, flat, wiry. Sheaths strongly compressed and sharply keeled, glabrous; blades 2–5 mm wide, flat to folded, bluish green. Panicles 3–10 cm long, with branches usually short, in pairs; spikelets 4–6 mm long, 3- to 6-flowered; first glume 1.5–2.5 mm long; second glume 2–3 mm long; lemmas 2–3 mm long, somewhat pubescent. Introduced; dry, poor, often stony soils; throughout the Prairie Provinces.

Poa cusickii Vasey

EARLY BLUE GRASS

Plants densely tufted, with erect culms 20–40 cm high. Sheaths compressed, sharply keeled, scabrous; blades 1–3 mm wide, flat to folded and bristle-like, erect. Panicles 3–8 cm long, dense, usually obovoid, pale; spikelets 7–9 mm long; glumes 3–4 mm long; lemmas 4–6 mm long, smooth or somewhat scabrous. Dry to moist prairie; throughout the Prairies and Parklands.

Poa epilis Scribn.

SKYLINE BLUE GRASS

Plants densely tufted, with erect culms 20–40 cm high. Sheaths compressed; blades 2–3 mm wide, folded or involute. Panicles 2–6 cm long, dense, oblong, usually purplish; spikelets about 5 mm long, 3-flowered; glumes about 5 mm long; lemmas 6 mm long, glabrous. Mountain meadows; Rocky Mountains.




Fig. 59. Canby blue grass, *Poa canbyi* (Scribn.) Piper.

Poa fendleriana (Steud.) Vasey

MUTTON GRASS

Plants tufted, with culms 30–50 cm high, erect, scabrous below the panicle. Sheaths scabrous; blades 1–2 mm wide, folded or involute, stiff, erect. Panicles 2–7 cm long, contracted; spikelets 6–8 mm long, 4- to 6-flowered; glumes about 3 and 4 mm long; lemmas 4–5 mm long, long pilose on lower keel and marginal nerves. Dry slopes, prairie; Prairies.

Poa glauca Vahl

Plants tufted, sometimes densely so, with erect culms 20–50 cm high. Sheaths glabrous; blades 1–2 mm wide, flat, blue green, glaucous. Panicles 3–10 cm long, narrow, often compact, with branches ascending, spreading in flower; spikelets 5–6 mm long, 2- or 3-flowered; glumes 2.5–4 mm long, subequal; lemmas 2.5–4 mm long, densely pubescent on lower half of keel and marginal nerves. Stony areas, sandy soils; Boreal forest, Rocky Mountains.

Poa glaucifolia Scribn. & Will.

GLAUCOUS BLUE GRASS

Plants with slender rhizomes; culms 30–60 cm high, loosely tufted, erect. Sheaths somewhat compressed, glabrous; blades 2–4 mm wide, flat to folded, glabrous, glaucous on both sides. Panicles 10–20 cm long, open, with branches ascending to spreading; spikelets 5–7 mm long, 2- to 4-flowered; glumes about 3.5 and 4.5 mm long; lemmas about 4 mm long, villous on lower half of keel and marginal nerves. Not common; moist, often somewhat alkali areas and meadows; Prairies, Parklands.

Poa gracillima Vasey

PACIFIC BLUE GRASS

Plants loosely tufted, with culms 30–60 cm high, mostly decumbent at the base. Sheaths somewhat scabrous; blades 0.5–1.5 mm wide, flat or folded, with the basal ones filiform. Panicles 3–10 cm long, pyramidal, with branches spreading, the lower ones often reflexed; spikelets 4–6 mm long; first glume 2.5–3 mm long; second glume 3–4 mm long; lemmas about 4 mm long, pubescent on the lower part of back. Slopes, riverbanks, lakeshores; Rocky Mountains.

Poa juncifolia Scribn.

ALKALI BLUE GRASS

Plants tufted, with erect culms 30–60 cm high. Sheaths somewhat flattened; blades 1–3 mm wide, involute, rather stiff. Panicles 10–20 cm long, narrow, with branches appressed; spikelets 7–10 mm long, 3- to 6-flowered; glumes about 4 mm long; lemmas about 4 mm long, glabrous or nearly so. Alkaline meadows; Prairies.

Poa leptocoma Trin.

BOG BLUE GRASS

Plants loosely tufted, with culms 20–50 cm high, often decumbent at the base. Sheaths slightly scabrous; blades 2–4 mm wide, short, flat, lax. Panicles 7–15 cm long, open, nodding at the tip, with branches very slender, ascending or spreading; spikelets 4–6 mm long, 2- to 4-flowered; glumes about 2.5 and 3 mm long; lemmas 3.5–4.5 mm long, pubescent on the back. Not common; bogs and wet meadows; Rocky Mountains.

Poa nemoralis L.

WOOD BLUE GRASS

Plants sod-forming, with culms 30–70 cm high, loosely tufted, erect, often decumbent at the base. Sheaths glabrous, smooth; blades to 2 mm wide, flat, lax. Panicles 5–10 cm long, loose, often nodding, with branches spreading in flower, later appressed, scabrous; spikelets 3–5 mm long, 1- to 6-flowered, light green; glumes about 2.5 and 3.5 mm long; lemmas 3–4 mm long, pubescent on the lower back. Meadows and open woods; Parklands, Boreal forests. Var. *interior* (Rydb.) Abbe & Butters (*P. interior* Rydb.), more densely tufted, stiffer, with the branches of the panicles more contracted, is very similar to var. *firmula* Gaud. of Europe and may be identical. Plants of this variety with reduced panicles are forma *rariflora* (Desf.) A. & G. In dry areas, sandy open forest, slopes; Boreal forest, Rocky Mountains.

Poa nervosa (Hook.) Vasey

WHEELER'S BLUE GRASS

Plants with long rhizomes; culms 30–60 cm high, erect, somewhat tufted. Sheaths often purplish below, the lower ones retrorsely pubescent; blades 2–4 mm wide, flat or folded. Panicles 5–10 cm long, open, nodding at the tip, with branches spreading, the lower ones often reflexed; spikelets 4–6 or sometimes 8 mm long; glumes subequal, about 2 mm long; lemmas 3–4 mm long, strongly nerved, pubescent to glabrous on lower back. Slopes, open woods; Rocky Mountains, Cypress Hills.

Poa palustris L.

FOWL BLUE GRASS

Plants loosely tufted, with culms 30–100 cm high, decumbent, purplish at base. Sheaths somewhat flattened, keeled, often somewhat scabrous; blades 2–4 mm wide, flat or loosely folded, lax, scabrous on both sides. Panicles 10–30 cm high, pyramidal or oblong, with branches spreading; spikelets 3–4 mm long, 3- or 4-flowered; glumes subequal, 2.5–3 mm long; lemmas 2.5–3 mm long, often bronzed at the tip, pubescent on lower back. Meadows, moist areas, lakeshores, and riverbanks; throughout Prairie Provinces.

Poa pattersonii Vasey

Plants densely tufted, with erect culms 5–20 cm high. Sheaths smooth; blades about 1 mm wide, folded, lax. Panicles 1–4 cm long, dense, purplish; spikelets 5–6 mm long, 4- or 5-flowered; glumes about 3 and 3.5 mm long; lemmas about 4 mm long, strongly pubescent on keel and marginal nerves. Rare; in alpine meadows and slopes; southern Rocky Mountains.

Poa pratensis L.

KENTUCKY BLUE GRASS

Plants with long rhizomes; culms to 100 cm high. Culms compressed and slightly keeled, glabrous, dark green; blades to 5 mm wide, often to 40 cm long, linear, dark green, the lower side glossy, soft. Panicles 5–15 cm long, contracted before flowering, with branches spreading during flowering; spikelets 3–6 mm long, 3- to 5-flowered, green- or purplish-tinged; glumes subequal, 3–3.5 mm long, the first 1-nerved, the second 3-nerved; lemmas 3.5–4 mm long, distinctly 5-nerved, densely short pubescent on the nerves, and strongly webbed at the base. Common throughout the Prairie Provinces. Sown in lawn mixtures, pastures; also found in meadows, moist prairies, and forest openings. Probably native, as well as introduced. Very variable, consisting of many

racess. Low plants, 15–20 cm high, with small, few-flowered panicles, and often rather grayish green, are var. *humilis* (Ehrh.) Griseb.; on dry ground. Plants 15–30 cm high, culms stiff, usually with a single culm leaf, and panicles small, stiff, with branches single or in pairs, are var. *arenaria* J. & W.; in dry pastures, sand dunes. The form described as *P. agassizensis* Boivin & Love seems to fit in both these varieties.

Poa rupicola Nash

TIMBERLINE BLUE GRASS

Plants densely tufted, with culms 10–20 cm high, erect, stiff. Sheaths smooth; blades 1–1.5 mm wide, erect. Panicles 2–5 cm long, narrow, with branches short, ascending to appressed; spikelets 4–5 mm long, mostly 3-flowered, purple; glumes about 3 mm long; lemmas about 3.5 mm long, villous on lower keel and marginal nerves. Slopes, openings, and meadows; Rocky Mountains.

Poa secunda Presl

SANDBERG'S BLUE GRASS

Plants densely tufted, with erect culms 10–30 cm high. Sheaths compressed, often somewhat scabrous; blades 1–2 mm wide, flat or folded, twisted to erect. Panicles 2–10 cm long, narrow, with branches short ascending to appressed; spikelets 4–6 mm long, pale green; glumes about 3–3.5 mm long; lemmas 3.5–4 mm long, pubescent on lower back. Apparently rare or lacking in Manitoba; dry grasslands; Prairies, Parklands.

Poa stenantha Trin.

Plants tufted, with erect culms 30–50 cm high. Sheaths smooth; blades 1–2 mm wide, flat or somewhat involute, lax. Panicles 5–15 cm long, nodding, with branches drooping; spikelets 6–8 mm long, 3- to 5-flowered; glumes about 2.5 mm and 3.5 mm long; lemmas about 5 mm long, pubescent on lower back. Moist meadows and openings; Rocky Mountains.

Poa trivialis L.

ROUGH-STALKED BLUE GRASS

Plants sod-forming, with culms erect or ascending, sometimes stoloniferous below. Sheaths compressed, sharply keeled; blades 2–5 mm wide, flat, lax, glossy green below. Panicles 10–20 cm long, oblong, often somewhat contracted, with branches scabrous, ascending-spreading; spikelets about 4 mm long, 2- to 5-flowered, usually green, sometimes bronze- or purplish-tinged; glumes about 2 and 3 mm long; lemmas 2.5–3 mm long, finely but distinctly 5-nerved, strongly webbed at the base. Introduced; occasionally sown in pastures; escaped in various locations.

Polypogon beard grass

Polypogon monspeliensis (L.) Desf.

RABBITFOOT GRASS

Plants annual, with culms 15–50 cm high, erect or decumbent. Sheaths scabrous; blades 4–6 mm wide, scabrous. Panicles 1–15 cm long, spike-like, dense, silky, yellowish when mature; glumes 2 mm long, with the apex somewhat lobed and the awn 6–8 mm long, pubescent on the back; lemmas about 1 mm long, short-awned. Introduced; established in various locations as a weedy plant in waste areas, dry banks, coulees, along streams.

Puccinellia salt-meadow grass

Low to medium, tufted, feathery paniced grasses of moist alkaline soils.

- 1. Plants stoloniferous, with the stolons bearing bulblets; lemmas entire, 3.5–4.5 mm long. *P. phryganodes*
Plants not stoloniferous, with the culms ascending or decumbent; lemmas 1.5–3 mm long. 2
- 2. Lemmas rounded or truncate at the tip; lower panicle branches reflexed at maturity. *P. distans*
Lemmas blunt, narrowed to a triangular tip; panicle branches not reflexed. *P. nuttalliana*

***Puccinellia distans* (L.) Parl.**

SLENDER SALT-MEADOW GRASS

Plants sod-forming, bluish green, with culms 15–60 cm high, mostly geniculate-ascending. Sheaths broad, smooth; blades 1–2 mm wide, linear, flat. Panicles 5–15 cm long, pyramidal, with branches scabrous, reflexed at maturity; spikelets 4–6 mm long, 4- to 7-flowered; glumes 1 and 2 mm long; lemmas 1.5–2 mm long, 5-nerved, often reddish-tinged with a hyaline or yellowish membranous margin. Introduced from Europe; not common; on alkaline soils around lakes.

***Puccinellia nuttalliana* (Schultes) Hitchc. (Fig. 60)**

NUTTALL'S SALT-MEADOW GRASS

Plants sod-forming or tufted, with culms 30–60 cm high, usually erect or ascending, slender. Sheaths smooth; blades 1–3 mm wide, flat or involute, glaucous. Panicles 10–20 cm long, open, pyramidal, with branches scabrous, spreading; spikelets 4–7 mm long, 3- to 6-flowered; first glume 1–1.5 mm long; second glume 1.5–2 mm long; lemmas 2–3 mm long, narrowed to an obtuse apex. Common; on moist to rather dry alkaline soils; throughout the Prairie Provinces.

***Puccinellia phryganodes* (Trin.) Scribn. & Merr.**

Plants sod-forming, with sterile culms stoloniferous, bearing bulblets at the nodes; fertile culms 20–40 cm high, erect or decumbent. Sheaths smooth; blades 1–3 mm wide, mostly folded, thick. Panicles 5–15 cm long, open; spikelets 8–12 mm long, 4- to 7-flowered; first glume 2–3 mm long; second glume 3–4 mm long; lemmas 3.5–4.5 mm long. Around sloughs and wet calcareous gravelly soils; Churchill.

Schedonnardus tumble grass

***Schedonnardus paniculatus* (Nutt.) Trel.**

TUMBLE GRASS

Plants spreading, tufted, with culms 20–40 cm high, erect or ascending. Sheaths compressed, sharply keeled; blades 1–2 mm wide, folded, twisted, and wavy. Spikes 2–5 cm long; spikelets about 4 mm long, narrow, distant. Inflorescence elongating at maturity into a spiral that breaks off and rolls with the wind. Dry, sandy, or infertile soils; Prairies.



Fig. 60. Nuttall's salt-meadow grass, *Puccinellia nuttalliana* (Schultes) Hitchc.

Schizachne purple oat grass

Schizachne purpurascens (Torr.) Swallen (Fig. 61)

PURPLE OAT GRASS

Plants with rhizomes; culms 50–100 cm high, loosely tufted, usually decumbent at the base. Sheaths round or slightly flattened; blades 2–6 mm wide, flat to loosely folded. Panicles 6–15 cm long, often secund, with branches more or less drooping, single or in pairs; spikelets 20–25 mm long; first glume 4–5 mm long; second glume 6–8 mm long, purplish; lemmas 8–10 mm long, with the awn 10–15 mm long. In woods throughout the Prairie Provinces; rare in Prairies.

Scolochloa spangletop

Scolochloa festucacea (Willd.) Link (Fig. 62)

SPANGLETOP

Plants with thick, long rhizomes; culms to 150 cm high, stout. Sheaths glabrous, prominently veined; blades 5–10 mm wide, flat to convolute. Panicles 15–20 cm long, loose, with branches fascicled, ascending; spikelets 6–9 mm long, 4- to 7-flowered; first glume 5–7 mm long; second glume 6–8 mm long; lemmas 5–7 mm long, villous at the base. Wet areas throughout the Prairie Provinces.

Secale rye

Secale cereale L.

RYE

Annual or winter annual; culms to 2 m high, usually densely pubescent above. Leaves flat, often glaucous, to 15 mm wide. Spikes to 15 cm long; spikelets usually 2-flowered; glumes awl-shaped, to 1 cm long; lemmas 3- to 5-nerved, to 18 mm long, with the awn 4–8 cm long. A cereal grain grown on light soils; occasionally established in roadsides and waste areas.

Setaria foxtail

Weedy annuals with cylindrical, dense, spike-like, bristly heads, found in waste places and as a weed in cultivated fields. Introduced from Europe but rapidly spreading.

Spikelets with more than 5 bristles, these mostly less than 3 times as long as the spikelet. *S. glauca*

Spikelets with fewer than 5 bristles, these 3–5 times as long as the spikelet. *S. viridis*

Setaria glauca (L.) Beauv.

YELLOW FOXTAIL

Plants annual, tufted, with culms 50 cm or more high, erect, occasionally branching above. Sheaths compressed, keeled, pale green; blades 5–12 mm wide, flat or loosely folded, twisted, soft, gray green. Panicles 5–10 cm long, spike-like, cylindrical; spikelets 3–3.5 mm long; first glume about 1.2 mm long, 3-nerved; second glume about 1.7 mm long, 5-nerved; lemmas about 3 mm long, rugose; bristles usually 6–8 or up to 20, 6–10 mm long. Not common; introduced; weedy in gardens and waste places.



Fig. 61. Purple oat grass, *Schizachne purpurascens* (Torr.) Swallen.




Fig. 62. Spangletop, *Scolochloa festucacea* (Willd.) Link.

Plants annual, tufted, with culms 20–50 cm high, erect or geniculate-ascending. Sheaths slightly compressed, glabrous or appressed-pubescent; blades 5–12 mm wide, flat, light green. Panicles 3–10 cm long; spikelets 2–2.5 mm long; first glume less than 1 mm long, 1-nerved; second glume about 2 mm long, 5-nerved; lemmas 2–2.5 mm long, smooth; bristles 1–3, below each spikelet, 8–10 mm long. Introduced; weedy in gardens, cultivated land, roadsides; throughout the Prairies and Parklands.

Sitanion squirreltail

Sitanion hystrix (Nutt.) J. G. Smith

SQUIRRELTAIL

Plants densely tufted, with culms 10–50 cm high, erect to spreading, stiff. Sheaths somewhat keeled above, softly pubescent or glabrous; blades 1–3 mm wide, flat to involute, finely pubescent on both sides. Spikes 2–7 cm long, the lower parts often included in the sheaths; spikelets about 15 mm long excluding the awns; glumes about 6 mm long excluding the awn, very narrow; lemmas about 10 mm long excluding the awn; awns of glumes and lemmas 3–10 cm long. Very rare; slopes, dry grassland, open areas; Prairies.

Sorghastrum Indian grass

Sorghastrum nutans (L.) Nash

INDIAN GRASS

Plants with long, scaly rhizomes; culms 60–150 cm high, loosely tufted, somewhat pubescent on the nodes. Sheaths round, distinctly veined; blades 5–10 mm wide, flat, to 50 cm long, dull green to glaucous. Panicles 10–25 cm long, narrow, yellowish, brownish at maturity; spikelets 6–8 mm long, with the awn 10–15 mm long; summit of branchlets, rachis joints, pedicels, and spikelets long grayish hirsute. Rare; in grassland; southeastern Parklands.

Spartina cord grass

Erect coarse grasses with scaly rhizomes found in moist areas throughout the Prairie Provinces. Spikelets crowded on one side of panicle branches, which are borne alternately on the main stem and closely parallel it. Leaves very coarse.

Lemmas awned, found only in eastern prairies. *S. pectinata*

Lemmas not awned. *S. gracilis*

Spartina gracilis Trin. (Fig. 64)

ALKALI CORD GRASS

Plants with tough rhizomes; culms 60–100 cm high, solitary or in small tufts. Sheaths glabrous, yellowish green, to 5 mm wide, very scabrous above. Inflorescence 6–15 cm long, with 4–8 closely appressed spikes 2–4 cm long; spikelets 6–8 mm long; first glume about 5 mm long; second glume about 10 mm long; lemmas about 10 mm long; glumes and lemmas ciliate on the keel. Common; in dry to wet alkali areas, and sandy soils; Prairies and Parklands.




Fig. 63. Green foxtail, *Setaria viridis* (L.) Beauv.




Fig. 64. Alkali cord grass, *Spartina gracilis* Trin.

Spartina pectinata Link

PRAIRIE CORD GRASS

Plants with long, tough rhizomes; culms to 2 m high, solitary or a few together. Sheaths glabrous, distinctly veined; blades 5–15 mm wide, to 60 cm long, flat to involute. Inflorescence 10–30 cm long, with 10–20 spikes 4–8 cm long, ascending to appressed; first glume 5–6 mm long, short-awned; second glume 8–12 mm long, with the awn 4–10 mm long; lemmas 7–9 mm long. Moist prairie, riverbanks, and lakeshores; southeastern Prairies and Parklands.

Sphenopholis wedge grass

Tall tufted grasses with narrow panicles and many shining spikelets.

Second glume much wider than lemma, wedge-shaped; panicle dense, spike-like, erect. *S. obtusata*

Second glume not much wider than lemma, obtuse or acute; panicle not dense or spike-like, drooping. *S. intermedia*

Sphenopholis intermedia Rydb.

SLENDER WEDGE GRASS

Plants in small tufts, with erect culms 30–60 cm high. Sheaths glabrous or somewhat pubescent; blades 2–5 mm wide, flat, soft. Panicles 8–15 cm long, to 3 cm in diam, more or less lobed, lustrous; first glume 1.5–2.5 mm long; second glume 2–3 mm long; lemmas 2.5–3 mm long. Moist soil, ravines, margins of woods, and meadows; throughout the Prairie Provinces.

Sphenopholis obtusata (Michx.) Scribn.

PRAIRIE WEDGE GRASS

Plants tufted, with erect culms 20–60 cm high. Sheaths glabrous or slightly scabrous, distinctly veined; blades 2–5 mm wide, flat, scabrous on both sides. Panicles 5–15 cm long, narrow, dense, mostly spike-like, sometimes lobed below, lustrous; first glume 1–2 mm long; second glume 1.5–2.5 mm long; lemmas 1.5–2.5 mm long. Moist areas, shrubbery, coulee bottoms, creek banks, and lakeshores; throughout the Prairie Provinces.

Sporobolus dropseed

Tufted plants with paniculate inflorescence, the seeds readily falling at maturity, giving the common name to the genus. Fairly palatable forage plants.

- 1. Plants annual; panicle contracted or spike-like. *S. neglectus*
- Plants perennial; panicle open and branches more or less spreading. 2
- 2. Spikelets less than 3.2 mm long. *S. cryptandrus*
- Spikelets more than 3.2 mm long. *S. heterolepis*

Plants tufted, with culms 30–100 cm high, solitary or few together, erect or spreading to prostrate. Sheaths prominently veined, often purplish at the base, with the margin ciliate; blades 2–5 mm wide, flat to involute, with conspicuous tufts of hairs at the base. Panicles 10–25 cm long, the base often included in the sheath, with branches spreading or reflexed; spikelets 2–2.5 mm long, grayish; first glume about 1 mm long; second glume 2–2.5 mm long; lemmas about 2.5 mm long. Prairies and open woods on sandy soils; Prairies and Parklands.

Sporobolus heterolepis A. Gray

Plants densely tufted, with culms 30–70 cm high, erect, slender. Sheaths somewhat flattened, glabrous or pubescent; blades 1–3 mm wide, to 45 cm long, flat to involute, erect or slightly drooping. Panicles 5–20 cm long, oblong to narrowly ovoid, with branches 3–6 cm long, ascending to appressed; spikelets 3–6 mm long, grayish; first glume 2–4 mm long; second glume 4–6 mm long; lemmas 5–5.5 mm long. Moist grasslands; southeastern Prairies and Parklands.

Sporobolus neglectus Nash

Plants annual, tufted, with culms 20–40 cm high, erect or spreading. Sheaths glabrous; blades 1–2 mm wide, with the upper culm leaves 1–2 cm long. Panicles 2–5 cm long, rarely exserted, often exceeded by the uppermost blade; spikelets 2.5–3 mm long; first glume 1.5–2.5 mm long; second glume 2–3 mm long; lemmas 2–3 mm long. Rare; dry open areas; Prairies.

Stipa needle grass

Medium- to tall-growing tufted grasses with narrow panicles of fairly large spikelets bearing twisted awns. Spikelets breaking off at maturity above glumes. Lemmas remaining attached to the seed characterized by a hard, sharp-pointed, hairy callus, and a long twisted awn. The needle-like lemmas injurious to the mouthparts and skin of grazing animals, particularly sheep. Except when the seeds are mature and still attached, the needle grasses are excellent forage plants, both for hay and pasture, and are one of the most important groups of forage plants on the western range.

- 1. Glumes 15–40 mm long; lemmas 8–20 mm long; awns 10–25 cm long. 2
- Glumes 8–10 mm long; lemmas 5–7 mm long; awns 2–3 cm long. 4
- 2. Awns indistinctly geniculate, with the upper part strongly curled and flexuous; glumes 15–30 mm long; lemmas 8–15 mm long. *S. comata*
- Awns distinctly twice geniculate. 3
- 3. Glumes 30–40 mm long; lemmas 15–25 mm long; awns to 25 cm long. *S. spartea*
- Glumes 20–30 mm long; lemmas 12–15 mm long; awns to 12 cm long. *S. spartea* var. *curtiseta*



Fig. 65. Sand dropseed, *Sporobolus cryptandrus* (Torr.) A. Gray.

4. Panicle open, with branches spreading and drooping; spikelets few at ends of branches. *S. richardsonii*
 Panicle narrow with appressed branches. 5
5. Callus blunt; collar and lower nodes of panicle villose. *S. viridula*
 Callus sharp-pointed; collar and lower nodes of panicle glabrous. *S. columbiana*

Stipa columbiana Macoun

COLUMBIA NEEDLE GRASS

Plants tufted, with culms 30–60 cm high, erect or ascending. Sheaths glabrous; blades 1–3 mm wide, to 25 cm long, usually involute. Panicles 10–25 cm long, narrow, with branches appressed-ascending; glumes about 1 cm long; lemmas 6–7 mm long, pubescent, with the callus sharp-pointed; awns 2–2.5 cm, twisted below, twice geniculate. Prairies and openings; Rocky Mountains, Peace River, Cypress Hills.

Stipa comata Trin. & Rupr. (Fig. 66)

SPEAR GRASS

Plants in small, dense tufts, with culms 30–60 cm high, erect. Sheaths round or somewhat compressed; blades 1–3 mm wide, flat to involute, prominently ridged above. Panicles 10–20 cm long, commonly included at the base; glumes 15–25 mm long; lemmas 8–12 mm long, at first pale, shiny brown at maturity, sparsely pubescent; awns 10–15 cm long, twisted below, curled toward the tip, flexuous. Very common; in grasslands of the Prairies and Parklands, and in the Peace River region. A very important forage species in the drier parts of the grasslands.

Stipa richardsonii Link

RICHARDSON NEEDLE GRASS

Plants tufted, with culms 50–80 cm high, usually erect. Sheaths slightly flattened, smooth or scabrous; blades 1–3 mm wide, involute, scabrous, indistinctly veined. Panicles 10–20 cm long, open, with branches slender, flexuous, spreading or drooping, bearing spikelets at the tips; glumes 8–10 mm long; lemmas about 5 mm long, pubescent, brown when mature; awns 25–35 mm long, weakly twice geniculate. Moist grasslands; Rocky Mountains, Cypress Hills, Parklands, Riding Mountain, Duck Mountain.

Stipa spartea Trin.

PORCUPINE GRASS

Plants in large tufts or tussocks, with erect culms to 1 m high. Sheaths round, with the outer margin usually ciliate; blades 3–5 mm wide, flat to convolute, ridged above. Panicles 15–20 cm long, narrow, nodding, with branches slender, each bearing one or two spikelets; glumes 3–4 cm long, tapering to a slender point; lemmas 15–25 mm long, brown, with the callus 7 mm long, villos; awns stout, 15–25 cm long, twice geniculate. Becoming rare; grassland and openings; southeastern Prairies and Parklands.

Stipa spartea Trin. var. *curtiseta* Hitchc.

WESTERN PORCUPINE GRASS

Very similar in stature to spear grass, but with wider, usually flat leaves; glumes 2–3 cm long; lemmas 12–15 mm long, brown; awns to 10 cm long, twice geniculate. Common; in moist prairie; throughout Prairies, Parklands, and Rocky Mountains; replacing spear grass as dominant in the moister sites.



Fig. 66. Spear grass, *Stipa comata* Trin. & Rupr.

Plants loosely tufted, with erect culms 50–100 cm high. Sheaths round, prominently veined, villous at the throat; blades 2–5 mm wide, to 25 cm long, flat. Panicles 10–20 cm long, narrow, with branches appressed-ascending; glumes 7–10 mm long; lemmas 5–6 mm long, plump, with the callus blunt, appressed-pubescent. Moderately dry to moist areas, in shrubbery, forest margins; throughout Prairies and Parklands. A variety of this species has been improved, and named green stipa grass, occasionally seeded for forage, mostly in comparative trials.

Trisetum trisetum

Tufted perennials, with open or contracted panicles; spikelets usually 2- or 3-flowered, shiny. Not common, of almost no forage value.

- 1. Lemmas awnless, or the awn less than 2 mm long, hidden by the glumes. *T. wolfii*
 Lemmas awned, with the awn bent and exerted. 2
- 2. Panicle open, nodding, with branches flexuous. *T. cernuum*
 Panicle contracted, with branches short, ascending. 3
- 3. Plants densely tufted; panicles 5–15 cm long, spike-like. *T. spicatum*
 Plants loosely tufted, with culms often solitary; panicles 10–25 cm long, narrow, loose. *T. canescens*

Trisetum canescens Buckl.

Plants loosely tufted, with culms 60–100 cm high, erect or decumbent at the base, often solitary. Sheaths, at least the lower one, sparsely to densely retrorse pubescent; blades 2–7 mm wide, flat, scabrous to pubescent. Panicles 10–25 cm long, narrow, usually loose, occasionally interrupted and spike-like, with branches appressed-ascending; spikelets about 8 mm long, 2- or 3-flowered; glumes 5–7 mm long; lemmas 5–6 mm long; rachilla hairs copious; awns to 12 mm long, geniculate. Moist meadows, woods, and openings; southern Rocky Mountains.

Trisetum cernuum Trin.

Plants loosely tufted, with culms 60–100 cm high, rather lax. Sheaths glabrous or sparsely pubescent; blades 6–12 mm wide, flat, lax, scabrous. Panicles 15–30 cm long, open, with branches flexuous, spreading or ascending; spikelets 6–12 mm long, usually with 3 florets; first glume 0.5–2 mm long, 1-nerved; second glume 3–4 mm long, 3-nerved; lemmas 5–6 mm long; callus hairs to 1 mm long, those of the rachilla to 2 mm long; awns 5–10 mm long, flexuous. Moist woods and openings; southern Rocky Mountains.



Fig. 67. Green needle grass, *Stipa viridula* Trin.

Trisetum spicatum (L.) Richt.

SPIKE TRISETUM

Plants tufted, with culms 10–50 cm high, erect, pubescent below the panicle. Sheaths retrorse-pubescent; blades 1–3 mm wide, flat or involute, pubescent to subglabrous. Panicles 3–10 cm long, dense; spikelets about 6 mm long, shiny; first glume 3–5 mm long, 1-nerved; second glume 3.5–5.5 mm long, 3-nerved; lemmas 3.5–5 mm long; awns to 6 mm long, inserted below the apex, flexuous. Not common; grasslands, open woods; Rocky Mountains, Cypress Hills.

Trisetum wolfii Vasey

AWNLESS TRISETUM

Plants loosely tufted, with culms 50–100 cm high, erect. Sheaths scabrous, or the lower sparsely pubescent; blades 2–4 mm wide, flat, scabrous or pilose above. Panicles 8–15 cm long, rather dense, with short branches appressed-ascending; spikelets 5–7 mm long, usually 2-flowered; glumes about 5 mm long, subequal; lemmas 4–5 mm long, awnless or with a short awn below the tip. Meadows and moist woods; southern Rocky Mountains, Cypress Hills.

Triticum wheat

Triticum aestivum L.

WHEAT

Annual plants, with stems to 1 m high. Leaves flat, to 2 cm wide. Spike to 12 cm long, dense; spikelets 2- to 5-flowered; glumes ovate, the upper part keeled, mucronate; lemmas glabrous to pubescent, awnless or awned, depending on the variety. The most important cereal grain, with many varieties, in the Prairie Provinces; occasionally established in roadsides and waste areas.

Zizania wild-rice

Zizania aquatica L.

ANNUAL WILD-RICE

Plants aquatic, with stout culms 1–2 m high. Sheaths long, with the ligule 10–15 mm long; blades 5–15 mm wide, flat. Panicles 30–50 cm long, with branches 10–20 cm long; spikelets unisexual; upper half of panicle pistillate, with branches appressed-ascending; lower half of panicle staminate, with branches spreading, spikelets pendulous, 6–11 mm long; pistillate spikelets with awns 1–6 cm long. Borders of streams and lakes; native in southeastern Boreal forest; introduced and established in several lakes in Boreal forests of Saskatchewan and Alberta.

CYPERACEAE—sedge family

Grass-like or rush-like annual or perennial plants, usually with solid stems. Root fibrous or of long running rootstocks. The long, narrow leaves with closed sheaths on three sides of the stem, or 3-ranked. Flowers either perfect or unisexual with no sepals or petals, their places being taken by bristles or scales (perigynia). Although usually associated with moist or marshy areas, many species occur in very arid localities.

1. Flowers perfect, with spikelets uniform; achenes not enclosed in a perigynium or bract. 2
 Flowers imperfect, with staminate and pistillate flowers in same or different spikelets; achenes enclosed in a perigynium or bract. 6
2. Scales of the spikelets 2-ranked, keeled; perianth lacking; spikelets in simple or compound terminal umbels. *Cyperus*
 Scales of the spikelets several-ranked, spirally arranged, not keeled; perianth present as bristles; spikelets solitary and terminal or partly lateral. 3
3. Base of the style persistent as a tubercle on top of the achene. 4
 Base of the style not persistent, the achene without a tubercle. 5
4. Culms leafless, with sheaths usually bladeless; spikelet solitary, terminal. *Eleocharis*
 Culms with bristle-like leaves; spikelets several. *Rhynchospora*
5. Perianth bristles 1–8, occasionally lacking, usually little longer or shorter than the achene. *Scirpus*
 Perianth bristles numerous, often 2–3 cm long, silky. *Eriophorum*
6. Achenes partly enclosed by a spathe-like scale. *Kobresia*
 Achenes entirely enclosed by a perigynium. *Carex*

Carex sedge

The genus *Carex* is very large, and its taxonomy is difficult. To distinguish between species, characters of the mature inflorescence are needed. The species can be divided into several groups of species that share common characters. However, determination within groups is often not as easy, because the characters are variable, hybridization between species is known to occur, and the opinions of experts differ on the validity of species and varieties.

Sedges are important in the Prairie Provinces. Some species, such as the awned sedge, beaked sedge, and water sedge, form a large part of the “slough hay,” harvested in many areas. In dry prairie, several species of sedges increase in abundance as a result of overgrazing, and thereby they help prevent soil erosion. In Boreal forest, sedges are often dominant in swamp vegetation, and are helpful in building peat soils. Also, in alpine tundra and meadows in the Rocky Mountains, sedges are often the dominant constituents of the vegetation.

Because of their importance, the sedges have been treated in detail in this publication. A total of 125 species and many varieties are described. The treatment followed is conservative, in that several closely related species are grouped into one large species. Keys are provided to separate the small species within these groups.

Well-developed mature fruits (perigynia) and inflorescences are needed for successful determinations. The various types of perigynia and inflorescences are shown in Figs. 68 and 69.




Fig. 68. Flowering and fruiting characters of sedges.




Fig. 69. Inflorescences of sedges: *a*, single spike; *b–k*, two or more spikes; *b–f*, mostly bisexual spikes; *b, c*, spikes approximate; *d*, spikes crowded; *e*, compound spikes; *f*, gynaecandrous spikes; *g–k*, terminal spikes staminate, lateral spikes pistillate; *g, h*, pistillate spikes more or less long-peduncled; *k*, pistillate spikes sessile.

Key to the Groups of *Carex*

- | | | |
|--|---------|---|
| 1. Spike one, usually bractless; leaves usually all basal. | Group 1 | |
| Spikes two or more, with bracts often present; stem leaves often present. | | 2 |
| 2. Spikes mostly bisexual, sessile; achenes lenticular; stigmas 2. | Group 2 | |
| Spikes of two kinds, the terminal spikes staminate, the lower spikes pistillate. | | 3 |
| 3. Achenes lenticular; stigmas 2. | Group 3 | |
| Achenes triangular; stigmas 3. | | 4 |
| 4. Perigynia with a conspicuously bidentate beak. | Group 4 | |
| Perigynia beakless, or the beak not conspicuously bidentate. | Group 5 | |

Key to Sections and Species in Group 1

- | | | |
|---|---|---|
| 1. Achenes lenticular; stigmas 2. | | 2 |
| Achenes triangular; stigmas 3. | | 4 |
| 2. Spikes usually completely staminate or completely pistillate; perigynia dark, lustrous, spreading to reflexed. | Section 3. Dioicae
(<i>C. gynocrates</i>) | |
| Spikes androgynous; perigynia ascending to spreading. | | 3 |
| 3. Perigynia stipitate, striate. | Section 1. Nardinae
(<i>C. nardina</i>) | |
| Perigynia sessile, not striate. | Section 2. Capitatae
(<i>C. capitata</i>) | |
| 4. Pistillate scales deciduous; at least the lower perigynia reflexed at maturity. | | 5 |
| Pistillate scales not deciduous; perigynia spreading to ascending at maturity. | | 6 |
| 5. Perigynia dark brown; style jointed to the achene. | Section 4. Callistachys | |
| Perigynia yellowish to light brown; style continuous with the achene. | Section 44. Orthocerates | |
| 6. Plants dioecious, rhizomatous; perigynia pubescent. | Section 23. Scirpinae
(<i>C. scirpoidea</i>) | |
| Plants monoecious; spikes androgynous. | | 7 |
| 7. Margins of the staminate scales united in the lower part. | | 8 |
| Margins of the staminate scales free to the base. | | 9 |

8. Pistillate scales leaf-like; perigynia clearly beaked.	Section 19. Phyllostachyae (<i>C. backii</i>)	
Pistillate scales not leaf-like; perigynia rounded at the tip, beakless.	Section 18. Politrichoideae (<i>C. leptalea</i>)	
9. Perigynia pubescent or puberulent; plants densely caespitose; leaves stiffly filiform.	Section 20. Filifoliae (<i>C. filifolia</i>)	
Perigynia not pubescent.		10
10. Perigynia stipitate, striate.	Section 1. Nardinae (<i>C. nardina</i>)	
Perigynia sessile, not striate.		11
11. Pistillate scales without hyaline margins, shorter than the perigynia; prairie species.	Section 21. Obtusatae (<i>C. obtusata</i>)	
Pistillate scales with hyaline margins, concealing the perigynia; mountain species.		12
12. Perigynia 3–4 mm long.	Section 25. Rupestres	
Perigynia 5–7 mm long.	Section 26. Firmiculmes (<i>C. geyeri</i>)	

Key to Sections and Species in Group 2

1. Culms single or few together from long creeping rhizomes or stolons.		2
Culms more or less densely tufted; rhizomes, if any, short.		6
2. Culms stoloniferous, producing shoots from the nodes of old decumbent stems.	Section 9. Chordorrhizae (<i>C. chordorrhiza</i>)	
Culms not stoloniferous; new shoots arising from rhizomes.		3
3. Spikes closely aggregated into a globose or ovoid head; bracts scarious; spikes androgynous.	Section 5. Foetidae (<i>C. maritima</i>)	
Spikes not closely aggregated; bracts not scarious.		4
4. Perigynia 4–6 mm long, wing-margined at the tip, with bidentate beak.	Section 8. Arenariae (<i>C. siccata</i>)	
Perigynia to 4 mm long, not wing-margined, not conspicuously bidentate beaked.		5

5. Ventral strip of sheath green, nerved to the top; collar with two dark warts; spikes 10–20 or more in a head. Section 7. Intermediae
(*C. sartwellii*)
- Ventral strip of sheath hyaline or brown, unveined; collar without warts; spikes usually less than 10 in a head. Section 6. Divisae
6. Spikes androgynous. 7
 Spikes gynaecandrous. 11
7. Spikes with only 2 or 3 perigynia; perigynia beakless, not flattened on the inside. Section 14. Heleonastes
(*C. disperma*)
- Spikes with more than 3 perigynia; perigynia beaked, flattened on the inside. 8
8. Spikes 4–10 in the inflorescence; all rachis nodes bearing a single spike. Section 10. Bracteosa
- Spikes usually more than 10 in the inflorescence; at least the lowest rachis node bearing 2 or more spikes. 9
9. Culms soft and weak, the angles sharply winged; perigynia widest at base, the beak about as long as the body; sheaths loose. Section 13. Vulpinae
- Culms stiff, not winged; perigynia widest above the middle, the beak about half as long as the body; sheaths tight. 10
10. Perigynia straw-colored; pistillate scales awned; ventral strip of sheaths cross-wrinkled. Section 11. Multiflorae
(*C. vulpinoidea*)
- Perigynia brown to black; pistillate scales not awned; ventral strip of sheaths smooth, with reddish spots. Section 12. Paniculatae
11. Perigynia winged along part or all of the margin; always with a more or less distinctly bidentate beak. Section 17. Ovale
- Perigynia not winged along the margin; the beak, if present, not distinctly bidentate. 12
12. Achenes filling the perigynia almost fully; perigynia white punctulate under magnification. Section 14. Heleonastes
- Achenes filling only the upper half or two-thirds of the perigynia; perigynia not punctulate. 13
13. Achene visible through the wall of the perigynium; perigynia more than 4 mm long; awn-like bracts present. Section 16. Deweyanae
(*C. deweyana*)

Achene not visible through the wall of the perigynium; perigynia less than 4 mm long; bracts only slightly developed.

Section 15. Stellulatae
(*C. muricata*)

Key to Sections and Species in Group 3

- | | | |
|--|---|---|
| 1. Lower pistillate spikes long-peduncled, pendulous. | Section 43. Cryptocarpae | |
| Lower pistillate spikes sessile or short-peduncled, erect or ascending. | | 2 |
| 2. Perigynia shiny, inflated, clearly ribbed, the beak pronounced; style continuous with the achene. | Section 48. Vesicariae
(<i>C. saxatilis</i>) | |
| Perigynia not shiny or inflated, the ribs obscure or absent, the beak not pronounced; style jointed to the achene. | | 3 |
| 3. Perigynia rounded, not beaked; bracts with a sheath. | | 4 |
| Perigynia flattened, with a short beak; bracts sheathless. | Section 42. Acutae | |
| 4. Sheaths of bracts long; perigynia serrulate toward the tip. | Section 27. Albae
(<i>C. rufina</i>) | |
| Sheaths of bracts short; perigynia not serrulate. | Section 28. Bicolores | |

Key to Sections and Species in Group 4

- | | | |
|--|---|---|
| 1. Perigynia pubescent or puberulent. | | 2 |
| Perigynia glabrous. | | 3 |
| 2. Perigynia rounded in cross section, not ribbed; lower bracts shorter than or equaling the inflorescence. | Section 22. Montanae | |
| Perigynia triangular in cross section, ribbed; lower bracts longer than the inflorescence. | Section 39. Hirtae | |
| 3. Style continuous with and usually strongly curved above the achene, persistent; leaves often strongly and clearly cross-veined. | | 4 |
| Style jointed to the achene, straight, withering; leaves slightly or not cross-veined. | | 8 |
| 4. Perigynia about 6 times as long as wide; lower bracts sheathing. | Section 45. Folliculatae
(<i>C. michauxiana</i>) | |
| Perigynia about 3–4 times as long as wide; lower bracts sheathless. | | 5 |

- | | | |
|---|---|----------|
| <p>5. Perigynia shiny, conspicuously 7- to 9-ribbed, usually contracted at base of beak.
Perigynia not shiny, 12- to 25-ribbed, though often obscurely, tapering to the beak.</p> | <p>Section 48. Vesicariae</p> | <p>6</p> |
| <p>6. Perigynia 9–18 mm long; pistillate spikes globose.
Perigynia to 10 mm long; pistillate spikes cylindrical.</p> | <p>Section 49. Lupulinae
(<i>C. intumescens</i>)</p> | <p>7</p> |
| <p>7. Lower pistillate spikes long-peduncled, pendulous.
Lower pistillate spikes short-peduncled, ascending or erect.</p> | <p>Section 46. Pseudo-Cyperae
Section 47. Paludosae</p> | |
| <p>8. Pistillate spikes long-peduncled, loosely flowered.
Pistillate spikes sessile or short-peduncled, densely and closely flowered.</p> | <p>Section 35. Longirostres
(<i>C. sprengelii</i>)
Section 36. Extensae</p> | |

Key to Sections and Species in Group 5

- | | | |
|---|---|----------------|
| <p>1. Margins of staminate scales united toward the base; pistillate scales leaf-like.
Margins of staminate scales free to the base; pistillate scales not leaf-like.</p> | <p>Section 19. Phyllostachyae
(<i>C. backii</i>)</p> | <p>2</p> |
| <p>2. Perigynia pubescent or puberulent.
Perigynia glabrous.</p> | | <p>3
6</p> |
| <p>3. Pistillate bracts developed, sheathed; perigynia 5–6 mm long, lanceolate.
Pistillate bracts lacking either sheath or blade; perigynia less than 5 mm long, ovoid.</p> | | <p>4
5</p> |
| <p>4. Spikes 3–8, the upper 3 or 4 closely aggregated; perigynia with a minute beak.
Spikes about 3, widely separated; perigynia with a beak about as long as the body.</p> | <p>Section 37. Ferrugineae
Section 33. Sylvaticae
(<i>C. assiniboinensis</i>)</p> | |
| <p>5. Pistillate bracts sheathless, blades more or less well developed; perigynia rounded in cross section.
Pistillate bracts long-sheathing, bladeless; perigynia triangular in cross section.</p> | <p>Section 22. Montanae
Section 24. Digitatae</p> | |
| <p>6. Leaves and sheaths pubescent or puberulent.</p> | <p>Section 38. Virescentes
(<i>C. torreyi</i>)</p> | |

Leaves and sheaths glabrous.		7
7. Lower pistillate bracts long-sheathing.		8
Lower pistillate bracts sheathless or almost sheathless.		13
8. Bracts bladeless; perigynia less than 2 mm long; leaves involute, to 0.5 mm wide.	Section 27. <i>Albae</i> (<i>C. eburnea</i>)	
Bracts with blades; perigynia longer than 2 mm; leaves wider than 0.5 mm.		9
9. Pistillate spikes long-peduncled, pendulous.		10
Pistillate spikes short-peduncled, erect to ascending.		11
10. Pistillate spikes 1–6 cm long; culms to 60 cm high; leaves 3–9 mm wide, deep green.	Section 32. <i>Gracillimae</i> (<i>C. gracillima</i>)	
Pistillate spikes usually less than 1 cm long; culms 10–20 cm high; leaves less than 2 mm wide, yellowish brown.	Section 34. <i>Capillares</i> (<i>C. capillaris</i>)	
11. Culms soft, somewhat winged; sheaths loose, dilated at summit; perigynia abruptly constricted and bent into a beak.	Section 30. <i>Laxiflorae</i> (<i>C. laxiflora</i>)	
Culms firm, not winged; sheaths tight; perigynia not contracted and bent into the beak.		12
12. Perigynia obscurely veined; veins less than 10; perigynia tapered at base, somewhat triangular in cross section.	Section 29. <i>Paniccae</i>	
Perigynia clearly veined; veins more than 10; perigynia rounded at base, rounded in cross section.	Section 31. <i>Granulares</i>	
13. Pistillate spikes long-peduncled, spreading or pendulous.	Section 33. <i>Sylvaticae</i> (<i>C. castanea</i>)	
Pistillate spike short-peduncled or sessile, ascending or erect.		14
14. Bracts of the pistillate spikes small or absent; pistillate spikes close together, about 15 mm long.		15
Bracts of the pistillate spikes well-developed; pistillate spikes not close together, longer than 15 mm.		16
15. Pistillate spikes oblong; culms longer than the leaves; plants cespitose.	Section 25. <i>Rupestres</i> (<i>C. glacialis</i>)	

- Pistillate spikes spherical; culms about as long as the leaves; plants rhizomatous. Section 21. Obtusatae (*C. supina*)
16. Perigynia 7–9 mm long, shiny, inflated, strongly 7- to 9-veined; style continuous with the achene. Section 48. Vesicariae (*C. oligosperma*)
- Perigynia less than 7 mm long, not shiny or inflated, obscurely veined; style jointed to the achene. 17
17. Perigynia with a beak about one-quarter as long as the body, not flattened, spreading at maturity. Section 36. Extensae
- Perigynia beakless or almost beakless, clearly flattened, ascending to erect. 18
18. Root system covered with a yellowish felt; terminal spike staminate; perigynia yellowish to greenish. Section 40. Limosae
- Root system without felt cover; terminal spike gynaecandrous or staminate; perigynia greenish to brown. Section 41. Atratae

Section 4. Callistachys

- Perigynia erect or spreading at maturity; leaves involute, 1 mm wide; staminate flowers few; plants cespitose. *C. pyrenaica*
- Perigynia deflexed at maturity; leaves flat, 1.5–2.5 mm wide; staminate flowers numerous; plants rhizomatous. *C. nigricans*

Section 6. Divisae

1. Inflorescence 1–2 times as long as broad; culms about 20 cm high; rhizomes brownish, 1–2 mm in diam. 2
- Inflorescence 2–5 times as long as broad; culms usually more than 20 cm high; rhizomes brown or black, 2–3 mm thick. *C. praegracilis*
2. Plants monoecious; perigynia 2.5–3 mm long, not concealed by the scales; the beak about 0.5 mm long. *C. stenophylla*
- Plants dioecious; perigynia 3.5–4 mm long, fully concealed by the scales; the beak about 1.5 mm long. *C. douglasii*

Section 10. Bracteosae

1. Spikes barely overlapping in the head, erect with a bract; beak of the perigynium about as long as the body. *C. hookerana*

- Spikes densely crowded in the head, only the lowest spike with a bract; beak of the perigynium shorter than the body. 2
2. Heads ovoid, about as long as broad; perigynia and scales brown. *C. hoodii*
- Heads elongate, more than twice as long as broad; perigynia and scales yellowish. 3
3. Culms firm, sharply triangular, not winged; perigynia about 4 mm long, the beak about 1 mm long. *C. gravida*
- Culms soft, the angles winged; perigynia about 3 mm long, the beak about 1.5 mm long. Section 13. Vulpinae
(*C. alopecoidea*)

Section 12. Paniculatae

- Perigynia dark brown, not concealed by the scales; sheaths with red spots. *C. diandra*
- Perigynia light brown to brown, concealed by the scales; sheaths without red spots. *C. prairea*

Section 13. Vulpinae

- Perigynia lanceolate, the beak about as long as the body, the base spongy. *C. stipata*
- Perigynia ovoid, the beak about half as long as the body, the base not spongy. *C. alopecoidea*

Section 14. Heleonastes

1. Spikes few, widely spaced in the inflorescence; perigynia 1-5 in a spike. 2
- Spikes usually several, at least the upper ones close together; perigynia usually more than 5 in a spike. 3
2. Spikes androgynous; bracts obsolete or that of the lowest spike about 1 cm long. *C. disperma*
- Spikes gynaeandrous; bract of the lowest spike as long as or longer than the inflorescence. *C. trisperma*
3. Beak of the perigynia merely indicated; scales silvery, more or less transparent. 4
- Beak of the perigynia at least 0.5 mm long; scales whitish to brown, not transparent. 5
4. Spikes closely crowded into a head at the summit of the stem; scales about as long as the perigynia. *C. tenuiflora*

- Spikes not crowded into a head, spaced along the stem; scales about half as long as the perigynia. *C. loliacea*
5. Spikes more or less crowded into a head at the summit of the stems. 6
- Spikes spaced in an elongated inflorescence. 7
6. Perigynia broadest at the base; dorsal veins pronounced and raised. *C. arcta*
- Perigynia broadest at the middle, tapering to the base; dorsal veins not pronounced and raised. *C. heleonastes*
7. Perigynia stipitate; the terminal spike with a long staminate base; scales reddish orange brown, as long as the perigynia. *C. mackenziei*
- Perigynia not stipitate; the terminal spike with few staminate flowers; scales whitish to brown, shorter than the perigynia. 8
8. Perigynia 5–10 in a spike; the beak about 1.5 mm long. *C. brunnescens*
- Perigynia 10–30 in a spike; the beak about 0.25 mm long. *C. curta*

Section 17. Ovales

1. Bracts leaf-like or considerably longer than the inflorescence or both. 2
- Bracts not leaf-like or longer than the inflorescence. 3
2. Bracts leaf-like and much longer than the inflorescence; perigynia about 5 times as long as broad. *C. sychnocephala*
- Bracts not leaf-like but much longer than the inflorescence; perigynia about 2–3 times as long as broad. *C. athrostachya*
3. Perigynia subulate, about 4–5 times as long as broad; spikes closely crowded into a head. *C. crawfordii*
- Perigynia lanceolate to ovate, 2–3 times as long as broad; spikes crowded to distant in the head. 4
4. The pistillate scales about as long as the perigynia, completely or almost completely concealing the perigynia. 5
- The pistillate scales shorter and narrower than the perigynia, exposing the beak and margins of the perigynia. 9

5. Beak of the perigynium flat, the teeth spreading, usually serrulate to the tip. 6
 Beak of the perigynium terete, the teeth, if developed, parallel, not serrulate to the tip. 8
6. Inflorescence flexuous, the spikes hardly or not at all overlapping. *C. argyrantha*
 Inflorescence stiffly erect, the spikes crowded or at least well overlapping. 7
7. Pistillate scales and perigynia closely appressed; bracts poorly developed. *C. xerantica*
 Pistillate scales and perigynia ascending, not appressed; lower bracts prominent. *C. adusta*
8. Perigynia 6–8 mm long, prominently veined on both sides. *C. petasata*
 Perigynia 4.5–6 mm long, obscurely veined or veinless. *C. phaeocephala*
9. Perigynia narrowly lanceolate, about 3 times as long as wide. 10
 Perigynia lanceolate to ovoid, 2–2.5 times as long as wide. 11
10. Spikes crowded into a head at the tip of the culm; perigynia 4–7 mm long, veined on both sides; leaves 1–3 mm wide. *C. scoparia*
 Spikes not crowded into a head, at least the lower spikes separated; perigynia 3–5 mm long, prominently veined ventrally, veinless or obscurely veined dorsally; leaves 3–7 mm wide. *C. tribuloides*
11. Perigynia broadly ovate, the body almost orbicular, rather abruptly beaked. *C. straminea*
 Perigynia elliptic to ovoid, not almost orbicular, tapering to the beak. 12
12. Beak of the perigynia terete, the teeth, if developed, parallel. 13
 Beak of the perigynia flattened, the teeth usually well-developed, spreading. 14
13. Spikes, at least the lower ones, well-separated in the head. *C. preslii*
 Spikes crowded in the head, hardly separated from each other. *C. macloviana*
14. Perigynia prominently 3- to 5-veined dorsally; pistillate scales pale green to light brown; inflorescence elongate, flexuous. *C. festucea*
 Perigynia not prominently veined dorsally; pistillate scales reddish to deep brown; inflorescence dense, stiff. *C. tincta*

Section 21. Obtusatae

- Head a solitary spike, staminate at the tip. *C. obtusata*
Head with 2 or more spikes, the upper one
entirely staminate or with a few pistillate
flowers at the base. *C. supina*

Section 22. Montanae

1. Pistillate spikes all on well-developed
culms, 1–4 dm high. 2
Pistillate spikes in part on very short
culms, more or less hidden among the
leaf bases. 4
2. Staminate spike to 20 mm long; pistillate
scales about as long as the perigynia;
perigynia subglobose. *C. pennsylvanica*
Staminate spike to 10 mm long; pistillate
scales much shorter than the perigynia;
perigynia ellipsoid. 3
3. Staminate spikes 5–10 mm long; pistillate
spikes 3 or 4, the lower ones pedun-
cled; perigynia 3–4 mm long; culms
1.5–3 dm high. *C. nigromarginata*
Staminate spikes 2–5 mm long; pistillate
spikes usually 2 or 3, sessile; perigynia
2.5–3 mm long; culms 0.5–1.5 dm high. *C. deflexa*
4. Culms bearing a staminate spike, occa-
sionally with one pistillate spike; bracts
poorly developed, shorter than the
inflorescence. 5
Culms bearing a staminate spike, usually
with 2 or 3 staminate spikes; bracts
well-developed, surpassing the
inflorescence. 6
5. Perigynia pubescent; blades 1–3 mm
wide. *C. umbellata*
Perigynia glabrous; blades 2–5 mm wide. *C. tonsa*
6. Staminate spike 2–5 mm long; perigynia
2.5–3 mm long, with a beak about 0.5
mm long. *C. deflexa*
Staminate spike 3–15 mm long; perigynia
3–3.5 mm long, with a beak about 1
mm long. *C. rossii*

Section 24. Digitatae

1. Lower pistillate spikes long-peduncled;
bracts usually with a short blade; pistil-
late scales short-awned. *C. pedunculata*

- Lower pistillate spikes short-peduncled or sessile; bracts bladeless or almost so; pistillate scales not awned. 2
2. Staminate spike 3–6 mm long, crowded by the pistillate spikes; pistillate scales light brownish. *C. concinna*
- Staminate spike 8–25 mm long, well overtopping the pistillate spikes; pistillate scales reddish or dark brown. 3
3. Lower pistillate spikes short-peduncled, often somewhat remote. *C. richardsonii*
- Lower pistillate spikes almost sessile, approximate and usually overlapping. *C. concinnoides*

Section 25. Rupestres

- Inflorescence a solitary androgynous spike. *C. rupestris*
- Inflorescence consisting of a sessile staminate spike, and one or more pistillate spikes, the lower ones peduncled. *C. glacialis*

Section 27. Albae

- Stigmas 3; achenes triangular in cross section. *C. eburnea*
- Stigmas 2; achenes lenticular in cross section. *C. rufina*

Section 28. Bicolores

- Terminal spike staminate; pistillate scales pale brown to whitish. *C. aurea*
- Terminal spike gynaeandrous; pistillate scales purplish brown to black. *C. bicolor*

Section 29. Paniceae

1. Perigynia beaked, with the beak bidentate, about one-quarter the length of the body; pistillate spikes loosely flowered. *C. vaginata*
- Perigynia beakless, or the beak very short, not bidentate; pistillate spikes rather closely flowered. 2
2. Pistillate spikes sessile or short-peduncled, ascending to erect; leaves whitish glaucous, usually folded or involute. *C. livida*
- Pistillate spikes, at least the lower ones, long-peduncled, ascending to spreading; leaves not very glaucous, flat. *C. tetanica*

Section 31. Granulares

- Staminate spike sessile or short-peduncled, not
overtopping the pistillate spikes. *C. granularis*
Staminate spike long-peduncled, overtopping
the pistillate spikes. *C. crawei*

Section 33. Sylvaticae

- Perigynia pubescent; pistillate spikes very
loosely flowered. *C. assiniboinensis*
Perigynia glabrous; pistillate spikes rather
closely flowered. *C. castanea*

Section 36. Extensae

- Perigynia 3.5–6 mm long, spreading to reflexed
in the spike, with the beak about half to
three-quarters the length of the body. *C. flava*
Perigynia 2–3.5 mm long, spreading to ascend-
ing in the spike, with the beak about one-
quarter to half the length of the body. *C. viridula*

Section 33. Ferrugineae

1. Perigynia beakless or almost beakless;
terminal spike entirely staminate. *C. atrofusca*
Perigynia tapering into a beak; terminal
spike pistillate above. 2
2. Perigynia 3.5–5 mm long, 1 mm wide;
blackish above, greenish or yellowish
below. *C. misandra*
Perigynia 4.5–6 mm long, 1.5–2 mm wide;
yellowish brown. *C. petricosa*

Section 39. Hirtae

1. Perigynia 4–7 mm long, prominently
veined; beak half as long as the body,
its teeth spreading; rhizomes long. *C. houghtonii*
Perigynia 2.5–4 mm long, the veins not
prominent, hidden under the pubes-
cence; beak less than half as long as the
body, its teeth slightly spreading; rhi-
zomes short. 2
2. Leaves to 2 mm wide, involute; perigynia
4–5 mm long; the body oval, somewhat
flattened. *C. lasiocarpa*
Leaves to 4 mm wide, flat; perigynia
2.5–3.5 mm long; the body almost orbi-
cular, terete. *C. lanuginosa*

Section 40. Limosae

1. Pistillate scales long acuminate, to twice as long as but much narrower than the perigynia. *C. magellanica*
 Pistillate scales ovate, about as long and wide as the perigynia. 2
2. Staminate spike short-peduncled; pistillate scales partly clasping and enclosing the perigynia. *C. rariflora*
 Staminate spike long-peduncled; pistillate scales not clasping or enclosing the perigynia. *C. limosa*

Section 41. Atratae

1. Culms arising from long rhizomes, the base without the previous year's dry leaves. *C. buxbaumii*
 Culms arising from a tuft of the previous year's dry leaves, or the lower sheaths not filamentose. 2
2. Lower pistillate spikes long-peduncled, the peduncles as long as or longer than the spikes. 3
 Lower pistillate spikes sessile or short-peduncled, the peduncles shorter than the spikes. 5
3. The terminal spike gynaeandrous. *C. mertensii*
 The terminal spike entirely staminate. 4
4. Perigynia round in cross section, not flattened; achenes not stipitate. *C. raynoldsii*
 Perigynia strongly flattened; achenes stipitate. *C. podocarpa*
5. Perigynia 2–3.5 mm long; pistillate scales 1.5–2.5 mm long. 6
 Perigynia 3.5–4 mm long; pistillate scales about 2.5 mm long. *C. atrata*
6. The terminal spike usually completely pistillate; scales much shorter than the perigynia; rhizomes short. *C. norwegica*
 The terminal spike staminate or gynaeandrous; scales about equaling the perigynia; rhizomes long. *C. parryana*

Section 42. Acutae

1. The terminal spike gynaeandrous; leaf blades 0.5–1.5 mm wide; pistillate scales shorter and narrower than the perigynia. *C. eleusinoides*

- The terminal spikes staminate; leaf blades
2–8 mm wide; pistillate scales about as
long and as wide as the perigynia. 2
2. Perigynia unveined or the veins obscure. 3
Perigynia clearly veined. 6
3. Culms in large, dense clumps; lowest new
leaf sheaths bladeless. *C. stricta*
Culms singly or in small tufts; lowest new
leaf sheaths with blades. 4
4. Staminate spikes usually several; culms in
tufts, together with sterile shoots. *C. aquatilis*
Staminate spike solitary; culms arising
singly or a few together from rhizomes. 5
5. Plants low, 1–4 dm high; pistillate scales
as wide as the perigynia. *C. bigelowii*
Plants taller, 3–10 dm high; pistillate
scales about half as wide as the
perigynia. *C. aperta*
6. Perigynia thick, leathery; beak 0.5–1 mm
long, bidentate. *C. nebraskensis*
Perigynia not thick and leathery; beak 0.1
mm long or less, blunt. 7
7. Perigynia prominently veined, stipitate;
pistillate scales with a green center
about one-fifth the width of the scale. *C. kelloggii*
Perigynia finely veined, hardly stipitate;
pistillate scales with a green center one-
third the width of the scale. *C. lenticularis*

Section 43. Cryptocarpae

1. Pistillate spikes short-peduncled, ascend-
ing to erect. *C. salina*
Pistillate spikes long-peduncled, spread-
ing to pendulous. 2
2. Staminate spikes spreading to pendulous;
pistillate scales tapering to a long awn. *C. paleacea*
Staminate spikes erect; pistillate scales
abruptly contracted to a long awn. *C. crinita*

Section 44. Orthocerates

- Perigynia 6–8 mm long; rachilla rudimentary;
culms with 2 or 3 leaves. *C. pauciflora*
Perigynia 4–5 mm long; rachilla well-devel-
oped; culms with 4–8 leaves. *C. microglochin*

Section 46. Pseudo-Cyperae

- Perigynia spreading or ascending at maturity,
thin-walled, scarcely stipitate. *C. hystricina*
Perigynia spreading to reflexed at maturity,
thick-walled, clearly stipitate. *C. pseudo-cyperus*

Section 47. Paludosae

1. Sheaths and base of leaf blades
pubescent. *C. atherodes*
Sheaths and blades glabrous. 2
2. Teeth of the perigynium 0.5 mm long. *C. lacustris*
Teeth of the perigynium 1–2 mm long. *C. laeviconica*

Section 48. Vesicariae

1. Stigmas 2; achenes lenticular. *C. saxatilis*
Stigmas 3; achenes triangular. 2
2. Pistillate spikes globose to very short
oblong with 3–15 perigynia; beak of
the perigynium short-bidentate or
edentate. *C. oligosperma*
Pistillate spikes oblong to cylindric with
15 to many perigynia; beak of the peri-
gynium usually prominently bidentate. 3
3. The lower perigynia reflexed at maturity;
lower bracts two or more times as long
as the inflorescence. *C. retrorsa*
The lower perigynia not reflexed at ma-
turity, ascending to spreading; lower
bracts slightly exceeding the
inflorescence. 4
4. Culms thick, spongy at base, and bluntly
angled; leaves and sheaths conspicu-
ously cross-veined. *C. rostrata*
Culms slender, not spongy at base, and
sharply angled; leaves and sheaths
cross-veined but not conspicuously so. 5
5. Perigynia 3–3.5 mm long, obscurely
veined; pistillate scales purplish black;
leaves 1–3 mm wide, involute. *C. rotundata*
Perigynia 4–10 mm long, clearly veined;
pistillate scales yellowish to purplish
brown; leaves 3–7 mm wide, flat. *C. vesicaria*

Carex adusta Boott

BROWNE SEDGE

Plants tufted, with culms 20–60 cm high, erect, much taller than the leaves; blades 3–4 mm wide. Inflorescence 2–3 cm long; spikes crowded, subglobose, 8–12 mm long; pistillate scales ovate, hiding the perigynia, brown, with the margin hyaline; perigynia 4–5 mm long, brown, ovate, sharp-edged below the middle. Not common; dry, usually sandy, soil; Prairies and Parklands.

Carex alopecoidea Tuck.

FOXTAIL SEDGE

Plants tufted, with culms 30–70 cm high, erect, soft, mostly as long as or shorter than the leaves; blades 3–4 mm wide. Inflorescence 2–5 cm long, simple or compound, with the upper spikes crowded, the lower ones distant; scales shorter than the perigynia, brown; perigynia 3–4 mm long, tapering into a beak nearly as long as the body, brown. Wet meadows, slough margins, banks of creeks and rivers; Prairies, Parklands, Boreal forest.

Carex aperta Boott

OPEN SEDGE

Plants with stout, woody rhizomes; culms 30–70 cm high, tufted, stiff, and sharply 3-sided; leaves 2–5 mm wide, flat to channeled. Inflorescence 15–20 cm long; terminal spike staminate, 2–3.5 cm long; pistillate spikes 1–5 cm long, usually 2 or 3, erect, sessile or on short peduncles, distant to approximate; scales mostly longer but narrower than the perigynia, reddish black with light midrib; perigynia 2.5–3.5 mm long, greenish to straw-colored. Moist areas; southern Rocky Mountains.

Carex aquatilis Wahl.

WATER SEDGE

Plants with long, scaly rhizomes; culms 10–80 cm high, densely tufted, often in large clumps; blades 3–8 mm wide, flat to channeled, light green or bluish green. Inflorescence 10–20 cm long; terminal spikes staminate, 1.5–3 cm long, usually 1–3; pistillate spikes 2–4 cm long, usually 2–6; scales usually narrower than the perigynia, brown with light green center; perigynia 2–4 mm long, purplish green or green, minutely beaked. Slough margins, marshes, wet meadows; throughout the Prairie Provinces. A rather variable species var. *altior* (Rydb.) Fern. is a larger and coarser plant; var. *stans* (Drej.) Boott, a smaller form. Both grade into the typical form. Water sedge is eaten by cattle, and forms part of slough hay.

Carex arcta Boott

NARROW SEDGE

Plants with short rhizomes; culms 20–40 cm high, densely tufted; blades 2–4 cm wide. Inflorescence 2–4 cm long, with spikes crowded, 5–7 or more; scales somewhat shorter and as wide as perigynia, pale brown; perigynia 2–2.5 mm long, dull brown or greenish brown, white punctulate, with the beak about one-quarter as long as the body. Not common; swamps, wet woods, muskeg; Boreal forest.

Carex argyrantha Tuck.

SILVERY-FLOWERED SEDGE

Plants densely tufted, with culms 30–70 cm high, equaling or exceeding the leaves; blades 3–6 mm wide, flat to somewhat involute. Inflorescence 5–10 cm long, flexuous, often nodding at the tip; spikes 6–12 mm long, with the

upper ones close, the lower distant; scales as long as, but narrower than, the perigynia, shiny greenish or light brown; perigynia 3.5–4.5 mm long, distinctly nerved. Moist grassland, open woods, clearings; throughout the Prairie Provinces. Includes *C. aenea* Fern.

Carex assiniboinensis Boott

ASSINIBOIA SEDGE

Plants loosely tufted, with culms 30–60 cm high, slender, weak; blades 1–3 mm wide. Inflorescence 10–20 cm long; terminal spike staminate; pistillate spikes 2–4 cm long, loosely flowered, spreading or drooping; scales equaling the perigynia in size, long acuminate; perigynia 5–8 mm long, narrowly lanceolate, densely short pubescent, with the beak slender, about as long as the body. Moist open woods; eastern Boreal forest, Parklands.

Carex atherodes Spreng. (Fig. 70)

AWNED SEDGE

Plants with creeping rhizomes; culms 50–120 cm high, loosely tufted, stout, erect; sheaths, especially the lower ones, pubescent; blades 4–12 mm wide, mostly pubescent toward the base; sheaths and blades rarely glabrous. Inflorescence to 25 cm long; terminal 2–6 spikes staminate, 2–4 cm long; pistillate spikes 4–10 cm long, usually 2–4, distant, erect, sessile or short-peduncled; scales shorter than the perigynia, pale brown; the green midrib prolonged into an awn; perigynia 6–10 mm long, prominently nerved, with the beak bearing long, often divergent or recurving teeth. Common; in slough margins, marshes, and wet places; throughout the Prairie Provinces. An important species, palatable to livestock, and a major part of slough hay.

Carex athrostachya Olney

LONG-BRACTED SEDGE

Plants with short rhizomes; erect culms 10–60 cm high, slender, about as tall as, or a little taller than, the leaves; blades 2–5 mm wide, mostly basal; inflorescence 1–2 cm long, and about the same width, with bracts to 6 cm long; spikes 4–10, crowded, gynaeandrous; scales a little shorter than the perigynia, reddish brown with green center; perigynia 3–5 mm long, straw-colored, nerved, the upper half serrulate. Not common; wet meadows; Prairies, Cypress Hills, Rocky Mountains.

Carex atrata L.

Plants with short rhizomes; culms 20–50 cm high, tufted; blades 2–5 mm wide, soft, much shorter than the culm. Inflorescence 3–10 cm long, with bracts leaf-like, the lowest one about equaling the inflorescence; terminal spike gynaeandrous, with the lower 2 or 3 pistillate, 1–3 cm long; at first erect, later spreading or drooping; scales about equaling the perigynia, dark brown to purplish black; perigynia 2.5–4 mm long, greenish to blackish. A large species, in which the following taxa have been recognized:

1. Lower pistillate spikes long-peduncled; perigynia greenish brown. *C. raymondii* Calder
- Lower pistillate spikes sessile or short-peduncled; perigynia greenish to purplish black. 2
2. Perigynia flattened. 3



A.C. Budd

Fig. 70. Awned sedge, *Carex atherodes* Spreng.

Perigynia somewhat inflated, not flattened. *C. atrosquama* Mack.

3. Perigynia granular on the surface, purplish black. *C. albo-nigra* Mack.

Perigynia smooth, yellowish green to brown. *C. epapillosa* Mack.

Of these, *C. raymondii* occurs in meadows throughout Boreal forest; the others, in meadows in the Rocky Mountains.

Carex atrofusca Schk.

Plants with rather stout rhizomes; culms 20–40 cm high, erect; blades 2–4 mm wide, flat. Inflorescence 3–6 cm long, with bracts 1–3 cm long; terminal spikelets entirely staminate, usually 1 or 2, 6–12 mm long, slender-peduncled; pistillate spikes 10–20 mm long, ascending to spreading, or later drooping on slender peduncles; scales shorter and narrower than the perigynia, blackish; perigynia 3–4 mm long, blackish, often with a green margin. Very rare; Churchill.

Carex aurea Nutt.

GOLDEN SEDGE

Plants with slender creeping rhizomes; culms 10–30 cm high, usually spreading or ascending; blades 2–4 mm wide. Inflorescence 5–10 cm long; blades of the lowest bracts exceeding the inflorescence; terminal spikelet gynaeandrous or staminate throughout, 5–15 mm long; pistillate spikes 5–15 mm long, usually 3–5, ascending to spreading on slender peduncles; scales as long as, but narrower than, the perigynia, straw-colored to brownish with green midrib; perigynia 2–3 mm long, elliptic to obovoid, at maturity golden orange. Wet meadows, springy places; throughout Parklands, Cypress Hills, Riding Mountain, Duck Mountain, Rocky Mountains, Boreal forest. Includes *C. garberi* Fern. var. *bifaria* Fern., having deep brown scales with conspicuous green midrib, and prominently veined perigynia.

Carex backii Boott

BACK'S SEDGE

Plants densely cespitose, with culms 5–35 cm high; blades 3–6 mm wide, often exceeding the culms. Inflorescence, without the scales, 1–3 cm long, the staminate portion of the spike 2–4 mm long, inconspicuous; scales foliaceous, to 35 mm long, prominently veined, broadened at the base, long acuminate; perigynia 5–6 mm long, tapering into a stout beak. Dry, open, sandy, or gravelly areas; Parklands, Rocky Mountains.

Carex bicolor Bell.

TWO-COLORED SEDGE

Plants with slender rhizomes; culms 1–12 cm high, very slender, flexuous, spreading; blades 1–3 mm wide, flat to channeled. Inflorescence 2–5 cm long, with the terminal spike gynaeandrous; pistillate spikes 5–10 mm long, usually 1–4, crowded; scales purplish black with green midrib; perigynia 1.5–3 mm long, whitish, beakless. Very rare; in muskeg and marl areas; Churchill.

Carex bigelowii Torr.

STIFF SEDGE

Plants with stout, scaly rhizomes; culms 10–40 cm high, solitary, or few tufted; blades 2–6 mm wide, flat, smooth. Inflorescence 5–15 cm long; the ter-

minal spike staminate, 5–20 mm long; peduncle 5–15 mm long; pistillate spikes 5–35 mm long, usually 2 or 3, erect, the lower ones with peduncles 5–20 mm long; the lowest bract shorter than the inflorescence; scales hiding the perigynia, dark brown, or with a narrow green midrib; perigynia 2–4 mm long, dull green or purple-tinged, occasionally completely purplish. Bogs, marshes, tundra; Boreal forest.

Carex brunnescens (Pers.) Poir.

BROWNISH SEDGE

Plants densely tufted, with culms 20–70 cm high, erect or spreading, much exceeding the leaves; blades 1–3 mm wide, flat or channeled. Inflorescence 3–6 cm long, with 5–8 spikes 4–6 mm long, short ovoid or subglobose, with 5–10 perigynia, the lower ones distant, the upper ones approximate; scales shorter than the perigynia, yellowish or tinged with brown; perigynia 2–2.5 mm long, lightly nerved, punctulate. Bogs, muskeg, and wet woods; Boreal forest.

Carex buxbaumii Wahl.

BROWN SEDGE

Plants with long, slender, brown rhizomes; culms 30–70 cm high, loosely tufted, erect, sharply 3-sided; sheaths reddish brown, becoming filamentose at the base; blades 2–3 mm wide, flat or channeled, gray green or blue green. Inflorescence 3–10 cm long, terminal spike gynaeandrous, 2–3 cm long; pistillate spikes 1–2 cm long, short-peduncled or subsessile; bract of the lowest spike leaf-like, the same length as, or somewhat longer than, the inflorescence; scales 3.5–4 mm long, awn-tipped, reddish brown, with green midrib; perigynia 3–4 mm long, light green, beakless or with two minute teeth. Swamps, wet meadows, and riverbanks; Boreal forest, Rocky Mountains.

Carex capillaris L.

HAIR-LIKE SEDGE

Plants densely tufted, with culms 5–40 cm high; blades 2–4 mm wide, flat, light green. Inflorescence 5–15 cm long; the terminal spike staminate, 4–8 mm long; pistillate spikes 5–15 mm long, usually 2 or 3, drooping on slender peduncles 1–4 cm long; scales 1–2 mm long, hyaline with a green midrib; perigynia 2.5–4 mm long, brown, with two strong veins. Springy areas, marshes, and bogs; Boreal forest. Large plants, 20–40 cm high, with leaves to 20 cm long have been named var. *elongata* Olney; plants 5–20 cm high with leaves 3–10 cm long are var. *capillaris*. However, intermediate sizes also occur.

Carex capitata L.

CAPITATE SEDGE

Plants with thin, hard, ascending rhizomes; culms 10–40 cm high, wiry, stiff; blades mostly less than 1 mm wide, channeled, stiffly erect, often as long as the culm. Inflorescence a single, androgynous, ovoid spike; scales much smaller than the perigynia, center brown with a strong midrib, and a wide hyaline margin; perigynia 2–3 mm long, straw-colored to brown, spreading. Meadows, bogs, open woods; Boreal forest, northern Rocky Mountains.

Carex castanea Wahl.

CHESTNUT SEDGE

Plants tufted, with culms 30–70 cm high, purplish at base, erect; blades 3–6 mm wide, softly pubescent. Inflorescence 4–10 cm long; terminal spike staminate, 1–2 cm long; pistillate spikes 1–2 cm long, usually 3, spreading or

drooping on slender peduncles 1–3 cm long; scales about equaling the perigynium, brownish; perigynia 4–6 mm long, with the beak slender, bidentate, light brown. Very rare; in damp woods; southeastern Boreal forest.

Carex chordorrhiza L. f.

PROSTRATE SEDGE

Plants sod-forming, with culms elongate, prostrate; fertile culms 10–30 cm high, arising from nodes of the old culms; blades 1–2 mm wide, 1–3 on fertile culms, the sterile shoots with several long leaves. Inflorescence with 3–8 spikes crowded in a head 5–15 mm long; spikes androgynous; scales equaling perigynia, straw-colored; perigynia 2.5–3.5 mm long, oblong-ovoid, plump, strongly veined. Bogs, muskeg, and lakeshores; Boreal forest.

Carex concinna R. Br.

BEAUTIFUL SEDGE

Plants with slender rhizomes; culms 10–25 cm high, slender, often recurved or curved ascending, loosely tufted; blades 2–4 mm wide, flat. Inflorescence 1–3 cm long, with the terminal spike staminate, 2–5 mm long, sessile; pistillate spikes 5–10 mm long, approximate, usually 2 or 3; scales about half as long as the perigynia, light to dark reddish brown, with the margins hyaline; perigynia 2.5–3 mm long, thinly pubescent, with a short beak. Damp woods, meadows, and riverbanks; Boreal forest, Rocky Mountains.

Carex concinnoides Mack.

LOW NORTHERN SEDGE

Plants with slender rhizomes; culms 10–35 cm high, slender, erect to curved; blades 3–5 mm wide, flat, stiff. Inflorescence 2–3 cm long, with the terminal spike 15–20 mm long, sessile; pistillate spikes 5–10 mm long, approximate, sessile; scales shorter than the perigynia, the center reddish brown with wide hyaline margins; perigynia 2.5–3 mm long, pubescent, green, with a short beak. In woods, on stony riverbanks; southern Rocky Mountains.

Carex crawei Dewey

CRAWE'S SEDGE

Plants with slender, elongate rhizomes; culms 5–30 cm high, one to several tufted, slender, stiffly erect; blades 2–5 mm wide, stiff. Inflorescence 5–20 cm long, with the terminal spike staminate, 5–20 mm long, on a peduncle 1–7 cm long; pistillate spikes 5–30 mm long, usually 1–4, ascending on peduncles 1–3 cm long, the upper ones often subsessile; scales much shorter than the perigynia, light brown, with a green center; perigynia 2.0–3.5 mm long, light green. Wet meadows, lakeshores, and moist woods; Parklands, Boreal forest.

Carex crawfordii Fern.

CRAWFORD'S SEDGE

Plants densely tufted, with culms 10–70 cm high, mostly barely exceeding the leaves, stiffly erect; blades 1–3 mm wide, erect, flat. Inflorescence 1–3 cm long, with 3–12 spikes, crowded in an oblong or ovoid head; spikes 6–10 mm long, gynaeandrous; scales lanceolate, much shorter than the perigynia, dull brown; perigynia lanceolate, 3–4.5 mm long, 0.7–1 mm wide, long-tapering into the beak. Meadows, swamps, margins of woods; Boreal forest, Rocky Mountains.

Carex crinita Lam.

LONG-HAIRED SEDGE

Plants densely cespitose, with culms 50–80 cm high, exceeding the leaves, 3-sided; blades 6–12 mm wide, flat. Inflorescence 10–20 cm long, with terminal

1–3 spikes staminate, to 5 cm long; pistillate spikes 3–10 cm long, usually 2–5, approximate or somewhat distant, drooping and nodding on slender peduncles; scales shorter and narrower than the perigynia, brownish with green midrib; perigynia 2–4.5 mm long, inflated, green, abruptly short-beaked. Very rare; southeastern Boreal forest.

Carex curta Good.

SHORT SEDGE

Plants with slender rhizomes; culms 20–50 cm high, densely tufted, sharply 3-sided; blades 2–3 mm wide, about as long as the culms, flat, rather stiff, gray green. Inflorescence 3–5 cm long, usually with 4–6 spikes, the upper ones approximate, the lower ones distant; spikes about 5 mm long, ovoid, gynaeandrous, with 10–20 or more flowers; scales shorter than the perigynia, straw-colored, with the midrib light green; perigynia 2–2.5 mm, yellowish green. Muskeg, bogs, and marshes; throughout Boreal forest. Syn.: *C. canescens* L.

Carex deflexa Hornem.

BENT SEDGE

Plants with rather stout horizontal or ascending rhizomes; culms 5–20 cm high, in leafy tufts, purplish at the base; blades 1–3 mm wide, soft, flat. Inflorescence 5–20 mm long, with the terminal spike staminate, 2–5 mm long, often hidden by the 2–4 pistillate spikes, these 5–10 mm long, approximate, the lower ones with a leaf-like bract 5–20 mm long; scales shorter than the perigynia, light brown, with midrib green; perigynia 2–3 mm long, stipitate, green, pubescent. Not common; open forest, clearings; Boreal forest.

Carex deweyana Schw.

DEWEY'S SEDGE

Plants densely tufted, with culms 30–80 cm high, erect to spreading, weak. Sheaths glabrous; blades 2–4 mm wide. Inflorescence 3–8 cm long, with 3–5 spikes, the lower one with a bract to 3 cm long; spikes sessile, the upper ones approximate, the lower ones remote; scales exceeding the perigynia, pale yellowish brown; perigynia 4–5.5 mm long, tapering to a beak more than half as long as the body, rough-margined. Moist or shady woods, margins of woods, meadows; Boreal forest.

Carex diandra Schrank

TWO-STAMENED SEDGE

Plants with short rhizomes; culms 40–80 cm high, in dense tufts, erect. Sheaths scabrous; leaves 2–4 mm wide, erect. Inflorescence 3–5 cm long, with 6–10 spikes, the lower ones somewhat distant, the upper ones approximate to crowded; scales equaling the perigynia, brownish; perigynia 2–3 mm long, shiny dark brown at maturity, tapering into a rough-margined beak, more than half as long as the body. Swamps, wet meadows, and lakeshores; Boreal forest, Cypress Hills, Riding Mountain.

Carex disperma Dewey

TWO-SEEDED SEDGE

Plants with slender, creeping rhizomes, sod-forming, culms 15–50 cm high, very slender, weak, spreading. Sheath smooth; blades 1–2 mm wide, flat, soft. Inflorescence 3–5 cm long, with 2–5 androgynous spikes, 3–6 mm long, distant; scales shorter than the perigynia, white hyaline with green midrib; perigynia 2–3 mm long, minutely beaked. Bogs, wet woods, springy places; throughout the Prairie Provinces.

Plants with slender, long-creeping rhizomes; culms 10–30 cm high, solitary or in small tufts. Sheaths smooth; blades 1–2 mm wide, erect. Inflorescence 2–3 cm long, ovoid, usually dioecious, but occasionally monoecious heads are found; spikes 8–12 mm long; scales as long as or slightly longer than the perigynia, light to dark brown, with green midrib; perigynia 3.5–4 mm long, with the beak more than half as long as the body. Dry grassland, sandhills; Prairies, Parklands.

Carex eburnea Boott

Plants with long, slender rhizomes; culms 10–30 cm high, densely tufted. Sheaths smooth; blades to 0.5 mm wide, flat or involute, erect. Inflorescence 2–6 cm long, with the staminate spike sessile, exceeded by the 2 or 3 long-peduncled pistillate spikes, all 3–6 mm long; scales much shorter than the perigynia, whitish to light brown with green midrib; perigynia 1.5–2 mm long, dark brown when ripe. Open woods and riverbanks; Boreal forest, Parklands.

Carex eleusinoides Turcz.

Plants with slender rhizomes; culms 15–30 cm high. Sheaths smooth; blades 0.5–1.5 mm wide, tufted at the base. Inflorescence 3–6 cm long, crowded, with the terminal spike gynaecandrous; the 2–3 pistillate spikes 7–10 mm long, sessile or subsessile, with the lowermost bract exceeding the inflorescence; scales much shorter and narrower than the perigynia, purplish black; perigynia 3.–3.5 mm long, stipitate, distinctly many nerved, white to light green. Very rare; northern Rocky Mountains.

Carex festucacea Schk.

Plants densely tufted, with culms 30–80 cm high, exceeding the leaves. Sheaths smooth; blades 2–7 mm wide. Inflorescence 1–6 cm long, with 5–10 spikes, 6–10 mm long, crowded or more or less distant below; scales shorter and narrower than the perigynia, tinged with brown, the margins hyaline; perigynia 2.5–4.5 mm long, ovate to obovate. Meadows, open woods, clearing, lakeshores; in Boreal forest, Parklands. A large species, including the following small species:

- 1. Perigynia distinctly veined; inflorescence elongate, flexuous. *C. tenera* Dewey
 Perigynia finely veined; inflorescence compact. 2
- 2. Perigynia 2.5–3.5 mm long; scales pale to dull brown; sheaths tight; blades 2–4 mm wide. *C. bebbii* Olney
 Perigynia 3–4.5 mm long; scales hyaline to light brown; sheaths loose; blades to 7 mm wide. *C. normalis* Mack.

Carex filifolia Nutt. (Fig. 71)

Plants densely tufted, with hard bases; culms 8–30 cm high, stiff, wiry. Sheaths smooth, reddish; blades 0.25 mm wide, needle-like, stiff. Inflorescence




Fig. 71. Thread-leaved sedge, *Carex filifolia* Nutt.

a solitary, androgynous spike, 1–3 cm long, erect; scales hiding the perigynia, the center dark brown, the broad margins hyaline; perigynia 3–4 mm long, straw-colored to light brown, puberulent. Dry grassland, eroded slopes, and hills; Prairies, Parklands. Palatable to livestock, and of moderate forage value.

Carex flava L.

YELLOW SEDGE

Plants with short rhizomes, sod-forming; culms 30–70 cm high, sharply 3-sided, stiff, as long as or slightly exceeding the leaves. Sheaths smooth, straw-colored to light brown; blades 4–5 mm wide, flat, lax. Inflorescence 3–6 cm long, with the terminal spike staminate, 5–20 mm long; pistillate spikes 2–4, 5–15 mm long, distant or approximate; bracts leaf-like, the lowest one exceeding the inflorescence; scales shorter than the perigynia, reddish brown with green midrib; perigynia 4.5–7 mm long, with the beak about as long as the body, yellow. Not common; wet meadows, riverbanks, and swamps; Boreal forest.

Carex geyeri Boott

GEYER'S SEDGE

Plants with thick, short, scaly rhizomes, to 3 mm thick; culms 10–40 cm high, rough. Sheaths loose; blades 1–3 mm wide, the lower ones short. Inflorescence a solitary androgynous spike; scales hiding the perigynia, brownish with hyaline margins; perigynia 4–6 mm long, light brown, beakless or nearly so. Dry woods, slopes; Rocky Mountains.

Carex glacialis Mack.

GLACIER SEDGE

Plants densely tufted, with erect culms 5–15 cm high. Sheaths scabrous; blades 1–1.5 mm wide, channeled, recurved. Inflorescence 1–1.5 cm long, with the terminal spike staminate, 2–6 mm long; 1–3 pistillate spikes, 2–6 mm long, the lowest short-peduncled, the upper sessile; scales slightly shorter than the perigynia, brownish to purplish black, with hyaline margins; perigynia 2–2.5 mm long, light to dark brown, with a hyaline beak. Very rare; barren tundra; Boreal forest.

Carex gracillima Schw. (Fig. 72)

SLENDER SEDGE

Plants tufted, with culms 40–80 cm high, slender, erect, purplish brown at the base. Sheaths smooth, tight; blades 3–10 mm wide, flat, thin. Inflorescence to 30 cm long, with the terminal spike gynaeandrous, 2–3 cm long; pistillate spikes 3–5 cm long, usually 3–5, long-peduncled, remote, the lower ones separated by as much as 15 cm, spreading or drooping; scales half to nearly as long as the perigynia, hyaline with green midrib; perigynia 2.5–4 mm long, distinctly nerved, green. Uncommon; moist woods; southeastern Parklands, Boreal forest.

Carex granularis Muhl.

GRANULAR SEDGE

Plants with short rhizomes; culms 30–80 cm high, tufted. Sheaths loose, glabrous; blades 4–10 mm wide, flat. Inflorescence 5–15 cm long, with the terminal spike staminate, often exceeded by the pistillate spikes, 1–2 cm long; pistillate spikes 15–20 mm long, usually 2–4, the upper ones subsessile, the lower ones progressively longer peduncled; bracts leaf-like, the upper ones exceeding the spikes; scales shorter than the perigynia, hyaline with a green



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

WRITES: *Carex gracillima* Schw.

LOCATION: Long Hill Community Pasture

COLL: J. Looman

DET:

FILE NO: 3104

DATE: 11 June 1958

ELEVATION: 450 m

ENVIRONMENT: Deep ravine,
open woods




Fig. 72. Slender sedge, *Carex gracillima* Schw.

midrib; perigynia 2.5–4 mm long, green, strongly ribbed, with a short bent beak. Damp meadows, springy areas; southeastern Parklands, Boreal forest.

Carex gravida Bailey

HEAVY SEDGE

Plants densely tufted, with culms 30–60 cm high, scabrous at the summit. Sheaths scabrous; blades 4–8 mm wide. Inflorescence 1–3 cm long, dense, oblong-ovoid, with 4–7 spikes; bracts shorter than the head; scales slightly shorter than the perigynia, brown; perigynia 4–5 mm long, yellowish brown, with the beak about one-third as long as the body. Rare; dry grasslands; Prairies.

Carex gynocrates Wormsk. (Fig. 73)

NORTHERN BOG SEDGE

Plants with slender, creeping rhizomes; culms 10–30 cm high, erect, solitary or a few tufted. Sheaths smooth; blades 0.5 mm wide, involute. Inflorescence a single spike, usually either staminate or pistillate, occasionally androgynous; scales shorter than the perigynia, brown, with hyaline margins and midrib; perigynia 3–3.5 mm long. Bogs, marshes, and muskeg; Boreal forest.

Carex heleonastes L. f.

HUDSON BAY SEDGE

Plants with short rhizomes; culms 10–30 cm high, tufted, slender. Sheaths smooth; blades 2 mm wide, flat, with the margins revolute. Inflorescence 3–5 cm long, with 1–4 spikes, approximate; spikes gynaeandrous, 3–5 mm long; scales shorter than the perigynia, with the center yellowish brown, the margins hyaline; perigynia 2–3.5 mm long, greenish, white punctate. Muskeg, meadows, and swamps; Boreal forest.

Carex hoodii Boott

HOOD'S SEDGE

Plants with short, stout rhizomes; culms 20–80 cm high, slender, sharply 3-sided. Sheaths glabrous, tight; blades 1.5–4 mm wide, flat or channeled. Inflorescence 1–2 cm long, ovoid to orbicular, with 4–8 spikes, crowded; spikes androgynous; scales hiding the perigynia or nearly so, shiny brown, with green midrib and hyaline margins; perigynia 3.5–5 mm long, broadly winged, with the beak about one-third as long as the body. Meadows and slopes; Rocky Mountains.

Carex hookerana Dewey

HOOKER'S SEDGE

Plants with short rhizomes; culms 10–40 cm high, erect, slender. Sheaths tight, glabrous; leaves 1–2.5 mm wide, flat, thin. Inflorescence 2–5 cm long, with 5–8 spikes, distant to approximate; spikes androgynous, each with a short bract; scales hiding the perigynia, pale brown, distinctly awned; perigynia 2.5–3.5 mm long, brownish. Rare; dry grasslands, openings, and clearings; Prairies, Parklands.

Carex houghtonii Torr.

SAND SEDGE

Plants with long-creeping, thick rhizomes; culms 30–60 cm high, loosely tufted. Sheaths glabrous, loose; blades 4–10 mm wide. Inflorescence 5–15 cm long; staminate spike terminal, usually solitary, occasionally with 1 or 2 small staminate spikes or 1 or 2 perigynia at the base; pistillate spikes 1–3, 1–4 cm



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Carex gynocrates* Wormsk.

LOCATION: 4 mi south of Descharme Creek

COLL: J. Looman

DET:

COLL. NO: 15937

DATE: 26 June 1971

ELEVATION: 625 m

HABITAT: Muskeg



Fig. 73. Northern bog sedge, *Carex gynocrates* Wormsk.

long, remote, sessile or short-peduncled, erect; perigynia 5–7 mm long, strongly many nerved, finely pubescent, with the beak about 2 mm long, bidentate. Sandy or rocky soils and open forests; Boreal forest.

Carex hystricina Muhl.

PORCUPINE SEDGE

Plants densely tufted, with erect culms 20–70 cm high. Sheaths smooth, loose; blades 2–10 mm wide; sheaths and blades often conspicuously cross-veined. Inflorescence 5–15 cm long, with the terminal spike staminate, 2–4 cm long, occasionally gynaecandrous; 1–4 pistillate spikes, 10–40 mm long, the upper ones approximate, the lowest one often 6 or 7 cm distant, peduncled, erect to spreading; perigynia 5–8 mm long, substipitate, ascending to spreading above, somewhat reflexed below, with the beak 2–2.5 mm long, slender, bidentate. Not common; swamps and wet meadows; Boreal forest.

Carex intumescens Rudge

SWOLLEN SEDGE

Plants in large tufts, with culms 30–70 cm high, erect to spreading, sharply triangular. Sheaths smooth; blades 3–8 mm wide. Inflorescence 3–5 cm long, with the terminal spike staminate, 15–25 mm long; pistillate spikes 1–3, globose, 10–20 mm in diam, crowded; perigynia 10–15 mm long, tapering into the bidentate beak. Rare; damp woods; southeastern Boreal forest.

Carex kelloggii Boott

KELLOGG'S SEDGE

Plants with slender, short rhizomes; culms 10–60 cm high, more or less densely tufted. Sheaths smooth, brownish at the base; blades 1–3 mm wide, equaling to exceeding the culms. Inflorescence 5–10 cm long, with the terminal spike staminate, 20–30 mm long, occasionally gynaecandrous; 3–5 pistillate spikes, 1–7 cm long, the upper ones sessile or subsessile, the lower ones short-peduncled; perigynia 1.5–3 mm long, flattened, light green; scales smaller than the perigynia, with the center dark reddish brown. Moist, springy areas; Rocky Mountains.

Carex lacustris Willd.

LAKESHORE SEDGE

Plants with long rhizomes; culms tufted, 50–125 cm high, stout, sharply triangular, rough on the edges. Sheaths reddish below, with the inner membrane disintegrating to the appearance of a ladder; blades 6–15 mm wide; sheaths and blades usually cross-veined. Inflorescence 15–35 cm long, with the terminal 2–4 spikes staminate, 1–8 cm long; pistillate spikes 2–4, usually distant, 3–10 cm long, erect, sessile or short-peduncled; perigynia 5–8 mm long, dull green, tapering into the beak. Occasionally staminate and pistillate spikes are compound, with 3–5 flowers on the rachillae. Marshes, swamps, and lake-shores; Boreal forest.

Carex laeviconica Dewey

SMOOTH-FRUITED SEDGE

Plants with rhizomes; culms 50–70 cm high, stout, loosely tufted. Sheaths glabrous, with the inner membrane disintegrating and laddering; blades 3–6 mm wide, flat. Inflorescence to 25 cm long, with the upper 2–6 spikes staminate, 2–10 cm long; pistillate spikes 2–4, distant, 3–7 cm long, erect, sessile or short-peduncled. Perigynia 5–8 mm long, strongly many nerved, tapering into the beak, with the teeth 1–2.5 mm long, straight. Swampy areas, slough margins; Prairies, Parklands.

Carex lanuginosa Michx.

WOOLLY SEDGE

Plants with slender, long-creeping rhizomes; culms 30–70 cm high, usually in small tufts, sharply triangular. Sheaths glabrous, reddish at the base, with the inner membrane laddering; blades 2–5 mm wide, flat or involute, often exceeding the culms. Inflorescence 5–20 cm long, with the terminal 1 or 2 spikes staminate, 1–5 cm long; pistillate spikes 1–3, distant, 15–40 mm long, erect, sessile or the lower spikes short-peduncled. Perigynia 2.5–3.5 mm long, densely pubescent, abruptly beaked, with the teeth about 0.5 mm long, erect. Slough margins, marshes, and wet places; throughout the Prairie Provinces. Syn.: *C. lasiocarpa* Ehrh. var. *latifolia* (Böck.) Gl.

Carex lasiocarpa Ehrh.

HAIRY-FRUITED SEDGE

Plants with stout long-creeping rhizomes, 2–3 mm thick; culms 50–120 cm high, tufted, bluntly triangular, scabrous above. Sheaths glabrous, loose, cross-veined, with the inner membrane laddering; blades 1–2 mm wide, folded or convolute, those of sterile shoots exceeding the flowering culms. Inflorescence 6–35 cm long, with the terminal 1–3 spikes staminate, 1–7 cm long, occasionally with a few perigynia at the base; pistillate spikes 2 or 3, distant, 1–3 cm long, often with a few staminate flowers at the tip, sessile or short-peduncled; perigynia 4–5 mm long, densely pubescent, tapering into the beak, with the teeth about 1 mm long. Bogs, lakeshores, and riverbanks; Boreal forest.

Carex laxiflora Lam. var. *varians* Bailey

PLEASING SEDGE

Plants tufted, with culms 20–50 cm high, somewhat winged. Sheaths glabrous, loose, expanded at the throat; blades 3–7 mm wide, flat. Inflorescence 5–15 cm long, with the terminal spike staminate, 1–2 cm long; pistillate spikes 2–4, the upper ones approximate to the pistillate spike, the lower distant by up to 10 cm; perigynia 2.5–4 mm long, nerveless to obscurely or distinctly 10- to 12-nerved, with the body abruptly contracted into an oblique short beak; scales hyaline with a green midrib, obtuse or with a short awn. Apparently very rare; moist woods; Boreal forest.

Carex lenticularis Michx.

LENS-FRUITED SEDGE

Plants with short rhizomes; culms 20–60 cm high, densely tufted. Sheaths glabrous, brown at the base; blades 2–3 mm wide, as long as or exceeding the culms. Inflorescence 5–12 cm long, with the terminal spike staminate, 1–3 cm long, occasionally gynaeandrous; pistillate spikes 3–5, crowded, 15–40 mm long; perigynia 2–3 mm long, glaucous green; scales smaller than the perigynia, with the midrib green, the margins purple brown. Rare; riverbanks, lakeshores, and marshes; Boreal forest, Rocky Mountains.

Carex leptalea Wahl.

BRISTLE-STALKED SEDGE

Plants with slender creeping rhizomes; culms 10–40 cm high, very slender, densely tufted. Sheaths smooth, tight; blades about 0.5 mm wide, shorter than to as long as the culms. Inflorescence a single spike, 5–15 cm long, androgynous; staminate scales blunt, with the margins overlapping around the rachis; perigynia 2.5–6 mm long, green, finely many nerved. Bogs and marshes; Boreal forests, Parklands.

Plants with stout creeping rhizomes, to 3 mm thick; rootlets clothed with a yellow to gray or brownish felt-like layer; culms 30–50 cm high, sharply 3-angled. Sheaths glabrous, fibrillose; blades 1–1.5 mm wide, stiff, channeled or folded. Inflorescence 3–6 cm long, with the terminal spikelet staminate, 1–2 cm long, erect; pistillate spikes 1 or 2, 10–15 mm long, on filiform pedicels, 10–25 mm long, spreading to drooping; perigynia 2.5–4 mm long, grayish green, distinctly 8- to 10-nerved; scales about as long as the perigynia, light or dark brown, with green midrib. Not common; in bogs, marshes, and muskeg; Boreal forest.

Carex livida (Wahl.) Willd.

Plants with slender rhizomes; culms 20–50 cm high, solitary. Sheaths glabrous; blades 1–3 mm wide, equaling or surpassing the leaves, strongly glaucous. Inflorescence 3–6 cm long, with the terminal spike staminate, rarely gynaeandrous 10–30 mm long; pistillate spikes 1–3, 5–15 mm long, sessile or short-peduncled; perigynium 3–5 mm long, glaucous green, finely many nerved; scales purplish with green midrib. Rare; bogs, marshes, and muskeg; Boreal forest.

Carex loliacea L.

Plants with slender rhizomes; culms 20–60 cm high, weak, tufted, bluntly 3-angled. Sheaths glabrous, tight; blades 1–2 mm wide, flat or channeled, shorter than to as long as the culms. Inflorescence 2–3 cm long, with 2–6 gynaeandrous spikelets, 3–5 mm long, at least the lower ones distant; perigynia 2–3 mm long, distinctly many veined; scales white hyaline, with midrib green, smaller than the perigynia. Bogs, wet places; Boreal forests.

Carex mackenziei Krecz.

Plants with slender rhizomes; erect culms 15–40 cm high. Sheaths scabrous; blades 1–3 mm wide. Inflorescence 2–5 cm long; spikes 3–6, about 1 cm long, distant, with the terminal spike gynaeandrous, about one-half staminate; perigynia 2.5–3.5 mm long, stipitate; scales as long as the perigynia, reddish brown. Brackish marshes; Hudson Bay.

Carex macloviana d’Urv.

Plants more or less densely tufted from short rhizomes, with erect culms 10–40 cm high. Sheaths glabrous, open; blades 2–6 mm wide. Inflorescence 5–20 mm long, with 3–8 spikes crowded into a head; perigynia 3.5–4.5 mm long, brown, finely nerved; scales shorter than the perigynia. Moist areas; throughout the Prairie Provinces. A large species in which several small species have been distinguished.

- 1. Perigynia much flattened, thin, scale-like
 except over the achene. 2
- Perigynia not very thin and scale-like,
 with the front convex, the back flat. 6
- 2. Scales and perigynia copper brown; peri-
 gynia obscurely veined. *C. macloviana*

- Scales and perigynia light, chestnut, or dull brown. 3
3. Perigynia veinless on the front. 4
 Perigynia veined on the front, at least toward the base. 5
4. Perigynia 4.5–6 mm long, light brown or greenish brown. *C. haydeniana* Olney
 Perigynia 2.5–3.5 mm long, olive brown to blackish brown. *C. limnophila* Hermann
5. Spikes distinguishable in the head; perigynia appressed, with the tip inconspicuous, margins rather wide. *C. festivella* Mack.
 Spikes hardly distinguishable in the head; perigynia spreading to ascending, with the tip conspicuous, margins narrow. *C. microptera* Mack.
6. Scales with distinct hyaline margins. 7
 Scales without distinct hyaline margins. *C. pachystachya* Cham.
7. Perigynia 3–3.5 mm long, with corky tissue around the achene. *C. incondita* Hermann
 Perigynia 3.5–4.5 mm long, no corky tissue. *C. macloviana*

Carex magellanica Lam. (Fig. 74)

BOG SEDGE

Plants with long slender rhizomes; the rootlets covered with a yellow, grayish, or brownish felt-like layer; culms 30–60 cm high, very slender. Sheaths glabrous, mostly brownish at the base; blades 2–4 mm wide, flat. Inflorescence 5–12 cm long, with the terminal spike staminate, 5–15 mm long; pistillate spikes 2–4, the lower ones distant, the upper approximate, 8–20 mm long, on slender peduncles 1–4 cm long, spreading to drooping; perigynia 3–3.5 mm long, glaucous green to dull brown, veinless to finely veined; scales equaling or exceeding, but narrower than, the perigynia, brown with green midrib, or all brown. Bogs and muskeg; Boreal forests. Syn.: *C. paupercula* Michx.

Carex maritima Gunn.

SEASIDE SEDGE

Plants with extensively creeping, tough, slender rhizomes; culms 5–15 cm high, mostly solitary, often curved at the summit. Sheaths glabrous, loose; blades 1–2 mm wide, flat or involute, often exceeding the culms, curved or curled. Inflorescence 5–20 mm long, with spikes 3–5, densely packed in a head, androgynous; perigynia 3.5–5 mm long, greenish to golden brown, finely veined or veinless; scales smaller than the perigynia, brown. Rare; gravelly or rocky areas; Boreal forest, Rocky Mountains.

Carex mertensii Prescott

PURPLE SEDGE

Plants with short, stout rhizomes; culms 30–100 cm high, densely tufted, sharply 3-angled, winged, erect. Sheaths glabrous, loose, purplish red below; blades 4–8 mm wide, flat or the margins revolute. Inflorescence 5–12 cm long, with the terminal spike staminate or gynaeandrous, lateral spikes gynaeand-




Fig. 74. Bog sedge, *Carex magellanica* Lam.

rous, with a few staminate flowers, 1–4 cm long, drooping on slender peduncles; perigynia 4.5–5 mm long, light green to pale brown, often purplish spotted; scales purplish brown, with light midrib and hyaline margins, smaller than the perigynia. Open slopes and forests; Rocky Mountains.

Carex michauxiana Böck.

LONG-FRUITED SEDGE

Plants rather densely tufted, with erect culms 20–60 cm high. Sheaths glabrous, loose; blades 2–4 mm wide. Inflorescence 2–10 cm long, with the terminal spike staminate, 5–15 mm long, often hidden by the upper pistillate spike; pistillate spikes 2–4, ovoid to suborbicular, 15–25 mm long, the lower ones distant; bracts exceeding the culms; perigynia 8–12 mm long, very slender, sharply veined; scales much smaller than the perigynia, brown. Bogs and wet meadows; southeastern Boreal forest.

Carex microglochin Wahl.

SHORT-AWNED SEDGE

Plants with slender creeping rhizomes; culms 8–25 cm high, erect, tufted. Sheaths glabrous, tight, brown below; blades about 0.5 mm wide, involute, stiffly erect. Inflorescence a solitary spike, 5–15 mm long, androgynous; perigynia 3–5 mm long, linear-lanceolate, light green to brownish, at first ascending, later spreading to reflexed; scales wider but shorter than the perigynia, early deciduous. Rare; alpine meadows; Rocky Mountains.

Carex misandra R. Br.

NODDING SEDGE

Plants with short rhizomes; culms 10–30 cm high, densely tufted, slender, erect. Sheaths glabrous, brownish below; blades 1–3 mm wide, densely clustered at base of culms. Inflorescence 3–6 cm long, nodding, with the terminal spike gynaecandrous, 5–10 mm long; pistillate spikes 2 or 3, approximate, 5–20 mm long, erect to nodding on slender peduncles; perigynia 3.5–5 mm long, purplish black above, straw-colored below; narrowly lanceolate; scales wider but shorter than the perigynia, blackish with hyaline margin and apex. Rare; tundra and alpine meadows; Boreal forest, Rocky Mountains.

Carex muricata L.

INLAND SEDGE

Plants densely tufted, with culms 20–70 cm high, slender, erect. Sheaths glabrous, tight; blades 1–4 mm wide. Inflorescence 3–5 cm long, with 3–7 spikes in an interrupted oblong head; the terminal spike gynaecandrous, lateral spikes pistillate or gynaecandrous; perigynia 2.5–5.0 mm long, tapering into a more or less distinctly bidentate beak; scales shorter than to about as long as the perigynium, pale brown with green midrib. Moist woods, meadows, and bogs; Boreal forest, Rocky Mountains. Includes *C. interior* Bailey and *C. angustior* Mack.

Carex nardina Fries

FRAGRANT SEDGE

Plants densely tufted, with culms 30–50 cm high, slender, erect. Sheaths smooth, tight, with the old sheaths persisting; blades 0.5–1 mm wide, stiff, shorter to longer than the culms. Inflorescence a single androgynous spike, 5–15 mm long; perigynia 3–4.5 mm long, striate, short-stipitate; scales about equaling the perigynia, brown, with greenish or straw-colored center. Not common; dry slopes, rocky areas; Rocky Mountains. Plants with culms mostly 15–30 cm high have been distinguished as var. *hepburnii* (Boott) Kük.

Plants with long creeping rhizomes; culms 20–100 cm high, tufted, coarse. Sheaths smooth; blades 3–8 mm wide, flat, usually cross-veined, glaucous green. Inflorescence 5–10 cm long, the upper 1 or 2 spikes staminate, 15–40 mm long; pistillate spikes 2–5, sessile or nearly so, 1–6 cm long, approximate; lowest bract commonly exceeding the inflorescence; perigynia 3–3.5 mm long, straw-colored, red-dotted, strongly veined; scales much smaller than the perigynia. Very rare; marshy ground; Prairies (Alberta).

Carex nigricans C. A. Mey.

BLACKENING SEDGE

Plants with stout, long-creeping rhizomes; culms 5–20 cm high, stiff. Sheaths smooth, the old ones persisting; blades 1–2.5 mm wide, flat or channeled. Inflorescence a single androgynous spike, 8–15 mm long; perigynia 3.5–4.5 mm long, at first appressed, later spreading to reflexed; stipitate, veinless; scales much shorter than the perigynia, dark brown. Exposed alpine meadows; Rocky Mountains.

Carex nigromarginata Schw. var. *elliptica* (Boott) Gl. BLACK-MARGINED SEDGE

Plants with short rhizomes; culms 5–50 cm high, tufted, very slender. Sheaths glabrous, tight; blades 1.5–3 mm wide, flat, soft, mostly shorter than the culms. Inflorescence 1–3 cm long, with the spikes approximate or crowded; terminal spike staminate, 5–10 mm long; pistillate spikes 1–4, with the lowest bract often exceeding the inflorescence; perigynia 3–4 mm long, pubescent, with the beak 0.5–1 mm, distinctly bidentate; scales much smaller than the perigynia, reddish brown with hyaline margins. Not common; open woods, margins of woods, and riverbanks; Parklands, Boreal forest.

Carex norvegica Retz.

NORWAY SEDGE

Plants with short, stout rhizomes; culms 20–60 cm high, erect or arched, loosely tufted. Sheaths glabrous; blades 1–3 mm wide, soft, flat. Inflorescence 10–35 mm long, usually with 3 spikes, crowded or the lower ones approximate, with the terminal spike pistillate or gynaeandrous, the lateral ones pistillate, 1–3 cm long, sessile or nearly so; perigynia 2–3.5 mm long, glaucous green, finely veined; scales deep purplish brown with white hyaline margins. Moist woods, lakeshores, and swamps; Boreal forest, Rocky Mountains.

Carex obtusata Lilj.

BLUNT SEDGE

Plants with slender, long-creeping, purplish black rhizomes; culms 6–20 cm high, slender, solitary or few together. Sheaths glabrous, usually reddish brown below; blades 1–1.5 mm wide, involute or flat. Inflorescence a single androgynous spike, 5–12 mm long, with 1–6 perigynia, 3–4 mm long, dark chestnut or blackish brown at maturity; scales smaller than the perigynia, pale brown. Locally common; dry to moist grassland; Prairies, Parklands.

Carex oligosperma Michx.

FEW-FRUITED SEDGE

Plants with long creeping rhizomes; culms 40–100 cm high, slender, stiffly erect. Sheaths glabrous, loose; blades 1–2 mm wide, involute, stiff, to 80 cm long. Inflorescence 6–10 cm long; the lowest bract exceeding the inflorescence; the terminal spike staminate, 1–5 cm long; the lateral spikes 1–3, pistillate, 1–2

cm long, ovoid or subcylindric, with 2–15 perigynia, 4–7 mm long, with the beak 1–2 mm long, bearing short teeth; scales smaller than the perigynia, pale brown to hyaline. Not common; bogs and sedge meadows; Boreal forest.

Carex paleacea Wahl.

CHAFFY SEDGE

Plants with long stolons; culms 10–60 cm high, forming small tussocks. Sheaths glabrous; blades 3–8 mm wide, flat. Inflorescence 6–15 cm long, the terminal spikes 1–3 staminate, 2–5 cm long; the lateral spikes 2–4, pistillate or gynaeandrous, 2–5 cm long, drooping on slender peduncles; perigynia 2.5–3 mm long, glaucous green, short-beaked; scales shorter than the perigynia, brown, with the pale midrib prolonged into an awn, up to 1 cm long. Very rare; salt marshes; Boreal forest.

Carex parryana Dewey

PARRY'S SEDGE

Plants with slender, long creeping rhizomes; culms 15–40 cm high, erect. Sheaths glabrous, brown to reddish below; blades 2–3 mm wide, flat or the margins revolute, erect. Inflorescence 4–8 cm long; the terminal spike gynaeandrous or staminate; the lateral spikes 2–4, pistillate, 1–3 cm long, approximate to crowded; perigynia 1.5–2.5 mm long, straw-colored or greenish, punctulate; scales equaling the perigynia, dark reddish brown, with green midrib. Rare; moist grasslands; Prairies, Parklands.

Carex pauciflora Lightf.

FEW-FLOWERED SEDGE

Plants with long, slender rhizomes; culms 10–40 cm high, few together or solitary. Sheaths glabrous; blades 1–2 mm wide, stiff, involute, shorter than the culm. Inflorescence a single androgynous spike, to 1 cm long; perigynia 6–7 mm long, narrowly lanceolate, long-tapering into a beak, spreading or reflexed; scales smaller than the perigynia, pale brown. Bogs and muskeg; Boreal forest.

Carex pedunculata Muhl.

STALKED SEDGE

Plants with short, thick rhizomes; culms 10–30 cm high, often barely exceeding the leaves. Sheaths glabrous, the lowest ones with a very short, stiff leaf, brownish; blades 2–4 mm wide, flat, thick. Inflorescence 4–10 cm long, with the terminal spike androgynous or staminate, 5–15 mm long; the lateral spikes 2 or 3, pistillate or sometimes androgynous, 1–3 cm long, with the upper ones ascending on short pedicels, the lower ones long-pedicel, spreading or ascending; perigynia 3.5–5 mm long, thinly pubescent, sharply angled, pale green; scales smaller than the perigynia, purplish brown, with the green midrib prolonged into a short awn. Rich woods; southeastern Boreal forest.

Carex pensylvanica Lam.

SUN-LOVING SEDGE

Plants with extensively creeping slender rhizomes; culms 10–30 cm high, usually exceeding the leaves, tufted. Sheaths glabrous, disintegrating; blades 1–3 mm wide, flat, erect. Inflorescence 15–50 mm long, with the terminal spike staminate, the lateral spikes 2 or 3, pistillate, 5–10 mm long, sessile or short-peduncled; perigynia 2–4 mm long, finely pubescent, abruptly narrowed to a bidentate beak, stipitate. Dry to moist grassland, open woods, and thickets;

throughout Prairies and Parkland. A variable species, the following varieties are present:

1. Perigynia 3–3.5 mm long, round; leaves stiff and scabrous. var. *digyna* Böck.
Perigynia 2–4 mm long, angled; leaves soft, not scabrous. 2
2. Perigynia 2–3 mm long, with the beak about 0.5 mm long. var. *pensylvanica*
Perigynia 3–4 long, with the beak more than 1 mm long. var. *distans* Peck

The var. *pensylvanica* occurs in southeastern Parklands and in a few locations in Boreal forest; var. *digyna* Böck. (*C. heliophila* Mack.) is the common form in grassland of Prairies and Parklands; var. *distans* Peck occurs occasionally in openings in Boreal forest.

Carex petasata Dewey

PASTURE SEDGE

Plants with short rhizomes; culms 20–80 cm high, densely tufted, smooth, mostly nodding at the top. Sheaths glabrous, tight; blades 2–5 mm wide, flat. Inflorescence 2–5 cm long, with 3–6 gynaeandrous spikes, approximate; spikes about 1 cm long; perigynia 4–8 mm long, oblong to lanceolate; scales equaling the perigynia, reddish brown with white hyaline margins. Meadows, open woods, and clearings; Parklands, Boreal forest. A large species, including the following:

1. Perigynia 4–4.5 mm long; blades 3–5 mm wide. *C. platylepis* Mack.
Perigynia 4.5–8 mm long; blades 2–3 mm wide. 2
2. Perigynia 6–8 mm long, prominently veined. *C. petasata*
Perigynia 4.5–6 mm long, obscurely veined. *C. praticola* Rydb.

Carex petricosa Dewey

STONE SEDGE

Plants with slender, long rhizomes; erect culms 10–30 cm high. Sheaths glabrous; blades 1–3 mm wide, stiff. Inflorescence 4–8 cm long, with 3–8 spikes; uppermost spikes androgynous or staminate, 7–20 mm long; lateral spikes 2 or 3, pistillate or androgynous, 10–30 mm long; perigynia 4–6 mm long, ciliate on the margins, yellowish brown to dark brown; scales shorter than the perigynia, reddish brown. Riverbanks, slopes, and alpine tundra; Rocky Mountains. Plants of the lower altitudes are usually larger, to 90 cm high, and have broad yellowish brown perigynia, 2–2.5 mm wide. These have been distinguished as var. *franklinii* (Boott) Boiv.

Carex phaeocephala Piper

HEAD-LIKE SEDGE

Plants with short, matted rhizomes, forming large clumps; culms 10–30 cm high, slender, scabrous above, stiff. Sheaths smooth, tight; blades 3–6 mm

wide, mostly basal. Inflorescence 10–25 mm long, with 2–5 gynaeandrous spikes, approximate to crowded; perigynia 4–6 mm long, oblong-ovate, straw-colored to dark brown; scales equaling the perigynia, dark brown with a light center and hyaline margins. Slopes and alpine meadows; Rocky Mountains.

Carex podocarpa R. Br.

ALPINE SEDGE

Plants with slender creeping rhizomes; culms 10–50 cm high, loosely tufted, bluntly triangular, more or less nodding above. Sheaths glabrous, reddish brown below; blades 2–4 mm wide, flat, deep green. Inflorescence 4–7 cm long, with the terminal spike staminate, 5–20 mm long; lateral spikes 2–4, pistillate or somewhat androgynous, 1–2 cm long, approximate, short pedunculate, ascending or spreading; perigynia 3.5–4.5 mm long, blackish brown; scales much smaller than the perigynia, brownish black. Alpine meadows, riverbanks; Rocky Mountains.

Carex praegracilis W. Boott

GRACEFUL SEDGE

Plants with blackish, long-creeping rhizomes; culms 15–60 cm high, sharply triangular, scabrous above. Sheaths scabrous, loose; blades 1–3 mm wide, flat or channeled, mostly basal. Inflorescence 1–5 cm long, with 5–10 androgynous spikes crowded into a dense head, or the lower ones approximate; spikes 5–10 mm long; perigynia 3–4 mm long, dull brownish black; scales larger than the perigynia, with the center chestnut brown, the margins hyaline. In marshes, around sloughs, wet meadows; throughout the Prairie Provinces.

Carex prairea Dewey

PRAIRIE SEDGE

Plants with short stout rhizomes; culms 50–100 cm high, erect. Sheaths smooth, brown below, the lower ones with short blades, the membrane continued 2–3 mm beyond base of blades, copper-colored; principal blades 2–3 mm wide, to 40 cm long. Inflorescence 3–8 cm long, compound, with the branches, at least the lower ones, bearing 2–4 androgynous spikes; perigynia 2.5–3 mm long, dull dark brown, tapering to a beak with serrulate margins; scales equaling the perigynia, reddish brown. Swamps and wet meadows; Parklands, Boreal forest.

Carex preslii Steud.

PRESL SEDGE

Plants with thick, corm-like, short rhizomes; culms 20–70 cm high, slender, scabrous, densely tufted. Sheaths smooth, very thin, with the membrane white hyaline, loose; blades 1–4 mm wide, flat. Inflorescence 10–20 mm long, with 2–4 gynaeandrous spikes aggregated into a head; perigynia 3–4.5 mm long, yellowish to greenish brown; scales smaller than the perigynia, olive brown to chestnut. Alpine slopes; Rocky Mountains.

Carex pseudo-cyperus L.

CYPERUS-LIKE SEDGE

Plants with short rhizomes, densely sod-forming; culms to 1 m high, often nodding at the top, sharply 3-angled, very scabrous. Sheaths scabrous, loose; blades 5–15 mm wide, light green, flat, scabrous. Inflorescence 10–20 cm long; the terminal spike staminate, 2–5 cm long; pistillate spikes 3–5, the upper ones approximate, the lower distant, 3–7 cm long, spreading or drooping on slender

peduncles, 3–10 cm long; perigynia 5–6 mm long, spreading to reflexed at maturity, with the beak 0.5–2 mm long, bidentate; scales 2–3 mm long, ovoid, with the midrib prolonged into an awn, 3–4 mm long. Swamps and bogs; Boreal forests.

Carex pyrenaica Wahl.

SPIKED SEDGE

Plants densely tufted, with culms 5–25 cm high, equaling or exceeding the leaves. Sheaths smooth, whitish, tight; blades 0.3–1.5 mm wide, folded. Inflorescence a single androgynous spike, 5–20 mm long; perigynia 3–4.5 mm long, glossy dark brown, stipitate, tapering into a hyaline beak; scales shorter but wider than the perigynia, brown to dark brown. Rocky or grassy alpine slopes; Rocky Mountains.

Carex rariflora (Wahl.) Smith

SCANT SEDGE

Plants with slender rhizomes, the rootlets covered with a yellow or grayish felt-like layer; culms 20–30 cm high, bluntly 3-angled. Sheaths glabrous; blades 1–3 mm wide. Inflorescence 4–8 cm long; the terminal spike staminate or somewhat gynaeandrous, 6–15 mm long; pistillate spikes 1–3, approximate, 10–15 mm long, short-peduncled; perigynia 3–4 mm long, glaucous green, beakless; scales about equaling the perigynia, dark purple. Bogs and muskeg; southeastern Boreal forest.

Carex raynoldsii Dewey (Fig. 75)

RAYNOLD'S SEDGE

Plants with short, stout rhizomes; culms 20–80 cm high, erect, sharply 3-angled. Sheaths glabrous; blades 2–5 mm wide, flat, to 50 cm long. Inflorescence 4–8 cm long; the terminal spike staminate, 1–2 cm long; pistillate spikes 2–4, approximate or the lower one somewhat distant, 1–2 cm long; perigynia 3.5–4.5 mm long, yellowish green or brown, distinctly several-veined; scales shorter than the perigynia, dark reddish brown or black. Meadows and open woods; Rocky Mountains, Cypress Hills.

Carex retrorsa Schw. (Fig. 76)

TURNED SEDGE

Plants with short, stout rhizomes; culms 40–100 cm high, densely tufted, stout, obtusely 3-angled. Sheaths glabrous, often cross-veined; blades 3–12 mm wide, to 40 cm long, flat, soft. Inflorescence 5–15 cm long; the upper 1 or 2 spikes staminate, 10–25 mm long, often hidden among the 3–8 pistillate spikes, 15–60 mm long, all aggregated, or the lower ones approximate; perigynia 7–10 mm long, tapering into a long bidentate beak, shiny, with many distinct veins, light green or straw-colored; scales smaller than the perigynia. Swampy areas, lakeshores, and wet meadows; Boreal forest.

Carex richardsonii R. Br.

RICHARDSON'S SEDGE

Plants with long rhizomes; culms 10–25 cm high, tufted, short pubescent. Sheaths glabrous, open; blades 1.5–2.5 mm wide, stiff, short, mostly basal. Terminal spike staminate, 10–25 mm long, peduncled; lateral 2 or 3 spikes pistillate, 1–2 cm long, short-peduncled, erect, approximate or somewhat distant; perigynia 2.5–3.5 mm long, pubescent, green; scales larger than the perigynia, purplish brown, with the margins hyaline. Dry woods, open areas; Boreal forest, Rocky Mountains.



Fig. 75. Raynold's sedge, *Carex raynoldsii* Dewey.



Fig. 76. Turned sedge, *Carex retrorsa* Schw.

Carex rossii Boott

ROSS' SEDGE

Plants with slender rhizomes; culms 10–30 cm high, erect, often exceeded by the leaves. Sheaths smooth, purplish brown below; blades 1–3 mm wide, stiffly erect or curly. Inflorescence about 2 cm long; the terminal spike staminate, to 15 mm long; lateral spikes pistillate, 5–10 mm long; basal spikes abundant; perigynia 3–4 mm long, green to light brown, pubescent, with the beak 1–2 mm long; scales equaling the perigynia, keeled. Dry slopes, rocky areas, and clearings; throughout Prairies, Parklands, Rocky Mountains, Boreal forest.

Carex rostrata Stokes

BEAKED SEDGE

Plants with short rhizomes, and long stolons; culms 50–100 cm high, stout, erect, usually exceeded by the leaves. Sheaths smooth, strongly cross-veined; blades 5–10 mm wide, usually cross-veined, flat. Inflorescence 10–30 cm long; the upper spikes 2–4, staminate, 15–50 mm long, well separated from the 2–5 pistillate spikes, 4–10 cm long, short-peduncled to sessile; perigynia 4–8 mm long, narrowed to a bidentate beak; scales narrower than the perigynia, light brown. Swamps, marshes, lakeshores, riverbanks; throughout Parklands, Boreal forest.

Carex rotundata Wahl.

ROUND SEDGE

Plants with short rhizomes; culms 20–60 cm high, sharply 3-angled, erect. Sheaths smooth, reddish brown below; blades 1–3 mm wide, involute. Inflorescence 5–15 cm long; the terminal spikes 2 or 3, staminate, 10–20 mm long; the lateral spikes 3 or 4, pistillate, 20–40 mm long, sessile or short pedunculate; perigynia 3–3.5 mm long, tapering to the bidentate beak, obscurely veined; scales smaller than the perigynia, purplish black. Peat marsh and tundra; eastern Boreal forest.

Carex rufina Drejer

REDDISH SEDGE

Plants with slender rhizomes; culms 10–25 cm high, loosely tufted. Sheaths glabrous, short; leaves 1–3 mm wide, often surpassing the culms. Inflorescence 5–10 cm long, with bracts long-sheathing; the terminal spike gynaeandrous, 5–10 mm long; the lateral spikes 2 or 3, pistillate, 5–15 mm long, ascending on elongate peduncles; perigynia 2–3 mm long, brownish green, serrulate above, with a beak 0.1–0.2 mm long; scales smaller than the perigynia, purplish brown. Very rare; arctic and Boreal tundra; Nueltin Lake, Manitoba.

Carex rupestris All.

ROCK SEDGE

Plants with slender rhizomes; culms 8–15 cm high, solitary or few together. Sheaths reddish brown below; blades 1–2 mm wide, to 15 cm long, usually curved, exceeding the culms. Inflorescence a single androgynous spike, 1–2 cm long; perigynia about 3 mm long; scales dark chestnut brown with hyaline margins, somewhat shorter but broader than, and partly enveloping, the perigynia. Rare; open, usually calcareous soils; Boreal forest.

Carex salina Wahl.

SALT SEDGE

Plants with slender rhizomes; culms 5–60 cm high, curving or erect. Sheaths glabrous; blades to 9 mm wide, flat or involute. Inflorescence 5–12 cm

long; the upper spikes 1–3, staminate, 2–3 cm long; the 1–3 pistillate spikes 5–80 mm long, short-peduncled to sessile; perigynia 2–3.5 mm long, pale green; scales somewhat longer than the perigynia, purple brown, with the green midrib prolonged into an awn. Salt marshes; Hudson Bay. Three forms are recognized:

1. Culms 5–15 cm high, slender, obtusely angled; blades 1–2.5 mm wide; pistillate spikes 5–15 mm long, 3–4 mm thick. var. *subspathacea* (Wormsk.) Tuck.
 Culms 10–60 cm high, stiff; blades 2–9 mm wide; pistillate spikes 1–8 cm long, 3–10 mm thick. 2
2. Culms 10–30 cm high, obtusely angled; blades 2–4 mm wide, with the margins revolute; pistillate spikes 1–3 cm long, 3–4 mm thick. var. *salina*
 Culms 15–60 cm high, rather sharply angled; blades 2–9 mm wide, flat; pistillate spikes 2–8 cm long, 4–10 mm thick. var. *kattegatensis* (Fries) Almq.

Carex sartwellii Dewey

SARTWELL'S SEDGE

Plants with slender rhizomes; culms 40–80 cm high, sharply triangular, stiff. Sheaths with a green, clearly veined membrane; blades 2–4 mm wide. Inflorescence 2–6 cm long, with 6–20 androgynous spikes 5–10 mm long; perigynia 2.5–3.5 mm long, brown, finely veined; scales smaller than the perigynia, reddish brown, with the midrib green, margin hyaline. Wet meadows, bogs, and margins of woods; Boreal forest.

Carex saxatilis L.

ROCKY-GROUND SEDGE

Plants with creeping purplish rhizomes; culms 20–80 cm high, decumbent at the base, leafy. Sheaths smooth; blades 2–5 mm wide, flat, scabrous toward the tip. Inflorescence 3–10 cm long; the upper 1 or 2 spikes staminate, 1–3 cm long; the 1–3 pistillate spikes 5–20 mm long; perigynia 3–5 mm long, purplish black; scales smaller than the perigynia, reddish black. Gravelly soil and muskeg; Boreal forest. Two varieties can be distinguished:

- Culms 20–40 cm high; pistillate spikes sessile or short-peduncled, approximate. var. *saxatilis*
 Culms to 80 cm high; pistillate spikes long-peduncled, drooping, distant. var. *major* Olney

Carex scirpoidea Michx.

RUSH-LIKE SEDGE

Plants with long, creeping rhizomes; culms 20–50 cm high, solitary or few together. Sheaths glabrous, loose, the lower ones reddish brown; blades 1–3 mm wide. Inflorescence a single spike, staminate or pistillate, 1–3 cm long; perigynia 2.5–3 mm long, densely pubescent, green; scales smaller than the perigynia, deep brown. Marshy areas, slopes, and meadows; throughout the Prairie Provinces.

Carex scoparia Schk.

BROOM SEDGE

Plants with short rhizomes; culms 20–100 cm high, loosely to densely tufted, sharply 3-angled. Sheaths smooth; blades 1–3 mm wide, 15–60 cm long. Inflorescence 25–40 mm long, with 3–12 gynaeandrous spikes 6–12 mm long, approximate or crowded into a head; perigynia 4–7 mm long, lanceolate, flat, greenish or straw-colored; scales smaller than the perigynia, light brown, with green midrib. Moist areas, swamp, muskeg; Boreal forest, a few locations in Parklands.

Carex siccata Dewey (Fig. 77)

HAY SEDGE

Plants with long, tough rhizomes; culms 15–80 cm high, solitary or few together. Sheaths smooth; blades 1–3 mm wide, often almost as long as the culms. Inflorescence 2–4 cm long, with 4–12 spikes; the uppermost spike androgynous, the middle spikes staminate, the lower ones pistillate; perigynia 3–6 mm long, nerved on both sides, winged in the upper half, narrowing into a bidentate beak; scales shorter than the perigynia, light brown with hyaline margins. Common; sandy areas, open pine woods, sometimes covering large areas; Boreal forest. Syn.: *C. foenea* Willd.

Carex sprengelii Dewey

SPRENGEL'S SEDGE

Plants with long, creeping rhizomes; culms 15–80 cm high, tufted. Sheaths smooth, loose; blades 2–4 mm wide, flat, soft. Inflorescence 10–20 cm long; the upper 1–3 spikes staminate, 1–2 cm long; the lateral 2–4 spikes pistillate, 2–4 cm long, distant, on long slender ascending or drooping peduncles; perigynia 5–6 mm long, with the body subglobose, contracted into a bidentate beak as long as, or longer than, the body; scales narrower than the perigynia, greenish white. Open woods, shrubbery, and moist semiopen areas; throughout the Prairie Provinces.

Carex stenophylla Wahl. ssp. *eleocharis* (Bailey) Hulten

LOW SEDGE

Plants with slender, long creeping rhizomes; culms 3–20 cm high, solitary or tufted. Sheaths glabrous, loose; blades 1–2 mm wide, channeled to involute. Inflorescence 5–20 mm long, with 3–5 androgynous spikes, closely aggregated in a head; perigynia 2.5–3.5 mm long, dark brown at maturity; scales wider than the perigynia. Often very abundant; dry grasslands; Prairies.

Carex stipata Muhl.

AWL-FRUITED SEDGE

Plants with short, thick rhizomes; culms 10–100 cm high, erect, soft, winged on the angles. Sheaths glabrous, loose; blades 2–8 mm wide, often equaling or surpassing the culms. Inflorescence 1.5–10 cm long, compound, with 5–15 spikes aggregated into a head, the lower spikes sometimes approximate, the upper ones crowded; perigynia 4–6 mm long, yellowish, sharply nerved on both sides; scales smaller than the perigynia, brownish. Moist woods, swamps, and bogs; Parklands, Boreal forest.

Carex straminea Willd.

STRAW-COLORED SEDGE

Plants with short, stout rhizomes; culms 30–80 cm high, densely tufted. Sheaths glabrous, brownish; blades 2–6 mm wide. Inflorescence 1–6 cm long,



Fig. 77. Hay sedge, *Carex siccata* Dewey.

with 3–10 gynaeceandrous spikes 7–10 mm long, approximate to somewhat distant; perigynia 3.5–7 mm long, suborbicular; scales smaller than the perigynia. Meadows, grasslands, open forest; Parklands, Boreal forest. A large species, which has been divided as follows:

1. Perigynia broadest above the middle; leaves 3–6 mm wide; sheaths loose. *C. cumulata* (Bailey) Mack.
Perigynia broadest below the middle; leaves 2–4 mm wide; sheaths tight. 2
2. Perigynia thin, 5–7 mm long, 3–5 mm broad, distinctly veined on both sides. *C. bicknellii* Britt.
Perigynia thick, firm, 3.5–5 mm long, 2–3.5 mm broad, obscurely veined or veinless on the inside. 3
3. Perigynia 4–5 mm long, strongly veined on the back, obscurely veined inside; wings thick; scales almost reaching tip of beak. *C. brevior* (Dewey) Mack.
Perigynia 3.5–4 mm long, obscurely veined on both sides; wings thin; scales reaching to base of beak. 4
4. Heads 1–3 cm long; spikes crowded; leaves 2–3 mm wide. *C. molesta* Mack.
Heads to 8 cm long; spikes more or less separated; leaves 3–4 mm wide. *C. merritt-fernaldii* Mack.

Carex stricta Lam.

ERECT SEDGE

Plants with long, scaly, rather stout rhizomes; culms 40–100 cm high, loose-tufted into large clumps. Sheaths brown or reddish below, tight; blades 3–5 mm wide, flat. Inflorescence 5–15 cm long; the upper 1–3 spikes staminate, 1–4 cm long; the 1–4 lateral spikes pistillate, or sometimes androgynous, 2–6 cm long, erect, sessile or the lower ones short-peduncled, approximate or somewhat distant; perigynia 1.5–3 mm long, light green to straw-colored; scales variable, from narrower and shorter to broader and longer than the perigynia, reddish brown to purple brown. Marshy areas; southeastern Boreal forest.

Carex supina Wahl. ssp. *spaniocarpa* (Steud.) Hulten

WEAK SEDGE

Plants with slender brown rhizomes; culms 8–30 cm high, sharply 3-angled, slender. Sheaths glabrous; blades 0.5–2 mm wide, channeled, crowded, often almost as high as the culms. Inflorescence 2–3 cm long; the terminal spike staminate, 10–15 mm long, pale; the 1–3 lateral spikes pistillate, 5 mm long, with 3–5 perigynia, approximate; perigynia 2.5–3 mm long, plump, lustrous, with a short beak; scales equaling the perigynia, light reddish brown. Rare; stony areas, beaches; southeastern Boreal forest, Parklands.

Carex sychnocephala Carey

LONG-BEAKED SEDGE

Plants densely tufted, with culms 10–50 cm high, erect or spreading. Sheaths smooth, tight; blades 2–5 mm wide, often exceeding the inflorescence,

flat. Inflorescence without the bracteal leaves 2–3 cm long; perigynia 5–6 mm long, lanceolate, green; scales smaller than the perigynia, hyaline with a green midrib. Wet meadows and lakeshores; Parklands, Boreal forest.

Carex tenuiflora Wahl.

THIN-FLOWERED SEDGE

Plants with very slender rhizomes; culms 20–60 cm high, slender. Sheaths smooth, loose; blades 1–2 mm wide, flat. Inflorescence 5–15 mm long, with 2–4 gynaecandrous spikes approximate to crowded; perigynia 3–3.5 mm long, greenish; scales equaling the perigynia, white hyaline with green midrib. Rare; wet wood, bogs, and muskeg; Boreal forest.

Carex tetanica Schk.

RIGID SEDGE

Plants with slender rhizomes; culms 30–60 cm high, tufted. Sheaths glabrous; blades 2–5 mm wide, flat. Inflorescence 5–10 cm long; the terminal spike staminate, 1–2 cm long, with the peduncle to 10 cm long; the 1–3 lateral spikes pistillate, 1–3 cm long, slender-peduncled, ascending; perigynia 3–4 mm long, light brown, finely veined; scales shorter than the perigynia, dark reddish brown with green midrib. Open grassland; southeastern Parklands, Boreal forest. Two varieties are represented:

Perigynia 2-ranked; the pistillate spikes 3–5 mm thick. var. *woodii* (Dewey) Wood

Perigynia 6-ranked; the pistillate spikes 5–7 mm thick. var. *meadii* (Dewey) Bailey

Carex tinctoria Fern.

TINGED SEDGE

Plants with short rhizomes; culms 40–80 cm high, slender, sharply triangular, tufted. Sheaths somewhat cross-veined, with the membrane wrinkled; blades 2–4 mm wide, flat, firm. Inflorescence 1–4 cm long, with 4–10 gynaecandrous spikes, approximate to closely crowded into a head; perigynia 3.5–4 mm long, brownish, sharply many nerved on the back, tapering into a serrate beak; scales about equaling the perigynia, reddish brown with green midrib and hyaline margins. Meadows and open woods; Parklands, Boreal forest. Presumably a hybrid of species in the *C. festucacea* group.

Carex tonsa (Fern.) Bickn.

BALD SEDGE

Plants with creeping rhizomes; culms poorly developed, hidden among the basal leaves. Sheaths reddish brown; blades 2–4 mm wide, spreading, curved. Inflorescence 10–15 mm long, hidden among the leaves; perigynia 1.5–3 mm long, glabrous or nearly so, yellowish brown, with the beak about 1.0–1.5 mm long, curved, bidentate; scales equaling the perigynia, pale brown with green midrib. Open woods, sandy areas; Boreal forest.

Carex torreyi Tuck.

TORREY'S SEDGE

Plants with short rhizomes; culms 25–50 cm high, slender, weak, sharply triangular, somewhat pubescent below the inflorescence. Sheaths soft pubescent, tight; blades 1.5–4 mm wide, flat, pilose. Inflorescence 3–5 cm long; the terminal spike staminate, 5–15 mm long; the 1–3 lateral spikes pistillate, 5–15 mm long, sessile or short-peduncled; perigynia 2.5–3.5 mm long, yellowish

green, strongly nerved, abruptly short-beaked; scales smaller than the perigynia, light brown with broad hyaline margins. Rare; in meadows and moist woods; Prairies, Parklands.

Carex tribuloides Wahl.

PRICKLY SEDGE

Plants tufted, with culms 30–70 cm high, stout. Sheaths glabrous, tight; blades 3–7 mm wide. Inflorescence 2–5 cm long, with 7–12 spikes 6–12 mm long, approximate or crowded; perigynia 3–5 mm long, straw-colored, obscurely veined on both sides; scales smaller than the perigynia, pale brown with hyaline margins. Wet areas, marshes, and shores; southeastern Boreal forest. Includes *C. projecta* Mack., *C. cristatella* Britt.

Carex trisperma Dewey

THREE-SEEDED SEDGE

Plants with slender rhizomes; culms 20–70 cm high, very slender and weak, often straggling. Sheaths glabrous, tight; blades to 30 cm long, 1–3 mm wide, flat. Inflorescence 3–6 cm long, with 1–3, but commonly 2, spikes 2–4 cm apart; the lower bract 2–4 cm long; spikes 3–5 mm long, gynaeandrous, with 1–5 perigynia 2.5–4.0 mm long, greenish brown; scales shorter than to equaling the perigynia, with the center green and margins hyaline. Wet woods, bogs, and marshes; Boreal forest.

Carex umbellata Schk.

UMBELLATE SEDGE

Plants with stout rhizomes; culms 3–20 cm high. Sheaths purplish or reddish brown below; blades 1–3 mm wide, stiff, usually exceeding the culms. Inflorescence of two kinds: 1 culm with 1 terminal spike staminate, 5–10 mm long, sometimes with a pistillate spike below it; and 1–3 very short culms bearing 1 pistillate spike 4–10 mm long; perigynia 1.5–3 mm long, pubescent, yellowish green, with a prominent, often curved, beak; scales equaling the perigynia, with the center green and the margins hyaline. Open woods on sandy or stony soils; Boreal forest, Rocky Mountains.

Carex vaginata Tausch

SHEATHED SEDGE

Plants with long, slender, creeping rhizomes; culms 20–60 cm high, solitary or few together. Sheaths glabrous, tight; blades 2–5 mm wide, flat. Inflorescence 10–15 cm long; the terminal spike staminate, 1–2 cm long, peduncled; lateral spikes 1–3, often androgynous, 10–25 mm long, short-peduncled, loosely spreading, or the lower ones drooping; perigynia 3–4 mm long, light brown, beaked; scales smaller than the perigynia, purplish brown. Wet woods, bogs, and muskeg; Boreal forest.

Carex vesicaria L.

BLISTER SEDGE

Plants with stout, creeping rhizomes; culms 50–100 cm high, tufted. Sheaths glabrous, cross-veined; blades 3–7 mm wide, equaling the culms, or those of sterile shoots exceeding them. Inflorescence 10–20 cm long; the 1–4 upper spikes staminate, 2–6 cm long; the lower spikes 1–3, but usually 2, pistillate, 3–5 cm long; perigynia 7–9 mm long, yellow green to light brown, inflated, strongly veined, with the beak about 2 mm long and having divergent teeth; scales smaller than the perigynia, light brown. Very rare; marshes and lakeshores; southeastern Boreal forest.

Plants more or less densely tufted, with erect culms 5–40 cm high. Sheaths glabrous, loose; blades 1–3 mm wide, usually yellowish green. Inflorescence 1–5 cm long; the terminal spike staminate, 5–15 mm long, peduncled, often barely exceeding the 2–6 pistillate spikes 5–10 mm long, crowded to approximate; the lowest bract much exceeding the inflorescence; perigynia 2–3.5 mm long, green or brownish green; scales about equaling the perigynia, yellowish brown. Wet areas, particularly calcareous or slightly alkali lakeshores; throughout the Prairie Provinces.

Plants with short, stout rhizomes; culms 30–80 cm high, densely tufted. Sheaths glabrous, with the membrane greenish, clearly veined, and cross-wrinkled; blades 2–4 mm wide, commonly exceeding the leaves. Inflorescence 4–8 cm long, compound, with many densely crowded spikes; bracts up to 5 cm long, mostly well exerted; perigynia 2–3 mm long, yellowish green; scales about equaling the perigynia. Wet areas, meadows, and bogs; Boreal forest.

Plants with short, thick rhizomes; culms 30–60 cm high, densely tufted. Sheaths glabrous; blades 2–3 mm wide, flat, stiff. Inflorescence 3–8 cm long, with 3–8 spikes 6–12 mm long, crowded toward the tip, approximate to distant below; perigynia 4–6 mm long, straw-colored to light brown; scales about equaling the perigynia, light brown, with the midrib green and the margins hyaline. Open woods, meadows, disturbed areas; Parklands, southern Boreal forest, Cypress Hills.

Cyperus nut-grass

Triangular-stemmed plants with long grass-like leaves, mostly basal. Flowers perfect and borne in spikelets or in an umbel-like inflorescence at the summit of the stem. Fruit a 3-angled achene.

- 1. Tip of the scales slender, recurved; spikelets greenish to pale brown, 2–10 mm long; annual with fibrous roots. *C. inflexus*
- Tip of the scales not recurved; spikelets yellowish to brownish, to 20 mm long; perennials with corm-like thickened roots. 2
- 2. Joints of the rachilla with conspicuous hyaline wings; scales 3.5–6 mm long, acute or mucronate. *C. strigosus*
- Joints of the rachilla sharp-edged, not winged; scales 2–4 mm long, awned or mucronate. 3
- 3. Upper scales with an awn 0.5–1 mm long; achenes 2–3 mm long, about 1 mm thick. *C. schweinitzii*

Upper scales obtuse or short mucronate;
 achenes 1.5–2 mm long, about 1.2 mm
 thick. *C. houghtonii*

Cyperus houghtonii Torr.

HOUGHTON'S NUT-GRASS

Plants with short rhizomes; culms 20–70 cm high, obtusely angled, smooth; leaves 2–4 mm wide. Heads loosely fan-shaped; bracts 2–4, exceeding the inflorescence; spikelets 5–20 mm long; scales almost orbicular, 2–3 mm long; achene 1–1.5 mm long, ellipsoid, dark brown. Rare; sandy areas; south-eastern Parklands, Boreal forest.

Cyperus inflexus Muhl.

AWNED NUT-GRASS

Plants annual, low, tufted, fragrant when bruised or dried; culms 1–15 cm high, very slender; leaves 1–2 mm wide. Heads dense; bracts 2–4, much exceeding the inflorescence; spikelets 5–10 mm long; scales 1.5–3 mm long, oblong, strongly nerved, narrowed to a long, recurving, slender tip; achene 0.6–1 mm long, 3-sided obovoid, pale brown. Rare; slough margins, Prairies.

Cyperus schweinitzii Torr.

SAND NUT-GRASS

Plants with hard, short, scaly rhizomes, with corm-like branches; culms 10–70 cm high, sharply angled; leaves 2–6 mm wide, firm. Heads mostly obovoid; bracts 3–6, exceeding the inflorescence; spikelets 6–15 mm long; scales 3.5–4.5 mm long, ovate, strongly nerved, with the midrib prolonged into a short awn; achene 2.5–3.5 mm long, 3-sided ellipsoid, light brown. Rare; sand-hills; Parklands.

Cyperus strigosus L.

STRAW-COLORED NUT-GRASS

Plants with a hard, short, corm-like rhizome; culms tufted, 10–60 cm high, stout; leaves 2–12 mm wide, flat, soft. Heads compound, condensed to somewhat open; bracts 2–6 or 7, exceeding the inflorescence; spikelets compressed, 5–10 mm long, yellowish; scales 3–4.5 mm long, oblong, strongly nerved; achene 1.5–2 mm, 3-sided, linear. Rare; moist meadows and swamps; eastern Parklands, southeastern Boreal forest.

Eleocharis spike-rush

Perennial, sedge-like, slough margin plants, from coarse, creeping roots, or annuals with fibrous roots. Leaves generally reduced to sheaths without blades and all basal. Flowers perfect and borne in a spike at summit of stem.

- 1. Achenes lenticular or biconvex; stigmas
 - 2. 2
 - Achenes triangular; stigmas 3. 6
- 2. Plants annual; the root system fibrous. *E. engelmannii*
 - Plants perennial; the root system rhizomatous. 3
- 3. Sterile basal scale 1 encircling the culm at base of spikelet. 4
 - Sterile basal scales 2 or 3 at base of spikelet. 5

4. Tubercle narrowly conical, the base less than half as wide as the achene; spikelets closely many-flowered. *E. calva*
 Tubercle deltoid-conical, the base more than half as wide as the achene; spikelets loosely few-flowered. *E. uniglumis*
5. Tubercle as broad as long; scales firm, stiffly acuminate. *E. smallii*
 Tubercle much longer than broad; scales soft, scarious at the tip. *E. palustris*
6. Tubercle confluent with the achene, forming a beak. 7
 Tubercle distinctly jointed to the achene, forming a cap. 8
7. Spikelets to 20 mm long, many-flowered, fusiform. *E. rostellata*
 Spikelets to 7 mm long, 2- to 8-flowered, ovoid. *E. pauciflora*
8. Culms to 2 dm high, loosely tufted, forming mats from slender rhizomes. *E. acicularis*
 Culms 2–10 dm high, not matted, arising from stout rhizomes. 9
9. Culms 4- to 8-angled; achene 0.6–1.1 mm long, including the tubercle. *E. elliptica*
 Culms flattened; achene 1–1.5 mm long, including the tubercle. *E. compressa*

Eleocharis acicularis (L.) R. & S.

NEEDLE SPIKE-RUSH

Plants with very slender rhizomes, tufted, forming dense mats; culms slender, 2–15 cm high; sheaths loose. Spikelets flattened, 2–7 mm long, 3- to 15-flowered; scales 1–2.5 mm long, membranous, with a green midrib; achenes 0.7–1.2 mm long, whitish, ellipsoid. Common; around sloughs, in mud flats, along shores, and other wet places; throughout the Prairie Provinces.

Eleocharis calva Torr.

BALD SPIKE-RUSH

Plants with slender, reddish rhizomes; culms 10–50 cm high, solitary or loosely tufted; sheaths reddish, tight. Spikelets 10–20 mm long, lanceolate, densely flowered; scales 2–3 mm long, oblong-ovate, membranous, pale brown; achenes 1–1.5 mm long, narrowly obovoid, light brown. Not common; marshes, lakeshores, mud flats; southeastern Parklands.

Eleocharis compressa Sulliv.

FLATTENED SPIKE-RUSH

Plants with thick, hard rhizomes, 2–4 mm thick; culms 15–50 cm high, flat, wiry; sheaths tight. Spikelets 5–15 mm long, ovoid to ellipsoid; scales 2–3 mm long, ovate, brown or purplish; achenes 1–1.5 mm long, bluntly 3-sided, golden brown. Marshy areas, sandy shores; Parklands.

Eleocharis elliptica Kunth

SLENDER SPIKE-RUSH

Plants with slender purplish rhizomes, 1–2 mm thick; culms 5–50 cm high, scattered or in small tufts; sheaths tight, reddish below. Spikelets 3–10 mm long, loosely flowered; scales 2–3.5 long, ovate, reddish brown; achenes about 1 mm long, yellow or orange, prominently cross-ridged. Wet areas, sand dunes, shores, meadows; southeastern Parklands.

Eleocharis engelmannii Steud.

ENGELMANN'S SPIKE-RUSH

Plants annual, with culms 10–50 cm high, densely tufted; sheaths loose. Spikelets 4–15 mm long, ovoid-cylindric; scales 1–1.5 mm long, elliptic-obovate, brownish; achenes 1–2 mm long, lustrous brown when ripe. Wet areas; Prairies, Parklands.

Eleocharis palustris (L.) R. & S. (Fig. 78)

CREEPING SPIKE-RUSH

Plants with slender, long, creeping, reddish rhizomes; culms 10–60 cm high, firm; sheaths loose, red or brownish. Spikelets 5–20 mm long; scales 3–4 mm long, oblong-ovate, reddish brown; achenes 1–1.5 mm long, obovoid, light brown. Common; in shallow water, mud flats, and lakeshores; throughout the Prairie Provinces.

Eleocharis pauciflora (Lightf.) Link var. *fernaldii* Svenson

FEW-FLOWERED SPIKE-RUSH

Plants with long, creeping rhizomes, often with scaly tubers; culms 10–30 cm high, in small tufts. Spikelets 3–6 mm long; scales 3–5 mm long, ovate, lustrous brown; achenes 2–2.5 mm long, net-veined, gray brown. Calcareous wet soils; Boreal forest.

Eleocharis rostellata Torr.

BEAKED SPIKE-RUSH

Plants with short caudex; culms 10–60 cm high, tufted; sheaths loose. Spikelets 6–12 mm long; scales 2–3 mm long, ovate, green with brown midrib; achenes 2–3 mm long, oblong-obovoid, olive brown. Wet soils; northwestern Boreal forest.

Eleocharis smallii Britt.

SMALL'S SPIKE-RUSH

Plants with long-creeping rhizomes; culms 10–60 cm high; sheaths loose, reddish. Spikelets 5–20 mm long, loosely flowered; scales 3–5 mm long, lanceolate to narrowly ovate; achenes 1–1.5 mm long, obovoid, brown. Mud flats, peaty swamps, and shores; throughout the Prairie Provinces.

Eleocharis uniglumis (Link) Schultes

ONE-GLUMED SPIKE-RUSH

Plants with slender reddish rhizomes; culms 10–50 cm high, loosely tufted, very slender; sheaths tight, reddish. Spikelets 1–1.5 cm long, loosely flowered; scales 2–3 mm long, thin; achenes 1–1.5 mm long, ellipsoid to obovoid. Wet areas, lakeshores; throughout the Prairie Provinces.




Fig. 78. Creeping spike-rush, *Eleocharis palustris* (L.) R. & S.

Eriophorum cotton-grass

1. Spikelets several, spreading or pendulous on peduncles; the involucre of one or more leafy bracts. 2
Spikelets solitary at the tip of the stem; the involucre lacking. 4
2. Involucre a single bract, shorter than the inflorescence; leaf blades 1–2 mm wide. *E. gracile*
Involucre 2 or more bracts, about as long as the inflorescence; leaf blades 2–8 mm wide. 3
3. Midrib of scales prominent, reaching the tip of the scales; upper leaf sheaths not dark at the collar. *E. viridi-carinatum*
Midrib of scales not prominent, not reaching the tip of the scales; upper leaf sheaths dark at the collar. *E. angustifolium*
4. Plants stoloniferous; the culms solitary or few together. 5
Plants not stoloniferous; the culms more or less densely tufted. 6
5. Scales lead gray to black, with narrow pale margins; bristles bright white. *E. scheuchzeri*
Scales brownish to black, with wide hyaline margins; bristles pale reddish brown. *E. chamissonis*
6. Scales with broad, whitish or hyaline margins, the basal ones reflexed; bristles bright white. *E. vaginatum*
Scales without broad margins, the basal ones not reflexed. 7
7. Bristles light brownish; upper sheaths scarcely inflated. *E. brachyantherum*
Bristles bright white; upper sheaths strongly inflated. *E. callitrix*

Eriophorum angustifolium Honck.

TALL COTTON-GRASS

Plants with short, stout rhizomes; culms 20–60 cm high, mostly solitary, soft blunt-angled. Sheaths smooth, reddish below, with blades 3–6 mm wide, flat to convolute, scabrous. Inflorescence with 2–10 spikelets; bracts 2–3, dark purple at the base; spikelets 1–2 cm long, divergent or drooping, with bristles 2–5 cm long, white; scales lead gray to brownish; achenes 2.5–3.5 mm long. Muskeg, swamps, wet meadows; Boreal forest.

Eriophorum brachyantherum Trautv.

CLOSE-SHEATHED COTTON-GRASS

Plants tufted, with culms 20–60 cm high, terete, slender. Sheaths smooth, with blades 1–2 mm wide, filiform, the upper sheaths without blades.

Inflorescence a solitary spikelet 1–1.5 cm long, with bristles 2–3 cm long, cream-colored; scales lead gray; achenes 2–2.5 mm long. Bogs and swamps; Boreal forest.

Eriophorum callitrix Cham.

BEAUTIFUL COTTON-GRASS

Plants tufted, with culms 10–25 cm high, stiff, stout. Sheaths smooth, with blades 2–4 mm wide, coarse; cauline sheaths with a short blade, the upper ones inflated. Inflorescence a single spikelet about 1 cm long, with bristles 2–3 cm long, bright white; scales grayish; achenes 2–2.5 mm long. Bogs and swamps; southern Boreal forest.

Eriophorum chamissonis C. A. Mey.

RUSSET COTTON-GRASS

Plants with short rhizomes and somewhat stoloniferous; culms 10–35 cm high, solitary or few together. Sheaths loose, the lower ones with blades 1–3 mm wide, channeled or involute; the upper sheath bladeless, tight. Inflorescence a single spikelet 1–1.5 cm long, with bristles 2.5–3 cm long, reddish brown; scales brownish to blackish, with white margins; achenes 2–2.5 mm long. A form with white bristles is f. *albidum* (Nyl.) Fern. Muskeg and swampy areas; Boreal forest.

Eriophorum gracile Koch

SLENDER COTTON-GRASS

Plants in small tufts, with culms 20–40 cm high, very slender, weak, mostly without basal leaves. Sheath of upper culm leaf 3–6 cm long, with the blade 1–3 cm long, 1–1.5 mm wide. Inflorescence with 2–5 spikelets, with a single bract 1–2 cm long, lead gray or blackish; spikelets spreading or nodding, 7–10 mm long, with bristles 1–1.5 cm long, bright white; scales lead gray to blackish; achenes 1.5–2 mm long. Muskeg, swamps, and bogs; Boreal forest.

Eriophorum scheuchzeri Hoppe

ONE-SPIKE COTTON-GRASS

Plants with rhizomes; culms 10–40 cm high, solitary, somewhat leafy at the base. Sheaths loose, the upper ones bladeless, black-tipped; the lower sheaths with blades 2–5 mm wide, channeled or involute. Inflorescence a solitary spikelet about 1 cm long, with bristles 1–1.5 cm long, bright white; scales lead gray to blackish, with narrow pale margins; achenes 1.5–2 mm long. Muskeg areas, sedge swamps; Boreal forest.

Eriophorum vaginatum L.

SHEATHED COTTON-GRASS

Plants densely tufted, forming large clumps, with culms 20–60 cm high, stiff, 3-sided. Sheaths glabrous, with blades stiff, filiform; culm leaves consisting of one or two inflated bladeless sheaths. Inflorescence a solitary spikelet 1–1.5 cm long, with bristles 2–2.5 cm long, bright white; scales lead gray to blackish, divergent or reflexed; achenes 2.5–3.5 mm long. In the Prairie Provinces, our plants are ssp. *spissum* (Fern.) Hulten, differing from the typical Eurasian plants in having longer and more inflated upper sheaths, larger anthers, and narrower achenes. Bogs and marshes; Boreal forest.

Eriophorum viridi-carinatum (Engelm.) Fern.

THIN-LEAVED COTTON-GRASS

Plants in small tufts, with culms 20–70 cm high, 3-sided, slender. Sheaths smooth; basal leaves numerous, with blades 2–6 mm wide, flat to channeled;

cauline leaves 2 or 3, short. Inflorescence with 3–15 spikelets; bracts 2–4, green or with a brownish base; spikelets 5–10 mm long, with bristles 1–2 cm long, cream-colored to grayish; scales olive green to lead gray, with the midrib prominent, reaching the tip, sometimes protruding; achenes 1.5–2 mm long. Muskeg, bogs, and marshes; Boreal forest.

Kobresia bog-sedge

Spikes linear, undivided; glumes 3–3.5 mm long; achenes about 2.5 mm long. *K. myosuroides*

Spikes ovoid, compound; glumes 2.5–3 mm long; achenes about 3 mm long. *K. simpliciuscula*

Kobresia myosuroides (Vill.) F. & P. BOG-SEDGE

Plants densely tufted, with culms 10–30 cm high, erect. Sheaths brown, becoming fibrillose and very conspicuous, with blades about 1 mm wide, filiform, erect, about equaling the culms. Inflorescence 1–3 cm long, linear, simple; upper spikelets staminate, the lower ones androgynous; scales 2–3 mm long; glumes 3–3.5 mm long, light brown, shiny, tightly enclosing the achenes, these 2.5 mm long, short stipitate, light brown. Rare; mountain slopes; Rocky Mountains.

Kobresia simpliciuscula (Wahl.) Mack. SIMPLE BOG-SEDGE

Plants densely tufted, with culms 10–30 cm high, erect. Sheaths cinnamon brown, with blades about 1 mm wide, filiform, involute; old sheaths and blades very conspicuous. Inflorescence 1–4 cm long, compound; the terminal spike staminate, the lateral ones androgynous or pistillate, 3–8 mm long; scales small, reddish brown; glumes 2.5–3 mm long, reddish brown, shiny, enclosing the achene. Rare; slopes and calcareous bogs; Rocky Mountains, Hudson Bay.

Rhynchospora beak-rush

Glomerules turbinate; uppermost bract about as long as the glomerules; bristles 10–12. *R. alba*

Glomerules ovoid or oblong; uppermost bract much longer than the glomerules; bristles 6. *R. capillacea*

Rhynchospora alba (L.) Vahl WHITE BEAK-RUSH

Plants densely tufted, with stems 15–30 cm high, slender. Leaves 0.5–2.5 mm wide, short, mostly basal. Inflorescence 8–15 mm long, at first white, later brownish; spikelets 3.5–5 mm long, usually 2-flowered; bristles 10–12, retrorsely barbed. Rare; bogs, wet places, muskeg; Boreal forest.

Rhynchospora capillacea Torr. SLENDER BEAK-RUSH

Plants tufted, with stems 10–40 cm high, very slender. Leaves about 1 mm wide, the upper ones commonly exceeding the inflorescence. Inflorescence 8–10 mm long, with up to 10 spikelets 5–7 mm long; bristles 6, retrorsely barbed. Rare; bogs, swamps, and wet sands; Boreal forest.

Scirpus bulrush

- 1. Involucre consisting of 1 scale-like bract; inflorescence appearing terminal. 2
 Involucre green, with the bract or bracts erect or leaf-like, spreading; inflorescence appearing lateral or terminal. 6
- 2. Culms round, smooth. 3
 Culms triangular, scabrous. 4
- 3. Plants densely cespitose; spikelet lanceolate; bristles 6. *S. caespitosus*
 Plants in small tufts, rhizomatous; spikelet ovate; bristles none. *S. pumilus*
- 4. Inflorescence of several spikelets. *S. rufus*
 Inflorescence a single spikelet. 5
- 5. Bristles 20–30 mm long. *S. hudsonianus*
 Bristles 2–3 mm long. *S. clintonii*
- 6. Involucre a single erect bract, appearing as a continuation of the stem; the inflorescence appearing lateral. 7
 Involucre of several leaf-like bracts, spreading; the inflorescence appearing terminal. 12
- 7. Culms terete; sheaths bladeless. 8
 Culms triangular; sheaths with blades. 9
- 8. Scales almost as long as the achenes, broadly ovate; culms spongy. *S. validus*
 Scales much longer than the achenes, narrowly ovate; culms firm. *S. acutus*
- 9. Spikelets arranged in a short spike. *S. rufus*
 Spikelets sessile, clustered at one point on the stem. 10
- 10. Achenes triangular; stigmas 3. *S. torreyi*
 Achenes plano-convex; stigmas 2. 11
- 11. Achene reticulate, blunt at the tip; leaves stiffly erect. *S. nevadensis*
 Achene smooth, minutely beaked; leaves usually spreading. *S. americanus*
- 12. Spikelets 10–50 mm long, short-peduncled or subsessile, not very numerous. 13
 Spikelets 3–10 mm long, long-peduncled in a much branched inflorescence, very numerous. 14
- 13. Spikelets in part sessile or subsessile, in part 1–4 at the tip of peduncles to 10 cm long; stigmas 3. *S. fluviatilis*

- Spikelets all sessile or subsessile, or in part sessile, in part on peduncles to 5 cm long; stigmas 2. *S. paludosus*
14. Bristles much longer than the scales, protruding. *S. cyperinus*
 Bristles shorter than the scales. 15
15. Achene triangular; stigmas 3; sheaths green. *S. atrovirens*
 Achene lenticular; stigmas 2; sheaths usually reddish purple. *S. microcarpus*

Scirpus acutus Muhl.

VISCID GREAT BULRUSH

Plants with extensive rhizomes; culms 50–200 cm high, round, olive green, hard, and firm; sheaths firm, often with a short blade. Panicle 5–15 cm long, with branches ascending or spreading, rarely pendulous; spikelets 1–2 cm long; scales red-dotted, viscid-pubescent above; achene shiny black at maturity. Sloughs, riverbanks, and lakeshores; Prairies, Parklands.

Scirpus americanus Pers.

THREE-SQUARE BULRUSH

Plants with long creeping, stout rhizomes; culms 20–100 cm high, solitary or few together, sharply 3-angled; leaves 2–8 mm wide, linear, channeled. Inflorescence 1–5 cm long, with 1–8 spikelets; bract 2–15 cm long; spikelets 5–20 mm long, ovoid to almost cylindrical, reddish brown. Wet, often brackish to moderately saline areas; Prairies, Parklands.

Scirpus atrovirens Willd.

GREEN BULRUSH

Plants tufted, with culms 30–100 cm high, bluntly 3-angled; leaves 5–15 mm wide. Inflorescence 5–20 cm long, umbelliform, with forked branches; spikelets 2–8 mm long, greenish brown; bracts 3–5, leafy, with the longest one much longer than the inflorescence. Bogs and marshes; Prairies, Parklands.

Scirpus caespitosus L. var. *callosus* Bigel.

TUFTED BULRUSH

Plants densely tufted, forming hard tussocks, with culms 10–30 cm high, wiry, filiform, round; basal sheaths bearing a blade to 15 mm long. Inflorescence a single spikelet 3.5–6 mm long, ovoid to lanceolate. Bogs, marshes, and tundra; Boreal forest.

Scirpus clintonii Gray

CLINTON'S BULRUSH

Plants densely tufted, with culms 10–30 cm high, very slender, sharply 3-angled; leaves 0.5–1 mm wide, short. Inflorescence a single spikelet 3–6 mm long, lanceolate to ovoid, pale brown; the outer scales with the midrib prolonged into an awn. Rocky Mountains.

Scirpus cyperinus (L.) Kunth

WOOL-GRASS

Plants tufted, with culms 50–100 cm high, obscurely 3-angled; leaves 2–5 mm wide, to 30 cm long. Inflorescence 5–20 cm long, with branches 3–10 cm long, ascending or spreading to drooping; spikelets 3–6 mm long, very numerous, ovoid, with the bristles exceeding the scale, elongating at maturity; bracts of the involucre 5–30 cm long. Wet meadows, swamps, and lakeshores; Boreal forests, rarely in Parklands.

Scirpus fluviatilis (Torr.) Gray

RIVER BULRUSH

Plants with long rhizomes having corm-like thickenings; culms 50–150 cm high, sharply 3-angled, leafy to near the inflorescence; leaves 5–15 mm wide, pale green, flat. Inflorescence 6–10 cm long, with 1–5 spikelets; bracts to 15 cm long; spikelets 15–40 mm long, brown. Along streams, lakeshores; Boreal forest.

Scirpus hudsonianus (Michx.) Fern.

ALPINE COTTON-GRASS

Plants with creeping rhizomes; culms 10–45 cm high, in small tufts, sharply angled. Inflorescence a single spikelet 4–7 mm long; the lowest scale with an awn 2–4 mm long, the other scales awnless; bristles 6, exerted in flowering, elongating to 1–3 cm at maturity, barbless. Marshes and boggy places; Boreal forest, Rocky Mountains.

Scirpus microcarpus Pers.

SMALL-FRUITED BULRUSH

Plants with a thick rhizome; culms 30–80 cm high, stout; leaves 4–15 mm wide, firm, flat. Inflorescence 5–20 cm long, with branches 3–15 cm long, the shorter ones ascending, the longer ones drooping; spikelets 3–6 mm long, numerous, dark green; bracts usually 3, the lower one exceeding the inflorescence. Damp areas, swamps, and bogs; Boreal forest, Parklands.

Scirpus nevadensis Wats.

NEVADA BULRUSH

Plants with creeping rhizomes; culms 30–50 cm high, solitary or few tufted, blunt-angled or almost terete. Inflorescence 1–5 cm long, with 2–10 spikelets; the bract 3–12 cm long; spikelets 5–20 mm long, ovoid to lanceolate, reddish brown. Not common; lakeshores and riverflats, often brackish; western Prairies, Parklands.

Scirpus paludosus Nels.

PRAIRIE BULRUSH

Plants with stout rhizomes, corm-like thickened internodes; culms 30–120 cm high, 5–20 mm thick at the base; leaves 5–15 mm wide, channeled. Inflorescence 5–10 cm long, with spikelets 10–25 mm long, sessile or more or less long-peduncled, ovoid, usually pale brown; the bracts 5–20 cm long. Common; in alkaline marshes and lakeshores; throughout the Prairie Provinces.

Scirpus pumilus Vahl

DWARF BULRUSH

Plants densely tufted on short rhizomes, with culms 10–20 cm high, sharply angled, wiry. Inflorescence a solitary spikelet, 3–5 mm long, ovoid, light brown. Bogs and marshes; Boreal forest, Parklands.

Scirpus rufus (Huds.) Schrad.

RED BULRUSH

Plants with extensively creeping rhizomes; culms 10–40 cm high, tufted, bluntly angled; leaves 1–3 mm wide, erect. Inflorescence a spike, 10–20 mm long, with two rows of spikelets, involucre a single bract, to 5 cm long; spikelets 5–10 mm long, 2- to 5-flowered, lustrous reddish brown; bristles lacking or small. Saline or brackish marshes; Boreal forest, Parklands. Syn.: *Blysmus rufus* (Huds.) Link.

Scirpus torreyi Olney

TORREY'S BULRUSH

Plants with a slender, lax, creeping rhizome; culms 40–100 cm high, 3-angled, solitary; leaves 5–10 mm wide, channeled, firm or lax. Inflorescence 3–5 cm long, spikelets 10–15 mm long, ovoid to cylindrical; bracts 3–15 cm long. Lakeshores and ponds; southeastern Boreal forest, Parklands.

Scirpus validus Vahl

GREAT BULRUSH

Plants with stout, extensively creeping, reddish rhizomes; culms 50–300 cm high, to 20 mm thick at the base, round, soft, easily compressed; sheaths usually bladeless. Inflorescence 5–10 cm long, with branches 1–7 cm long, the longer ones repeatedly branching, drooping; spikelets 3–7 mm long, brown. Very common; often forming extensive borders along sloughs and lakeshores; throughout the Prairie Provinces.

LEMNACEAE—duckweed family

Aquatic perennials with small disk-like, leafy fronds floating on or in the water, with one or several rootlets hanging from the lower surface of the frond. Flowers unisexual, minute, and borne on the surface of the fronds; fruit a minute achene with a thin-walled covering. Duckweeds are found in still water throughout the Prairie Provinces.

- Rootlets solitary, one to a frond. *Lemna*
- Rootlets several to a frond. *Spirodela*

Lemna duckweed

- Fronds short-stalked or stalkless, floating. *L. minor*
- Fronds long-stalked, usually submerged. *L. trisulca*

Lemna minor L.

LESSER DUCKWEED

Small, floating fronds 3–5 mm across, found floating in large quantities in still water. Common; throughout the Prairie Provinces in suitable locations.

Lemna trisulca L.

IVY-LEAVED DUCKWEED

Small oblong or lanceolate fronds tapering to a narrow stalk, and found floating at various depths in the water. Fairly common; throughout the Prairie Provinces.

Spirodela larger duckweed

Spirodela polyrhiza (L.) Schleid.

LARGER DUCKWEED

Found floating solitary or in small groups on the surface of water. Has 5–11 rootlets hanging from beneath fronds. Fairly common; in Parklands, Boreal forest.

ARACEAE—arum family

Plants with flowers closely borne in a dense spike called a spadix and with a large leaf-like bract called a spathe at the back of, or sometimes partly enclosing, the spadix.

Leaf blades linear; spathe green and leaf-like. *Acorus*

Leaf blades ovate and cordate; spathe white
and petal-like. *Calla*

Acorus calamus

Acorus calamus L.

SWEET FLAG

Perennial herbs growing from stout rhizomes to 40–100 cm high with long, narrow sword-like leaves. Inflorescence a spike-like spadix 4–10 cm long, borne at an angle with the stem and with a leaf-like spathe projecting 3–10 cm beyond the inflorescence. Plant has a peculiar taste and odor. Found in swamps and along water courses; Boreal forest.

Calla water-arum

Calla palustris L.

WATER CALLA

Low-growing perennial herb from long rhizomes, rooting at the nodes. The long-stalked leaves are usually broadly ovate, cordate-based, 5–10 cm long on stalks 7–20 cm long. Inflorescence, borne on a long stalk, a spike-like spadix 1.5–2.5 cm long, backed by an oval white spathe 2.5–7.0 cm long. Fruits red berries, each containing a few seeds. Found in swamps and shallow water; Boreal forest.

COMMELINACEAE—spiderwort family

Tradescantia spiderwort

Tradescantia occidentalis (Britt.) Smyth.

WESTERN SPIDERWORT

Plants with slender stems, 20–60 cm high; the leaves linear, less than 1 cm wide, involute. Cymes solitary, terminal; sepals 6–10 mm long; petals 10–15 mm long, rose to blue; pedicels and sepals glandular pubescent. Very rare; dry grassland; southeastern Parkland.

JUNCACEAE—rush family

Grass-like plants growing from either fibrous roots or rhizomes, and sometimes bearing round or stem-like leaves. Flowers perfect, regular, with 3 sepals and 3 petals, both very similar and scale-like. Flowers borne on top of the stem, but the projecting stem-like bract often makes the flowers appear to be on the side of the stem. The fruit, a capsule, splitting at maturity.

- Capsule containing many small seeds; plants glabrous. *Juncus*
- Capsule containing 3 large seeds; plants with pubescent leaves. *Luzula*

Juncus rush

- 1. Bract of the inflorescence terete, appearing as a continuation of the stem; inflorescence appearing lateral. Group 1
- Bract of the inflorescence flat or channeled; inflorescence appearing terminal or lateral. 2
- 2. Leaves not septate, terete, or flat with the flat side toward the stem. Group 2
- Leaves septate, terete, or flat with the edge of the leaf toward the stem. Group 3

Group 1

- 1. Inflorescence few-flowered, usually only 2 or 3 flowers; seeds with tail-like appendages. 2
- Inflorescence usually with several to many flowers; seeds without tail-like appendages. 3
- 2. Blade of the uppermost leaf sheath well-developed; capsule acute. *J. parryi*
- Blade of the uppermost leaf sheath greatly reduced; capsule retuse. *J. drummondii*
- 3. Seeds about 0.5 mm long; flowers greenish; rhizomes slender. *J. filiformis*
- Seeds about 1 mm long; flowers brown to blackish; rhizomes stout. 4
- 4. Inflorescence many-flowered; anthers 2–4 times as long as the filament. *J. balticus*
- Inflorescence few-flowered; anthers about as long as the filaments. *J. arcticus*

Group 2

- 1. Flowers borne singly in the inflorescence, each one with a pair of bractlets. 2
- Flowers borne in heads with several flowers. 7
- 2. Plants annual; inflorescence diffuse, about half the height of the plant. *J. bufonius*
- Plants perennial; inflorescence much less than half the height of the plant. 3

3. Sepals obtuse, the tips curved inward; leaf sheaths extending halfway up the culm; rhizomes spreading. *J. compressus*
 Sepals acute, the tips not curved inward; leaf sheaths confined to the lower one-third of the culm; rhizomes erect. 4
4. Seeds with a tail-like appendage at each end; leaves somewhat channeled. *J. vaseyi*
 Seeds without appendages; leaves flat or involute. 5
5. Capsule completely 3-celled. *J. confusus*
 Capsule not completely 3-celled, with the partitions reaching less than halfway. 6
6. Perianth 4–6 mm long; leaf blades mostly less than half as long as the culm; auricles short, yellowish. *J. dudleyi*
 Perianth 3–4.5 mm long; leaf blades mostly more than half as long as the culm; auricles long, whitish. *J. tenuis*
7. Seeds without tail-like appendages. *J. longistylis*
 Seeds with tail-like appendages. 8
8. Plants rhizomatous; heads 1–3, each with 6–8 flowers. *J. castaneus*
 Plants tufted; heads solitary, with 1–6 flowers. 9
9. Perianth dark brown to blackish; the bract leaf-like, as long as or slightly longer than the inflorescence. *J. biglumis*
 Perianth whitish to pinkish or light brown; the bract scale-like. *J. albescens*

Group 3

1. Leaves flat, with the edge toward the stem; the septa incomplete. 2
 Leaves terete or occasionally somewhat flattened; the septa complete. 4
2. Seeds with tail-like appendages. *J. tracyi*
 Seeds without tail-like appendages. 3
3. Stamens 3; stems strongly flattened; sheaths not auricled. *J. ensifolius*
 Stamens 6; stems not strongly flattened; sheaths mostly auricled. *J. saximontanus*
4. Heads spherical, short-peduncled; stamens 6; capsules lanceolate. 5
 Heads at most hemispherical, long-peduncled; stamens 3 or 6; capsules oblong. 6

5. Sepals longer than the petals; heads 10–15 mm in diam with 25–80 flowers. *J. torreyi*
 Sepals as long as the petals; heads 7–10 mm in diam with 10–20 flowers. *J. nodosus*
6. Seeds with tail-like appendages. 7
 Seeds without tail-like appendages. 9
7. Heads usually solitary, densely many-flowered; perianth about 4 mm long, dark brown. *J. mertensianus*
 Heads few to many in a more or less diffuse inflorescence; perianth 2.5–5 mm long, color various. 8
8. Perianth about 2.5 mm long; seeds short-tailed, 0.7–1.2 mm long, the body longer than the tails. *J. brevicaudatus*
 Perianth 4–5 mm long; seeds long-tailed, 1.2–2 mm long, the body shorter than the tails. *J. canadensis*
9. Branches of inflorescence erect or stiffly ascending; plants in small tufts from slender rhizomes; capsule straw-colored or light brown. *J. alpinus*
 Branches of inflorescence loosely ascending or spreading; plants densely tufted from short rhizomes; capsule dark shiny brown. *J. articulatus*

Juncus albescens (Lange) Fern.

WHITE RUSH

Plants loosely tufted, with stems 5–15 cm high, very slender; leaves 1–5 cm long, narrow. Inflorescence a single head, 1- to 5-flowered; lower bract long-acuminate or awned as long as or longer than the first flower; perianth whitish or pinkish; sepals 3–4 mm long; capsule about 3.5 mm long; seeds 1.5–2 mm long, smooth, brown, with tails shorter than the body. Moist areas; Rocky Mountains.

Juncus alpinus Vill.

ALPINE RUSH

Plants with creeping rhizomes; stems 15–50 cm high, in small tufts; stem leaves 1 or 2, seldom reaching the inflorescence, narrow. Inflorescence a cyme 5–15 cm long; glomerules 3- to 12-flowered, obpyramidal; perianth segments 2–3 mm long; capsule 2.5–3.5 mm long. Wet meadows, bogs, and lakeshores; Parklands, Boreal forest.

Juncus arcticus Willd.

ARCTIC RUSH

Plants with stout rhizomes; stems 10–50 cm high, stout. Inflorescence a cyme, few-flowered, with branches mostly simple; the bract appearing like a continuation of the stem; perianth segments 2.5–3.5 mm long, dark brown; capsule about 3 mm long; seeds about 1 mm long. Sedge tundra; Boreal forest.

Juncus articulatus L.

JOINTED RUSH

Plants tufted, with slender stems 20–60 cm high; stem leaves usually 2–4, mostly 3–8 cm long, strongly septate. Inflorescence a cyme 3–10 cm long, with branches ascending to divergent; glomerules 3- to 11-flowered; perianth segments 2–3 mm long, brown; capsule 2.5–4 mm long, chestnut or purple brown. Bogs, wet meadows, lakeshores; Boreal forest.

Juncus balticus Willd.

BALTIC RUSH

Plants with long-creeping, thick rhizomes; stems 20–60 cm high, in more or less regular rows from the rhizomes, or singly; basal sheath 8–15 cm long, bladeless. Inflorescence appearing lateral, cymose or densely head-like, mostly 2–8 cm long, occasionally to 12 cm long; perianth 4–5 mm long, purplish brown; capsule about 5 mm long, chestnut brown; seeds 0.8–1 mm long. Wet meadows, slough margins, sandhills, lakeshores, often in great abundance. A variable species in which several varieties have been distinguished. Slender plants with a small dense inflorescence have been named var. *montanus* Engelm. (Fig. 79); stout plants with a more open inflorescence, 4–8 or 12 cm long, are var. *littoralis* Engelm. Both varieties occur throughout the Prairie Provinces.

Juncus biglumis L.

TWO-GLUMED RUSH

Plants tufted, with stems 5–20 cm high; leaves basal, filiform. Inflorescence a single head; the lowest bract mostly well-developed; perianth 3–3.5 mm long; capsule 3.5 mm long, purplish black; seeds about 1.5 mm long, with white tails. Tundra; Hudson Bay, Boreal forest.

Juncus brevicaudatus (Engelm.) Fern.

SHORT-TAIL RUSH

Plants densely tufted, with stems 10–50 cm high, slender; leaves 1–2 mm in diam, erect. Inflorescence 3–12 cm long, narrow, with branches ascending to erect; glomerules few to many, each 2- to 7-flowered; perianth segments 2.5–3 mm long, 3-nerved; capsule 3.5–4.5 mm long; seeds about 1 mm long, with short tails. Marshes and lakeshores; Boreal forest, Parklands.

Juncus bufonius L.

TOAD RUSH

Plants annual, with stems 3–20 cm high, tufted, erect or spreading; leaves about 1 mm wide. Inflorescence 2–10 cm long, spreading, freely branching; flowers singly or in twos or threes; perianth segments unequal, with sepals 3.5–6.5 mm long, petals 3–5.5 mm long. Common in low, wet ground; throughout the Prairie Provinces.

Juncus canadensis J. Gay

CANADA RUSH

Plants tufted, with stems 40–100 cm high, stout and rigid; leaves 1.5–2.5 mm in diam, erect. Inflorescence 2–20 cm long, varying from compact to loosely branching; glomerules few to many, 2- to 10-flowered; perianth segments 2.5–4 mm long; capsule 3.5–5 mm long, brown; seeds about 1 mm long, with white tails. Rare; marshes and wet meadows; northeastern Parkland and Boreal forest.



Fig. 79. Baltic rush, *Juncus balticus* Willd. var. *montanus* Engelm.

Juncus castaneus Sm.

CHESTNUT RUSH

Plants with slender rhizomes; stems 20–40 cm high, erect; leaves 1–2 mm wide, basal. Inflorescence 2–5 cm long; the involucral leaf 2–8 cm long; glomerules 1–3, the lowest ones sessile or nearly so, the upper one peduncled, 10–18 mm in diam; perianth segments 5–7 mm long, chestnut brown; capsule 6–9 mm long, with a conspicuous beak; seeds 2.5–4 mm long, with long slender tails. Arctic and alpine meadows and tundra; Boreal forest, Rocky Mountains.

Juncus compressus Jacq.

FLATTENED RUSH

Plants with slender rhizomes; stems 20–60 cm high, erect, tufted; leaves 1 mm wide, erect. Inflorescence 2–10 cm long, cymose, with branches ascending, bearing many-flowered glomerules; perianth segments 2–2.5 mm long, purple brown; capsule 2–3 mm long; seeds about 1 mm long. Introduced; wet ground, salt marshes; Manitoba.

Juncus confusus Cov.

FEW-FLOWERED RUSH

Plants tufted, with stems 30–50 cm high, slender; leaves filiform. Inflorescence 5–20 mm long, compact, pale; perianth segments 3.5–4 mm long, brown with a yellowish midrib and narrow margins; capsule 3–3.5 mm long; seeds about 1 mm long. Moist grassland, open woods, and meadows; southern Rocky Mountains.

Juncus drummondii E. Mey.

DRUMMOND'S RUSH

Plants densely tufted, with the base matted, persistent; stems 10–30 cm high, erect; sheaths short, bladeless, or the blade bristle-like. Inflorescence 1- to 5-flowered; the bract appearing like a continuation of the stem; perianth segments 5–7 mm long, brown with a light midrib; capsule 5–7 mm long, triangular; seeds about 2 mm long, tailed at both ends. Moist places; Rocky Mountains.

Juncus dudleyi Wieg.

DUDLEY'S RUSH

Plants densely or loosely tufted, with stems 30–60 cm high; leaves basal, flat, 10–30 cm long; auricles of sheaths rounded, papery, yellowish. Inflorescence 2–5 cm long, with the few branches loosely ascending; perianth segments 3.5–5.5 mm long; capsule 3–4 mm long; seeds about 0.4 mm long. Moist or dry soil, waste areas; throughout the Prairie Provinces. Syn.: *J. tenuis* Willd. var. *dudleyi* (Wieg.) Hermann.

Juncus ensifolius Wikstr.

EQUITANT-LEAVED RUSH

Plants tufted, with stems 30–60 cm high, flattened, sharply 2-keeled; leaves 2–6 mm wide, flat. Inflorescence 15–70 mm long, with 2–6 heads, about 1 cm thick, many-flowered, brownish; perianth segments about 3 mm long, dark brown; capsule about 3 mm long; often somewhat exceeding the perianth, dark brown; seeds about 0.3 mm long. Riverbanks, lakeshores; Rocky Mountains, Cypress Hills.

Juncus filiformis L.

THREAD RUSH

Plants with slender rhizomes; stems 10–40 cm high, arising from the rhizomes in rows, or in small tufts; leaves short, thick. Inflorescence 2–4 cm long,

sparingly branched, few-flowered; involucre leaf erect, to 20 cm long; perianth segments 2.5–4.5 mm long, greenish, with hyaline margins; capsules 2.5–4 mm long; seeds about 0.3 mm long. Lakeshores, bogs, and alpine meadows; Boreal forest, rare in northeastern Parkland.

Juncus longistylis Torr.

LONG-STYLED RUSH

Plants with slender rhizomes; stems solitary, 20–60 cm high; leaves mostly basal, 1–4 mm wide, flat; stem leaves short, flat or involute. Inflorescence 2–3 cm long, with 1–5 heads 8–15 mm across, 2- to 8-flowered; perianth segments 4.5–6 mm long, greenish brown; capsule 4–5 mm long, brownish; seed about 0.5 mm long, tailed at both ends. Fairly common; wet meadows, lakeshores, and riverbanks; Prairies, Parklands, Boreal forest.

Juncus mertensianus Bong.

SLENDER-STEMMED RUSH

Plants with matted rhizomes; stems 10–40 cm high, densely tufted; leaves 1–2 mm wide, compressed, septate. Inflorescence usually a solitary head, occasionally 2 or 3 heads, dark brown, almost spherical; perianth segments 3.5–4.5 mm long, dark brown; capsule 3.5–4.5 mm long, dark brown; seeds 1 mm long, reticulate. Rare; wet ground, meadows, and slopes; Rocky Mountains, Cypress Hills.

Juncus nodosus L.

KNOTTED RUSH

Plants with slender rhizomes; stems 20–60 cm high, erect; leaves to 10 cm long, and 1.5 mm thick, septate. Inflorescence a cyme 2–5 cm long, contracted or more or less open with 2–20 globose heads, 7–12 mm in diam, 10- to 25-flowered; perianth segments 2.5–4 mm long, reddish brown; capsule 3.5–4.5 mm long; seeds 1.5 mm long. Bogs, marshes, and lakeshores; Prairies, Parklands, Boreal forest.

Juncus parryi Engelm.

PARRY'S RUSH

Plants densely tufted, with stems 10–30 cm high, slender; leaves only from the upper sheaths. Inflorescence a single head, 1- to 3-flowered; perianth segments 6–7 mm long; capsule about 7 mm long; seeds about 0.5 mm long, tailed at both ends. Mountain slopes and meadows; Rocky Mountains.

Juncus saximontanus A. Nels.

ROCKY MOUNTAIN RUSH

Plants with stout creeping rhizomes; stems 20–50 cm high, compressed; leaves 2–4 mm wide, flat, with the edge toward the stem, 10–20 cm long. Inflorescence 6–10 cm long, with 2–12 heads, 4–6 mm across, few- to many-flowered; perianth segments 2.5–3.5 mm long, dark brown; capsule about 3 mm long; seeds 0.5 mm long, reticulate. Marshes and wet areas; Rocky Mountains, Cypress Hills.

Juncus tenuis Willd.

SLENDER RUSH

Plants loosely or densely tufted, with stems 10–60 cm high, mostly erect; leaves mostly basal, 10–25 cm long, flat or involute; the sheaths conspicuously long auricled. Inflorescence a cyme 2–8 cm long, commonly exceeded by the involucre leaf, compact to open; perianth segments 3–5 mm long; capsule 3–4 mm long, straw-colored; seeds about 0.5 mm long. Moist to wet areas; Prairies, Parklands, Boreal forest.

Plants with slender tuberous-thickened rhizomes; stems 40–80 cm high, stout, rigid, erect; leaves 1–3 mm thick, 10–30 cm long; sheaths with auricles 2.5–3.5 mm long. Inflorescence 2–4 cm long, with 2 to many heads, usually crowded; heads 10–15 mm in diam, globose; 25- to 100-flowered; perianth segments unequal, with the petals 4.5–5 mm long, the sepals 3.5–5 mm, linear-lanceolate; capsule 4.5–5.5 mm long, narrowly lanceolate; seeds about 0.3 mm long. Wet areas; Prairies, Parklands.

Plants with stout rhizomes; stems 30–60 cm high, stout, compressed; leaves 5–20 cm long, 2–4 mm wide. Inflorescence 4–8 cm long, with 5–9 heads, 8–10 mm in diam; perianth segments 3–4 mm long, light brown; capsule about 3–3.5 mm long, brown; seeds 0.5 mm long, reticulate. Rare; wet areas; northern Rocky Mountains.

Plants with short rhizomes; stems 30–80 cm high, erect, tufted; basal leaves to 30 cm long, terete or nearly so. Inflorescence 1–4 cm long, compact; perianth segments 3.5–4.5 mm long, brown; capsule 4–5.5 mm long, truncate, straw-colored; seeds 0.5–0.8 mm long, conspicuously tailed at both ends. Moist soil, meadows, and lakeshores; Prairies, Parklands, Boreal forest.

Luzula wood-rush

Similar to *Juncus* but capsule 1-celled with 3 seeds; foliage more or less pilose.

- 1. Flowers solitary or in pairs at the end of branches of the inflorescence. 2
Flowers few to many in glomerules. 5
- 2. Seeds with a tail-like appendage about as long as the body. *L. pilosa*
Seeds without a tail-like appendage. 3
- 3. Perianth about 3 mm long; panicle branches ascending. *L. glabrata*
Perianth about 1–2 mm long; panicle branches spreading to drooping. 4
- 4. Leaves 4–10 mm wide; usually 3–6 leaves on the plant, shiny green. *L. parviflora*
Leaves 2–5 mm wide; usually 1 or 2 leaves on the plant, dull green. *L. wahlenbergii*
- 5. Seeds without a tail-like appendage, but with a tuft of minute hairs. *L. confusa*
Seeds with a tail-like appendage, without hairs. 6
- 6. Appendage well-developed; perianth segments 2.5–3.5 mm long. *L. campestris*

Appendage of seeds small; perianth segments 2–2.5 mm long. 7

7. Sepals ciliate on the margins; leaves 1–4 mm wide, subulate at the tip. *L. spicata*

Sepals with entire or lacerate margins; leaves 2–6 mm wide, blunt, callus-tipped. *L. groenlandica*

Luzula campestris (L.) DC.

FIELD WOOD-RUSH

Plants tufted, with stems 20–40 cm high; leaves 2–6 mm wide, flat; cauline leaves 2 or 3, short. Inflorescence 3–5 cm long, with slender ascending peduncles, and one or more sessile glomerules; perianth segments 2.5–3.5 mm long; capsule 3.5 mm long; seeds 1–1.3 mm long, with the appendage 0.4–0.5 mm. Dry to moist woods; Boreal forest, Cypress Hills.

Luzula confusa Lindeb.

NORTHERN WOOD-RUSH

Plants densely tufted, with stems 10–30 cm high; leaves 1–3 mm wide; stem leaves 2 or 3, short. Inflorescence 3–5 cm long, with 2–5 glomerules on slender peduncles, or occasionally the glomerules crowded into a short spike; perianth segments 2.5–3 mm long, brown; capsule 3 mm long, red brown; seeds 1–1.2 mm long, with a tuft of minute hairs. Gravel ridges and barrens; Hudson Bay, Boreal forest, also reported from Rocky Mountains.

Luzula glabrata (Hoppe) Desv.

SMOOTH WOOD-RUSH

Plants with scaly rhizomes; stems 20–50 cm high; leaves 4–10 mm wide, with stem leaves well-developed. Inflorescence 6–10 cm long, with the branches ascending or spreading; perianth segments 3–3.5 mm long, dark brown; capsule about 3 mm long, almost black; seeds 1–1.3 mm long, with a tuft of hairs. Meadows and slopes; southern Rocky Mountains.

Luzula groenlandica Boecher

GREENLAND WOOD-RUSH

Plants tufted; stems 20–50 cm high; leaves 3–7 mm wide, flat, with stem leaves 2 or 3, short. Inflorescence 3–5 cm long; perianth segments 2–2.5 mm long; capsule 2.5–3 mm long, dark brown; seeds about 1 mm long, with the appendage short. Tundra; Hudson Bay.

Luzula parviflora (Ehrh.) Desv.

SMALL-FLOWERED WOOD-RUSH

Plants with creeping rhizomes; stems 20–60 cm high, erect; leaves 5–10 mm wide. Inflorescence 5–12 cm long, with branches loosely spreading to drooping; perianth segments 1.5–2.5 mm long, pale brown; capsule 2–2.5 mm long, dark brown; seeds 1–1.4 mm long, with a tuft of long hairs. Infertile soil, open forests; Boreal forest, Rocky Mountains.

Luzula pilosa (L.) Willd. var. *saltuensis* (Fern.) Boiv.

HAIRY WOOD-RUSH

Plants tufted from short rhizomes; stems 10–40 cm high; leaves 3–10 mm wide, and up to 30 cm long, with stem leaves 2–4, shorter and narrower than the basal leaves. Inflorescence 3–6 cm long, with branches loosely spreading; perianth segments 2.5–4.5 mm long, the center chestnut brown, with narrow margins; capsule 3–4.5 mm long, brown; seeds about 1 mm long, with the appendage about equaling the body. Open woods; Boreal forest.

Plants densely tufted, with stems 10–30 cm high, slender; leaves 1–4 mm wide; stem leaves 1–3, short. Inflorescence 1–3 cm long, spike-like, often nodding, with the glomerules dense, sessile; perianth segments 2–2.5 mm, bristle-pointed, brown; capsule 2.5 mm long, red brown; seeds 1–1.2 mm long, with a short appendage. Arctic and alpine meadows; Boreal forest, Rocky Mountains.

Plants with rhizomes; stems 20–60 cm high, erect; leaves 3–8 mm wide; stem leaves well-developed, thick, dull. Inflorescence 5–10 cm long, with the slender branches spreading and drooping; perianth segments 1.5–2.5 mm long, pale; capsule 2–2.5 mm long, dark brown; seeds 1–1.3 mm long, gray brown to yellow with a tuft of long hairs. Meadows and open forests; southern Rocky Mountains.

LILIACEAE—lily family

Perennial herbs growing from bulbs, corms, or rootstocks. With a few exceptions, the leaves are parallel-veined. Except in the *Maianthemum* genus, the flowers have 3 sepals and 3 petals, all colored, and 6 stamens in 2 whorls. The fruit is a capsule or a berry.

- 1. Leaves reduced to thin, dry scales; branchlets filiform; flowers small, greenish. *Asparagus*
Leaves normal, not reduced to scales. 2
- 2. Plants climbing with tendrils; leaves roundish, net-veined, flowers in axillary umbels. *Smilax*
Plants not climbing. 3
- 3. Leaves linear, many times as long as wide. 4
Leaves variously shaped, not more than five times as long as wide. 12
- 4. Flowers orange, large, 5–7 cm long. *Lilium*
Flowers not orange, smaller. 5
- 5. Inflorescence an umbel, head-like cluster, or few-flowered. 6
Inflorescence a many-flowered raceme. 7
- 6. Inflorescence few-flowered; flowers yellowish white, large, to 25 mm long. *Calochortus*
Inflorescence an umbel or head-like cluster; flowers 5–15 mm long. *Allium*
- 7. Flowers purplish blue, to 2 cm long. *Camassia*
Flowers not purplish blue. 8

8. Flowers greenish or reddish brown, 10–15 mm long, nodding on slender pedicels. *Stenanthium*
 Flowers some shade of white, in erect racemes. 9
9. Flowers large, 3–5 cm long; leaves with a rigid point at the tip. *Yucca*
 Flowers smaller, to 1.5 cm long; leaves not with a rigid point. 10
10. Racemes large, densely flowered; leaves rough; stems to 1.2 m high. *Xerophyllum*
 Racemes slender, loosely flowered; leaves smooth; plants not more than about 0.6 m high. 11
11. Leaves keeled, V-shaped in cross section; flowers 5–15 mm long. *Zygadenus*
 Leaves flat; flowers to 5 mm long. *Tofieldia*
12. Flowers solitary or in umbels, terminal. 13
 Flowers axillary, in panicles or in racemes. 17
13. Stems leafy. 14
 Stems leafless. 16
14. Flowers white or pinkish. *Trillium*
 Flowers orange yellow or straw yellow. 15
15. Flowers nodding; plants with a deep-seated bulb. *Fritillaria*
 Flowers erect or nodding; plants with rhizomes. *Uvularia*
16. Flowers bright yellow, nodding; fruit a capsule; leaves glabrous. *Erythronium*
 Flowers white or greenish white; fruit a bluish berry; leaves pilose beneath. *Clintonia*
17. Flowers in a large, diffuse panicle. *Veratrum*
 Flowers axillary or in racemes, or at tips of branches. 18
18. Flowers 1–4 at the tip of the stem or branches; fruit a red or orange berry. *Disporum*
 Flowers axillary or in racemes. 19
19. Flowers axillary. 20
 Flowers in racemes. 21
20. Perianth segments free at the tip only. *Polygonatum*
 Perianth segments free to the base. *Streptopus*
21. Plants small, usually with only 2 or 3 leaves; raceme small. *Maianthemum*
 Plants large, with leafy stems; racemes simple or paniculate. *Smilacina*

Allium onion

Strong-scented herbs growing from a bulbous root, which is generally covered with a fibrous coat. Leaves linear and narrow, flowers brightly colored and borne in umbels. Fruit a capsule containing dark seeds.

- 1. Flower stalks shorter than individual flowers; leaves circular in cross section and hollow. *A. schoenoprasum*
- Flower stalks longer than individual flowers; leaves not hollow. 2
- 2. Umbel nodding or drooping. *A. cernuum*
- Umbel erect. 3
- 3. Flowers rose-colored. *A. stellatum*
- Flowers white or pale pink. *A. textile*

Allium cernuum Roth

NODDING ONION

Growing from coarse-necked bulbs, usually on a short rhizome; flower heads rose or rarely white. Found occasionally; in grassy openings; Parklands.

Allium schoenoprasum L. var. *sibiricum* (L.) Hartm.

WILD CHIVES

Perennial from small bulbs, 20–50 cm high, with many hollow leaves up to 3 mm thick and circular in cross section. Inflorescence a dense, compact umbel 2–5 cm across; flower stalks very short and flowers bright rose pink. Not common; moist areas, meadows, and coulees; Rocky Mountains, Boreal forests.

Allium stellatum Fraser

PINK-FLOWERED ONION

Erect umbels of pink or rose flowers from a membranous-coated bulb. Fairly common; in wooded lands; throughout Parklands.

Allium textile Nels. & Macbr. (Fig. 80)

PRAIRIE ONION

A low plant, from an onion-like bulb having a net-like coating and a very pungent odor. From 8–25 cm high, with umbels of white or pale pink flowers, which blossom early in season. Common; on dry prairie; throughout Prairies, Parklands.

Asparagus asparagus

Asparagus officinalis L.

ASPARAGUS

Plants with succulent rhizomes; stems 60–150 cm high; branches filiform; leaves scaly. Flowers small, yellowish green, axillary, with the pedicels jointed; fruit a red berry. Introduced, occasionally escaped from cultivation.

Calochortus mariposa lily

Calochortus apiculatus Baker

MARIPOSA LILY

Plants with ovoid bulbs and membranous scales; stems 10–30 cm high; leaf solitary, basal, 5–15 mm wide. Flowers 1–4, yellowish white, 3–5 cm across; the sepals shorter than the petals; stamens 6; stigmas 3; fruit a 3-angled capsule. Dry slopes, open woods; southern Rocky Mountains.




Fig. 80. Prairie onion, *Allium textile* Nels. & Macbr.

Camassia camas

Camassia quamash (Pursh) Greene

COMMON CAMAS

Herbaceous perennial plants from an ovoid bulb 1–3 cm broad, the leaves mostly basal, narrow, and grass-like, 20–40 cm long. Blue flowers with 6 petal-like segments 10–25 mm long, borne in a raceme at the summit of scapose stems 30–60 cm high. Moist meadows; in southern Rocky Mountains.

Clintonia

Flowers white, usually solitary. *C. uniflora*

Flowers yellowish green, in 3- to 8-flowered umbel. *C. borealis*

Clintonia borealis (Ait.) Raf.

BLUEBEAD LILY

Plants with slender rootstocks; scapes 15–40 cm high; leaves 2–5, to 30 cm long, obovate to oblong-elliptic, glossy green. Flowers on pedicels 1–3 cm long; perianth segments 15–18 mm long, greenish yellow, nodding; fruit a blue berry, 8 mm in diam. Moist woods; southeastern Boreal forest.

Clintonia uniflora (Schult.) Kunth

CORN LILY

Plants with slender rootstocks; scapes 6–20 cm high; leaves 2–5, to 20 cm long, oblanceolate, pubescent below. Flowers showy; perianth segments about 1 cm long, white, villose; fruit a deep blue berry 8–10 mm in diam. Moist woods; southern Rocky Mountains.

Disporum fairybells

Stigma 3-lobed; flowers solitary or in pairs. *D. trachycarpum*

Stigma entire; flowers in clusters of 1–4. *D. hookeri*

Disporum hookeri (Torr.) Britt. var. *oreganum* (Wats.) Q. Jones

OREGON FAIRYBELLS

Plants 30–80 cm high, with leaves ovate to oblong-lanceolate, 3–9 cm long, 2–6 cm wide, cordate-based. Flowers 1–4 in a cluster, 10–12 mm long, creamy white or tinged with green; stigma entire; berry ovoid, 8–16 mm in diam, 6-seeded. Moist woods; southern Rocky Mountains.

Disporum trachycarpum (S. Wats.) B. & H. (Fig. 81A)

FAIRYBELLS

Plants 30–60 cm high, with leaves ovate to oblong-lanceolate, 3–8 cm long, 2–5 cm wide, short pubescent, rounded or subcordate at the base. Flowers solitary or in pairs, occasionally 3 together, 8–14 mm long, creamy white; stigma 3-lobed; berry depressed globose, 8–10 mm in diam, 4- to 18-seeded. Moist woods, ravines, and coulees; throughout Prairies, Parklands, southern parts of Boreal forest.




Fig. 81. A, Fairybells, *Disporum trachycarpum* (S. Wats.) B. & H.; B, two-leaved Solomon's-seal, *Maianthemum canadense* Desf. var. *interius* Fern.; C, clasping-leaved twistedstalk, *Streptopus amplexifolius* (L.) DC.

Erythronium fawn lily

Erythronium grandiflorum Pursh (Fig. 82)

GLACIER LILY

Plants with deep-seated corm, arising from a rootstock; scape 20–30 cm high. Leaves 7–15 cm long, oblong-lanceolate. Flowers solitary or in a 2- to 6-flowered raceme; perianth segments 3–5 cm long, bright yellow, recurved; capsule 3–5 cm long. Open forests, hillsides, and grassy slopes; southern Rocky Mountains.

Fritillaria fritillary

Fritillaria pudica (Pursh) Spreng.

YELLOWBELL

A short plant, 10–30 cm high, from a very scaly bulb. Leaves about 8 cm long and up to 6 mm wide, somewhat whorled about halfway up the stem. Flower, nodding and single, bell-shaped, yellow or orange, on the summit of the stem, about 15–20 mm high. Very rare; on moist banks; southern Rocky Mountains. Syn.: *Ochrocodon pudicus* (Pursh) Rydb.

Lilium lily

Lilium philadelphicum L.

WOOD LILY

Very showy, erect plants 20–60 cm high, from whitish, scaly bulblets. Leaves linear, in whorls. Flowers showy, 8 cm long, sepals and petals red or orange with black spots, 1–5 flowers on a plant. Found in the eastern part of the Prairie Provinces, but westward giving place to the western red lily, *L. philadelphicum* L. var. *andinum* (Nutt.) Ker (Fig. 83), having the lower leaves alternate and the upper ones whorled. The variety is found in moist meadows throughout the entire area, but is becoming scarcer with the advance of settlement and overpicking. Plants found in the Foothills region have wider, lanceolate leaves and were formerly considered a separate species.

Maianthemum wild lily-of-the-valley

Maianthemum canadense Desf. var. *interius* Fern. (Fig. 81B)

TWO-LEAVED SOLOMON'S-SEAL


Low, erect plant, 5–15 cm high, with 2, sometimes 3, ovate leaves, cordate-based, borne alternately on the stem. Flowers small, white, in a rather dense raceme. Berries pale red, speckled, about 5 mm in diam. Fairly common; in rich, moist woods; throughout the Prairie Provinces.

Polygonatum Solomon's-seal

Polygonatum canaliculatum (Muhl.) Pursh

COMMON SOLOMON'S-SEAL

Coarse plants, from jointed rootstocks, 30–100 cm high. Leaves large, alternate, up to 15 cm long and to 10 cm wide. White, cylindric flowers suspended in little bunches below the stem, about 15 mm long. Berries dark blue, 8–12 mm in diam. Fairly common; in open woodlands in the eastern part. Syn.: *P. commutatum* (R. & S.) Dietr.



HERBARIUM OF THE RESEARCH STATION
 SWIFT CURRENT, SASK.

SPECIES: *Erythronium grandiflorum* Pursh.

LOCATION: 111119, Riverbank, Sask.

DATE: 21. 10. 1950

COLLECTOR: J. R. ...

AT: ...

REMARKS: ...



Fig. 82. Glacier lily, *Erythronium grandiflorum* Pursh.



Fig. 83. Western red lily, *Lilium philadelphicum* L. var. *andinum* (Nutt.) Ker.

Smilacina false spikenard

Plants from rootstocks, with alternate, ovate to lanceolate leaves, and small, perfect, greenish white flowers borne in racemes or panicles. Fruits small berries.

1. Inflorescence a branched, dense panicle. *S. racemosa*
Inflorescence an unbranched raceme. 2
2. Leaves 6–12; berries green. *S. stellata*
Leaves 2–4; berries dark red. *S. trifolia*

Smilacina racemosa (L.) Desf.

Large-leaved plants, 30–90 cm high. Leaves elliptic or oval, up to 15 cm long and to 7 cm wide, parallel-veined, without stalks, except lower ones, which may have very short stalks. White flowers in a densely branched panicle at the end of the stem. Berries red, purple-specked, up to 6 mm across. Not common; found in moist woods; throughout southeastern Boreal forest, Riding Mountain. Replaced in western Boreal forest, Parklands, Rocky Mountains, and Cypress Hills by *S. racemosa* (L.) Desf. var. *amplexicaulis* (Nutt.) S. Wats., having all leaves sessile and clasping.

Smilacina stellata (L.) Desf.

STAR-FLOWERED SOLOMON'S-SEAL

Erect, low plants, 15–50 cm high. Leaves arranged on opposite sides of the stem, folded at the base, overlapping each other in early stages, but spreading and flattening out later. Flowers small, white, in a spike-like raceme on the stem. Fruits green berries, each with 6 black stripes. Very common; in moist soil, meadows, woods, and low areas; throughout the Prairie Provinces. Readily eaten by livestock.

Smilacina trifolia (L.) Desf.

THREE-LEAVED SOLOMON'S-SEAL

Short, slender plants from a thin rootstock, growing to 5–30 cm high and usually bearing 3 alternate, oblong-lanceolate leaves without stalks, but often sheathing the stem at their base. Flowers few, in a terminal raceme. Berries dark red, up to 6 mm in diam. Wet woods and bogs; Boreal forest.

Smilax carrionflower

Unlike most monocotyledons, this genus has netted-veined leaves. The parallel veins, however, are much more distinct and conspicuous than the others.

Smilax herbacea L. var. *lasioneura* (Hook.) DC.

CARRIONFLOWER

Plants climbing by tendrils, up to 1.5 m long. Leaves alternate, long-stalked, oval to ovate, cordate-based, up to 8 cm long and to 6 cm across. Flowers small, greenish, in many-flowered umbels. Fruits purplish berries. These plants derive their common name from the carrion-like odor of the flowers, which attracts flies that aid in fertilization. Fairly common; in shady, moist woodlands; throughout Prairies, Parklands.

Stenanthium stenanthium

Stenanthium occidentale A. Gray BRONZEBELLS

Bulbous-rooted herbs, with grass-like, linear, mostly basal leaves, 15–30 cm long. Flowers greenish purple, drooping, 10–15 mm long, in a raceme on a stem 30–50 cm high. Moist woods and slopes; Rocky Mountains.

Streptopus twistedstalk

Leaves clasping the stem; flowers greenish white. *S. amplexifolius*

Leaves sessile, not clasping the stem; flowers pink or rose. *S. roseus* var. *perspectus*

Streptopus amplexifolius (L.) DC. (Fig. 81C) CLASPING-LEAVED TWISTEDSTALK

Herbs, from rootstock, 30–100 cm high. Leaves ovate to lanceolate, 5–10 cm long and 3–6 cm wide, clasping the stem at their bases. Greenish white flowers, 8–12 mm long, on long twisted stalks, usually in pairs beneath the axils of the leaves. Fruits red berries, 10–15 mm long, on a long twisted stalk. Represented by var. *americanus* Schultes, as described. Moist woods, ravines, and thickets; Parklands, Boreal forest. Locally, var. *denticulatus* Fassett, with the leaf margins finely denticulate, has been found.

Streptopus roseus Michx. var. *perspectus* Fassett ROSE MANDARIN

Herbs with matted rootstocks; stems 30–80 cm high, simple or branched, finely pubescent. Leaves lanceolate, broadly rounded at the base, sessile, 5–9 cm long, 2–4 cm wide. Flowers about 1 cm long, rose or purplish, nodding. Not common; moist woods and thickets; southeastern Parklands, Boreal forest.

Tofieldia asphodel

Stem not hairy; flowers single in a short raceme. *T. pusilla*

Stem slightly hairy near top and with sticky glands; flowers in bunches of 3 in a short raceme. *T. glutinosa*

Tofieldia glutinosa (Michx.) Pers. STICKY ASPHODEL

Plants with slender rootstocks; stems 10–50 cm high, viscid above; basal leaves 5–20 cm long, 3–8 mm wide. Inflorescence 2–5 cm long; flowers white, with segments about 4 mm long; capsules 5–6 mm long, ovoid. Marshes, bogs, and riverbanks; Boreal forest, Rocky Mountains.

Tofieldia pusilla (Michx.) Pers. BOG ASPHODEL

Plants with short rootstocks; stems 5–20 cm high, slender, smooth; basal leaves 2–6 cm long, 1–3 mm wide. Inflorescence 5–20 mm long; flowers greenish white 1.5–2.5 mm long; capsules 2.5–3 mm long. Marshes, bogs, and riverbanks; Boreal forest, Rocky Mountains.

Trillium wakerobin

Flowers on recurved peduncles, to below the leaves. *T. cernuum* var. *macranthum*
Flowers on erect peduncles, above the leaves. *T. ovatum*

Trillium cernuum L. var. *macranthum* Eam. & Wieg. NODDING WAKEROBIN

Plants with short rootstocks; stems 20–40 cm high, slender; leaves rhombic-ovate, 6–10 cm long, 5–12 cm broad; flowers on peduncles 1–4 cm long, reflexed to below the leaves; perianth segment 15–25 mm long; petals white. Rare; moist woods; eastern Boreal forest.

Trillium ovatum Pursh WESTERN WAKEROBIN

Plants with short, stout rootstocks; stems 30–50 cm high; leaves rhombic-ovate, 6–10 cm long, 5–9 cm wide. Flowers on an erect peduncle 25–75 mm long; perianth segments 25–50 mm long; petals white, turning deep rose. Very rare; moist woods; southern Rocky Mountains.

Uvularia bellwort

Uvularia sessilifolia L. SMALL BELLWORT

Slender, smooth-stemmed herbs, 15–30 cm high, with stalkless, lanceolate-oblong leaves, 3–8 cm long. Flowers borne singly 12–30 mm long, pale greenish yellow. Fruit sharply 3-angled, a capsule about 25 mm long. Very rare; in shady woodlands; southeastern Boreal forest.

Veratrum false hellebore

Veratrum eschscholtzii Gray GREEN FALSE HELLEBORE

Plants with a short, thick rootstock; stems stout, very leafy, somewhat pubescent, 1–2 m high. Leaves oval to elliptic, 10–30 cm long, acute, short-petioled to sessile or clasping, strongly veined, and often plaited. Flowers numerous in a panicle 20–50 cm long; perianth greenish or yellowish green, 15–20 mm across; capsule ovoid, 20–25 mm long, to 12 mm thick, with many large flat seeds. Moist forests and wet places; southern Rocky Mountains.

Xerophyllum turkey-beard

Xerophyllum tenax (Pursh) Nutt. BEAR-GRASS

Plants in dense clumps; stems slender 20–150 cm high, with scattered ascending leaves; basal leaves 50–80 cm long, 3–6 mm wide, rigid, flat, with rough margins. Inflorescence a dense raceme 30 cm or more long; flowers creamy white, with the segments 6–9 mm long; fruit a small capsule 5–7 mm long. Rare; dry mountain slopes; southern Rocky Mountains.

Yucca yucca

Yucca glauca Nutt. YUCCA, SOAPWEED

Coarse plants with short, woody stems and stiff, narrow basal leaves with sharp, hard tips. Flowers cream-colored or greenish white, with petals and se-

pals 25–50 mm long, in a raceme surmounting a stem 20–60 cm high. Fruits capsules containing numerous black seeds in layers. This plant is fertilized only by the yucca moth, which feeds on it, thereby making the insect and the plant interdependent. Very rare; on dry slopes; Prairies.

Zygadenus camas

Plants with an onion-like bulbous root and long, linear, grass-like leaves, somewhat flattened in cross section. Flowers perfect, with sepals and petals very similar, in racemes or panicles. Fruit an ovoid capsule containing many seeds.

Flowers pale yellow, about 5 mm long. *Z. gramineus*
 Flowers greenish yellow or straw-colored,
 about 10 mm long. *Z. elegans*

Zygadenus elegans Pursh SMOOTH CAMAS

Grows 30–60 cm high; leaves pale green; flowers grayish white to greenish, in an open raceme. Very common; in moist places, saline meadows; throughout the Prairie Provinces. Syn.: *Anticlea elegans* (Pursh) Rydb.

Zygadenus gramineus Rydb. (Fig. 84) DEATH CAMAS

A low-growing early herb from an onion-like bulb, which is usually 6–15 cm below the soil surface. Leaves linear and grass-like, 10–20 cm high. **Very poisonous** to sheep and somewhat poisonous to cattle. Common; in draws, around grassy sloughs and uplands; throughout the south central and southwestern Prairies. Because of its very early spring growth, it is sought after by livestock, and, wherever dense stands occur, poisoning may result. Syn.: *Toxicoscordion gramineum* Rydb.

AMARYLLIDACEAE—amaryllis family

Hypoxis star-grass

Hypoxis hirsuta (L.) Coville YELLOW STAR-GRASS

Plants 10–20 cm high, from a bulbous root. Leaves very narrow, grass-like, slightly hairy. Flowers perfect, 3 or 4 to an umbel, with 3 sepals and 3 petals, greenish outside and yellow inside. Flowers 5–10 mm long; fruit a narrow capsule. Not common; found growing among grass in eastern Parklands.

IRIDACEAE—iris family

Flowers large, irregular; leaves to 3 cm wide;
 plants with stout rootstocks. *Iris*
 Flowers small, regular or nearly so; leaves to 5
 mm wide; plants with fibrous roots. *Sisyrinchium*



Fig. 84. Death camas,
Zygadenus gramineus
Rydb.

Iris flag

Flowers yellow. *I. pseudacorus*

Flowers blue violet. *I. versicolor*

Iris pseudacorus L.

WATER FLAG

Stems 50–60 cm high, with leaves stiff, erect, 1–2 cm broad, equaling or exceeding the stems. Perianth 7–9 cm across, bright yellow to cream. Capsule 5–8 cm long, 6-angled. Introduced, occasionally escaped; southeastern Parklands, Boreal forest.

Iris versicolor L.

BLUE FLAG

Stems 20–80 cm high, arising from a thick, creeping rhizome; leaves erect, arching, 1–2 cm broad, shorter than to equaling the stems. Perianth 6–8 cm across, pale to dark blue or violet. Capsule 3–6 cm long, 3-angled. Marshes, swamps, and lakeshores; southeastern Parklands, Boreal forest.

Sisyrinchium blue-eyed grass

Perennial plants from rootstocks, with 2-edged stems. Leaves narrowly linear or grass-like and borne on two sides of the stem. The flowers perfect, with 3 sepals and 3 petals, all alike, blue, and with sharp-pointed tips. Fruit a round capsule.

Sisyrinchium montanum Greene

COMMON BLUE-EYED GRASS

Stems 10–30 cm high with narrow-winged edges. Flowers bright blue, with petals 3 mm long. Common, often found in large colonies; in meadows and moist places; throughout Prairies and Parklands. This, the commonest of the species, was formerly called *Sisyrinchium angustifolium* Miller.

ORCHIDACEAE—orchid family

Perennial herbs, with roots often fleshy. Leaves entire, usually sheathing the stems. Flowers irregular, usually with a large lip. *Cypripedium* has 2 stamens but the other genera have a single stamen, with the pollen grains united into 4–8 masses called pollinia. A difficult, widely distributed family with pretty flowers.

- 1. Flowers large, to 8 cm long; lip inflated and mocassin-shaped; fertile anthers
 - 2. *Cypripedium*
- Flowers mostly smaller; lip concave or flat, not mocassin-shaped; fertile anther 1. 2
- 2. Flower solitary; plants with a single basal leaf; scape with sheathing bracts. 3
- Flowers 2 to many, in racemes or spikes. 4
- 3. Flower purplish, 15–20 mm long; leaf oval to round-ovate, 2–6 cm long. *Calypso*
- Flower rose purplish, 20–30 mm long; leaf linear, 10–20 cm long. *Arethusa*
- 4. Plants without chlorophyll; leaves reduced to scales. *Corallorhiza*
- Plants with chlorophyll; leaves normal. 5
- 5. Flowers distinctly spurred. 6
- Flowers not spurred. 7
- 6. Leaves 2, oval, basal; lip white, spotted with purple; sepals and petals roseate. *Orchis*
- Leaves 1 to several; flowers uniformly greenish or white. *Habenaria*
- 7. Leaves 2, sessile near the middle of the stem. *Listera*
- Leaves 1 to several, basal or scattered. 8
- 8. Plants with a subbasal, solitary leaf. 9
- Plants with 2 to several leaves. 10

9. Leaf linear to narrowly oblong; flowers 25–50 mm long, pink purple. *Calopogon*
 Leaf oval to elliptic; flowers small, 2–3 mm long, greenish. *Malaxis*
10. Leaves somewhat fleshy, strongly reticulate, ovate to obovate, basal. *Goodyera*
 Leaves not fleshy, not strongly reticulate, linear or lanceolate. 11
11. Leaves 2, strongly keeled, sheathing the stem at the base; flowers greenish. *Liparis*
 Leaves several, not strongly keeled, basal or scattered on the stem; flowers creamy white. *Spiranthes*

Arethusa arethusa

Arethusa bulbosa L.

DRAGON'S-MOUTH

Plants with solid bulbs, 10–30 cm high; the leaf 2–4 mm wide, nearly equaling the scape. Flower with a pair of small bracts at the base; sepals and petals lanceolate, magenta; lip pinkish white with purple and yellow spots and streaks. Very rare; bogs and wet meadows; southeastern Boreal forest.

Calopogon grass-pink

Calopogon pulchellus (Salisb.) R. Br.

PURPLE GRASS-PINK

Plants with solid bulbs, 30–90 cm high; the leaf to 40 cm long, grass-like. Inflorescence a loose raceme with 3–15 rose purple flowers, 15–20 mm long. Very rare; bogs and swamps; southeastern Boreal forest.

Calypso Venus-slipper

Calypso bulbosa (L.) Oakes (Fig. 85A)

VENUS-SLIPPER

Growing from bulbous corm, with stem 8–15 cm high. One round-ovate leaf at base and a single pink flower 15–20 mm long with large pink sac having purple lines and inner tuft of yellow hairs. An exceedingly delicate and beautiful flower. In cool woods; Boreal forest, Cypress Hills. Syn.: *Cytherea bulbosa* (L.) House.

Corallorhiza coralroot

Plants growing on dead and decaying organic matter; therefore, found only in wooded areas. Lacking green color; stems scaly, pinkish yellow; roots coral-like.

1. Flowers small, not longer than 10 mm. *C. trifida*
 Flowers longer than 10 mm. 2
2. Flowers conspicuously striped; lip entire. *C. striata*
 Flowers more spotted than striped; lip 3-lobed. *C. maculata*




Fig. 85. Orchids: *A*, Venus-slipper, *Calypso bulbosa* (L.) Oakes; *B*, small yellow lady's-slipper, *Cypripedium calceolus* L. var. *parviflorum* (Salisb.) Fern.; *C*, green-flowered bog orchid, *Habenaria hyperborea* (L.) R. Br.; *D*, northern twayblade, *Listera borealis* Morong; *E*, round-leaved orchid, *Orchis rotundifolia* Banks.

Corallorhiza maculata Raf.

LARGE CORALROOT

Plant with stem 20–50 cm high, purplish, having scaly leaves. Flowers 12–20 mm long, reddish purple; lip white, spotted with red. Not common; in shady woods; throughout the Prairie Provinces.

Corallorhiza striata Lindl.

STRIPED CORALROOT

Plant with coarse, stout stem 20–50 cm high. Flowers dark purple, with darker purple stripes. Very rare; in shady woodlands; southeastern Boreal forest, Cypress Hills.

Corallorhiza trifida Chat.

EARLY CORALROOT

Small species with slender stem 10–30 cm high, and small greenish flowers. Not common; in shaded woods; throughout the Prairie Provinces.

Cypripedium lady's-slipper


Perennial plants with thick fibrous roots and erect stems, bearing two to several large leaves and 1–3 large flowers. All are rare or very rare; the most colorful orchids in the Prairie Provinces.

- 1. Lower sepals separate and spreading. *C. arietinum*
 Lower sepals united, usually 2-veined or
 2-toothed. 2
- 2. Flowering stem a naked scape, with 2
 leaves basal, subopposite. *C. acaule*
 Flowering stem leafy, with 2–6 leaves
 along the stem. 3
- 3. Sepals and petals white; 25–40 mm long;
 lip white with rose purple streaks. *C. reginae*
 Sepals greenish, brownish, or yellowish. 4
- 4. Sepals yellowish; petals greenish yellow to
 brownish; lip yellow. *C. calceolus*
 Sepals greenish or brownish; lip not
 yellow. 5
- 5. Flower solitary; sepals and petals green-
 ish, crimson-striped. *C. candidum*
 Flowers 1–3 on stem; petals brownish or
 white. 6
- 6. Sepals and petals greenish or purplish
 brown, 40–60 mm long. *C. montanum*
 Sepals green, with the lateral petals white,
 15–20 mm long. *C. passerinum*

Cypripedium acaule Ait. (Fig. 86)

STEMLESS LADY'S-SLIPPER

Scape leafless, 20–40 cm high, with 2 basal leaves, 10–20 cm long, elliptic, thinly pubescent. Flower solitary; sepals and lateral petals yellowish green to greenish brown, 30–50 mm long, lip 30–60 mm long, pink with reddish veins. Rare; dry to moist woods, bogs, and swamps; Boreal forest.



HERBARIUM OF THE RESEARCH STATION
 WEST POINT, N.Y.
 PLANT: *Cypripedium acaule*, var. *puberulum* (L.)
 JONES, 1910. (See also No. 1114)
 D. ATTEN: (See also provincial book
 1114, p. 100, 1911)
 ALLEN: (See also No. 1114, p. 100, 1911)
 BOSTON: (See also No. 1114, p. 100, 1911)

Fig. 86. Stemless lady's-slipper, *Cypripedium acaule* Ait.

Cypripedium arietinum R. Br.

RAM'S-HEAD LADY'S-SLIPPER

Stem with 3–5 sessile leaves above and 2 or 3 sheathing scales below, 10–40 cm high; leaves 5–10 cm long, lanceolate, finely ciliate. Flower solitary; sepals and lateral petals greenish brown, 15–25 mm long; lip whitish, veined with red, 10–20 mm long. Very rare; moist coniferous woods; southeastern Boreal forest.

Cypripedium calceolus L.

YELLOW LADY'S-SLIPPER

Plants 20–40 cm high, with leafy stems; the leaves more or less sheathing 6–20 cm long, elliptic. Flowers 1–2, with an erect leaf-like bract at the base; sepals and petals greenish yellow to purplish brown, the lip yellow, usually more or less purple-veined. Small yellow lady's-slipper, *C. calceolus* var. *parviflorum* (Salisb.) Fern. (Fig. 85B), has sepals 3–5 cm long, the lip 20–35 mm long; moist woods; Boreal forest. *C. calceolus* var. *pubescens* (Willd.) Correll has sepals 5–8 cm long, and the lip 35–60 mm long. Both varieties are becoming increasingly rare; moist woods; southeastern Boreal forest.

Cypripedium candidum Muhl.

SMALL WHITE LADY'S-SLIPPER

A somewhat glandular or sticky-haired plant 10–25 cm high. The ovate to lanceolate leaves 6–12 cm long. The single flower has a white lip with purplish stripes inside. Scarce; but has been found in moist spots; southeastern Boreal forest.

Cypripedium montanum Dougl.

MOUNTAIN LADY'S-SLIPPER

Plants 20–50 cm high, more or less glandular pubescent throughout; leaves 4–6, ovate to lanceolate 5–16 cm long. Flowers with brownish green sepals and petals, 4–6 cm long, the lip 2–3 cm long, white, purple-veined. Moist woods; southern Rocky Mountains.

Cypripedium passerinum Richardson

NORTHERN LADY'S-SLIPPER

Differs from other lady's-slippers by having rounded sepals and smaller flowers, with the lip 15–20 mm long, white or pale lilac with purple spots. Rare; moist coniferous woods; Cypress Hills, Boreal forest.

Cypripedium reginae Walt.

SHOWY LADY'S-SLIPPER

A very showy species, 30–60 cm high; leaves oval to elliptic, 8–20 cm long, covered with soft hairs; flowers white, with a large, inflated sac-like lip 30–40 mm long, with reddish purple stripes. Rare; swampy woodlands; southeastern Boreal forest, Riding Mountain.

Goodyera rattlesnake-plantain

Plants with fleshy rootstocks and a basal clump of dark green mottled leaves and greenish white flowers borne on long stem.

Stem 10–20 cm high; lip of flower decidedly inflated; flowers to 5 mm long. *G. repens*

Stem 30–45 cm high; lip of flower scarcely inflated, margins turned inward; flowers to 10 mm long. *G. oblongifolia*

Plants 30–45 cm high, with ovate-lanceolate to narrowly elliptic leaves to 8 cm long, 20–25 mm wide. Raceme 6–12 cm long; flowers 8–10 mm long, with the lip 6–7 mm long, the body semiglobose. Very rare; dry to moist woods; Boreal forest, Rocky Mountains, Cypress Hills.

Goodyera repens (L.) Br.

LESSER RATTLESNAKE-PLANTAIN

Plants 10–30 cm high, with ovate to oblong leaves 15–30 mm long, 8–12 mm wide. Raceme 3–6 cm long; flowers 3.5–5 mm long, with the lip 3–3.5 mm long, the body deeply pouch-like. Rare; dry to moist woods; Boreal forest, Rocky Mountains, Cypress Hills, Riding Mountain. The var. *repens* has the leaves reticulate green-veined; plants with the leaves white reticulate distinguished as var. *ophioides* Fern.

Habenaria bog orchid

- 1. Leaves basal. 2
 Leaves cauline. 5
- 2. Leaf solitary, 5–15 cm long, 1–5 cm wide;
 scape usually bractless. *H. obtusata*
 More than one basal leaf. 3
- 3. Leaves lanceolate or oblanceolate, 6–12
 cm long, 1–3 cm wide; spur 5 mm long. *H. unalascensis*
 Leaves broadly ovate or rotund, to 15 cm
 long, and often as wide; spur longer
 than 10 mm. 4
- 4. Stem bractless; spur 10–25 mm long. *H. hookeri*
 Stem with several small bracts; spur
 25–45 mm long. *H. orbiculata*
- 5. Lip 3-toothed, with the central tooth
 short; spur 2–3 mm long. *H. viridis* var. *bracteata*
 Lip entire; spur 4–8 mm long. 6
- 6. Spur 4–5 mm long, pouch-like, often
 purplish. *H. saccata*
 Spur 5–8 mm long, cylindrical. 7
- 7. Flowers white, with the lip abruptly wid-
 ened at base. *H. dilatata*
 Flowers greenish, with the lip gradually
 widened toward base. *H. hyperborea*

Habenaria dilatata (Pursh) Hook.

WHITE BOG ORCHID

Plant with slender, leafy stem 30–60 cm high; flowers small, white, about 10–15 mm long, in a spike-like raceme. Rare; in bogs; Boreal forest. Syn.: *Limnorchis dilatata* (Pursh) Rydb.

Habenaria hookeri Torr.

HOOKER'S BOG ORCHID

Plant with 2 basal leaves, 20–40 cm high; the leaves broadly elliptic to rotund; flowers yellowish green. Very rare; moist woods; southeastern Boreal forest.

Habenaria hyperborea (L.) R. Br. (Fig. 85C) GREEN-FLOWERED BOG ORCHID

Fairly stout, leafy-stemmed plant 20–60 cm high; flowers greenish, in spike-like raceme. In moist woodlands, meadows, and stream banks; Boreal forest, Riding Mountain, Cypress Hills. Syn.: *Limnorchis viridiflora* (Cham.) Rydb.

Habenaria obtusata (Pursh) Richardson SMALL NORTHERN BOG ORCHID

Slender plant without stem leaves, 10–25 cm high. Solitary, obovate basal leaf and loose raceme of greenish yellow flowers. Moist woods, marshes, and bogs; Boreal forest. Syn.: *Lysiella obtusata* (Pursh) Rydb.

Habenaria orbiculata (Pursh) Torr. ROUND-LEAVED BOG ORCHID

Plants with 2 basal leaves; scape 30–60 cm high; leaves broadly elliptic to rotund; flowers greenish white, about 2 cm long, with the spike to 20 cm long. Rare; moist woods and bogs; Boreal forest.

Habenaria saccata Greene SLENDER BOG ORCHID

Plants with leafy stems 20–50 cm high; leaves 4–12 cm long, lanceolate or oblanceolate; flowers green, tinged with purple or brown, with the lip 4–7 mm long, the spur 3–5 mm long, pouch-shaped, green, often purplish-tinged. Rare; wet meadows, bogs, and forests; southern Rocky Mountains.

Habenaria unalascensis (Spreng.) Wats. ALASKA BOG ORCHID

Plants with basal leaves; scape 20–60 cm high, with a few scale-like leaves; basal leaves 1–4, narrowly oblanceolate; flowers green, in a spike 10–30 cm long, with the spur 3–5 mm long. Rare; moist or dry woods and meadows; Rocky Mountains, Boreal forest.

Habenaria viridis (L.) R. Br. var. *bracteata* (Muhl.) Gray LONG-BRACTED ORCHID

Stout, leafy-stemmed plant 15–60 cm high, with very conspicuous green bracts, the lower ones at least twice as long as the greenish flowers. In moist meadows and open woods; throughout the Prairie Provinces. Syn.: *H. bracteata* (Muhl.) R. Br.

Liparis twayblade

Liparis loeselii (L.) Rich. TWAYBLADE

Stem strongly ribbed, 5–20 cm high, with 2 lanceolate leaves at the base; flowers few, small, greenish, in a raceme. Rare; in bogs and moist woods; Boreal forest.

Listera twayblade

Low-bog or moist-soil plants with 2 almost opposite leaves about halfway up the stem.

1. Lip cleft to below middle; lobes linear; leaves cordate to deltoid. *L. cordata*

- Lip shallowly cleft; lobes oblong to ovate; leaves ovate to elliptic. 2
2. Lip nearly oblong, narrowed at the middle, auriculate at base. *L. borealis*
- Lip widening toward the tip, not auriculate at base. 3
3. Lip 8–10 mm long, minutely toothed on the margins near the base. *L. convallarioides*
- Lip 4–6 mm long, prominently toothed on the margins near the base. *L. caurina*

Listera borealis Morong (Fig. 85D)

NORTHERN TWAYBLADE

Plants 5–25 cm high, with the leaves ovate to elliptic, 1–5 cm long. Flowers pale to yellow green, with deep green veins in sepals and petals. Rare; mossy woods, moist meadows and slopes; Rocky Mountains, Cypress Hills.

Listera caurina Piper

WESTERN TWAYBLADE

Plants 10–20 cm high, with the leaves broadly ovate to orbicular, 3–6 cm long; stems glandular puberulent above. Flowers greenish or yellowish. Rare; moist woods; southern Rocky Mountains.

Listera convallarioides (Sw.) Torr.

BROAD-LIPPED TWAYBLADE

Plants 10–20 cm high, with the leaves broadly oval, 3–5 cm long; stems and inflorescence glandular pubescent. Flowers green or yellow green. Rare; meadows, bogs, and moist woods; Boreal forest.

Listera cordata (L.) R. Br.

HEART-LEAVED TWAYBLADE

Plants 10–20 cm high, with the leaves broadly round-ovate, 15–30 mm long, truncate to cordate at the base. Flowers green, tinged with purple. Very rare; moist woods and swamps; Boreal forest.

Malaxis adder's-mouth

1. Stem leaves 2 or more; lip erect. *M. paludosa*
 Stem leaf solitary; lip deflected. 2
2. Lip not lobed, pointed. *M. monophylla* var. *brachypoda*
 Lip deeply lobed. *M. unifolia*

Malaxis monophylla (L.) Sw. var. *brachypoda* (Gray) Morris & Eames

ADDER'S-MOUTH

A slender bog plant 5–20 cm high, from an ovoid corm or solid bulb-like base. The single leaf clasping, oval to elliptic, to 8 cm long; greenish yellow flowers in a raceme, very small, about 3 mm long. Rare; Boreal forest.

Malaxis paludosa (L.) Sw.

BOG ADDER'S-MOUTH

Plants 5–15 cm high, with the 2–5 leaves ovate-lanceolate, 1–3 cm long; flowers yellowish green. Rare; bogs; Boreal forest.

Malaxis unifolia Michx.

GREEN ADDER'S-MOUTH

Scape 10–30 cm high, with the single leaf sessile, oval or elliptic, 3–6 cm long, 1–3 cm wide; flowers greenish. Rare; damp woods and bogs; southeastern Boreal forest.

Orchis orchid

Orchis rotundifolia Banks (Fig. 85E)

ROUND-LEAVED ORCHID

An orchid 20–25 cm high, with 1 oval to almost round leaf near the base, and often with 1 or 2 sheathing scales below it. Leaf 3–8 cm long and 2–5 cm wide. Flowers on stem 2–5 cm long, rose, 10–15 mm long, with white lips, spotted with purple. In moist woodlands; especially in Boreal forests and in Cypress Hills.

Spiranthes lady's-tresses

Flowers in 1 spiral. *S. gracilis*

Flowers in 2 or 3 spirals. *S. romanzoffiana*

Spiranthes gracilis (Bigel.) Beck

SLENDER LADY'S-TRESSES

An orchid 15–40 cm high. Leaves mostly basal, short-stalked, ovate or elliptic, and usually withering before the flowers appear. Rare; open woods and bogs; Boreal forest.

Spiranthes romanzoffiana Cham.

HOODED LADY'S-TRESSES

An orchid 15–40 cm high, with the lower leaves linear to linear-lanceolate; flowers in 3 spirals, delicately fragrant. In swampy places; throughout the Prairie Provinces.

Class: DICOTYLEDONEAE

Seedlings with 2 seedling leaves (cotyledons); stems with a central pith or, if woody, the wood arranged in annual layers; the leaves netted-veined; and the flower parts usually in fours and fives, or multiples of four and five.

SALICACEAE—willow family

Deciduous (that is, shedding their leaves in fall, not evergreen) trees or shrubs. Leaves alternate on stem. Flowers in aments (catkins) without sepals and petals, these being replaced by glands or a cup-like disk. Male and female flowers produced on separate plants. The numerous seeds bear a tuft of hairs at the apex, which aids in dissemination.

Winter buds covered by several scales; bracts below flowers fringed; stamens usually more than 10; a cup-shaped disk below each flower.

Populus

Winter buds covered by one scale; bracts below flowers entire, not fringed; stamens 2–10; one or more glands below each flower. *Salix*

Populus poplar

Fairly tall trees with either smooth or furrowed bark and light-colored, soft, straight-grained wood. Leaves usually broad and pointed, with petioles sometimes as long as the leaf itself. Flowers unisexual, those of each sex in long catkins on separate trees. Fruit capsule containing small seeds, each bearing a tuft of white hairs.

The genus *Populus* consists of species that appear to hybridize freely, which has resulted in several forms, intermediate between the parents. Because of the considerable variability in the size and form of the leaves in the species, many species have been described that are now considered to be varieties or hybrids.

- 1. Petioles distinctly flattened. 2
 Petioles terete. 4
- 2. Leaves coarsely toothed. *P. grandidentata*
 Leaves not coarsely toothed. 3
- 3. Leaves roundish in outline, with the margins finely crenate. *P. tremuloides*
 Leaves deltoid in outline, with the margins coarsely dentate. *P. deltoides*
- 4. Leaves lanceolate to linear-lanceolate. *P. angustifolia*
 Leaves ovate or roundish. 5
- 5. Leaves roundish, coarsely toothed, white below. *P. alba*
 Leaves ovate, dark green above, light green or glaucous below. *P. balsamifera*

Populus alba L. WHITE POPLAR

Trees up to 30 m high, with bark grayish, furrowed; branches brownish; branchlets coarse, glabrous, grayish brown; leaves 4–6 cm long, 3–4 cm broad, oval to round in outline, with margins coarsely toothed, green above, white tomentose below. Aments appearing before the leaves, on leafless peduncles. Capsules 5–7 mm long, sessile. Male flowers with 6–10 stamens. Introduced from Europe, and planted in shelterbelts.

Populus angustifolia James NARROW-LEAVED COTTONWOOD

Trees to 15 m high, with bark greenish; branches greenish brown; branchlets green, coarse, glabrous; leaves 5–12 cm long, lanceolate to long-ovate, acute to acuminate, base cuneate to subcordate, with margins finely crenate, dark green above, pale green below. Aments appearing with the leaves, drooping. Capsules 5–7 mm long, sessile. Male flowers with 6–10 stamens. Along rivers in Prairies.

Populus balsamifera L.

BALSAM POPLAR

Large trees, to 25 m high, with bark at first grayish white, becoming dark gray and furrowed in age; branches grayish brown; twigs coarse, light gray; leaves 5–15 cm long, ovate to ovate-lanceolate, acute to acuminate, base obtuse to cuneate or subcordate, with margins minutely crenulate to subentire, dark green and somewhat shiny above, light yellow green to rusty below. Aments appearing before the leaves, on drooping peduncles. Capsules 6–10 mm long. Stamens 12–30. Along rivers and lakes, in coulees and ravines, throughout the Prairie Provinces, but becoming more common in Boreal forest. This species is very variable in leaf shape and size; it includes several forms and varieties, sometimes described as separate species: *Populus candicans* Michx., *P. tacamahaca* Mill., *P. trichocarpa* T. & G., and *P. gileadensis* Rouleau.

Populus deltoides Marsh.

COTTONWOOD

Large trees 15–25 m high, with bark greenish gray, furrowed; branches grayish brown; twigs greenish gray; leaves 5–10 cm long, broadly deltoid, acuminate, base often somewhat cordate, with margins coarsely serrate-crenate, shiny green above, lighter green below. Aments appearing before the leaves, on drooping peduncles. Capsules 6–10 mm long. Stamens about 60. Along rivers and lakeshores, sand dunes, Prairies. Western plants are classified as var. *occidentalis* Rydb. based on the coarser serration; in eastern specimens the number of teeth on one side of the leaves is about 50–60% higher than that in the western specimens.

Populus grandidentata Michx.

LARGE-TOOTHED ASPEN

Medium-sized trees with bark greenish gray, becoming brown and furrowed in age; branches gray brown; twigs gray; leaves 6–10 cm long, broadly ovate; with margins usually having 5–10 large rounded teeth on each side; densely white tomentose when young, later thinly pubescent. Aments appearing before the leaves. Capsules 5–7 mm long. Male flowers with 5–12 stamens. Southeastern Boreal forest.

Populus tremuloides Michx.

ASPEN POPLAR

Slender trees, to 30 m high, with bark grayish white, furrowed in lower part of stem in age; branches brown; twigs yellow green; leaves 3–10 cm long, broadly ovate to orbicular, abruptly pointed, base truncate to subcordate, with margins finely crenate or serrate to subentire, dark green or dark yellow green above, pale green below. Aments appearing before the leaves, on drooping peduncles. Capsules 4–6 mm long. Male flowers with 5–12 stamens. Very common; in depressions and other moist areas; in the Prairies, widespread in the Parklands and Boreal forest.

Salix willow

The willows are very difficult to identify; the flowers or fruits of both sexes, and also in many instances the mature leaves, are needed for positive identification. Even then, several of the diagnostic characters are not constant as a result of hybridization between species, which may occur quite often.

Nevertheless, it is possible to determine typical species with a high degree of accuracy, and come close with most of the more or less intermediate forms. Determinations should be made on fertile material using the female plants. Matching of the male plants is then possible on the basis of the characters common to both sexes, for example, color of bark, leaf characters, and characters of the catkins. In the determination of the female plants, the following characters are used:

- position of catkin: sessile or on a leafy branchlet (also for male plants);
- capsules: pubescent or glabrous, size, sessile or pedicellate;
- length of pedicel: longer or shorter than the bract;
- bracts: light or dark in color.

With the use of well-developed material, that is, with mature capsules and, if possible, expanded leaves also, most specimens can be determined accurately based on the combinations of these four characters.

Sterile specimens can usually be determined with a fair degree of accuracy by anyone who has experience in working with willows. New shoots often bear leaves that are not typical in size, and therefore it is best always to use mature leaves in all determinations. In Fig. 87, the main characters of both fruiting and sterile material, used in the keys, are illustrated.

Key to species of *Salix* by fruiting characters

1. Capsules pubescent. 2
 Capsules glabrous. 23
2. Bracts of the capsules light in color, yellowish to light brownish or purplish. 3
 Bracts of the capsules dark in color, brown to black. 8
3. Plants very small, with creeping stems mostly underground; underside of leaves reticulate-veined. *S. reticulata*
 Plants not creeping, with stem at least partly upright. 4
4. Leaves narrowly linear, the length-to-width ratio 10:1 or more; leaf margins remotely dentate. *S. interior*
 Leaves not narrowly linear, the length-to-width ratio seldom more than 5:1. 5
5. Capsules on long pedicels; catkins loose. 6
 Capsules with short pedicels; catkins compact. 7
6. Shrubs with stems to 2 m high; leaves oblanceolate or ellipsoid, dull green, mostly somewhat pubescent, rugose-veiny below. *S. bebbiana*
 Shrubs with stems seldom higher than 1 m; leaves narrowly elliptic oblanceolate, green above, glaucous below, not rugose-veiny. *S. pedicellaris*




Fig. 87. Main characters of willows. Catkins: *a*, on leafy branchlet; *b*, sessile. Capsules: *c*, pedicel longer than bract; *d*, pedicel as long as bract; *e*, pedicel shorter than bract. Male flower: *f*. Leaves: *g*, acuminate; *h*, acute; *k*, rounded; *l*, irregularly veined; *m*, remotely dentate; *n*, finely serrate.

7. Undersurface of leaves densely silky-pubescent; leaves sessile; catkins to 2 cm long. *S. brachycarpa*
 Undersurface of leaves glaucous, pubescent but becoming glabrous; leaves petioled; catkins to 7 cm long. *S. glauca*
8. Leaf margins distinctly toothed. 9
 Leaf margins entire or indistinctly toothed. 12
9. Leaves deep green and rugose above, silvery silky and strongly veined below. *S. vestita*
 Leaves not as above. 10
10. Leaves shiny green above, glaucous and silky pubescent below. *S. arbusculoides*
 Leaves glabrous below. 11
11. Leaves glabrous on both sides, with underside light green or glaucous, strongly reticulate-veined. *S. maccalliana*
 Leaves pubescent above, glabrous below, not noticeably reticulate-veined. *S. barclayi*
12. Leaves lanceolate, densely white tomentose below, dark green and pubescent above; leaf margins entire, involute. *S. candida*
 Leaves not as above. 13
13. Shrubs strongly depressed, with stems partly or entirely underground. 14
 Shrubs not strongly depressed, with stems mostly upright. 15
14. Capsules silky-villose, grayish; leaves green above, yellowish below. *S. arctica*
 Capsules thinly pubescent, purplish; leaves shiny green above, glaucous below. *S. arctophila*
15. Catkins on leafy peduncles. 16
 Catkins not on leafy peduncles. 17
16. Leaves densely pubescent below. *S. sitchensis*
 Leaves glabrous below. *S. petiolaris*
17. Leaves pubescent on both sides, more or less cordate at the base. *S. barrattiana*
 Leaves glabrous or nearly so above, not cordate at base. 18
18. Young twigs and branches densely tomentose. *S. alaxensis*
 Young twigs and branches not densely tomentose. 19

19. Leaves pubescent below; branchlets yellowish to brown.	20
Leaves not pubescent below, mostly glaucous; branchlets reddish or yellowish.	21
20. Leaves narrowly lanceolate, the length-to-width ratio about 6:1; capsules 4–6 mm long.	<i>S. pellita</i>
Leaves oblanceolate or obovate, the length-to-width ratio less than 5:1; capsules 7–9 mm long.	<i>S. humilis</i>
21. Leaves lanceolate, the length-to-width ratio about 6:1; capsules 4–6 mm long; twigs yellow.	<i>S. pellita</i> var. <i>psila</i>
Leaves elliptic or elliptic-oblanceolate, the length-to-width ratio about 3:1; capsules 6–12 mm long; twigs reddish.	22
22. Lateral veins of leaves regular, parallel; capsules 6–7 mm long, subsessile.	<i>S. planifolia</i>
Lateral veins of leaves irregularly spaced; capsules 7–12 mm long, clearly pedicellate.	<i>S. discolor</i>
23. Bracts of the capsules dark brown to black.	24
Bracts of the capsules yellow to light brown or purplish.	31
24. Catkins sessile or subsessile, leafless.	25
Catkins on leafy peduncles.	26
25. Leaves ovate to orbicular, strongly veined below, entire to somewhat glandular dentate.	<i>S. calcicola</i>
Leaves elliptic-ovate to narrowly ovate, reticulate-veined and glaucous below, distinctly serrate.	<i>S. monticola</i>
26. Leaves pubescent.	27
Leaves glabrous.	28
27. Leaves green, pubescent on both sides.	<i>S. commutata</i>
Leaves green, pubescent above, glabrous and glaucous below.	<i>S. barclayi</i>
28. Pedicels of capsules much longer than bracts.	29
Pedicels of capsules as long as or shorter than bracts.	30
29. Young twigs dark brown or yellow brown; leaves lanceolate to obovate, rounded to cordate at base.	<i>S. mackenzieana</i>
Young twigs yellow or reddish, shiny; leaves ovate to lanceolate, fragrant when crushed.	<i>S. pyrifolia</i>

30. Leaves green and reticulate-veined on both sides; stipules usually lacking.	<i>S. myrtillifolia</i>
Leaves yellowish green above, glaucous below; stipules usually present.	<i>S. lutea</i>
31. Shrubs depressed; stems underground with only leaves and catkins showing.	<i>S. herbacea</i>
Shrubs not depressed; stems upright.	32
32. Leaves narrowly linear, length-to-width ratio about 10:1; leaf margins remotely dentate.	<i>S. interior</i>
Leaves not narrowly linear, length-to-width ratio 5:1 or less.	33
33. Foliage fragrant when young or crushed.	34
Foliage not fragrant.	35
34. Leaves firm, dark glossy green above; petioles with glands at leaf base; pedicels of capsules shorter than bracts.	<i>S. pentandra</i>
Leaves thin, dull or slightly shiny above; petioles without glands; pedicels of capsules longer than bracts.	<i>S. pyrifolia</i>
35. Young twigs fragile, breaking off at the base in strong wind or on touch.	<i>S. fragilis</i>
Young twigs not fragile.	36
36. Leaves pubescent, especially below.	<i>S. alba</i>
Leaves glabrous.	37
37. Petioles with glands at leaf base.	38
Petioles glandless.	39
38. Catkins appearing at the same time as the leaves; capsules 4–7 mm long.	<i>S. lucida</i>
Catkins appearing later than the leaves; capsules 7–12 mm long.	<i>S. serissima</i>
39. Leaves usually large, to 12 cm long, closely and finely serrate.	<i>S. amygdaloides</i>
Leaves usually small, to 6 cm long, often with revolute margins.	<i>S. pedicellaris</i>

Key to species of *Salix* by leaf characters

1. Leaves densely pubescent beneath.	2
Leaves glabrous or only sparsely pubescent beneath.	15
2. Leaf margins toothed or entire, revolute.	3
Leaf margins toothed or entire, not revolute.	6

3. Length-to-width ratio of leaves about 3:1. 4
 Length-to-width ratio of leaves about 6:1. 5
4. Young twigs densely felty tomentose; leaf margins entire or minutely glandular serrulate. *S. alaxensis*
 Young twigs not densely tomentose; leaf margins entire or coarsely glandular serrate. *S. bebbiana*
5. Leaf margins entire; pubescence often interspersed with brown hairs. *S. candida*
 Leaf margins obscurely glandular serrulate; pubescence silvery. *S. pellita*
6. Length-to-width ratio of leaves 10:1 or more. *S. interior*
 Length-to-width ratio of leaves usually not more than about 5:1. 7
7. Leaves glabrous or glabrate above. 8
 Leaves clearly pubescent above. 10
8. Leaves obovate, often almost orbicular, length-to-width ratio about 1.5:1, deep green and rugose above. *S. vestita*
 Leaves elliptic to oblanceolate, length-to-width ratio about 4:1, not rugose above. 9
9. Length-to-width ratio of leaves usually about 3:1. *S. discolor*
 Length-to-width ratio of leaves usually about 5:1. *S. humilis*
10. Leaf margins finely glandular serrulate. *S. alba*
 Leaf margins entire or obscurely dentate. 11
11. Leaves sessile, the petiole usually less than 1 mm long. *S. brachycarpa*
 Leaves with well-developed petioles. 12
12. Twigs and young branches grayish or light brown. 13
 Twigs and young branches dark brown to black. 14
13. Leaf scars on older branches very prominent; young branches yellowish brown. *S. barrattiana*
 Leaf scars on older branches not prominent; young branches grayish brown. *S. glauca*
14. Leaves as pubescent above as below, pubescence tomentose. *S. commutata*
 Leaves much less pubescent above than below, pubescence silky. *S. sitchensis*

15. Length-to-width ratio of leaves 10:1 or more.	<i>S. interior</i>
Length-to-width ratio of leaves usually not more than 6:1.	16
16. Leaf margins clearly toothed.	17
Leaf margins entire or obscurely toothed.	31
17. Petioles glandular at leaf base.	18
Petioles glandless.	21
18. Twigs very fragile, readily breaking in strong wind or on touch.	<i>S. fragilis</i>
Twigs not fragile.	19
19. Leaves very dark glossy green above, lighter and bluish green below, acute to short acuminate; introduced small tree.	<i>S. pentandra</i>
Leaves not very glossy above.	20
20. Leaves green below, long acuminate.	<i>S. lucida</i>
Leaves glaucous below, short acuminate.	<i>S. serissima</i>
21. Twigs drooping or arching; leaves long acuminate.	<i>S. amygdaloides</i>
Twigs not drooping; leaves not long acuminate.	22
22. Leaves ovate to orbicular; branchlets stout, with conspicuous leaf scars.	<i>S. calcicola</i>
Leaves lanceolate to oblong; branchlets more slender, not with conspicuous leaf scars.	23
23. Length-to-width ratio of leaves about 6:1.	24
Length-to-width ratio of leaves less than 5:1.	25
24. Leaf margins coarsely serrate; leaves glabrous below.	<i>S. petiolaris</i>
Leaf margins closely and shallowly serrate; leaves somewhat silky pubescent below.	<i>S. arbusculoides</i>
25. Leaves thin, aromatic when young or crushed.	<i>S. pyrifolia</i>
Leaves not thin, not aromatic.	26
26. Plants small shrubs, usually less than 1 m high; stems often decumbent.	<i>S. myrtillifolia</i>
Plants usually more than 1 m high; stems usually upright.	27
27. Leaves pubescent above, especially on the veins, glabrous and glaucous below.	<i>S. barclayi</i>
Leaves glabrous on both sides.	28

28. Length-to-width ratio of leaves about 4:1;
leaves rather glossy green above,
midrib yellow. *S. maccalliana*
Length-to-width ratio of leaves about
2.5:1. 29
29. Leaves elliptic-lanceolate, acute. *S. monticola*
Leaves ovate-lanceolate, acuminate. 30
30. Twigs yellow to yellowish gray; leaves
yellowish green above, glaucous
below. *S. lutea*
Twigs dark brown; leaves dark green
above, pale green below. *S. mackenzieana*
31. Plants low, usually creeping shrubs with
entirely or partly buried stems. 32
Plants usually shrubs with mostly upright
stems; 50 cm high or higher. 38
32. Leaf petioles short, usually less than 5
mm long. 33
Leaf petioles more than 5 mm long. 35
33. Plants glabrous throughout; leaves mostly
less than 3 cm long, reticulate below. *S. herbacea*
Plants not glabrous throughout. 34
34. Leaves pubescent below, rugose above;
stipules none. *S. vestita*
Leaves glabrous, glaucous and strongly
veined below; stipules often present. *S. calcicola*
35. Leaves conspicuously reticulate-veined,
oval to oblong, 1–3 cm long, usually
glabrous. *S. reticulata*
Leaves not conspicuously reticulate-
veined, obovate to lanceolate, to 6 cm
long, often pubescent. 36
36. Leaves dull green above, glaucous below,
often more or less pubescent on both
sides; stems usually upright. *S. glauca*
Leaves somewhat shiny green above, usu-
ally glabrous above; stems usually
creeping. 37
37. Leaves mostly obovate, strongly glaucous
below, somewhat rugose above. *S. arctophila*
Leaves mostly oblanceolate, glaucous
below, strongly veined and often
obscurely reticulate, especially below. *S. arctica*
38. Leaves glabrous on both sides. 39
Leaves more or less pubescent. 44
39. Leaves linear-lanceolate, the tips acute. 40
Leaves elliptic to ovate. 41

40. Leaves obscurely veined below. *S. pellita*
 Leaves clearly reticulate-veined below. *S. pedicellaris*
41. Leaves yellowish green above, pale green
 or glaucous below; usually some leaves
 serrate or undulate. *S. lutea*
 Leaves not yellowish green; leaf margins
 entire or obscurely and remotely
 crenate. 42
42. Lateral veins regular, parallel, rather
 closely spaced. *S. planifolia*
 Lateral veins irregular, not closely spaced. 43
43. Stipules usually present, large, clasping
 most of the stem. *S. monticola*
 Stipules usually absent, when present,
 small, hardly exceeding width of
 petiole base. *S. discolor*
44. Leaves usually more or less pubescent on
 both sides. 45
 Leaves glabrous above, more or less
 pubescent below, particularly on the
 veins. *S. discolor*
45. Leaves rugose below, glaucous. *S. bebbiana*
 Leaves glaucous below, clearly veined but
 not rugose. *S. glauca*

Salix alaxensis (Anderss.) Cov.

ALASKA WILLOW

A shrub to 4 m high; bark dark to chestnut brown, more or less persistently gray villous tomentose; branchlets densely white or yellow tomentose. Leaves 5–11 cm long, 1.5–3.5 cm broad, narrowly ovate to oblong, acute, with base cuneate; margins revolute, entire; bright green above, densely white pubescent below. Catkins appearing before the leaves; sessile. Capsules 4–5 mm long, densely puberulent; scales black; pedicels shorter than the scales. Male flowers with 2 stamens. Along streams and lakeshores; in the Rocky Mountains south to Jasper National Park.

Salix alba L.

WHITE WILLOW

A tree up to 20 m high; bark grayish; branches ascending, appearing silvery gray. Leaves 5–10 cm long, 1–1.5 cm broad, lanceolate, acuminate, with base cuneate; margins finely serrate; appressed white silky pubescent on both sides. Catkins appearing at the same time as the leaves; terminating short leafy peduncles. Capsules 3–5 mm long, glabrous; scales yellowish green; pedicels shorter than the scales. Male flowers with 2 stamens. A Eurasian species, often planted in shelterbelts.

Salix amygdaloides Anderss.

PEACH-LEAVED WILLOW

A shrub or tree 3–15 m high; bark rough, dark reddish brown; branches gray brown; twigs yellowish to reddish brown, drooping. Leaves 5–12 cm long,

1.5–3 cm broad, ovate-lanceolate, acuminate, tapering to base; margins closely serrate; light to yellowish green above, glaucous below, glabrous on both sides. Catkins appearing at the same time as the leaves; terminating short leafy peduncles. Capsules 4–7 mm long, glabrous; scales yellow, deciduous; pedicels 2–3 mm long. Male flowers with 4–7 stamens. Not common; along rivers and streams; Prairies.

Salix arbusculoides Anderss.

SHRUBBY WILLOW

Shrubs or occasionally small trees 1–6 m high; bark reddish brown; branches reddish brown; bark peeling; branchlets velvety. Leaves 5–8 cm long, 1–2 cm broad, narrowly ovate to elliptic, acute, with base acute; margins glandular serrulate; glossy and glabrous above, sparsely pubescent to almost glabrous below. Catkins appearing shortly before or at the same time as the leaves; subsessile, with bract-like leaves at base. Capsules 4–5 mm long, sparsely pubescent; scales dark brown; pedicels shorter than the scales. Male flowers with 2 stamens. Along rivers and in muskeg areas; Boreal forest.

Salix arctica Pall.

ARCTIC WILLOW

Dwarf shrubs, usually prostrate, occasionally to 50 cm high; branches stout to long, rooting at some nodes, yellowish to chestnut brown; branchlets yellowish green to brown, often villous when young. Leaves 2–7 cm long, 1–2.5 cm broad, oblanceolate to elliptic, acute to obtuse, with base acute; margins entire; mostly glabrous above, glabrous or very sparsely pubescent, somewhat reticulately veined below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 6–9 mm long, sparsely to densely pubescent; scales black; pedicels shorter than the scales. Male flowers with 2 stamens. Alpine tundra and alpine meadows; Rocky Mountains.

Salix arctophila Cock.

TRAILING WILLOW

Dwarf shrubs, usually trailing along the ground; branches chestnut brown to greenish gray brown; branchlets yellowish green. Leaves 2–4 cm long, 1–2 cm broad, obovate to broadly elliptic, obtuse to acute, with base acute; margins entire to shallowly crenulate, glandular; glabrous, often glossy green above, very sparsely pubescent or glabrous, glaucous below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 5–6 mm long, thinly pubescent to subglabrous; scales black; pedicels shorter than the scales. Male flowers with 2 stamens. Tundra; Boreal forest.

Salix barclayi Anderss.

BARCLAY'S WILLOW

Shrubs 1–3 m high; branches dark brown; branchlets yellowish green at first villous, later subglabrous. Leaves 3–8 cm long, 1.5–4 cm broad, acuminate to acute, with base acute to subcordate; margins glandular serrulate to subentire; the upper side pubescent along the midrib, glabrous and glaucous below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 5–7 mm long, glabrous; scales dark brown or brown only in the upper half; pedicels shorter than to as long as the scales. Male flowers with 2 stamens. Along creeks, rivers, and lakeshores; Rocky Mountains.

Low shrub to 1 m high; branches reddish brown, usually pubescent with conspicuous leaf scars; branchlets short, with short internodes, villous. Leaves 4–8 cm long, 1–2 cm broad, elliptic, acute, with base acute; margins entire to obscurely serrulate; sparsely pubescent above, densely white pubescent below. Catkins appearing before the leaves; sessile. Capsules 4–6 mm long, densely pubescent; scales black; pedicels shorter than scales. Male flowers with 2 stamens. Alpine meadows and slopes; Rocky Mountains.

Salix bebbiana Sarg.

BEAKED WILLOW

Shrubs to small trees 1–10 m high; branches reddish brown; branchlets light brown, at first densely pubescent. Leaves 2–7 cm long, 1–2.5 cm broad, acute to obtuse, with base acute to obtuse; margins entire to crenate, somewhat revolute; thinly pubescent to subglabrous above; glaucous, rugose, subglabrous to pubescent below. Catkins appearing shortly before or at the same time as the leaves; terminating short leafy peduncles. Capsules 5–9 mm long, pubescent to subglabrous; scales yellow, in part deciduous; pedicels as long as or longer than the scales. Male flowers with 2 stamens. One of the most common willows; around sloughs, along rivers and lakeshores, and in woods; throughout the Prairie Provinces.

Salix brachycarpa Nutt. (Fig. 88)

SHORT-CAPSULED WILLOW

Small shrubs, usually 0.5–1.0 m high, occasionally to 3 m high; branches stout, grayish to reddish brown, the outer layers of bark peeling; branchlets pubescent. Leaves 1–3 cm long, 0.5–1.5 cm broad, obovate to elliptic, obtuse to acute, with base acute; petioles very short. Catkins appearing at the same time as the leaves; terminating short leafy peduncles; seldom more than 2 cm long. Capsules 3–5 mm long, pubescent; bracts yellowish; pedicels shorter than the bracts. Male flowers with 2 stamens. Sand dunes, gravel bars, and marsh areas; in various parts of the Prairie Provinces, and in the Rocky Mountains.

Salix calcicola Fern. & Wieg.

LIME WILLOW

Shrubs 0.5–2 m high; branches pubescent, becoming glabrate with age; branchlets stout with inconspicuous leaf scars, pubescent. Leaves 2–5 cm long, 1.5–3.5 cm broad, ovate to almost orbicular, with base rounded to cordate; margins subentire to glandular denticulate, acute; rugose green above, glaucous below. Catkins appearing before the leaves; sessile. Capsules 7–9 mm long, glabrous; scales dark brown to black; pedicels shorter than the scales. Male flowers with 2 stamens. Tundra; Hudson Bay; rare in Rocky Mountains.

Salix candida Fluegge

HOARY WILLOW

Shrubs to 3 m high; branches dark brown; branchlets densely tomentose to sparsely pubescent. Leaves 5–8 cm long, 0.7–2 cm broad, narrowly elliptic; margins entire to subentire, revolute; dull green, somewhat tomentose, with sunken veins above; densely white tomentose below; acute; base acute. Catkins appearing at the same time as the leaves; terminating short leafy peduncles. Capsules 6–8 mm long, white tomentose; scales brown; pedicels shorter than the scales. Male flowers with 2 stamens. In wet, usually somewhat saline, swampy areas; throughout the Prairie Provinces.



Fig. 88. Short-capsuled willow, *Salix brachycarpa* Nutt.

Low shrub 0.2–1.2 m high; branches dark brown, usually pubescent; branchlets densely white tomentose. Leaves 2.5–5.5 cm long, 1.5–3.5 cm broad, elliptic to ovate-elliptic; margin subentire or entire to serrulate; pubescent on both sides. Catkins appearing after the leaves; terminating leafy peduncles. Capsules 4–6 mm long, glabrous; scales brown to dark brown, or brown only in upper half; pedicel shorter than to as long as the scales. Male flowers with 2 stamens. Gravel beds along rivers and lakes; in Rocky Mountains.

Salix discolor Muhl.

PUSSY WILLOW, DIAMOND WILLOW

Large shrubs or small trees up to 6–7 m high; bark grayish brown; branches reddish brown; branchlets rather stout yellowish to reddish brown, at first pubescent, later glabrous. Leaves 5–8 cm long, 2–3.5 cm broad, acute to short acuminate, with base acute to obtuse; margin entire to undulate-crenate; dark green, often somewhat glossy above, glaucous, with strongly raised veins below; somewhat pubescent to glabrous on both sides; veins branching before reaching margin. Catkins appearing long before the leaves; sessile, often with 2 or 3 small bracts. Capsules 6–10 mm long, densely puberulent; scales black; pedicels shorter than to as long as the scales. Male flowers with 2 stamens. One of the most common and earliest flowering willows; around sloughs and lakes, along riverbanks, and in woods and swamps; throughout the Prairie Provinces.

Salix fragilis L.

BRITTLE WILLOW

Tree up to 20 m high; bark rough and fissured on old trees; branches grayish brown; branchlets yellow to reddish, easily breaking at base. Leaves 7–15 cm long, 2–4 cm broad, lanceolate; margin glandular serrate; dark green above, glaucous below. Catkins appearing at the same time as the leaves; terminating leafy peduncle. Capsules 4–6 mm long, glabrous; scales yellow, deciduous; pedicels shorter than the scales. Male flowers with 2 stamens. Introduced from Eurasia, and planted in shelterbelts.

Salix glauca L.

SMOOTH WILLOW

Shrubs to about 1 m or occasionally 2 m high; branches dull grayish; branchlets grayish tomentose. Leaves 2.5–10 cm long, 1–3.5 cm broad, obovate to elliptic; margins entire; dark green, pubescent to glabrous above, glaucous and usually villous below; acute to acuminate or obtuse, with base acute to obtuse. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 3–5 mm long, pubescent; scales yellow to light brown; pedicels shorter than the scales. Male flowers with 2 stamens. Along rivers, in muskeg; Rocky Mountains, Boreal forest.

Salix herbacea L.

Small, creeping shrubs; stems and branches buried, glabrous; branchlets chestnut brown, glabrous. Leaves 1–3 cm long, 1–2.5 cm broad, oval to suborbicular; margins serrate or crenate-serrate; glabrous on both sides. Catkins appearing later than the leaves; terminating leafy peduncles. Capsules 3–6 mm long, glabrous; scales brown to dark brown; pedicels shorter than the scales. Male flowers with 2 stamens. Tundra; Hudson Bay.

Salix humilis Marsh.

GRAY WILLOW

Shrubs 1–3 m high; branches grayish brown; branchlets pubescent to subglabrous, yellowish. Leaves 3–12 cm long, 1–3 cm broad, oblanceolate to ovate; margins entire to sparsely dentate; acute to short acuminate; base acute; dark green above, glaucous; somewhat rugose and pubescent below. Catkins appearing before the leaves; sessile to subsessile. Capsules 7–9 mm long, pubescent; scales black; pedicels shorter than to as long as the scales. Male flowers with 2 stamens. Openings around sloughs and meadows; Boreal forest.

Salix interior Rowlee

SANDBAR WILLOW

Shrubs 0.5–4 m high, with extensively creeping root systems; stems and branches grayish brown; branchlets reddish, at first sparsely pubescent, becoming glabrous. Leaves very variable, 3–15 cm long, 0.3–1 cm broad, narrowly oblong to linear; margins distantly denticulate to subentire, with teeth 0.5–1 cm apart; green and glabrous above, pale green, pubescent to subglabrous below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 5–8 mm long, glabrous to somewhat pubescent; scales yellowish, deciduous; pedicels shorter than the scales. Male flowers with 2 stamens. This species belongs to the group Longifoliae, which has several highly variable species. *S. interior* is also highly variable, and has been subdivided into several varieties and species on the basis of leaf length and width, pubescence, and other characters. The form with very narrow leaves, 3–4 mm broad, is var. *pedicellata* (Anderss.) Ball; the form with small, permanently pubescent leaves is var. *wheeleri* Rowlee (*S. exigua* Nutt.). The species is also considered to be *S. fluviatilis* Nutt. with the two previously mentioned varieties then named var. *sericans* (Nees) Boiv. with the forma *hindsiana* (Berth.) Boiv.

The “typical” plants occur around sloughs and lakes, along creeks, rivers, and canals throughout the Prairie Provinces; var. *pedicellata* and var. *wheeleri* appear to be more restricted to drier locations and sand dune areas.

Salix lucida Muhl.

SHINING WILLOW

Shrubs or small trees 1–6 m high; bark brown; branches brown, sometimes sparsely pubescent; branchlets reddish brown, sparsely pubescent. Leaves 5–15 cm long, 1.5–4 cm broad, usually ovate, long acuminate; base acute to obtuse or cordate, with several glands at apex of petiole; margin glandular serrate; dark green and often glossy above, pale green to glaucous below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 5–7 mm long, glabrous; scales yellow, deciduous; pedicels shorter than the scales. Western specimens usually have the leaves glaucous below (*S. lasiandra* Berth.), eastern ones have the leaves light green. Along rivers and in meadows; throughout the Prairie Provinces.

Salix lutea Nutt.

YELLOW WILLOW

Shrubs 2–5 m high; branches grayish brown; branchlets reddish brown. Leaves 4–10 cm long, 1.5–4 cm broad, lanceolate, acute to short acuminate, with base obtuse to somewhat cordate; margins serrulate to entire; yellowish

green above, glaucous below. Catkins appearing at the same time as or shortly before the leaves; subsessile on a short leafy peduncle. Capsules 4–5 mm long, glabrous; scales brown; pedicels as long as or longer than the scales. Male flowers with 2 stamens. Along streams and lakeshores; throughout the Prairie Provinces.

Salix maccalliana Rowlee

VELVET-FRUITED WILLOW

Shrubs 1–3 m high; branches reddish to dark brown; branchlets reddish brown. Leaves 4.5–7 cm long, 0.6–2 cm broad, elliptic to oblong; margins entire to glandular serrate or crenulate; acute to short acuminate; base acute to obtuse; dark green and glossy above, pale green, somewhat reticulate-veined below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 7–10 mm long, pubescent; scales brown, occasionally only in the upper half; pedicels shorter than to as long as the scales. Male flowers with 2 stamens. Lakeshores and muskeg; throughout the Prairie Provinces.

Salix mackenzieana (Hook.) Barratt

MACKENZIE WILLOW

Shrubs or small trees 2–5 m high; branches grayish brown; branchlets chestnut brown. Leaves 6–10 cm long, 2–3.5 cm broad, ovate-lanceolate; margins glandular serrulate; acute to long acuminate; base obtuse to cordate; somewhat glossy green above, glaucous and reticulate below. Catkins appearing shortly before the leaves; terminating leafy peduncles. Capsules 4–6 mm long, glabrous; scales brown; pedicel longer than the scales. Male flowers with 2 stamens. Along creeks and rivers, on lakeshores; Boreal forest.

Salix monticola Bebb

MOUNTAIN WILLOW

Shrubs 1–3 m high; branches dark reddish brown; branchlets yellowish. Leaves 4–6 cm long, 1.5–3 cm broad, elliptic to ovate-elliptic; margins glandular serrulate to crenate; acute to short acuminate; base obtuse to subcordate; green above, glaucous below. Catkins appearing before the leaves; sessile to subsessile; sometimes with one or two small bracts at base. Capsules 6–8 mm long, glabrous; scales dark brown; pedicels smaller than to as long as the scales. Male flowers with 2 stamens. Lakeshores, muskeg, and marshy areas; throughout the Prairie Provinces.

Salix myrtillofolia Anderss.

MYRTLE-LEAVED WILLOW

Low shrubs usually less than 1 m high; branches grayish brown; branchlets greenish to reddish brown. Leaves 2–6 cm long, 0.8–2.5 cm broad, elliptic to obovate; margins fine glandular serrulate to crenate; green, often glossy above, pale green, often somewhat reticulate below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 4.5–7 mm long, glabrous; scales black or black-tipped only; pedicels shorter than to as long as the scales. Male flowers with 2 stamens. In muskeg, along wet lakeshores, and riverbanks; throughout Boreal forest.

Salix pedicellaris Pursh

BOG WILLOW

Low shrubs 0.2–1.5 m high; branches grayish brown; branchlets reddish yellow to brownish. Leaves 2–6 cm long, 5–15 mm broad, elliptic to narrowly

obovate; margins entire often revolute; acute to obtuse; base obtuse to acute; dull green and somewhat glaucous above, glaucous below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 5–6.5 mm long, glabrous; scales light brown; pedicels longer than scales. Male flowers with 2 stamens. In var. *athabascensis* (Raup) Boiv. the capsules and leaves are somewhat pubescent. Muskeg and marshy areas; in Boreal forest.

Salix pellita Anderss.

SATIN WILLOW

Shrubs or small trees 3–5 m high; branches brown; branchlets yellowish or greenish brown, brittle. Leaves 4–12 cm long, 0.8–2.0 cm broad, linear-lanceolate; acuminate; base acute to obtuse; margins entire to obscurely serrate, revolute; glabrous above, densely silky pubescent below. Catkins appearing shortly before or at the same time as the leaves; subsessile, often with 2 or 3 bracts at base. Capsules 4–6 mm long, pubescent; scales black; pedicels shorter than the scales. Male flowers with 2 stamens. Lakeshores, riverbanks, swamps, and muskeg; in Boreal forest. In forma *psila* Schn., the leaves are glabrescent and glaucous below. Local in the area where the species occurs. In Alberta specimens occur in which the pubescence is shorter and more uniform; these have been named var. *angustifolia* (Bebb) Boiv., but are considered to be a distinct species, *S. drummondiana* Barr., by some taxonomists.

Salix pentandra L.

BAY-LEAVED WILLOW

Usually small trees up to 7 m high; branches brown; branchlets reddish brown, shiny. Leaves 4–12 cm long, 2–4 cm broad, ovate to elliptic; long acuminate; base obtuse; dark green and usually very glossy above, pale green below; margins glandular serrulate. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 5–7 mm long, glabrous; scales yellow, deciduous; pedicels shorter than the scales. Male flowers with 5 stamens. A Eurasian species, introduced and planted as an ornamental.

Salix petiolaris Sm.

BASKET WILLOW

Shrubs or small trees 2–7 m high; branches reddish brown; branchlets at first yellowish, later reddish. Leaves 4–12 cm long, 0.8–2.5 cm broad, narrowly lanceolate; acute to acuminate; base acute; margins subentire to rather closely glandular serrulate; dark green above, glaucous below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 6–8 mm long, pubescent; scales brown; pedicels as long as or longer than the scales. Male flowers with 2 stamens. Meadows, lakeshores, and along streams; throughout the Prairie Provinces.

Salix planifolia Pursh

FLAT-LEAVED WILLOW

Shrubs 1–4 m high; branches dark grayish to reddish brown; branchlets greenish brown, often at first pubescent, later glabrous. Leaves 3–6 cm long, 1–2 cm broad, elliptic to lanceolate; acute; base acute; margins subentire to glandular serrulate; glossy green with somewhat sunken veins above; glaucous below. Catkins appearing before the leaves; sessile, sometimes with 2 or 3 small bracts at the base. Capsules 5–6 mm long, pubescent; bracts dark brown or black; pedicels shorter than the bracts. Male flowers with 2 stamens. One of the most common species; around sloughs and lakes, along streams, in

swampy areas; throughout the Prairie Provinces. This species is considered to be *S. phylicifolia* L. or *S. phylicifolia* ssp. *planifolia* (Pursh) Hiitonen by some authors. *S. phylicifolia*, however, has catkins terminating leafy peduncles.

Salix pyrifolia Anderss.

BALSAM WILLOW

Shrubs usually to 4 m high; branches brown; branchlets greenish yellow when young, becoming reddish brown. Leaves 4–12 cm long, 2–4 cm broad; ovate to lanceolate-oblong; margins glandular serrulate to crenate; acute to acuminate, with the apex often asymmetric; base obtuse to cordate; dull green above, glaucous and finely reticulate below; leaves thin and somewhat translucent; often purplish when young, fragrant, especially when crushed. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 5–9 mm long, glabrous; scales light brown or purplish; pedicels longer than the scales. Male flowers with 2 stamens. Sloughs and lakeshores, muskeg, swamps, and riverbanks; Boreal forest.

Salix reticulata L.

SNOW WILLOW

Dwarf shrubs, with stem buried; branches short, light brown; branchlets greenish yellow to light brown. Leaves 1–6 cm long, 0.8–5 cm broad; orbicular to suborbicular; obtuse to somewhat retuse; base obtuse to cordate; margins subentire to somewhat crenate, revolute; glossy dark green above, with the veins sunken; pale green, conspicuously reticulately veined, thinly pubescent to glabrous below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 4–5 mm long, pubescent; scales yellowish to reddish; pedicels shorter than the scales. Male flowers with 2 stamens. Tundra; in the Boreal forest of northern Manitoba and Saskatchewan. A somewhat smaller plant with only the branchlets clearly above ground occurs in alpine tundra in the Rocky Mountains; this plant is considered as var. *navalis* (Hook.) Anderss. or *S. nivalis* Hook. by some authors.

Salix serissima (Bailey) Fern.

AUTUMN WILLOW

Shrubs to 4 or 5 m high; stems grayish brown; branches brown to olive brown; branchlets yellowish brown, shiny. Leaves 4–8 cm long, 1–3 cm broad, lanceolate to elliptic-lanceolate; acute to short acuminate; base acute; margins fine glandular serrulate; green and often glossy above, lighter green and subglaucous below. Catkins appearing after the leaves, maturing in late summer; terminating leafy peduncles. Capsules 7–12 mm long, glabrous; scales yellow, deciduous; pedicels shorter than to as long as the scales. Male flowers with 5 stamens. Swampy areas; Boreal forest.

Salix sitchensis Sanson

SITKA WILLOW

Shrubs 0.5–3 m high; stems grayish; branches dark to grayish brown; branchlets densely pubescent when young, later glabrescent. Leaves 3–10 cm long, 1.5–5 cm broad, narrowly elliptic to obovate; obtuse, often with an asymmetric acuminate tip; base acute; margins subentire, distantly glandular; dull green, sparsely pubescent above, densely silky pubescent below. Catkins appearing at the same time as the leaves; terminating leafy peduncles. Capsules 3–5.5 mm long, pubescent; scales dark brown, or with a dark brown apex only; pedicels shorter than the scales. Male flowers with 1 stamen. Banks of streams and lakeshores; Rocky Mountains.

Depressed shrubs 0.25–0.75 m high. Branches brown with conspicuous leaf scars; branchlets brown, pubescent. Leaves 2–6 cm long, 1.5–5 cm broad, oval to broadly oblong or suborbicular; obtuse to retuse; base obtuse to subcordate; margins subentire, often somewhat revolute; deep green, rugose above, densely long pubescent and glaucous below. Catkins appearing after the leaves; subsessile or on naked peduncles. Capsules 4–5 mm long, pubescent; scales dark brown; pedicels shorter than the scales. Wet, rocky, usually shaded habitats on lakeshores in Boreal forest, and subalpine locations in the Rocky Mountains.

MYRICACEAE—bayberry family

Myrica sweet gale

Myrica gale L.

SWEET GALE, BOG-MYRTLE

A shrub to 100–120 cm high, with wedge-shaped leaves up to 5 cm long and 1–2 cm wide, which have a pleasing odor when bruised. Flowers in catkins before leaves appear, unisexual, the male and female inflorescence usually on separate plants. Plants bearing flowers of one sex one year have been known to produce flowers of the opposite sex the next year. Fruit in the form of a small nutlet, coated with a resinous wax and having two wing-like scales, borne in small heads. Found along stream banks and in swamps on acid soil; Boreal forest.

BETULACEAE—birch family

Trees or shrubs with unisexual flowers, both sexes borne on the same plant. Flowers in catkins, fruit a nutlet borne in a cone-like head or a nut enclosed in bracts.

- 1. Nuts wingless, enclosed in a leafy involucre. 2
- Nuts mostly winged, not enclosed in a leafy involucre. 3
- 2. Pistillate flowers 2–4; inflorescence capitate; involucre not inflated; nut large, acorn-like. *Corylus*
- Pistillate flowers several; inflorescence ament; involucre inflated; nut small, achene-like. *Ostrya*
- 3. Fruiting bracts woody, persistent; nutlets narrow-winged or with leathery margins; stamens 4. *Alnus*
- Fruiting bracts thin, deciduous; nutlets mostly broadly winged; stamens 2. *Betula*

Alnus alder

- Fruit thin-margined; flowering before the leaves. *A. incana*
- Fruit distinctly winged; flowering after the leaves. *A. viridis* var. *sinuata*

Alnus incana (L.) Moench SPECKLED ALDER

A tall shrub or small tree with elliptic leaves, dark green above and paler below, usually with some hairiness on the veins on the underside. Flowers developing before leaves appear; fruit not winged. Common; along streams, river valleys, and other habitats; Boreal forest, less common in Parkland. Syn.: *Alnus rugosa* (Du Roi) Spreng. var. *americana* (Regel) Fern.; *A. tenuifolia* Nutt.

Alnus viridis (Chaix) DC. var. *sinuata* Regel GREEN ALDER

A shrub 2–3 m high with oval leaves, which are often sticky on the underside. Fruit with thin wings on both sides. Fairly abundant; in dry sandy, coniferous woods; Boreal forest. Syn.: *Alnus crispa* (Ait.) Pursh.

Betula birch

- 1. Trees with white, papery bark. *B. papyrifera*
Shrubs or small trees with reddish or brown bark. 2
- 2. Wing of seed broader than body of seed; tall shrub or small tree; leaves sharply double-toothed. *B. occidentalis*
Wing of seed narrower than body of seed; low bushes or shrubs. 3
- 3. Young twigs and branches with scattered hairs, slightly glandular; leaves tapering at base; fruit with wing half as wide as body. *B. glandulifera*
Twigs and branches not hairy, densely glandular; leaves rounded at base; fruit with very narrow wing. *B. nana* var. *sibirica*

Betula glandulifera (Regel) Butler SWAMP BIRCH

A shrub, with leaves 20–30 mm long, dark green above and yellowish or reddish green beneath. Twigs and undersides of young leaves with very fine hairs. Seed with a distinct wing half the width of the seed. Around swamps and bogs; in Boreal forest and Rocky Mountains. Sometimes considered *B. pumila* L. var. *glandulifera* Regel.

Betula nana L. var. *sibirica* Led. SCRUB BIRCH

A shrub 30–150 cm high, with glandular twigs and small rounded leaves 5–20 mm long. Twigs and leaves without any hairiness, but with resinous glands. Seed with a very narrow wing, sometimes almost lacking. Fairly common; in marshes, sloughs, and boggy places; Boreal forest, Rocky Mountains. An arctic-alpine species. Syn.: *Betula glandulosa* Michx.

A small tree or large shrub up to 10 m high, growing in many-stemmed clumps, with ovate or almost round leaves 2–5 cm long. Inflorescence a cylindrical catkin; fruit borne in cylindrical, cone-like heads; seeds with broad wings. Common in low places, stream banks, depressions in sandhills; throughout the Prairie Provinces. Syn.: *B. fontinalis* Sarg.

Betula papyrifera Marsh.

WHITE BIRCH, CANOE BIRCH

A tree with white bark, up to 15 m high in favorable locations. Leaves ovate to rhomboid, serrate, dark green above. Along rivers, in openings, and somewhat moister sites; commonly in cutover areas. Three forms can be distinguished:

- 1. Leaves glabrous; twigs densely glandular. var. *neoalaskana* (Sarg.) Raup
 Leaves pubescent along the nerves below; twigs not densely glandular. 2
- 2. Pubescence consisting of tufts of hairs in the axils of the veins; leaves rounded to truncate at the base. var. *papyrifera*
 Pubescence more or less velvety below, often also pubescent above; leaves cordate at the base. var. *cordifolia* (Regel) Fern.

Corylus hazelnut

Shrubs or small trees with much-branched stems and smooth bark. Leaves broadly oval, to 10 cm long. Flowers in catkins, which are produced before leaves develop, the slender red stigmas quite conspicuous. Fruit a true nut enclosed in two leaf-like bracts.

- Bracts not much united, barely covering nut. *C. americana*
- Bracts united and produced into long beak, completely enclosing nut, and extending about 3 cm beyond nut. *C. cornuta*

Corylus americana Walt. (Fig. 89A)

AMERICAN HAZELNUT

Shrub 1–3 m high, with ovate leaves covered with pinkish hairs. Nuts rather compressed and enclosed in two distinct leafy bracts, which are slightly longer than the nut. Common throughout eastern Boreal forest and Parkland, but unusual farther west.

Corylus cornuta Marsh. (Fig. 89B)

BEAKED HAZELNUT

Similar in growth to *C. americana*, but with leaves sometimes hairless or with flattened sparse hairs. Nuts, which are very little compressed, enclosed in the united bracts, extending about 3 cm beyond the nut, forming a kind of beak. Fairly plentiful in woodlands and moist hillsides; throughout Parklands and Boreal forest.




Fig. 89. A, American hazelnut, *Corylus americana* Walt.; B, beaked hazelnut, *Corylus cornuta* Marsh.

Ostrya ironwood

Ostrya virginiana (Mill.) K. Koch

HORN HOPBEAM

Trees to 20 m high; the leaves narrowly to broadly oblong or ovate, short acuminate, serrate. Catkins 3–5 cm long, at maturity resembling a hops strobile, the involucral bracts inflated, 1–3 cm long, loosely enclosing the fruit. Very rare; dry to moist woods; southeastern Boreal forest.

FAGACEAE—beech family

Quercus oak

Quercus macrocarpa Michx.

BUR OAK

Trees with gray, flaky bark and very hard wood, 15 m high in eastern locations, but a small scrubby tree farther west. Leaves bright, shiny green above and grayish white and slightly woolly beneath, deeply cut with rounded lobes. Flowers unisexual, in slender catkins. Fruit an acorn resting in a shallow fringed cup. Common in southeastern Parkland, but farther west only found in the valleys of the rivers tributary to the Assiniboine River in Saskatchewan.

ULMACEAE—elm family

Leaves with 2 prominent veins beside the midrib; fruit a drupe with large stone. *Celtis*

Leaves with a single midrib; fruit a thin, broadly winged samara. *Ulmus*

Celtis nettletree

Celtis occidentalis L.

HACKBERRY

Trees to 15 m high, with prominently ridged bark. Leaves 6–12 cm long, conspicuously serrate. Fruit ellipsoid, 7–13 cm long, dark red to nearly black. Very rare; moist woods; southeastern Boreal forest.

Ulmus elm

Ulmus americana L.

AMERICAN ELM

A tall tree with reddish bark and smooth twigs. Leaves oval, with soft hairs beneath and rough above, veins very prominent. Flowers perfect, in little bunches early in spring before the leaves appear. Fruit a one-seeded, flat samara, or winged fruit. A common tree in southeastern Parkland, rare along rivers in Prairie, but in various locations planted and escaped.

CANNABINACEAE—hemp family

Humulus hop

Humulus lupulus L.

COMMON HOP

A perennial twining climber, often 3–6 m long; leaves usually large, palmately 3- to 7-lobed, with cordate bases, opposite, 2–8 cm across, and bearing many tiny whitish or yellowish glandular spots on the undersides. Upper leaves near the inflorescence often not divided, merely toothed. Stems rough to the touch, with fine, stiff, reflexed hairs. Male flowers in loose panicles, green, with 5 sepals. Female flowers on separate plants, in catkin-like or cone-like heads in the axils of the leaves, with broad greenish or yellowish imbricated bracts covering the fruits. These clustered fruits (hops) vary from 2 to 5 cm in length. Fairly common in moist places in southeastern Parklands, particularly in river valleys, and scattered throughout the Prairies. Contact with this plant causes a form of dermatitis in some individuals.

URTICACEAE—nettle family

Plants often with stinging hairs and greenish flowers, which are borne in clusters at the junction of the stem and the leaf stalk.

1. Plants without stinging hairs. *Parietaria*
Plants with stinging hairs. 2
2. Leaves alternate; sepals 5. *Laportea*
Leaves opposite; sepals 4. *Urtica*

Laportea wood nettle

Laportea canadensis (L.) Gaud.

WOOD NETTLE

A perennial with alternate and toothed leaves with stinging hairs. Fairly common; in rich woodlands; southeastern Parklands and Boreal forest.

Parietaria pellitory

Parietaria pensylvanica Muhl.

AMERICAN PELLITORY

A weak-stemmed, hairy, annual plant, 10–50 cm high, with opposite, lanceolate, 3-nerved leaves 2–8 cm long. Greenish flowers in dense clusters around the leaf axils. Very rare; shaded areas; southeastern Parklands.

Urtica nettle

Plants perennial, with extensive rootstocks;

stipules 5–15 mm long. *U. dioica* var. *procera*

Plants annual; stipules 1–3 mm long. *U. urens*

Urtica dioica L. var. *procera* (Muhl.) Wedd. (Fig. 90)

STINGING NETTLE

A perennial with very coarse rootstocks, which have pink offshoots. Plants have tall, straight stems, usually square in cross section, and with ovate to lanceolate, serrate (toothed), opposite leaves bearing stinging hairs. Greenish flowers borne in clusters at the junction of stem and leaf stalk. Very common around sloughs, moist places, and bushes; throughout the Prairie Provinces. Syn.: *U. gracilis* Ait.

Urtica urens L.

ENGLISH NETTLE

Annual plants with erect stems 10–50 cm high, simple or branched, with numerous stinging hairs. Leaves elliptic to ovate, deeply coarse serrate; flower clusters oblong. A rare weed; in gardens; Prairies and Parklands.

SANTALACEAE—sandalwood family

Smooth, hairless perennials with rootstocks, partly parasitic on the roots of other plants. Flowers perfect, without petals, but with 5 sepals, which are united to form a bell-like tube. Fruit a 1-seeded drupe (dry or fleshy fruit with a hard nut-like seed in the center).

Comandra comandra

Flowers axillary. *C. livida*

Flowers in terminal or subterminal clusters. *C. umbellata*

Comandra livida Rich.

NORTHERN COMANDRA

A slender, erect plant, 10–30 cm high, with oval leaves 1–3 cm long on short stalks. Flowers in the axils of the leaves. Fruit a spherical red drupe about 3 mm in diam, edible. Not very common; but found in moist woods; Boreal forest, Riding Mountain, Rocky Mountains. Syn.: *Geocaulon lividum* (Richards.) Fern.

Comandra umbellata (L.) Nutt. (Fig. 91)

BASTARD TOADFLAX

An erect plant, from a white creeping rootstock, 6–30 cm high, usually in bunches of several plants from the same rootstock. Leaves linear or linear-lanceolate, 10–25 mm long, without stalks, borne alternately on the stems. Flowers at the summit of stems, greenish white to pinkish, small, about 5 mm long. Fruit ovoid, 3–8 mm long. This plant is often attacked by a tiny insect that causes small round galls about 6 mm in diam that are sometimes mistaken for the fruit.

Three forms can be distinguished:

1. Leaves thin, distinctly veined; calyx lobes 2–3 mm long; fruit 3–6 mm across. var. *umbellata*
 Leaves thick, not distinctly veined; calyx lobes 3–4 mm long; fruit 5–8 mm across. 2
2. Plants usually small, to 15 or 20 cm high; panicle branches short; the panicle ovoid. var. *angustifolia* (DC.) Torr.



Fig. 90. Stinging nettle, *Urtica dioica* L. var. *procera* (Muhl.) Wedd.



Fig. 91. Bastard toadflax, *Comandra umbellata* (L.) Nutt.

Plants usually 20–40 cm high; panicle branches long; the panicle more corymbiform. var. *pallida* (DC.) G. N. Jones

The var. *umbellata*, Richards comandra, is the common form in openings, open forests, and disturbed areas throughout Boreal forest (syn.: *C. richardsiana* Fern.); var. *angustifolia*, pale comandra, is very common in dry grasslands of the Prairies (syn.: *C. pallida* A. A.); var. *pallida* occurs in grassland in southern Rocky Mountains (syn.: *C. pallida* A. DC.). Many plants, such as those of the Peace River and Parkland areas, are intermediate between these forms, but they approach one form more closely than the others.

LORANTHACEAE—mistletoe family

Arceuthobium dwarf mistletoe

Stems 5–10 cm high; staminate flowers terminal. *A. americanum*

Stems 0.5–2 cm high; all flowers axillary. *A. pusillum*

Arceuthobium americanum Nutt. AMERICAN MISTLETOE

A parasitic plant on conifers, mostly *Pinus*, often causing witches'-broom. Stems fragile, greenish yellow, often much-branched; leaves opposite, scale-like. Fruit 1-seeded, berry-like, 2–3 mm long, ejecting the seed explosively. Boreal forest.

Arceuthobium pusillum Peck DWARF MISTLETOE

A parasitic plant, mostly on black spruce, more rarely on white spruce or tamarack. Stems fragile, greenish brown, simple or sparingly branched; leaves scale-like. Fruit about 2 mm long. Eastern Boreal forest.

ARISTOLOCHIACEAE—birthwort family

Asarum wild ginger

Asarum canadense L. WILD GINGER

Plants with slender, branched rootstocks and 2 large, rotund or reniform leaves with a deeply cordate base, pubescent especially on the petioles. Flower solitary, with the petals minute or absent. Fruit a capsule with large wrinkled seeds. Rare; woods; southeastern Boreal forest.

POLYGONACEAE—buckwheat family

Herbs with mostly alternate, entire leaves and, except in *Eriogonum*, with sheathing stipules called ocrea. Flowers small, without petals, but with a perianth of 4–6 more or less united sepals. Achenes either triangular or lens-shaped.

1. Leaves without stipules; a whorl of bracts below flower clusters; stamens 9. *Eriogonum*
 Leaves with stipules; without whorl of bracts below flowers; stamens less than 9. 2
2. Sepals 4 or 6; stigmas tufted. 4
 Sepals 5, more or less equal; stigmas not tufted. 3
3. Seeds generally enclosed by somewhat enlarged calyx, but, if protruding from calyx, the leaves are long and narrow. *Polygonum*
 Seeds protruding from calyx; leaves broadly arrow-shaped. *Fagopyrum*
4. Sepals 6, the three inner ones large, 15–20 mm, enclosing the fruit; leaves lanceolate. *Rumex*
 Sepals 4, the inner ones erect, 4–6 mm; leaves reniform. *Oxyria*

Eriogonum umbrellaplant

1. Plants annual; inflorescence a diffusely branched raceme. *E. cernuum*
 Plants perennial; inflorescence a more or less dense umbel. 2
2. Inflorescence subtended by leaf-like bracts. 3
 Inflorescence subtended by small, reduced bracts. 4
3. Involucral bracts linear, ascending. *E. flavum*
 Involucral bracts lanceolate or oblong, spreading or reflexed. *E. umbellatum*
4. Leaves oval, obovate or orbicular, about 1 cm long. *E. ovalifolium*
 Leaves lanceolate or linear-lanceolate. 5
5. Leaves in dense crowded rosettes; flowers yellow or reddish. *E. androsaceum*
 Leaves in loose rosettes; flowers whitish or pinkish. *E. multiceps*

Eriogonum androsaceum Benth.

CUSHION UMBRELLAPLANT

Plant forming small, compact cushions; scapes 2–10 cm high. Leaves 1–2 cm long, oblanceolate to spatulate, densely villose, later becoming glabrate. Inflorescence often a single small umbel; flowers 4–5 mm long. Not common; exposed, often rocky, areas; southern Rocky Mountains.

Eriogonum cernuum Nutt.

NODDING UMBRELLAPLANT

Low annual plants up to 30 cm high, with a whorl of small leaves near base of stem. Leaves almost circular, 1–2 cm in diam, white woolly beneath.

Flowers white or rose, borne in small clusters about 5 mm wide on stems in a much-branched inflorescence, usually drooping. Not common but abundant locally; found on badlands, abandoned fields, dry and sandy soil; throughout southwestern Prairies.

Eriogonum flavum Nutt. (Fig. 92)

YELLOW UMBRELLAPLANT

A low perennial with a very coarse, scaly, dark brown, woody, tufted root. Leaves all basal, usually linear-oblong or spatulate, 2–5 cm long, green and slightly hairy above, densely white woolly beneath. Flowers pale yellow, in umbel-like clusters at head of stems. Common; on dry and eroded hillsides, badlands, and canyons; throughout Prairies.

Eriogonum multiceps Nees

BRANCHED ERIOGONUM

A low perennial, rarely over 20 cm high, with branched stems. Leaves very narrow, to 5 cm long; the whole plant, leaves, and stem densely white woolly. Whitish or pale pink flowers in small clusters at the summits of the stems. Uncommon; on eroded banks and badlands; Prairies.

Eriogonum ovalifolium Nutt.

SILVERPLANT

A low plant with the caudex closely branched; leaves densely white tomentose. Scapes 5–15 cm high; inflorescence usually a single dense umbel, with very small bracts. Not common; dry hillsides and plains; southern Rocky Mountains.

Eriogonum umbellatum Torr.

UMBRELLAPLANT

Plant with short woody branches forming loose mats; leaves clustered in numerous rosettes, 3–5 cm long, glabrous above, dense white tomentose below. Scapes 10–30 cm high, pubescent below; umbel with rays 1–6 cm long; bracts 10–25 mm long. Exposed sites, summits, and slopes; southern Rocky Mountains. Syn.: *E. subalpinum* Greene.

Fagopyrum buckwheat

Fagopyrum tataricum (L.) Gaertn. (Fig. 93)

TARTARY BUCKWHEAT

An annual weed 50–100 cm high, with somewhat triangular leaves often as broad as long, 3–10 cm long and wide. Flowers small, white, in bunches on the flowering stems, arising from the junction of the leaf stalks and stems. Seeds about the same size as wheat kernels. This weed, introduced from Asia, is a serious pest in northern and central Alberta; rare in Boreal forest. Common cultivated buckwheat, *F. esculentum* Moench, has larger white or pinkish flowers, and smooth shiny seeds. Rarely found as a weed.

Oxyria mountain sorrel

Oxyria digyna (L.) Hill

MOUNTAIN SORREL

Plant with rootstock and stout root; stems 5–30 cm high. Leaves mostly basal, alternate, long-petioled; stipule sheath loose. Flowers whorled in a dense raceme 4–6 mm long; achenes lens-shaped. Mountain meadows, rock slopes; Rocky Mountains.



Fig. 92. Yellow umbrellaplant, *Eriogonum flavum* Nutt.



Fig. 93. Tartary buckwheat, *Fagopyrum tataricum* (L.) Gaertn.

Polygonum knotweed

- 1. Plants twining; leaves broad, hastate (arrow-shaped) at base. Section: *Bilderdykia*
Plants not twining; leaves linear, lanceolate, or oblong. 2
- 2. Flowers solitary or in clusters in axils of leaves; leaves small. Section: *Avicularia*
Flowers in dense spikes, either terminal or in axils of leaves; leaves large. 3
- 3. Basal leaves with long stalks, stem leaves smaller and stalkless; stem single and unbranched. Section: *Bistorta*
Leaves all similar; stem much-branched. Section: *Persicaria*

Polygonum (Sect.: *Avicularia*) doorweed

Annual low-growing plants, often prostrate on ground, with tough stems and alternate leaves, usually narrow. Sheathing stipules usually fairly conspicuous and often white and translucent. Flowers small, borne in clusters in axils of leaves, and having a perianth of five partly united sepals. Seeds 3-angled achenes, black or brown. Plants appear to thrive when they are trampled.

- 1. Inflorescence crowded at summit of stem; bracts leaf-like. 2
Inflorescence not crowded at summit of stem; flower cluster axillary. 3
- 2. Bracts broadly white-margined; achenes jet black. *P. confertiflorum*
Bracts mostly not white-margined; achenes brown. *P. kelloggii*
- 3. Fruit reflexed or pendulous; achenes black. 4
Fruit erect or ascending; achenes brownish. 6
- 4. Calyx 3-4 mm long; plants 10-40 cm high. *P. douglasii*
Calyx 2-3 mm long; plants 5-15 cm high. 5
- 5. Leaves linear or linear-lanceolate; bracts awl-shaped. *P. engelmannii*
Leaves ovate to oblanceolate; bracts oblong or lanceolate. *P. austiniae*
- 6. Leaves oval or blunt; length-to-width ratio 2.5:1. 7
Leaves linear to lanceolate; length-to-width ratio 4:1 or more. 8
- 7. Plants erect, 3-15 cm high; flower clusters crowded at the tips of branches. *P. minimum*
Plants prostrate or decumbent, much larger; flower clusters scattered along the stems. *P. achoreum*

- 8. Plants erect or ascending, usually quite high. 9
Plants usually prostrate, depressed. 10
- 9. Leaves 10–25 mm wide, margins white. *P. erectum*
Leaves 5–15 mm wide, margins not white. *P. ramosissimum*
- 10. Petioles included in ocreae; margins of perianth segments white or pinkish; achene 2.5–3.5 mm. *P. aviculare*
Petioles exserted; margins of perianth segment bright pink; achene 3.5–4.5 mm. *P. boreale*

Polygonum (Sect.: *Bilderdykia*) wild buckwheat

- 1. Plants annual. *P. convolvulus*
Plants perennial. 2
- 2. Ocreae reflexed-bristly at base; stems sharply angled; fruit not winged. *P. cilinode*
Ocreae smooth; stems nearly terete; fruit broadly winged. *P. scandens*

Polygonum (Sect.: *Bistorta*) bistort

- Basal leaves tapering gradually to stalk; inflorescence not bulblet-bearing. *P. bistortoides*
- Basal leaves cordate or blunt-based; inflorescence bearing bulblets at lower part. *P. viviparum*

Polygonum (Sect.: *Persicaria*) smartweed

The smartweeds, especially the perennial species, are a rather confusing group of plants: they can have three phases, each phase differing from the others by so much that in some instances they have been classed as different species instead of phases of the same species. The aquatic phase is found submerged in water, is usually without hairs, and has broad leaves. The paludose (or marshy) phase is found where water has recently receded, and the plants grow in wet mud. The terrestrial phase is found where the soil has dried somewhat; the leaves are usually narrow and the plant often more or less hairy. The flowers of smartweeds are borne in conspicuous racemes, usually white and pink or greenish, and the perennial species are usually the dominant plant in the location they occupy. Most species are palatable and nutritious. Heavy pasturing may cause bighead or yellows in sheep.

- 1. Plants perennial, with extensively creeping rootstocks. *P. amphibium*
Plants annual. 2

- 2. Leaves 3–6 cm long, oblong-sagittate, cordate at base. *P. sagittatum*
 Leaves not sagittate, cuneate at base. 3
- 3. Perianth covered with brownish glands. 4
 Perianth not glandular. 5
- 4. Cleistogamous flowers present in the leaf axils. *P. hydropiper*
 Cleistogamous flowers not present. *P. punctatum*
- 5. Ocreae ciliate; peduncles not glandular. *P. persicaria*
 Ocreae not ciliate; peduncles and often perianth bearing yellow sessile glands. 6
- 6. Achene 1.5–2.0 mm wide; leaves glabrous below. *P. lapathifolium*
 Achene 2.0–3.0 mm wide; leaves sparingly pubescent below. *P. scabrum*

Polygonum achoreum Blake

STRIATE KNOTWEED

An annual growing to about 40 cm high, with oval to elliptic bluish green leaves 6–25 mm long. Common around yards and waste places, appearing to thrive on abuse. If kept mowed, it takes the place of a lawn, and helps to prevent soil erosion by wind and water. In waste places and farmyards; throughout the Prairie Provinces.

Polygonum amphibium L.

SWAMP PERSICARIA

Perennial with far-creeping blackish, branching rootstocks. Leaves 5–15 cm long, oblong-ovate, truncate, cordate or cuneate at the base, glabrous or somewhat pubescent. Ocreae often with foliaceous collar when young. Spikes 1–3 cm long. Very common; in standing water; throughout the Prairie Provinces. Syn.: *P. coccineum* Muhl. forma *natans* (Wiegand) Stanford; *P. natans* A. Eaton forma *genuinum* Stanford. The terrestrial form, with larger inflorescence, spikes 5–10 cm long, leaves often densely pubescent, and ocreae with a broad foliaceous collar, is *P. amphibium* L. var. *emersum* Michx. Very common; in mud flats, slough margins, and lakeshores; throughout the Prairie Provinces. Syn.: *P. coccineum* Muhl. forma *terrestre* (Willd.) Stanford; *P. natans* A. Eaton forma *hartwrightii* (Gray) Stanford; *P. amphibium* L. var. *stipulaceum* (Coleman) Fern.

Polygonum austiniiae Greene

Plants erect, 4–12 cm high, branched from the base. Lower leaves oval to obovate or lanceolate, petioled; upper leaves reduced, sessile. Flowers in the leaf axils, solitary or in pairs, reflexed. Achenes 2.5–3 mm long, shiny black. Dry, usually disturbed ground; southern Rocky Mountains.

Polygonum aviculare L. (Fig. 94)

DOORWEED

A semiprostrate annual plant, 10–50 cm long. Leaves fairly broad, oblong, 1.5–6 cm long, pale bluish green. The ocreae (or stipules) silvery and translucent. Very common; in waste places; throughout the Prairie Provinces.




Fig. 94. Doorweed, *Polygonum aviculare* L.

A perennial plant growing from a horizontal fleshy rootstock to 10–40 cm high. Basal leaves 6–15 cm long, lanceolate to oblong, on a stalk about as long as the blade. Stem leaves much shorter and without stalks. Inflorescence a dense, oblong raceme of white or pinkish flowers, borne at the head of the stem. A plant of swamps, stream beds, and wet places; Rocky Mountains, Cypress Hills.

Polygonum boreale (Lange) Small

NORTHERN DOORWEED

Stems prostrate or decumbent to erect, 10–50 cm long. Leaves 3–5 cm long, obovate; petioles well exerted from the ocreae. Perianth green with conspicuous pink margins. Gravel banks; Hudson Bay.

Polygonum cilinode Michx.

BINDWEED

Stems to 2 m long, twining, trailing, sometimes erect; pubescent. Leaves triangular-ovate, deeply cordate. Racemes long-peduncled, 4–10 cm long, perianth white; achene glossy black, 1.5–2 mm, with the outer calyx rarely narrowly winged. Clearings and rock outcrops; eastern Boreal forest.

Polygonum confertiflorum Nutt.

SMALL KNOTWEED

Plants annual, stems erect, 3–15 cm high, glabrous, simple or branched. Leaves 1–3 cm long, linear-lanceolate. Flower clusters crowded into short terminal spikes; perianth segments with a conspicuous white margin; achene jet black. Rare or overlooked; dry, disturbed areas; Cypress Hills.

Polygonum convolvulus L.

WILD BUCKWHEAT

An annual, tap-rooted weed, with weak twining stems and pale green arrow-shaped leaves, 1–5 cm long, and pointed at the tip. Flowers greenish or pink, in small racemes in the axils of the leaves or at the end of the stem. Seeds black, granular, three-angled, and often enclosed in a green or brown coating, and about 3 mm long. Introduced from Europe, but very common in cultivated land and waste places; throughout the Prairie Provinces.

Polygonum douglasii Greene

DOUGLAS KNOTWEED

An erect annual to 45 cm high; narrow leaves 1–5 cm long. Seeds oblong, narrow, reflexed or pendulous on the stem. Not very common; various locations; southwestern Prairies.

Polygonum engelmannii Greene

SLENDER KNOTWEED

An erect annual 5–15 cm high; stems slender, often diffusely branched at base; linear to lanceolate leaves, and axillary clusters of 1–4 flowers. Moist banks and slopes; Rocky Mountains.

Polygonum erectum L.

ERECT KNOTWEED

Plants with erect or decumbent stems 10–50 cm high; leaves elliptic to elliptic-lanceolate, finely crenulate and white-margined, 1–4 cm long, 0.5–2 cm broad. Achenes of two kinds: dark brown, shiny, punctate, about 2.5 mm long; or dull brown, 3–3.5 mm long. Waste areas and lakeshores; southeastern Parklands and Boreal forest.

Polygonum hydropiper L.

WATER-PEPPER

Stems 10–50 cm long, erect or spreading, often reddish. Leaves narrowly lanceolate to ovate-lanceolate, 2–5 cm long; ocreae short ciliate. Racemes 3–6 cm long; perianth greenish; achenes dull dark brown, 2–3 mm long. Rare; wet places; southeastern Boreal forest.

Polygonum kelloggii Greene

DWARF KNOTWEED

A small annual 3–15 cm high, erect, simple or branched; leaves 1–3 cm long, linear to linear-lanceolate; flower clusters crowded at the ends of branches; achenes dark brown. Rare; moist areas; southwestern Prairies.

Polygonum lapathifolium L.

PALE PERSICARIA

A variable annual plant, with lanceolate to ovate leaves, sometimes deep green and hairless, but more often pubescent beneath. The type with pubescent leaves was formerly listed as a separate species, *P. tomentosum* Schrank. Although hirsute and whitish underneath when young, mature plants appear glabrous. Flowers in erect spikes, usually pale pink to white; those growing under moist conditions usually paler than those of the drier locations. Very common along slough margins and low areas; occasionally in cultivated fields; throughout most of the Prairie Provinces.

Polygonum minimum S. Wats.

LEAST KNOTWEED

Small annuals 3–15 cm high, erect or spreading, branched at the base, leafy. Leaves oval to obovate, 5–15 mm long. Flower clusters of 2 or 3, axillary, perianth white-margined; achenes 2–2.5 mm long. Dry ground; southern Rocky Mountains.

Polygonum persicaria L. (Fig. 95)

LADY'S-THUMB

Large annuals, with stems 30–80 cm high, erect or ascending. Leaves narrowly lanceolate, 3–7 cm long; ocreae short-ciliate, minutely pubescent. Spikes dense, 1–5 cm long; perianth pink or reddish; achenes black, 2–2.5 mm long. Not common; wet areas; throughout the Prairie Provinces.

Polygonum punctatum Ell. var. *confertiflorum* (Meisn.) Fassett

WATER SMARTWEED

Large annuals, with stems to 60 cm high, simple or branched, erect or ascending. Leaves to 15 cm long, 2 cm wide, narrowly lanceolate. Racemes to 10 cm long, much interrupted; perianth greenish, glandular; achenes 2.5–3 mm long. Wet areas, lakeshores, waste places; eastern Parklands and Boreal forest.

Polygonum ramosissimum Michx.

BUSHY KNOTWEED

A yellowish green erect plant 15–90 cm high, with a much-branched stem and lanceolate to linear-oblong leaves to 3 cm long. Found occasionally; in sandy soil, river bottoms, and lakeshores; throughout Prairies and Parklands.

Polygonum sagittatum L.

ARROW-LEAVED TEAR-THUMB

Slender annuals, with stems 10–50 cm high, weak, reclining, with reflexed barbs. Leaves lanceolate-sagittate, deeply cordate at base. Inflorescence head-like, terminal on stems or branches; perianth pink; achene dark brown or black, 2–3 mm long. Rare; wet areas; southeastern Boreal forest.



Fig. 95. Lady's-thumb, *Polygonum persicaria* L.

Large annuals, with stems 30–80 cm high, simple or branching, erect to spreading-ascending. Leaves lanceolate to oblong-lanceolate, 3–8 cm long, white pubescent below, especially the lower ones. The upper leaves glandular-dotted. Spikes erect, usually greenish, very glandular below; achenes 2.5–3.5 mm long. Waste areas, lakeshores; western Parklands and Boreal forest.

Polygonum scandens L.

Twining perennials, with sharply angled stems to 2 m long. Leaves oblong-ovate, deeply cordate at base. Inflorescences from most upper leaf axils 5–10 cm long; perianth white; fruit broadly winged, the outer sepals expanding, to 15 mm long; achene glossy black, 3–5 mm long. Rare; shores, clearings, and margins of woods; southeastern Parklands and Boreal forest.

Polygonum viviparum L.

A perennial plant 5–25 cm high, from a scaly corm-like rootstock, with cordate-based basal leaves 5–10 cm long. Stem leaves narrow and tapering to the base, which is enclosed in a membranous sheath or ocreae. Inflorescence a dense raceme of whitish or pinkish flowers terminating the stem, but the lower flowers replaced by small bulbs. An alpine species; Rocky Mountains.

Rumex dock

Mostly perennial plants with thick roots, leaves of varying shapes and sizes. Flowers, perfect or unisexual, consisting of 6 sepals, the 3 inner ones enlarging and forming wings or valves enclosing the fruit. These valves sometimes entire and sometimes fringed with bristle-like teeth. In some species one or all three of the valves may bear a protuberance called a tubercle, useful as a character for identification. Usually in wet or marshy places; throughout the Prairie Provinces, but at least one species is a plant of the sandhills.

- 1. Valves greatly enlarging in fruit, becoming bright pink. *R. venosus*
- Valves not greatly enlarging. 2
- 2. Flowers unisexual. 3
- Flowers perfect. 5
- 3. Leaves cuneate at base, long-petioled. *R. paucifolius*
- Leaves hastate or sagittate at base. 4
- 4. Valves much larger than achene. *R. acetosa*
- Valves equaling the achene, or a little larger. *R. acetosella*
- 5. Valves distinctly lobed or toothed on the margin. 6
- Valves entire or nearly so. 7
- 6. Inflorescence continuous; valves 2.5–3 mm. *R. maritimus* var. *fueginus*
- Inflorescence with distant whorls; valves 4–5 mm. *R. dentatus*

- 7. Pedicels not jointed near the middle. 8
 Pedicels jointed. 9
- 8. Valves all with a grain-like body. *R. orbiculatus*
 Valves without grains. *R. occidentalis*
- 9. Fruit with 1–3 large grains. 10
 Fruit without grains, or with only 1 small
 grain. 12
- 10. Plants with prostrate or ascending stems;
 most stem leaves with axillary shoots. *R. salicifolius*
 Plants with erect stems; axillary shoots
 absent or few. 11
- 11. Valves entire, 3–6 mm long; leaf margins
 strongly undulate. *R. crispus*
 Valves with small teeth, 3.5–4 mm long;
 leaf margins undulate, but not strongly
 so. *R. stenophyllus*
- 12. Basal and lower stem leaves almost as
 wide as long, deeply cordate at base. *R. confertus*
 Basal and lower stem leaves much longer
 than wide. 13
- 13. Lower stem leaves linear-lanceolate,
 length-to-width ratio about 8:1; valves
 3.5–5 mm long, less than 5 mm wide. *R. fennicus*
 Lower stem leaves broadly lanceolate,
 length-to-width ratio about 4:1; valves
 4.5–6 mm long, more than 5 mm wide. *R. longifolius*

Rumex acetosa L.

SOUR DOCK

An erect plant sometimes over 60 cm high, with broadly lanceolate arrow-shaped leaves 2–12 cm long. Used as food; escaped from gardens in many parts of the Prairie Provinces.

Rumex acetosella L. (Fig. 96)

SHEEP SORREL

A low plant not over 30 cm high, with leaves narrowly hastate or spear-shaped, 2–10 cm long. In this species sepals do not enlarge in fruit. An introduced weed but only rarely found; in dry or sandy places and on acid soils.

Rumex confertus Willd.

Fall perennial, 60–100 cm high, with the basal leaves triangular cordate, to 20 cm wide; the petiole shorter than the blade. Valves about 6 mm long, 8 mm wide. Very rare; waste areas; eastern Parklands.

Rumex crispus L.

CURLED DOCK

An introduced weed, to over 1 m high in favorable localities, with leaves crinkled at the edge, the lower ones 15–30 cm long, the upper ones smaller. Fairly common; in moist places; eastern Prairies and Parklands.



Fig. 96. Sheep sorrel, *Rumex acetosella* L.

Rumex dentatus L.

TOOTHED DOCK

Plants annual, 20–70 cm high, with the basal leaves small, 2–3 times as long as wide, truncate to subcordate at the base. Valves 4–6 mm long, 2–3 mm wide, with teeth 3–6 mm long. Rare; weed in cultivated land; southwestern Alberta.

Rumex fennicus Murb.

FIELD DOCK

An introduced perennial growing to more than 1 m high, with lanceolate leaves 15–25 cm long, tapering toward the stalk. Erect, dense fruiting heads turn brown in late summer and are very conspicuous in fields and along roadsides. A tiny swelling on the stalk of fruiting valves about a third of the way up from the base distinguishes this species from western dock, which it superficially resembles. Originally introduced from Europe but has spread rapidly across the south central part of the Prairies and Parklands. Now a common plant on almost every type of soil; in upland fields and moist areas. Syn.: *R. domesticus* Hartm. var. *pseudonatronatus* Borb.

Rumex longifolius DC.

LONG-LEAVED DOCK

Stout perennial plants 60–120 cm high, with leaves 20–40 cm long, 10–15 cm wide. Valves 4.5–5.5 mm long, 5.5–6.5 mm wide. Rare; weed in moist areas; eastern Parklands and Boreal forest.

Rumex maritimus L. var. *fueginus* (Phil.) Dusen

GOLDEN DOCK

An annual species to 60 cm high, from a very fleshy root, and usually diffusely branched, with leaves lanceolate, 6–25 cm long, pale green. Stems covered with very short hairs. Fruiting heads conspicuous, golden yellow, very dense. Valves of calyx have a spine-like bristly margin, which gives the heads a prickly appearance and makes this species easy to distinguish. Found in moist and saline places, lake flats, and lakeshores, and often the dominating plant in such sites, sometimes in masses covering 1000 m² or more. Common in suitable locations; throughout the Prairie Provinces.

Rumex occidentalis S. Wats.

WESTERN DOCK

A tall, coarse, erect plant to 1 m high, with leaves broadly oblong or lanceolate, up to 30 cm long, often cordate or heart-shaped at the base. Fruiting heads dense and very conspicuous. Common; in wet or moist places; throughout western part of the Prairie Provinces.

Rumex orbiculatus Gray

GREAT WATER DOCK

Stout, erect plants to 1.5 m high, with leaves lanceolate, acute or rounded at base. Valves 5–8 mm long and wide. Rare; swamps and shallow water; Boreal forest.

Rumex paucifolius Nutt.

ALPINE SHEEP SORREL

Perennial plants with simple or branched taproot; stems 20–70 cm high; leaves mostly basal, 4–10 cm long, cuneate at the base. Valves 2.5–3 mm long, suborbicular, becoming reddish. Mountain meadows and streams; southern Rocky Mountains.

A narrow-leaved species, erect or sprawling, in low and moist places. Leaves pale green, 5–10 cm long and 10–25 mm wide. Fruiting heads narrow and usually not so dense as in most docks. Common in moist or saline places, roadside ditches, and similar sites; throughout the Prairie Provinces. Syn.: *R. triangulivalvis* (Danser) Rech. f.; *R. mexicanus* Meisn.

An introduced perennial, 1–2 m high, with narrowly lanceolate leaves to 15 cm long, narrowed at both ends and crinkly margined. Inflorescence tall, narrow, and somewhat leafy at the base. Fruiting valves toothed on the margins and bearing a tubercle on each valve; a tiny swollen joint on the lower part of each valve stalk. This plant resembles field dock in many ways. Becoming increasingly common; in southern part of Prairies.

A perennial species with running, woody rootstocks and branched stems. Sheathing stipules whitish and papery, very conspicuous. Leaves pale green, 5–15 cm long, and nearly half as wide. Calyx valves conspicuous in fruit, large, sometimes to 3 cm broad, almost round, bright red to pink. Often picked and dried to form winter bouquets. Common in suitable localities, becoming dominant in some places; on sandy soil, sand dunes, and often on railway grades and roadsides; often an early invader of a shifting sand dune; throughout Prairies and Parklands.

CHENOPODIACEAE—goosefoot family

A large and varied family consisting of both annual and perennial herbs and shrubs. Leaves usually alternate, without stipules, and often white mealy. Flowers have no petals but 2–5 sepals, and may be either perfect or unisexual. Many species are weedy and most produce vast quantities of seed. Some are edible, and nearly all are palatable to livestock, but one species is **poisonous**.

- 1. Leaves and branches opposite; fleshy-stemmed plants; leaves reduced to scales; flowers sunk into stem. *Salicornia*
- Leaves alternate. 2
- 2. Plants with spiny branches or spine-tipped leaves; calyx forming wing around mature fruit. 3
- Plants neither spiny nor with spine-tipped leaves. 4
- 3. Shrubs with spiny branches. *Sarcobatus*
- Annual herbs with fleshy spine-tipped leaves. *Salsola*
- 4. Flowers unisexual, female flowers enclosed in two bracts that enlarge after flowering. 5



A. F. Best

Fig. 97. Sand dock, *Rumex venosus* Pursh.

Flowers perfect.	9
5. Fruiting bracts covered with silvery, silky hairs; low wooly shrubs.	<i>Eurotia</i>
Fruiting bracts not covered with silky hairs.	6
6. Fruiting bracts not united; seed coat often winged at apex.	<i>Axyris</i>
Fruiting bract united, at least at base; seed coat not winged at apex.	7
7. Fruit with 2 teeth at apex; seed coat membranous.	<i>Suckleya</i>
Fruit not 2-toothed at apex; seed coat leathery.	8
8. Female flowers with no calyx or sepals, merely a pair of bracts.	<i>Atriplex</i>
Female flowers with 2 or 3 translucent sepals, which are much shorter than the bracts.	<i>Endolepis</i>
9. Calyx with only one sepal; 1–3 stamens.	10
Calyx with 3–5 sepals; 1–5 stamens.	11
10. Leaves spear-shaped; stems fleshy; flowers in clusters in leaf axils; seeds round and small.	<i>Monolepis</i>
Leaves narrow and linear; flowers single in axils of leaf-like bracts; seeds flat and large.	<i>Corispermum</i>
11. Mature calyx with wings, or minute spines.	12
Mature calyx without wings or spines.	14
12. Mature calyx with a tiny spine on each lobe.	<i>Bassia</i>
Mature calyx without spines, but winged.	13
13. Leaves flat, wavy-edged, ovate.	<i>Cycloloma</i>
Leaves linear, usually turning red when mature.	<i>Kochia</i>
14. Leaves circular in cross section, fleshy.	<i>Suaeda</i>
Leaves not circular in cross section.	<i>Chenopodium</i>

Atriplex atriplex

Annual or perennial plants with unisexual flowers. Male flowers usually with 3–5 sepals and the same number of stamens and no bracts; female flowers usually without sepals but with a pair of bracts.

1. Plant a small shrub, with pale leaves covered with small silvery scales; leaves without stalks. *A. nuttallii*

- Plant a herb, not shrubby but sometimes slightly woody at base. 2
2. Foliage green or greenish, sometimes deep red on both surfaces, sparsely mealy and, therefore, sometimes grayish when young. 3
- Foliage gray or whitish with a fine scurfiness, at least on the lower surface. 4
3. Bracts rounded ovate, over 6 mm broad; two kinds of female flowers, some with bracts and no sepals, others with sepals and no bracts. *A. hortensis*
- Bracts not orbicular, not over 6 mm broad; all female flowers with bracts and no sepals. *A. patula*
4. Plants perennial, woody at the base. *A. nuttallii*
- Plants annual. 5
5. Leaves more or less toothed or hastate; fruiting bracts toothed. *A. argentea*
- Leaves entire, strongly veined; fruiting bracts entire above the middle. *A. powellii*

Atriplex argentea Nutt.

SILVERY ATRIPLEX

A bushy annual growing to 40–80 cm high, but usually much smaller; stem angled. Leaves usually deltoid or triangular, 6–25 mm long, grayish green and scurfy. Male flowers usually in terminal spikes, and female flowers in clusters in axils of leaves. Very common in saline flats and similar sites in southern Prairies, but apparently rare elsewhere.

Atriplex hortensis L. (Fig. 98)

GARDEN ATRIPLEX

A fairly tall, erect annual plant to 60–120 cm high. Leaves of various shapes, cordate to triangular and ovate. Peculiarly interesting, because it bears two kinds of female flowers on the same inflorescence. One kind has no calyx but two large bracts, becoming quite large at maturity, over 6 mm; these flowers produce large, flat, pale brown seeds, about 3–10 mm in diam, which germinate a few days after being sown. The other kind of female flower has a small 3- to 5-lobed calyx but no bracts, and produces small, black, shiny seeds about 2 mm in diam, which sometimes remain dormant in the soil for many months before germination. Inflorescence often large, sometimes up to 30 cm long. A garden escape, it has become very common around towns and cities; in many parts of the Prairie Provinces. A red-leaved variety, *atrosanguinea* Hort., is often found escaped from gardens.

Atriplex nuttallii S. Wats.

NUTTALL'S ATRIPLEX

A perennial shrub or subshrub with a very deep rooting system. May be nearly prostrate, or with branches to 75 cm high. Leaves and stem pale green, with a fine scurfiness; leaves linear-oblong to spatulate or obovate, 2–5 cm long. Almost certainly dioecious (male flowers on one plant and female flowers on another). Readily eaten by livestock and considered a useful pasture plant



Fig. 98. Garden atriplex, *Atriplex hortensis* L.

because of its high mineral content and palatability. Found on badlands, eroded soils, and alluvial flats; throughout southern Prairies.

Atriplex patula L.

ORACHE

An annual plant 30–90 cm high; stem coarse, erect, often marked with vertical grooves or channels. There are two varieties: the common orache, *A. patula* L.; and the halberd-leaved orache, *A. patula* L. var. *hastata* (L.) Gray. In the common orache the leaves taper to their stalks, but in the halberd-leaved orache the base of the leaf is flatter and abruptly narrowed to the stalk. Both varieties common; in saline meadows and waste places; particularly in southwestern Prairies. Often mistaken for lamb’s-quarters.

Atriplex powellii S. Wats.

Annual plants 20–50 cm high, freely branched; leaves 1–2 cm long, silvery white. Flowers clustered in the axils; fruiting bracts about 5 mm broad, toothed below. Rare; alkali flats; southwestern Prairies.

Axyris axyris

Axyris amaranthoides L. (Fig. 99)

RUSSIAN PIGWEED

An erect, bushy annual weed 20–60 cm high. Stem much-branched, with ascending branches. Leaves lanceolate, pale green, to 8 cm long. Flowers borne in dense, leafy clusters; male flowers toward the ends of the stems. The whole plant turns a pale straw color at maturity. Seeds are of two types on the same plant: one type bearing a 2-lobed, long oval-shaped membranous wing at the top; and the other type almost round with no wings. The long winged seeds germinate readily, but the round wingless ones have a long period of dormancy. This introduced annual is very common in shelterbelts, on cultivated fields and abandoned lands, and in waste places; throughout the Prairie Provinces. It can be found often among bushes far from any cultivated land.

Bassia bassia

Bassia hyssopifolia (Pall.) Kuntze

FIVE-HOOK BASSIA

A slender-stemmed, weedy, branching annual 15–60 cm high. Leaves linear, 10–25 mm long, grayish green, usually closely appressed to the stem. Inflorescence in form of narrow interrupted spikes of tiny, inconspicuous flowers. Fruit not much larger than the head of a pin but enclosed with 5 tiny bracts, each bearing a small hooked spine. Very plentiful in some saline areas in southwestern Prairies, appears to be spreading eastward along railway grades.

Chenopodium goosefoot

Annual plants, usually weedy, with alternate, often farinose or mealy leaves. Flowers perfect, with no bracts, calyx of 2–5 sepals enclosing the fruit.

- 1. Calyx fleshy, red, and globular, resembling small strawberry. *C. capitatum*
- Calyx not fleshy, red, or strawberry-like. 2




Fig. 99. Russian pigweed, *Axyris amaranthoides* L.

- 2. Stamens 1 or 2; calyx only slightly fleshy and reddish. *C. rubrum*
 Stamens 5; calyx not fleshy. 3
- 3. Leaves with large, divaricate lobes, not mealy, shiny green; seeds about 2 mm in diam. *C. hybridum* var. *gigantospermum*
 Leaves entire or wavy-edged; seeds not over 1 mm in diam. 4
- 4. Calyx lobes not keeled; panicles of flowers in axils of leaves and shorter than the leaves. *C. glaucum* var. *pulchrum*
 Calyx lobes keeled; upper panicles of flowers longer than the leaves. 5
- 5. Seeds separating readily from their coating. 6
 Coating rather firmly attached to the seeds. *C. album*
- 6. Leaves ovate to triangular or hastate, many-nerved. 7
 Leaves linear, entire, 1- to 3-nerved. 8
- 7. Leaves triangular or hastate, to ovate, almost as broad at the base as long. *C. fremontii*
 Leaves ovate to oblong-lanceolate, rounded to cuneate at the base. *C. polyspermum*
- 8. Leaves grayish mealy, especially below. *C. leptophyllum*
 Leaves pale green, almost glabrous, not mealy. *C. subglabrum*

Chenopodium album L. (Fig. 100)

LAMB'S-QUARTERS

A rank annual weed 20–80 cm high. Leaves variable, usually wavy-margined and roughly ovate with a mealy coating, up to 7 cm long. Stems usually longitudinally grooved, often with reddish lines and blotches. Inflorescence usually of a bluish tinge, in dense panicles in leaf axils and at summit of stem. Although this species has been introduced from Europe, it is probably also a native. Young plants of this species are often eaten in place of spinach. One of the commonest weeds; found in waste places, roadsides, and gardens; throughout the Prairie Provinces. Syn.: *C. lanceolatum* Muhl.; *C. dacoticum* Standl.

Chenopodium capitatum (L.) Aschers.

STRAWBERRY BLITE

An annual herb with stem either erect or spreading, to 40 cm high. Leaves pale green, roughly triangular, 3–7 cm long. Inflorescence in form of small round clusters at intervals on the stem, turning red, resembling small strawberries. Found on rocky or stony soil and around bluffs and woodlands; throughout the Prairie Provinces, particularly in Boreal forest.




Fig. 100. Lamb's-quarters, *Chenopodium album* L.

Chenopodium fremontii S. Wats. (Fig. 101)

FREMONT'S GOOSEFOOT

An annual with rather slender, erect stems, usually with longitudinal dark green lines. Grows to about 60 cm high, and has broadly triangular leaves up to 5 cm long, pale green on both sides. Inflorescence open, interrupted, and not dense, usually appearing rather straggly. Common among bushes and bluffs; throughout the Prairies and Parklands.

Chenopodium glaucum L. var. *pulchrum* Aellen

SALINE GOOSEFOOT

A low prostrate plant 10–40 cm long, with rather fleshy, often reddish stem. Leaves usually small, varying from somewhat triangular to oval or oblong, sinuately dentate or lobed, resembling very small oak leaves, 10–25 mm long, and mealy on underside. Inflorescence in small spikes in axils of leaves. Common in moist, saline locations; throughout Prairies and Parklands. Syn.: *C. salinum* Standl.

Chenopodium hybridum L. var. *gigantospermum* (Aellen) Rouleau

MAPLE-LEAVED GOOSEFOOT

A tall annual 50–120 cm high. Leaves bright green, rather shiny, to 15 cm long, usually having 2–4 large sharp-pointed lobes on either margin and a long pointed apex. Inflorescence somewhat similar to that of *C. fremontii*, with which it is often associated, loose and interrupted, in long open panicles. Fairly common in shady wooded places; throughout the Prairie Provinces.

Chenopodium leptophyllum Nutt.

NARROW-LEAVED GOOSEFOOT

A tall erect annual with a somewhat mealy stem, striate or longitudinally grooved with alternate yellow and green lines. Leaves linear or linear-lanceolate, pale green and mealy above and densely mealy below, usually entire, and to 2–4 cm long and 6 mm wide. Inflorescence in small spikes in upper leaf axils and at summit of stem, densely mealy. Very plentiful on dry, sandy soil; in southwest Prairie Provinces, especially in the sandhills.

Chenopodium polyspermum L.

MANY-SEEDED GOOSEFOOT

An erect or procumbent annual to 1 m high, with glabrous stems and leaves. Leaves ovate-elliptic, entire or a slight tooth above the base. Inflorescence long, lax with many axillary cymes. Rare; an introduced weed; Parklands.

Chenopodium rubrum L.

RED GOOSEFOOT

A tall erect plant to 75 cm high, with ascending branches. Leaves not mealy, thick, and dark green, pointed at both ends, coarsely toothed, 3–10 cm long. Flower clusters in leafy spikes in leaf axils, fruit turning dull red when ripe. Very common in saline, moist soil; throughout the Prairie Provinces.

Chenopodium subglabrum (Wats.) Nelson

An erect annual 20–80 cm high, with lanceolate or oblong leaves, glabrous and green above, 3–6 mm long. Inflorescence diffuse; the stem leaves mealy. Rare or overlooked; on sandy soils; Prairies.




Fig. 101. Fremont's goosefoot, *Chenopodium fremontii* S. Wats.

Corispermum bugseed

Annual plants with very branched stems and narrow, linear leaves; flowers solitary in the axils of the leaf-like bracts and the seeds flattened, giving rise to the common name.

- Plants not hairy; fruit with a distinct wing around it. *C. hyssopifolium*
- Plants slightly hairy; fruit scarcely winged, merely acute-margined. *C. orientale* var. *emarginatum*

Corispermum hyssopifolium L. BUGSEED

A much-branched annual 20–50 cm high, with narrow, linear, pale green leaves about 3–5 cm long. Inflorescence in small clusters at axils of leaves, each separate flower having a small leaf-like bract. Seeds medium brown, about 3 mm across, flat on one side and slightly convex on the other, with a decided wing around them. Not very common; but has been found on sandy soil in several locations; in southwest and south central Prairies. Syn.: *C. marginale* Rydb.

Corispermum orientale Lam. var. *emarginatum* (Rydb.) Macbr. (Fig. 102) VILLOSE BUGSEED

An annual, very similar to *C. hyssopifolium*, but with stems and leaves having fine, white, stellate or star-shaped hairs. Seeds similar to the preceding but without a wing. Plant eaten by livestock; also a good sand binder. Fairly plentiful in sandy soil; especially the sandhills of southwest. Syn.: *C. villosum* Rybd.

Cycloloma winged pigweed

Cycloloma atriplicifolium (Spreng.) Coulter WINGED PIGWEED

A much-branched annual 15–50 cm high, sometimes with cobwebby hairiness on stem and leaves. Leaves up to 6 cm long, lanceolate, with a toothed margin. Inflorescence borne on open panicles; each fruit having a broad membranous wing below it. When mature, the plant becomes purplish and breaks off, blowing with the wind and scattering its seed. Rare; but has been found on sandy soils; in Prairies.

Endolepis endolepis

Endolepis suckleyi Torr. ENDOLEPIS

A low erect annual to 30 cm high, with short leaves 15–25 mm long, lanceolate and with 1 nerve. Stem and midrib often purplish. Flowers in clusters on short spikes in axils of leaves and at ends of stem. Not common, but plentiful locally; found on saline flats, eroded clay slopes, and badlands; throughout southern Prairies. Syn.: *Atriplex dioica* (Nutt.) Macbr.



Fig. 102. Villose bugseed, *Corispermum orientale* Lam. var. *emarginatum* (Rydb.) Macbr.



Fig. 103. Winterfat, *Eurotia lanata* (Pursh) Moq.

Eurotia winterfat

Eurotia lanata (Pursh) Moq. (Fig. 103)

WINTERFAT

A perennial subshrub or herb 15–50 cm high. Plant covered with fine, star-like white hairs, which become reddish as the plant ages. Leaves 1–5 cm long, linear, with rolled margins. Flowers unisexual, both male and female flowers being on the same plant, the male flower clusters above the female clusters. Female flowers and fruit enclosed in 2 bracts, which are united almost to the top and have 2 horns on the top. Whole female inflorescence clothed with long, silky, white hairs, making the plants very conspicuous. Much relished by livestock as an excellent forage plant. Very useful on winter range because its protein and mineral content remain high throughout the fall and winter. Common on dry prairies and on heavy soils; throughout Prairies.

Kochia kochia

Kochia scoparia (L.) Schrad.

SUMMER-CYPRESS

An annual plant, escaped from gardens, about 60 cm high, erect, and of regular pyramidal or ovoid shape. Leaves linear and closely compacted, pale green when young, but purplish red when mature. The plant produces many seeds, which germinate readily and form a mat of seedlings the following year. Used as an ornamental in gardens because of its symmetrical shape and its red coloring in the fall, but its prolific seeding habits have caused it to become a weed in the vicinity of most towns and cities in Prairies and Parklands.

Monolepis monolepis

Monolepis nuttalliana (R. & S.) Greene

SPEAR-LEAVED GOOSEFOOT

A prostrate annual with fleshy reddish stems and hastate or spear-shaped leaves, which form a rosette in the plant's early stage. Spreading on the soil, 20–50 cm across. Inflorescence in small clusters in axils of leaves, produced early in season. One of the first weeds to start in the spring on fallow and in waste places, where its fleshy taproots and succulent stems cause a great drain on moisture reserves. Common around gopher holes; a native that has become weedy and, though originally found on saline soils, is now found on cultivated land; throughout Prairies and Parklands.

Salicornia samphire

Salicornia rubra A. Nels.

RED SAMPHIRE

A low annual with no true leaves, their places being taken by scales at the nodes of the stems. Stems circular in cross section (terete), the branches opposite. Flowers very minute, sunk into the tissue of the stems. Plant turning a bright crimson at maturity and giving a reddish color to dry sloughs in late summer and early autumn. Very common in strongly saline sloughs; throughout Prairies and Parklands. Syn.: *S. europaea* L. ssp. *rubra* (Nels.) Breitung; *S. europaea* L. var. *prona* (Lunell) Boiv.

Salsola glasswort

Salsola kali L. var. *tenuifolia* Tausch.

RUSSIAN-THISTLE

An introduced annual weed 10–60 cm high. Early leaves dark green, thread-like, about 2 cm long; later leaves shorter and broader, coming to a sharp hard point. Flowers borne in axils of upper leaves, with a membranous wing on calyx around seed. As plants age they become dry and the spiny tips of the leaves and bracts harden, making the whole plant prickly. Plants usually become reddish and at maturity break off at the ground and drift across the country with the winds, scattering their seeds. Palatable to livestock, but highly laxative. Very common; in fields, waste places; throughout Prairies; less common in Parklands. Syn.: *S. pestifer* A. Nels.

Sarcobatus greasewood

Sarcobatus vermiculatus (Hook.) Torr. (Fig. 104)

GREASEWOOD

A much-branched, shrubby perennial, with spiny branches, in some localities 1.5–2 m high. Stems usually almost white. Leaves pale yellowish green, linear, about 2–4 cm long. Male flowers borne in small cylindrical spikes at ends of stems, female flowers borne singly in axils of leaves. A broad membranous wing forms on calyx around fruit. If eaten in large amounts, greasewood is rather **poisonous** to livestock, especially to lambs during the spring and summer. It contains sodium and potassium oxalates and during dry seasons these salts occur in a concentrated form. Around strongly saline sloughs and flats; throughout Prairies.

Suaeda sea-blite

Suaeda depressa (Pursh) S. Wats.

WESTERN SEA-BLITE

A low-growing annual or perennial, with very dark green, rather fleshy, narrow leaves. There are two forms: the species *depressa* and its variety *erecta* or erect sea-blite. In the *depressa* form the plant is low and spreading and the leaves are 10–30 mm long. The var. *erecta* is upright, with longer leaves 15–40 mm long. Flowers dark greenish, clustered in axils of upper leaves. At maturity plants turn very dark, almost black. Common in saline areas around sloughs and saline flats; throughout Prairies and Parklands. The var. *erecta* S. Wats. is probably a little more plentiful than the species. Syn.: *S. maritima* L. var. *americana* (Pers.) Boiv.

Suckleya suckleya

Suckleya suckleyana (Torr.) Rydb. (Fig. 105)

POISON SUCKLEYA

A somewhat prostrate, much-branched annual, with succulent stems up to 30 cm long. Leaves stalked, alternate, round to diamond-shaped, up to 3 cm long, irregularly blunt-toothed along the upper margins. Flowers unisexual, the male in the axils of the upper leaves, the female in the lower leaf axils; female flowers enclosed in a couple of stiff winged bracts. Fruit 5–6 mm long with an abruptly pointed end. This plant is **poisonous** to livestock because it contains hydrocyanic acid. Not plentiful; however, it has been found in moist and saline areas; in the south central Prairies.



Fig. 104. Greasewood, *Sarcobatus vermiculatus* (Hook.) Torr.




Fig. 105. Poison suckleya, *Suckleya suckleyana* (Torr.) Rydb.

AMARANTHACEAE—amaranth family

Amaranthus amaranth

Coarse, annual, weedy plants, somewhat similar to the goosefoot family, but having flowers enclosed in three dry and persistent bracts.

- 1. Plants tall; upper flowers in dense terminal spikes. *A. retroflexus*
 Plants low, much-branched; flowers in small clusters in axils of leaves. 2
- 2. Plants erect and bushy; leaves pale green and not shiny; sepals 3, much shorter than the somewhat spine-tipped bracts. *A. albus*
 Plants prostrate, with succulent reddish stems; leaves dark green and rather shiny; sepals 4 or 5, almost as long as the blunt-tipped bracts. *A. graecizans*

Amaranthus albus L.

TUMBLEWEED

A bushy, much-branched, whitish-stemmed annual 25–60 cm high, with pale green, dull, spatulate leaves 10–35 mm long. The end of the midrib of the leaf is usually projected as a tiny spine at the end of the leaf, a condition often found in other species of amaranth, but more pronounced in this species. Flowers in small clusters in leaf axils; seed black and shiny, enclosed in a small utricle, or envelope, the top of which falls off at maturity. A native plant and an early invader of newly broken ground, fireguards, and rough ground. When dry, this plant breaks off and blows across the prairie, dispersing its seeds. Now a common weed; on waste ground, in gardens, and along roadsides; throughout the Prairies and Parklands. This species has usually been called *A. graecizans*, but that name was intended by Linnaeus for the prostrate amaranth.

Amaranthus graecizans L. (Fig. 106)

PROSTRATE AMARANTH

A coarse, prostrate, weedy annual, with reddish, fleshy stems forming mats 15–60 cm in diam. Leaves usually spatulate, the broadest part beyond the middle, 5–25 mm long, dark, shiny green. Flowers borne in leaf axils. Not considered a native of the area, but has come in from the southwest, where it is a native of dry sites in the intermountain areas. A very common garden weed; throughout Prairies and Parklands. This species has been known as *A. blitoides* S. Wats.

Amaranthus retroflexus L.

RED-ROOT PIGWEED

A coarse, rough, erect annual with reddish-colored roots. Often growing to 1 m high and having rough, angular stems somewhat hairy near the top. Leaves usually ovate, on fairly long stalks, rough to the touch, and 6–10 cm long. Inflorescence harsh and rough, borne in dense spikes in leaf axils and in a large terminal spike at summit of stem. Sepals dry and parchment-like with spiny tips, the 3 bracts around each flower also spine-tipped, giving the inflorescence a decidedly rough appearance. Seeds shiny, black, and in a small




Fig. 106. Prostrate amaranth, *Amaranthus graecizans* L.

utricle, similar to other amaranths. A native of the subtropical states of USA, but a common weed over all North America. Very common; in waste places, roadsides, gardens, and fields; throughout Prairies and Parklands.

NYCTAGINACEAE—four-o'clock family

Plants annual; bracts free. *Abronia*
 Plants perennial; bracts united into a tube. *Mirabilis*

Abronia sand verbena

Abronia micrantha Torr.

SAND PUFFS

Annual plants with decumbent stems, 10–30 cm high, branching at base, often somewhat pubescent. Leaves opposite, petioled, 2–4 cm long, prominently veined. Inflorescence a head supported by 4–6 bracts; perianth a corolla-like salverform tube with 5 petal-like lobes. Fruit with wings 10–15 mm, rarely 20 mm, broad. Rare; sand dunes in Prairies.

Mirabilis umbrellawort

Perennial herbs with opposite, entire leaves without stipules. Stems usually swollen at nodes. Flowers regular and perfect, with no petals, but sepals brightly colored and petal-like, and generally united into a bell-like or funnel-form tube. Flowers usually in clusters of 3–5 with a saucer-like involucre bract beneath them.

1. Leaves cordate, heart-shaped, with distinct stalks. *M. nyctaginea*
 Leaves not cordate, but ovate-lanceolate or linear, mostly without stalks or the lower ones with short stalks. 2
2. Stem more or less hairy, sticky. *M. hirsuta*
 Stem not at all hairy below, but fine, sticky-haired above. *M. linearis*

Mirabilis hirsuta (Pursh) MacM.

HAIRY UMBRELLAWORT

A slender erect plant 30–60 cm high, with glandular hairs on the stem. Leaves usually lanceolate or linear-lanceolate and hairy, 2–7 cm long. Another form equally common has the lower part of the stem and leaves almost devoid of hairs, except just under the nodes of the stem, and slightly narrower leaves. This form has sometimes been designated *Oxybaphus pilosus*, a separate species. Most authorities, however, do not separate the forms. Common; on sandy soils; throughout Prairies and Parklands. Syn.: *Oxybaphus hirsutus* (Pursh) Sweet.

Mirabilis linearis (Pursh) Heimerl

LINEAR-LEAVED UMBRELLAWORT

Erect, with very narrow, single-nerved leaves, 15–65 mm long. It was reported from two locations in the southwestern area by the early botanist Macoun. However, its range is south of the Prairie Provinces and its appear-

ance north of the International Boundary is unusual. Syn.: *Oxybaphus linearis* (Pursh) Robinson.

Mirabilis nyctaginea (Michx.) MacM.

HEART-LEAVED UMBRELLAWORT

A tall erect species 40–90 cm high, with an almost hairless stem and large cordate or heart-shaped leaves, 5–10 cm long and 2–7 cm wide. Inflorescence usually deep reddish, with a greenish involucre below the flowers. Fairly common; in rich soils in southeastern Parklands and Boreal forest, gradually spreading westward along the railway tracks. Several large clumps have been located on the main line of the Canadian Pacific Railway in southern Saskatchewan, and, where found, it appears to be the dominant plant on the cinder fill of the railway grade. Syn.: *Oxybaphus nyctagineus* (Michx.) Sweet.

PORTULACACEAE—purslane family

Rather succulent plants with perfect flowers, 2 sepals, and 5 petals. The fruit is a capsule, which opens with 2 or 3 valves at the top or with the top falling off like a lid (circumscissile).

1. Capsule opening by valves. *Claytonia*
Capsule circumscissile at maturity. 2
2. Ovary partly inferior; prostrate annual
weed. *Portulaca*
Ovary superior; perennial with taproot
and bracteate scape. *Lewisia*

Claytonia springbeauty

1. Leaves alternate. 2
Leaves opposite. 3
2. Plants annual; leaves linear. *C. linearis*
Plants perennial; leaves ovate to lanceo-
late. *C. parvifolia*
3. Plants annual; stem leaves numerous. *C. fontana*
Plants perennial. 4
4. Plants with a globose corm; stem leaves 2. *C. caroliniana* var. *lanceolata*
Plants with a thick taproot; leaves basal. *C. megarrhiza*

Claytonia caroliniana Michx. var. *lanceolata* (Pursh) Wats.

LANCE-LEAVED SPRINGBEAUTY

A low-growing early spring flowering plant, growing from a corm or tuber-like globose root 6–10 cm below the soil surface. In some plants several stems arise from a single corm. Occasionally a plant bears one or two basal, stalked leaves, but they are usually absent. Two stalkless (sessile) opposite leaves are borne on the stem, lanceolate in shape, 15–50 mm long, with three distinct veins. Flowers have petals about 1 cm long, white with pink lines or

faintly pinkish. A western species, common along the margin between grassland and wooded areas in the Foothills, but eastward has been collected only in the Cypress Hills, where it is found flowering in large masses soon after snow melts in spring. Found along edges of woodlands and around margins of clearings.

Claytonia fontana (L.) Davis

INDIAN LETTUCE

Low annual with weak branching stems, often rooting at the nodes. Leaves opposite, 5–15 mm long. Inflorescence leafy, the flowers nodding. Rare; wet places; Hudson Bay.

Claytonia linearis Dougl.

LINEAR-LEAVED SPRINGBEAUTY

A low-branching annual, growing from fibrous roots to 6–10 cm high. Leaves linear or almost thread-like, 10–30 mm long. Flowers nodding, 2–7 in a raceme, with pale pinkish petals about 3 mm long. A western species, found in very early spring in higher meadows in the Cypress Hills and southern Rocky Mountains. Syn.: *Montia linearis* (Dougl.) Greene.

Claytonia megarrhiza (Gray) Parry

ALPINE SPRINGBEAUTY

Plants usually not more than 10 cm high; basal leaves numerous, fleshy, spatulate or obovate, 1–8 cm long; flowering stalks numerous, flowers 10–15 mm across, white or pink. Rocky places, scree fields; southern Rocky Mountains.

Claytonia parvifolia Moc.

SMALL-LEAVED SPRINGBEAUTY

Perennial plants with branched rootstocks, often developing stolons. Basal leaves 10–35 mm long, obovate, long-petioled, more or less fleshy; stem leaves reduced, alternate. Inflorescence racemose, with 1–10 flowers, 10–15 mm across, white or pink. Moist areas, mountain meadows; southern Rocky Mountains.

Lewisia bitter-root

Lewisia pygmaea (Gray) Rob.

DWARF BITTER-ROOT

Perennial with a thick fleshy taproot bearing several stems. Basal leaves linear, 3–6 cm long, 1.5–3 mm wide; stems 15–50 mm long, with a pair of opposite bracts. Flowers 1–3, about 15 mm across, white or pinkish. Rare; alpine slopes; southern Rocky Mountains.

Portulaca purslane

Portulaca oleracea L.

PURSLANE

A succulent, prostrate annual, sometimes making a mat up to 40 cm across. Stems reddish, very thick, fleshy, juicy, and hairless. Leaves alternate, dark shiny green, spatulate or obovate, 5–25 mm long, thick, and fleshy. Flowers borne singly, without stalks, in the axils of the leaves, and usually open only in bright sunshine, bright yellow, about 6 mm in diam, with 2 sepals and 5 petals. Seeds numerous and minute, contained in a pointed capsule, or pyxis, the top of which breaks off, releasing the seeds. An introduced weed, common in gardens; throughout the Prairie Provinces.

CARYOPHYLLACEAE—pink family

Herbs with stems swollen at the joints and entire, opposite leaves. Flowers usually with 4 or 5 petals, sometimes none, and 4 or 5 sepals. The fruit is a capsule opening by valves at the top. A difficult family to identify, often requiring the fruiting capsule for positive identification.

1. Plants with woody base; leaves with spiny tips; fruit with a single seed. *Paronychia*
 Plants not with woody base; fruit many-seeded. 2
2. Sepals united part way, forming a tube. 3
 Sepals entirely separate. 8
3. Calyx with 1–3 pairs of bracts below. *Dianthus*
 Calyx without bracts. 4
4. Styles 2 only. 5
 Styles more than 2. 6
5. Calyx becoming inflated and wing-angled; flowers few, pink. *Saponaria*
 Calyx not inflated or angled; flowers very numerous. *Gypsophila*
6. Sepals with long, leaf-like lobes, over 20 mm long; styles 5, opposite the petals; flowers dark purple. *Agrostemma*
 Lobes of sepals not over 20 mm long; styles alternate with the petals. 7
7. Styles usually 3; capsule usually divided into partitions at base. *Silene*
 Styles 5; capsule 1-celled to base. *Lychnis*
8. Plants with small, ovate, papery stipules at junction of stem and leaf, and stem and branch. 9
 Plants without stipules. 10
9. Styles 5; leaves in whorls. *Spergula*
 Styles 3; leaves opposite. *Spergularia*
10. Capsule opening with twice as many valves or teeth as there are styles; petals deeply 2-cleft. 11
 Capsule opening with as many entire (or later 2-cleft) valves as there are styles; petals entire or merely notched at apex. 12
11. Capsule short, ovate or oblong, usually opening with 6 valves; styles usually 3. *Stellaria*
 Capsule long, cylindrical, often curved, usually opening with 10 valves; styles usually 5. *Cerastium*

12. Flowers 5-merous. *Sagina*
 Flowers with 5 sepals, 3 styles, and 3
 valves. *Arenaria*

Agrostemma corn cockle

Agrostemma githago L. (Fig. 107)

PURPLE COCKLE

An erect hairy annual plant with a taproot, 30–75 cm high, with hairy linear leaves up to 10 cm long. Flowers borne singly at head of stems with hairy sepals united at base, but with long lobes often up to 2.5 cm, much longer than the petals. Petals purple and flowers 25–40 mm in diam. Seed about 3 mm in diam, somewhat flattened, black, and roughened with rows of minute protuberances. Seeds **poisonous** to chickens. A weed introduced from Europe, but not common; in grainfields throughout the Prairie Provinces.

Arenaria sandwort

Low tufted herbs with opposite leaves. Flowers white, with 4 or 5 sepals and 4 or 5 entire or slightly notched petals.

1. Leaves oval to oblong. 2
 Leaves linear or filiform. 6
2. Bracts of the inflorescence small, scarious. 3
 Bracts of the inflorescence large, leaf-like. 4
3. Leaves pubescent below along the
 midrib; sepals rounded. *A. lateriflora*
 Leaves glabrous below; sepals acuminate. *A. macrophylla*
4. Plants fleshy, glabrous. *A. peploides* var. *diffusa*
 Plants not fleshy, at least the pedicels
 puberulent. 5
5. Plants perennial; stems 3–10 cm high,
 puberulent; leaves glabrous. *A. humifusa*
 Plants annual; stems 5–20 cm high,
 puberulent; leaves puberulent. *A. serpyllifolia*
6. Sepals rounded or obtuse at the tip. 7
 Sepals distinctly acute to acuminate. 9
7. Leaves 3–8 mm long, about 1 mm wide. *A. laricifolia* var. *occulta*
 Leaves 1–6 cm long, linear to filiform. 8
8. Plants glabrous. *A. congesta* var. *lithophila*
 Plants glandular puberulent, at least in
 the inflorescence. *A. capillaris* var. *americana*
9. Leaves clearly spine-tipped, recurved
 spreading. *A. nuttallii*
 Leaves not spine-tipped. 10
10. Flowers solitary, terminal. 11
 Flowers in cymose inflorescence, 2 or
 more together. 12



Fig. 107. Purple cockle, *Agrostemma githago* L.

11. Inflorescence cymose, open. *A. stricta* ssp. *dawsonensis*
 Inflorescence with flowers solitary on
 leafy stems. *A. rossii* var. *columbiana*
12. Plants usually glabrous; stem leaves often
 with axillary branches. *A. stricta* ssp. *dawsonensis*
 Plants usually glandular puberulent, at
 least in the inflorescence; axillary
 branches few or none. *A. verna*

Arenaria capillaris Poir. var. *americana* (Maguire) Davis

Plants tufted, with a slender caudex; stems 5–15 cm high. Leaves 2–7 cm long, filiform, mostly erect. Inflorescence a few-flowered cyme, with the pedicels 5–20 mm long, sepals 3–4 mm long, petals 6–8 mm long. Alpine meadows; southern Rocky Mountains.

Arenaria congesta Nutt. var. *lithophila* (Rydb.) Maguire

ROCKY-GROUND SANDWORT

An erect perennial to 30 cm high, with tufted basal leaves and a few stem leaves. Leaves linear and thread-like, 1–5 cm long. Flowers in open clusters on tops of stems, with white petals 4–6 mm long, and straw-colored sepals. Very local, but found on rocky benchland and slopes; Cypress Hills, Wood Mountain. Syn.: *A. lithophila* Rydb.

Arenaria humifusa Wahl.

Plants tufted, mat-forming; stems 3–10 cm high, slender, few-flowered; leaves 2–10 mm long; sepals 3–5 mm long; petals 5–7 mm long. Sand and gravel bars; Boreal forest.

Arenaria laricifolia (L.) Rob. var. *occulta* (Ser.) Boiv.

Plants with more or less woody, branched caudex, mat-forming; leaves 3–7 mm long, 0.5–1 mm wide; flowering stems ascending, puberulent, often glandular; 1- to 6-flowered; sepals 3–5 mm long, glandular puberulent; petals 4–7 mm long. Stony slopes, scree fields; Rocky Mountains. Syn.: *Minuartia laricifolia* (L.) S. & T.; *A. obtusiloba* (Rydb.) Fern.

Arenaria lateriflora L.

BLUNT-LEAVED SANDWORT

An erect perennial with thin, weak, slightly hairy stems and oval to oblong, thin, pale green leaves. Flowers about 6–12 mm across, usually borne in pairs, and petals about twice the length of sepals; sometimes 4 sepals and petals, and sometimes 5 of each. Common; in moist woodlands; throughout Prairies and Parklands. Syn.: *Moehringia lateriflora* (L.) Fenzl.

Arenaria macrophylla Hook.

LARGE-LEAVED SANDWORT

Resembling the preceding species in habit and pubescence. Leaves mainly 2–5 cm long, 3–8 mm wide; sepals 3–6 mm long; petals 5–10 mm long. Rare; rocky areas; Boreal forest, Saskatchewan.

Arenaria nuttallii Pax

Plants with a deep taproot, many-stemmed, loosely matted stems 6–12 cm high, glandular pubescent; leaves 6–10 mm long; inflorescence a few- to many-

flowered cyme, with the pedicels 5–15 mm long, sepals 4–5 mm long, petals shorter than sepals. Slopes; southern Rocky Mountains.

Arenaria peploides L. var. *diffusa* Hornem.

SEA-PURSLANE

Plants glabrous; stems trailing, succulent, rooting at the nodes. Leaves 5–20 mm long, 5–10 mm wide; flowers solitary in leaf axils, and in few-flowered terminal cymes; capsule 6–10 mm wide, globose. Beaches; along Hudson Bay.

Arenaria rossii R. Br. var. *columbiana* Raup

Plants densely tufted, with the flowering stems spreading. Leaves 2–5 mm long, awl-shaped, with axils of upper leaves having short branches; flowers solitary, with the pedicels 5–15 mm long, sepals 1.5–2.5 mm long, petals slightly shorter than sepals. Rare; alpine slopes; southern Rocky Mountains.

Arenaria serpyllifolia L.

THYME-LEAVED SANDWORT

A much-branched annual with a slightly downy stem 8–20 cm high. Leaves small and ovate, 3–8 mm long; flowers very small, usually 6–10 mm across, and borne on summit of stem. Introduced from Europe and very rare; a weed in cultivated fields.

Arenaria stricta Michx. ssp. *dawsonensis* (Britton) Maguire

DAWSON SANDWORT

An annual, branched from the base, 10–30 cm high, with thread-like leaves usually 6–12 mm long; petals no longer than the sepals, which are 3-nerved and 2.5–4 mm long. Lakeshores, river flats; Boreal forest, Rocky Mountains.

Arenaria verna L.

EARLY SANDWORT

Plants loosely tufted; stems 2–15 cm high, usually glandular pubescent above. Leaves 5–20 mm long, linear-lanceolate to oval-shaped; cymes 1- to 6-flowered; pedicels 2–25 mm long; sepals 3 mm long; petals 2–4.5 mm long. Slopes, meadows, and rock slides; Rocky Mountains, rare in Cypress Hills.

Cerastium chickweed

Low-growing annuals or perennials with opposite leaves; white flowers with cleft petals and long capsule, usually opening with 10 valves.

1. Annuals; capsule 2–3 times as long as sepals. *C. nutans*
Perennials; capsules once or twice as long as sepals. 2
2. Axillary shoots numerous, at the axils of most leaves. 3
Axillary shoots few or lacking. 4
3. Plants glandular pubescent throughout. *C. arvense*
Plants densely white tomentose, not glandular. *C. tomentosum*
4. Stems usually glandular pubescent, with the leaves stiffly hirsute, ciliate. *C. vulgatum*

Stems and leaves pilose or glandular
pilose. *C. alpinum*

Cerastium alpinum L. ALPINE CHICKWEED

Perennial plants 5–20 cm high; leaves about 10 mm long, 5 mm wide, soft long pubescent, sometimes also glandular. Flowers on peduncles 1–4 cm long, 20–25 mm across, with the petals about twice as long as the sepals. Alpine slopes and scree fields; southern Rocky Mountains, Boreal forest.

Cerastium arvense L. FIELD CHICKWEED

A low-growing perennial species 10–30 cm high, often with the stems prostrate at the base. Stems covered with hairs pointing downward; white flowers 15–20 mm across, blooming very early in the season. A very common spring flower; on prairie; throughout the Prairie Provinces.

Cerastium nutans Raf. LONG-STALKED CHICKWEED

A bright green erect annual to 25 cm high, with leaves 6–30 mm long. Fruiting capsules much longer than sepals, usually decidedly curved at maturity. Fairly plentiful; in moist woodlands; throughout Boreal forest, Rocky Mountains.

Cerastium tomentosum L. SNOW-IN-SUMMER

Perennial mat-forming plants, densely white lanate, up to 45 cm high, with leaves 10–30 mm long, 2–5 mm wide, the margins somewhat revolute. Inflorescence an elongated cyme; flowers about 30 mm across. Introduced as an ornamental; escaped into various locations.

Cerastium vulgatum L. MOUSE-EARED CHICKWEED

A biennial or perennial plant 10–40 cm high, with glutinous hairy stems and oblong-spatulate leaves 10–25 mm long. Fairly common; in woodlands and fields; Boreal forest.

Dianthus pink

1. Calyx pubescent. *D. deltoides*
Calyx glabrous. 2

2. Leaves 5–20 mm wide, obtuse; flowers in
heads. *D. barbatus*
Leaves 0.5–1 mm wide, acute; flowers soli-
tary. *D. sylvestris*

Dianthus barbatus L. SWEET WILLIAM

Perennial plants to 60 cm high. Leaves lanceolate, the lower ones obtuse; flowers white or pink, several crowded in a head. Introduced as an ornamental plant; occasionally escaped and persistent in moist places.

Dianthus deltoides L. STEPPEN PINK

Perennial plants 20–45 cm high. Leaves linear-lanceolate, short pubescent; flowers solitary, terminating stems or peduncles, reddish purple. Probably introduced; found only once, in Meadow Lake Provincial Park.

Perennial plants 10–30 cm high. Leaves filiform; flowers solitary on stems or peduncles, pink, with petals fringed. Introduced as ornamental; occasionally escaped.

Gypsophila baby's-breath

- 1. Annual plants 30–50 cm high. *G. elegans*
 Perennial plants 50–150 cm high. 2
- 2. Inflorescence usually glandular pubescent; calyx 3–3.5 mm long. *G. acutifolia*
 Inflorescence usually glabrous; calyx 1.5–2 mm long. *G. paniculata*

***Gypsophila acutifolia* Steven** STICKY BABY'S-BREATH

Perennial plants with long-creeping rootstocks, up to 150 cm high. Inflorescence a rather dense panicle, with the pedicels 1–4 mm long; flowers white. Introduced ornamental; occasionally escaped and established as a roadside weed.

***Gypsophila elegans* Bieb.** ANNUAL BABY'S-BREATH

An annual, much-branched plant, with narrow lanceolate leaves and many white flowers up to 12 mm across, with petals much longer than sepals. Escaped from gardens, sometimes found growing in waste places.

***Gypsophila paniculata* L.** BABY'S-BREATH

A perennial, much-branched plant, with linear-lanceolate leaves and many white flowers about 6 mm across, with petals about the same length as the sepals. Escaped from gardens, locally abundant in waste areas, roadsides, and gravel pits.

Lychnis champion

- 1. Petals large, much protruding beyond the calyx; calyx enlarging in fruit; flowers of different sexes on separate plants. *L. alba*
 Petals small, not or little protruding beyond the calyx; calyx not or little enlarging; flowers perfect. 2
- 2. Flowers nodding, usually solitary, rarely 2 or 3. *L. apetala*
 Flowers erect, usually 3 or more, on stiffly ascending pedicels. 3
- 3. Calyx not inflated, tightly enclosing the capsule; petals included or barely exerted. *L. drummondii*
 Calyx somewhat inflated; petals conspicuously exerted. *L. affinis*

A perennial, with stems 5–30 cm high, stiffly erect, more or less glandular viscid; flowers usually 3, on erect pedicels; calyx 8–10 mm long, glandular, somewhat inflated, with petals white, conspicuously exerted. Rare; rock crevices; Hudson Bay, Rocky Mountains.

Lychnis alba Mill.

WHITE COCKLE

A biennial or short-lived perennial 30–75 cm high, much-branched, and with many sticky-haired stems. Leaves opposite, ovate-oblong, 2–8 cm long; flowers white, about 2 cm in diam, with stamen-bearing flowers on some plants and pistillate flowers on others. Fruiting capsule becoming enlarged and swollen at maturity and bearing 10 teeth at the top. Resembles night-flowering catch-fly, which, however, has perfect flowers. Becoming increasingly common; in western parts of the Prairie Provinces.

Lychnis apetala L.

NODDING COCKLE

Small perennials 5–15 cm high; stems pubescent, often reddish viscid above. Flowers nodding; calyx 10–15 mm long; petals included or barely exerted, purple. Plants with the calyx conspicuously inflated, and petals not exerted are var. *arctica* (Fries) Cody; those with the calyx not inflated, and petals clearly exerted are ssp. *attenuata* (Farr) Mag. Alpine areas; Rocky Mountains, Hudson Bay.

Lychnis drummondii Wats.

DRUMMOND'S COCKLE

A tall, slender, erect perennial to 70 cm high with a sticky, hairy stem and opposite linear leaves up to 10 cm long. Flowers borne at head of stem on fairly long stalks, usually white or purplish and only slightly exceeding the sepals. Sepals about 12 mm long and joined to form a somewhat cylindric tube, usually pale yellow with green lines. Not plentiful but very widespread; on open prairie, especially on sandy soils; throughout Prairies and Parklands. Syn.: *Wahlbergella drummondii* (Wats.) Rydb.

Paronychia whitlowwort

Paronychia sessiliflora Nutt.

LOW WHITLOWWORT

A woody-based perennial, forming dense cushions, usually 20–75 mm high and 7–15 cm in diam. Leaves linear, spine pointed and very short, 5–6 mm long, bright green, and so closely overlapping that the stems or branches are usually hidden. Flowers yellow, 3–4 mm high, and dotted singly at intervals on the plant. Fairly common; on dry hillsides and eroded places; throughout Prairies. On casual inspection, this species may be mistaken for moss phlox, *Phlox hoodii*, although moss phlox is more open. Syn.: *P. depressa* Nutt.

Sagina pearlwort

- 1. Plants annual. *S. decumbens*
- Plants perennial. 2

- 2. Stem leaves at the upper 2 nodes distinctly shorter than lower ones, and with axillary bulb-like leaf bundles. *S. nodosa*
- Stem leaves all about the same size. 3
- 3. Plants tufted; sepals with purple margins. *S. caespitosa*
- Plants mat-forming; sepals with white margins. *S. saginoides*

Sagina caespitosa (J. Vahl) Lange CUSHION PEARLWORT

Perennial plants growing in small cushions. Flowers singly or in pairs, 10–12 mm across. Known only from Northern Manitoba (Baralson Lake).

Sagina decumbens (Ell.) T. & G. SPREADING PEARLWORT

Annual plants usually less than 10 cm high; stems ascending or decumbent. Leaves linear to awl-shaped; petals the same length as or shorter than the sepals, or lacking. Rare weed; introduced from the eastern USA.

Sagina nodosa (L.) Fenzl PEARLWORT

Perennial plants 5–15 cm high; stems erect, ascending, or decumbent. Lower stem leaves 5–20 mm, the upper ones scale-like, with bulb-like sterile shoots replacing flowers. Rocky or sandy shores and beaches; Boreal forest.

Sagina saginoides (L.) Karsten MOUNTAIN PEARLWORT

Perennial plants usually less than 10 cm high; rosette leaves, when present, to 20 mm long, the cauline leaves 5–10 mm. Flowers mostly solitary or occasionally in pairs; the petals about the same length as the sepals. Moist slopes and shores; Rocky Mountains.

Saponaria soapwort

Saponaria vaccaria L. (Fig. 108) COW COCKLE

A smooth, hairless annual plant 30–70 cm high. Leaves grayish green, smooth, entire, clasping, borne opposite each other in pairs on the stem, ovate-lanceolate, 2–8 cm long. Flowers in loose corymbose cymes at head of stem. Ovate calyx formed by the united sepals, about 12 mm long and 10 mm across, with the 5 pale pink petals forming a flat corolla about 12 mm across. Seeds borne in an ovoid capsule, round, dull black, about 2 mm in diam. Introduced from Europe, now common; in grainfields and on roadsides; throughout the southern part of the Prairie Provinces. Seeds are often in “Wild Flower Garden” packets available commercially. Syn.: *Vaccaria vulgaris* Host.

Silene catchfly

Annual, biennial, or perennial plants with opposite, entire leaves and perfect flowers. Seeds contained in a capsule.

- 1. Annual or biennial plants. 2
- Perennial plants. 4




Fig. 108. Cow cockle, *Saponaria vaccaria* L.

- 2. Plants not sticky. *S. cserei*
 Plants sticky, at least around the nodes of
 the stem. 3
- 3. Plants almost without hairs; flowers pink-
 ish white and about 5 mm across. *S. antirrhina*
 Plants sticky and hairy; flowers white and
 6–10 mm across. *S. noctiflora*
- 4. Plants densely matted or cushion-
 forming. *S. acaulis* var. *excapa*
 Plants with elongate, erect or decumbent
 stems. 5
- 5. Calyx more or less glandular pubescent. 6
 Calyx glabrous, or if pubescent, not glan-
 dular. 7
- 6. Plants rather straggly, weak-stemmed;
 calyx 5–8 mm. *S. menziesii*
 Plants with stems 20–50 cm high; calyx
 12–16 mm. *S. parryi*
- 7. Calyx to 8 mm long, with 10 simple veins. *S. sibirica*
 Calyx 10–15 mm long, with 20 main veins,
 strongly reticulate. *S. cucubalus*

Silene acaulis L. var. *excapa* (All.) DC.

MOSS CAMPION

Cushion plants or mat-forming 3–6 cm high; leaves 4–12 mm long, linear-lanceolate. Flowers solitary, often imperfect, purplish. Alpine meadows, scree fields; Rocky Mountains.

Silene antirrhina L.

SLEEPY CATCHFLY

An erect annual 40–50 cm high, with erect branches. Stem usually sticky near the nodes, sometimes slightly hairy. Leaves lanceolate, 2–5 cm long. Flowers pink, very small, about 5 mm across. Not common; may be found in sandy areas; Parklands, Boreal forest.

Silene cserei Baumgarten

SMOOTH CATCHFLY

A biennial plant 10–70 cm high somewhat resembling bladder campion but with a taproot. Leaves elliptic-lanceolate, stalkless, glaucous and thick, 5–10 cm long, and borne oppositely on the stem. Flowers numerous with pinkish slightly inflated calyx and white petals cleft at the tip and borne in whorls around the stem. At maturity, calyx is somewhat inflated and ovoid, about 12 mm long. Becoming increasingly abundant; along railway grades; throughout Prairies and Parklands.

Silene cucubalus Wibel

BLADDER CAMPION

A tall erect perennial with hairless stems 15–60 cm high. Leaves opposite, lanceolate and smooth, 2–8 cm long. The flowers, 10–20 mm across, borne at the summit of the stem in loose open panicles; sepals united to form a bladder-like calyx 10–12 mm long; petals white and 2-cleft. An introduced and very persistent weed. Not common, but generally plentiful where found. Syn.: *S. vulgaris* (Moench) Garcke.

Silene menziesii Hook.

MENZIES CATCHFLY

A straggly weak-stemmed perennial 15–40 cm high, with oval to lanceolate leaves 2–5 cm long. Flowers white with cleft petals, about 10 mm long. Found sparingly; in Boreal forest, Rocky Mountains, and has been reported from the Cypress Hills.

Silene noctiflora L.

NIGHT-FLOWERING CATCHFLY

A stout, very sticky, hairy, erect annual weed 30–90 cm high. Basal leaves short-stalked, 5–12 cm long, oblanceolate. Upper leaves stalkless, lanceolate, 2–8 cm long. Leaves hairy and slightly sticky. Sepals united to form an oval tubular calyx up to 12 mm long, with light and dark green upright stripes. White petals deeply cleft, opening only at night. An introduced weed; in grainfields; throughout southern Prairie Provinces.

Silene parryi (Wats.) Hitchc. & Mag.

MACOUN'S CAMPION

A perennial with erect stems 10–30 cm high. Leaves 5–7 cm long, linear-lanceolate; stem and leaves finely puberulent and densely glandular above. Flowers usually 3–7, with the pedicels of lateral flowers 2–3 cm long; calyx 8–10 mm long, prominently 10-nerved, glandular pubescent. Grassy slopes; southern Rocky Mountains.

Silene sibirica (L.) Pers.

SIBERIAN CAMPION

Perennial plants with erect subglabrous stems 40–60 cm high. Basal leaves oblong-linear; stem leaves all with sterile shoots in the axils. Inflorescence interrupted, the flowers in verticils, unisexual; calyx 4–5 mm long. Rare, an introduced weed.

Spergula spurry

Spergula arvensis L.

CORN SPURRY

An introduced annual, branching plant 15–40 cm high, with very narrow leaves 20–30 mm long, borne in whorls around the stem. White flowers about 6 mm across, in branching cymes at end of stem. Introduced, not common; sometimes found in fields and on roadsides; western Prairies.

Spergularia sand spurry

Spergularia marina (L.) Griseb. var. *leiosperma* (Kindb.) Gurke

SALT-MARSH SAND SPURRY

A low, annual, clustered plant, much-branched, about 10–20 cm high, with small ovate stipules at the bases of the opposite, linear, fleshy leaves. Leaves circular in cross section, 6–20 mm long. Flowers pink and numerous; petals shorter than sepals, which are about 3 mm long. This is a rather rare plant; found in the margins of saline sloughs; throughout Prairies and Parklands. Syn.: *S. salina* J. & C. Presl.

Stellaria stitchwort

Mostly low-growing or straggling plants with small white flowers, 5 petals so deeply cleft that often they look like 10 petals.

1. Plants glandular pubescent throughout. *S. americana*
Plants glabrous. 2
2. Plants annual; basal leaves ovate or
rhombic-ovate, petioled. *S. media*
Plants perennial; all leaves sessile. 3
3. Bracts of the inflorescence, at least the
upper ones, membranous. 4
Bracts of the inflorescence all green, not
membranous. 8
4. Upper part of stems and leaf margins
papillate, rough. *S. longifolia*
Upper part of stems and leaf margins
smooth. 5
5. Bracts and leaves or sepals ciliate. 6
Bracts, leaves, and sepals not ciliate. 7
6. Bracts and leaves ciliate; leaves to 15 mm
long. *S. calycantha*
Bracts and sepals ciliate; leaves to 10 mm
long. *S. ciliatisepala*
7. Inflorescence subumbellate; petals
absent. *S. umbellata*
Inflorescence cymose; petals present,
larger than sepals. *S. longipes*
8. Leaves ovate-lanceolate, length-to-width
ratio less than 4:1. 9
Leaves linear- or elliptic-lanceolate,
length-to-width ratio more than 4:1. 10
9. Sepals 1.5–2.5 mm long, obtuse; leaves
about 5 mm long. *S. obtusa*
Sepals 2.5–3.5 mm long, acute; leaves
5–10 mm long. *S. crispa*
10. Petals shorter than the sepals, or lacking. *S. calycantha*
Petals the same length as or longer than
the sepals. 11
11. Petals cleft more than halfway, almost to
base. 12
Petals cleft less than halfway. 13
12. Leaves mostly 6–20 mm long, 2–6 mm
wide; sepals 2–3 mm long. *S. crassifolia*
Leaves mostly 2–8 mm long, 1–4 mm
wide; sepals 3.5–6 mm long. *S. humifusa*
13. Plants usually less than 10 cm high; inter-
nodes short; leaves thick, rather rigid,
green. *S. laeta*
Plants usually 10–20 cm high; internodes
long; leaves not thick or rigid, glau-
cous. *S. longipes*

Stellaria americana Porter

Plants with slender rootstocks, mat-forming; stems leafy, 10–20 cm long; leaves 1–3 cm long; cymes leafy; sepals 3–4 mm long; petals exceeding the sepals. Rocky alpine slopes; southern Rocky Mountains.

Stellaria calycantha (Ledeb.) Bong.

NORTHERN STITCHWORT

A weak-stemmed trailing plant 15–50 cm long, with linear-lanceolate leaves 10–35 mm long. Not common; in shady woodlands; Rocky Mountains, Boreal forest, Cypress Hills. Syn.: *S. borealis* Bigel.

Stellaria ciliatisepala Trautv.

Plants with short creeping rootstocks; stems to 20 cm high; leaves to 10 mm long. Inflorescence few-flowered, or rarely a single flower. Open, rocky areas; Boreal forest.

Stellaria crassifolia Ehrh.

FLESHY STITCHWORT

A small rather weak-stemmed plant 5–25 mm high, with short, fleshy oblong-lanceolate leaves 5–15 mm long. White flowers 4–6 mm across, not numerous. Uncommon; may be looked for in wet and very shaded places.

Stellaria crispa Cham. & Schl.

Weak-stemmed plants with creeping rootstocks; leaves ovate, sessile or subsessile; flowers axillary, solitary; sepals 3–4 mm long; petals minute or absent. Moist places in mountains; southern Rocky Mountains.

Stellaria humifusa Rottb.

Mat-forming plants with creeping rootstocks; often reproducing by vegetative buds in axils of leaves; flowers 8–16 mm across. Sandy beaches; Hudson Bay.

Stellaria laeta Richardson

Mat-forming plants with creeping rootstocks, and a few-flowered inflorescence. The sepals ciliate, puberulent; the stem often puberulent. Sandy beaches; Hudson Bay, southern Rocky Mountains.

Stellaria longifolia Muhl.

LONG-LEAVED STITCHWORT

A weak-stemmed semierect plant 20–40 cm high, with bright green opposite linear leaves, tapered at both ends, 10–60 mm long. Numerous flowers 6–9 mm across, borne at the summit of the stem, with the petals deeply cleft and longer than the sepals. Fairly common; woodlands, moist and shady places; throughout the Prairie Provinces.

Stellaria longipes Goldie

LONG-STALKED STITCHWORT

An erect plant 6–30 cm high, with short lanceolate leaves 10–25 mm long, broadest near the base and tapering upward. Flowers few, on long stalks at the head of the stem, the 2-cleft white petals slightly longer than the sepals. Petals not usually so deeply cleft as in the preceding species. Common; in moist places and woodlands; throughout the Prairie Provinces.

Stellaria media (L.) Cyrill.

COMMON CHICKWEED

A prostrate-growing trailing annual, with lines of fine white hairs on the stems. Leaves broadly ovate, 5–25 mm long, on short stalks. Flowers white, about 6 mm across, and deeply cleft. An introduced European weed, which has become very common; on lawns and in gardens; throughout the Prairie Provinces.

Stellaria obtusa Engelm.

Matted plants with many prostrate branching stems 5–15 cm long; flowers solitary in leaf axils; sepals 2–3 mm long; petals minute or absent. Very rare; wet areas; southern Rocky Mountains.

Stellaria umbellata Turcz.

Erect or ascending plants with creeping rootstocks, 10–30 cm high. Leaves 5–20 mm long, linear-lanceolate, slim; flowers in upper leaf axils and in terminal umbellate cymes; sepals 2–5 mm long; petals minute or absent. Rare; moist areas in mountains; southern Rocky Mountains.

CERATOPHYLLACEAE—hornwort family

Ceratophyllum hornwort

Ceratophyllum demersum L.

HORNWORT

A completely submersed aquatic plant with thread-like leaves, 2 or 3 times forked, 5–25 mm long, arranged in whorls of 6–9 leaves at intervals on the stem. Flowers without stalks, unisexual, and borne singly in the axils of the leaves. Fruit an achene about 5 mm long, with a persistent style about 6 mm long. May be mistaken for the water milfoil, *Myriophyllum*, from which it is distinguished by its 2- or 3-forked leaves, the leaves of *Myriophyllum* being pinnate and not forked. Fairly common; in ponds and still water; eastern Prairies and Parklands, rare along northern fringe of Parklands and in Boreal forest.

NYMPHAEACEAE—water-lily family

Flowers yellow, with sepals petal-like, larger than the petals. *Nuphar*

Flowers white, with green sepals and large petals. *Nymphaea*

Nuphar yellow pond-lily

Leaves 5–10 cm long; flowers 15–20 mm across. *N. microphyllum*

Leaves 10–25 cm long; flowers 40–50 mm across. *N. variegatum*

Nuphar microphyllum (Pers.) Fern.

SMALL POND-LILY

Leaves submersed or floating, oval or elliptic in outline, deeply cordate at the base. Flowers yellow within; anthers 1.5–3 mm long, shorter than the filaments; the stigmatic disk red; fruit about 15 mm long. Ponds and lakes; eastern Parklands and Boreal forest.

Nuphar variegatum Engelm.

YELLOW POND-LILY

Perennial aquatic plant with stout creeping rootstock. Large single cordate leaves that float on the surface borne on long slender stalks, arising from the rootstock. Leaves 10–25 cm long and 8–15 cm broad. Occasionally a few entirely submersed leaves, thin and membranous. Flowers reddish within; anthers 4.5–7 mm, shorter than the filaments; stigmatic disk green; fruit about 4 cm long. Ponds and lakes; Parklands, Boreal forest, Cypress Hills.

Nymphaea water-lily

Leaves rotund in outline, 10–20 cm across;

flowers 5–8 cm across, fragrant. *N. odorata*

Leaves elliptic in outline, 7–12 cm long, 4–8

cm wide; flowers 2–5 cm across, odorless. *N. tetragona* ssp. *leibergii*

Nymphaea odorata Ait.

FRAGRANT WATER-LILY

Leaves commonly purple or red below, deeply cordate, with the petiole inserted almost at the middle of the leaf. Flowers with 17–32 petals, opening in the morning. Lakes and ponds; southeastern Boreal forest.

Nymphaea tetragona Georgi ssp. *leibergii* (Morong) Porsild SMALL WATER-LILY

Leaves with a V-shaped sinus. Flowers with 8–17 petals, opening in the afternoon. Ponds, lakes, and marshes; Boreal forest.

RANUNCULACEAE—crowfoot family

A large and variable family, with leaves alternate, except in *Clematis*; sepals 3–15, often colored, and resembling petals. Petals sometimes absent, but, when present, the same number as sepals. Many stamens. Fruit in the form of achenes, follicles (dry pods), or berries.

1. Climbing plants with opposite leaves;
without petals but with colored sepals;
fruit with persistent feathery style. *Clematis*
- Plants not climbing; leaves either basal or
alternate. 2
2. Flowers irregular or spurred. 3
- Flowers regular, not spurred. 6
3. Flowers irregular, not spurred. *Aconitum*
- Flowers spurred. 4

4. Flowers with 1 spur. *Delphinium*
 Flowers with 5 spurs. 5
5. Plants tall; leaves compound; flowers
 15–25 mm long. *Aquilegia*
 Plants small; leaves simple; flowers 3–5
 mm long. *Myosurus*
6. Fruit berry-like. *Actaea*
 Fruit not berry-like. 7
7. Fruit dry pods (or follicles). 8
 Fruit achenes. 10
8. Leaves simple, entire or incised. 9
 Leaves compound with 3 leaflets. *Coptis*
9. Leaves entire or toothed; flowers yellow. *Caltha*
 Leaves palmately incised; flowers white. *Trollius*
10. Leaves all basal, 3-lobed, with the lobes
 entire, rounded. *Hepatica*
 Leaves not all basal; stem leaves present;
 leaf segments mostly toothed, acute. 11
11. Petals usually present. *Ranunculus*
 Petals absent, but sepals colored and
 petal-like. 12
12. Leaves all basal except involucre of leafy
 bracts some distance below
 inflorescence. *Anemone*
 Leaves not basal; without involucral
 bracts below inflorescence. *Thalictrum*

Aconitum monkshood

Aconitum delphinifolium DC.

MONKSHOOD

Perennial plants 30–70 cm high, with short tubers. Stems finely pubescent; leaves glabrous, palmately lobed, with the segments lanceolate. Inflorescence a short raceme; flowers deep blue to purple. **Poisonous** to cattle. Mountain meadows; Rocky Mountains.

Actaea baneberry

Actaea rubra (Ait.) Willd.

RED BANE BERRY

Erect perennial herbs 30–100 cm high, with large compound leaves. Flowers small, white, in dense clusters at the ends of the stems, with the sepals falling off when the flower opens. Fruits large and berry-like, 6–10 mm long, and clustered in a raceme. **Poisonous** to humans, especially children. Fruit of the common typical species is bright red, but there is also a fairly common white-berried form, *Actaea rubra* forma *neglecta* (Gilman) Robins, white baneberry. Both baneberries common; in rich woodlands and in shady, wooded ravines; throughout the Prairie Provinces.

Perennial plants from bulb-like taproots, with basal leaves long-stalked and palmately divided or dissected. Flowering stalks bearing a whorl of involucreal leaves (large bracts resembling leaves) borne part way up the flowering stem. Flowers borne at the summits of long stems and consisting of 5 or 6 colored sepals but no petals. Fruits achenes borne in globular or cylindrical heads.

1. Styles long and feathery at maturity; sepals 15–40 mm long. 2
 Styles short; sepals 5–25 mm long. 3
2. Flowers purplish or bluish; bracts sessile; basal leaves expanding after flowering. *A. patens* var. *wolfgangiana*
 Flowers white; bracts short-petioled; basal leaves fully expanded at flowering. *A. occidentalis*
3. Carpels and achenes not woolly; glabrous or somewhat pubescent. 4
 Carpels and achenes densely woolly. 7
4. Involucreal leaves petioled; achenes fusiform, with the style short, hooked. *A. nemorosa* var. *bifolia*
 Involucreal leaves sessile or subsessile; achenes flattened. 5
5. Sepals yellow; styles very long in fruit, reflexed. *A. richardsonii*
 Sepals white; styles straight or hooked. 6
6. Achenes stipitate, glabrous; styles hooked, short. *A. narcissiflora*
 Achenes sessile, pubescent; styles straight, long. *A. canadensis*
7. Involucreal leaves long-petioled; heads of achenes cylindrical or narrowly ovoid. 8
 Involucreal leaves sessile or subsessile; heads of achenes globose or ovoid. 9
8. Sepals 8–10 mm long, greenish white; fruiting heads long cylindrical. *A. cylindrica*
 Sepals 10–20 mm long, white; fruiting heads oblong or oblong-ovoid. *A. virginiana*
9. Leaves with 3 broad wedge-shaped segments. *A. parviflora*
 Leaves 2–4 times ternate, with the segments linear or lanceolate. 10
10. Leaves 3–4 times ternate; mostly 1-flowered, with mature styles 4–6 mm long. *A. drummondii*
 Leaves 2–4 times ternate; mostly 1- to 4-flowered, with mature styles 1–2 mm long. *A. multifida* and var. *richardsiana*

A hairy-stemmed plant to 30 cm high, with several 5- to 7-parted basal leaves. In this species the flowering stems divide, and fresh stems appear, each bearing a whorl of involucre leaves and a flower at the end. Flowers white, 25–30 mm across. Fruiting head globular. One of the commonest anemones; found in large patches at the edges of woodlands, low moist places, and hollows; throughout the Prairie Provinces.

Anemone cylindrica A. Gray

LONG-FRUITED ANEMONE

Plants with a long slender stem 15–50 cm high, branching at the involucre leaves into 2–6 flowering stems, each bearing a 5-sepaled greenish white flower almost 20 mm across. Fruiting head long, cylindric, often 20 mm high, and densely woolly. Common; on moist prairie; throughout Prairies and Parklands.

Anemone drummondii Wats.

DRUMMOND'S ANEMONE

Plants with a stout woody caudex, silky hirsute stems 10–25 cm high, and long-petioled basal leaves. Flower usually solitary, appressed pubescent. Mountain meadows and alpine tundra; Rocky Mountains.

Anemone multifida Poir.

CUT-LEAVED ANEMONE

Erect plants, usually with several silky-haired purplish stems 15–60 cm high. Flowering stalks 1–7, one usually having no involucre leaves. Leaves cleft several times into very narrow lobes. Flowers varying from reddish purple to white or yellowish green and the fruiting heads globular and very woolly. Common; in moist spots; throughout the Prairie Provinces. A var., *richardsiana* Fern., with longer sepals, usually whitish, but sometimes bright red. Fairly common; in moister spots; throughout the Prairie Provinces.

Anemone narcissiflora L.

NARCISSUS ANEMONE

Plants with stout rootstocks; the stems pubescent, 20–40 cm high. Leaves deeply palmately divided, long-petioled. Inflorescence umbellate, 3- to 8-flowered. Mountain meadows; Rocky Mountains. Occurrence in Alberta not yet definitely established.

Anemone nemorosa L. var. *bifolia* (Farwell) Boiv.

WOOD ANEMONE

Plants with long creeping rootstocks; sterile leaves 3 or 5 foliate; flowering stems 10–20 cm high, usually with a single flower, and a single verticil of 3-petioled stem leaves. Moist woods; Parklands and Boreal forest. Syn.: *Anemone quinquefolia* L. var. *interior* Fern.

Anemone occidentalis Wats.

CHALICEFLOWER

Plants 10–60 cm high, at first densely villose, later often glabrate. Basal leaves few, 3-parted, with numerous linear-lanceolate segments; involucre leaves short-petioled. Flowering stalk elongating in fruit, the styles 2–4 cm long. Mountain meadows and slopes; Rocky Mountains. Syn.: *Pulsatilla occidentalis* (Wats.) Freyn.

Plants with slender rootstocks; stems 10–30 cm high, sparingly pubescent. Basal leaves long-petioled; the segments crenately lobed into broad, obtuse divisions; peduncle solitary, with a single flower. Woods, meadows, and slopes; Boreal forest, Rocky Mountains.

Anemone patens L. var. *wolfgangiana* (Bess.) Koch (Fig. 109) CROCUS ANEMONE

An early spring flowering plant, from a thick woody taproot. Flower appears before leaves on an erect silky stem 8–20 cm high. Basal leaves much divided, on long stalks, with the involucral leaves also being cleft, but without stalks. Flowers mauve or pale blue, sometimes white, 4–6 cm across, usually the first flower to appear on the prairies in spring. Fruiting stems tall, often over 30 cm high, bearing many achenes, each having a long persistent tail-like feathery style about 3 cm long. Common on open prairie and hills; throughout southern and western parts of the Prairie Provinces, and particularly conspicuous on burned-over prairie and railway rights-of-way. Sheep may be **poisoned** by this plant and their digestive system may be impaired by its felty hairs. Very common on overgrazed pasture, where dense stands during the early spring indicate an overgrazed condition. Syn.: *Pulsatilla ludoviciana* (Nutt.) Heller; *P. patens* (L.) Miller ssp. *teklae* (Zamels) Zamels.

Anemone richardsonii Hook.

Plants with slender yellow or brownish rootstocks; stems slender, delicate, 6–15 cm high. Basal leaves solitary, rotund or reniform in outline, 5-cleft into cuneate incised segments; involucral leaves 3, sessile or subsessile. Flowers sulfur yellow. Marshes, wet woods, and bogs; Boreal forest, Rocky Mountains.

Anemone virginiana L.

A tall erect species 30–80 cm high, with hairy stems. Flowers greenish white, with sepals about 10 mm long. Head of fruit is ovoid or cylindrical and woolly. A very variable species, found occasionally; in woods and shady places; in eastern part of the Prairie Provinces, but rare elsewhere. Some authorities consider the western plant to be *A. riparia* Fern.

Aquilegia columbine

Erect, branching plants from rootstocks, with slightly hairy and glandular stems. Leaves compound and divided into 3 leaflets. Flowers with 5 sepals colored and petal-like, also 5 colored petals extending at the base forming a long tubular spur. Fruit a follicle or dry capsule containing numerous seeds.

- 1. Plants small, usually less than 20 cm high; leaves all basal; usually 1-flowered. *A. jonesii*
- Plants 20–80 cm high, with stem leaves; several-flowered. 2
- 2. Blade of petals longer than the spurs; flowers blue or purple. *A. brevistyla*
- Blade of petals shorter than the spurs. 3
- 3. Flowers yellow; spurs of petals slightly hooked. *A. flavescens*



Fig. 109. *Crocus anemone*, *Anemone patens* L. var. *wolfgangiana* (Bess.) Koch.

Flowers scarlet; spurs of petals almost straight. 4

4. Sepals erect, shorter than the spur. *A. canadensis*
Sepals spreading, longer than the spur. *A. formosa*

Aquilegia brevistyla Hook. SMALL-FLOWERED COLUMBINE

A small plant 20–50 cm high; flowers blue or purple, nodding, with sepals about 15–20 mm long. Rare; in woodlands; Boreal forest, Rocky Mountains.

Aquilegia canadensis L. WILD COLUMBINE

A stout erect plant 30–80 cm high; flowers large, nodding, with scarlet sepals and sometimes yellow petals. Stamens and styles much longer than sepals and protruding conspicuously from the flower. Fairly common; in open woodlands; eastern Boreal forest, reported from Qu'Appelle Valley in Saskatchewan.

Aquilegia flavescens S. Wats. YELLOW COLUMBINE

A slender branching plant 30–80 cm high, with yellow or yellowish white flowers. In woodlands; Rocky Mountains.

Aquilegia formosa Fisch. CRIMSON COLUMBINE

A stout glabrous or sparingly pubescent plant 30–80 cm high, with sepals about 2 cm long; sepals and spurs crimson; the lamina of the petals yellow. Open woods; southern Rocky Mountains.

Aquilegia jonesii Parry BLUE COLUMBINE

Plants with a short caudex; the leaves crowded, finely pubescent, and often glaucous. Flowering stem with a single erect flower about 1.5 cm long; sepals blue; petals often whitish. Exposed slopes, rocky areas; southern Rocky Mountains.

Caltha marsh-marigold

1. Stems erect; bearing a single leaf or none;
flowers 1 or 2. *C. leptosepala*
Stems ascending, decumbent or floating;
bearing several leaves; flowers several. 2
2. Sepals white or pink; flowers not over 15
mm across; plants often floating. *C. natans*
Sepals yellow; flowers over 15 mm across;
plants usually rooted in mud. *C. palustris*

Caltha leptosepala DC. MOUNTAIN-MARIGOLD

Plants 10–40 cm high, from a short erect rootstock. Basal leaves long-petioled; blades oval to oblong-oval, cordate at base, with the margins toothed. Flowers 2.5–3.5 cm across, white or bluish. Alpine meadows; Rocky Mountains.

Plants of lakes and small ponds, with cordate to kidney-shaped leaves and stems rooting at the nodes. Flowers 10–15 mm across, with white or pinkish petal-like sepals; fruiting clusters about 6 mm across. Quite rare; has been found floating or rooted in the mud in woodland lakes and slow streams; Boreal forest.

Caltha palustris L. (Fig. 110)

Stout marsh plants with smooth hollow stems arising from coarse fleshy roots. Leaves round or kidney-shaped and heart-shaped at the base, the basal ones with long stalks and the upper ones stalkless. Flowers bright yellow, 20–30 mm across, with 5–9 sepals but no petals. Fruits many-seeded follicles. Common in wet places, more common in slightly moving water; Parklands and Boreal forest.

Clematis virgin’s-bower

Climbing vines with more or less woody stems and opposite leaves. Plants climb by clasping the supporting plant with the stalks of their leaves and not by means of tendrils. Flowers have colored sepals but no obvious petals; fruit an achene with a long, persistent, feathery style.

- 1. Flowers solitary. 2
Flowers in panicles or corymbs. 3
- 2. Flowers blue; leaves with 3 leaflets. *C. verticellaris* var. *columbiana*
Flowers yellow; leaves with 5 leaflets. *C. tangutica*
- 3. Flowers yellowish or greenish; leaves with
3 leaflets. *C. virginiana*
Flowers white; leaves with 5–7 leaflets. *C. ligusticifolia*

Clematis ligusticifolia Nutt.

A climbing plant, with opposite leaves divided into 5-stalked leaflets arranged pinnately on stalk, each leaflet pointed at tip, 2–8 cm long. Flowers borne in clusters, each flower about 10–15 mm across with white sepals and no petals. Fruit a cluster of achenes, each with a persistent feathery style. Common; in coulees and ravines, climbing over bushes and shrubs; Prairies.

Clematis tangutica (Max.) Korsh.

Plant with greenish yellow flowers 3–5 cm across. Leaflets 2–5 cm long, coarsely toothed. Introduced ornamental, locally escaped and established; in coulees and shrubbery.

Clematis verticellaris DC. var. *columbiana* (Nutt.) A. Gray

A climbing plant, with opposite leaves divided into 3 long-stalked leaflets, each 2–8 cm long, pointed at the tip. Flowers 5–10 cm across, having 4 (sometimes 5) pale blue petal-like sepals. Petals reduced, resembling sterile stamens. Achenes borne in clusters, each with a feathery style about 5 cm long. In shady woodlands; in Cypress Hills and Foothills region. Syn.: *C. columbiana* (Nutt.) T. & G.




Fig. 110. Marsh-marigold, *Caltha palustris* L.

Clematis virginiana L.

VIRGIN'S-BOWER

Plants climbing, with stems 3–4 m long; leaflets usually 3, ovate, commonly coarsely toothed. Panicles from many of the leaf axils many-flowered. Woods; eastern Parklands and Boreal forest.

Coptis goldthread

Coptis trifolia (L.) Salisb.

GOLDTHREAD

Low, evergreen, perennial herbs 5–15 cm high. The leaves all basal, long-stalked, each bearing 3 leaflets. The single flowers about 6 mm across, on a long slender stem, and bearing 5–7 white sepals with yellowish bases, and smaller club-shaped petals, each with a tiny drop of nectar in its summit. Fruit a cluster of follicles, each about 6 mm long. Called goldthread because of its fine yellow rootstocks. In moist spots; Boreal forest.

Delphinium larkspur

Perennial herbs with alternate much-divided or lobed leaves. Flowers in racemes, perfect, irregular, with 5 petal-like colored sepals, the upper one extended at the base into a spur. Petals 4, the upper pair extended into spurs projecting into the sepal spur, the lower pair small with short claws. Fruit a many-seeded follicle.

Plants tall and erect, 50–150 cm high; racemes of flowers spike-like; lower flower stalks shorter than flowers. *D. glaucum*

Plants low, 15–50 cm high; racemes of flowers loose and spreading; lower flower stalks longer than flowers. *D. bicolor*

Delphinium bicolor Nutt. (Fig. 111)

LOW LARKSPUR

A perennial from a thick, fleshy, fibrous root, with hairy stems 15–50 cm. Leaves finely hairy, much cleft and dissected, on long stalks. Flowers dark blue, with a long blue spur at the rear. Flowers 15–35 mm across, borne on long stalks in a loose terminal raceme. Blooming in early spring, May, or June. Fruit a dry follicle 15–20 mm high, brownish, containing many seeds. Not common, but locally very abundant; in large patches in openings in woodlands, hillsides, and sheltered areas; on Cypress loam soils throughout the southwest, particularly in the Cypress Hills and Wood Mountains, also in the Rocky Mountains. Very **poisonous** to cattle and causing some heavy losses, but apparently harmless to sheep.

Delphinium glaucum Wats. (Fig. 112)

TALL LARKSPUR

A tall erect plant 30–150 cm high, with smooth stem and deeply cut, dissected slightly hairy leaves. Dark blue flowers borne on very short stalks in a long close spike-like raceme. Fairly abundant; throughout the Foothills region. Very **poisonous** to cattle. Growing in association with open stands of aspen poplar and willows in the Rocky Mountains and western Boreal forest. Syn.: *D. brownii* Rydb.; *D. canmoreense* Rydb.



HERBARIUM - THE UNIVERSITY OF CALIFORNIA
JAN 10 1891



Fig. 111. Low larkspur, *Delphinium bicolor* Nutt.



Fig. 112. Tall larkspur, *Delphinium glaucum* Wats.

Hepatica hepatica

Hepatica nobilis Miller

LIVERLEAF

Plants with short thick brown rootstocks; scapes to 15 cm high. Leaves overwintering and persisting until flowering, reniform in outline, 5–15 cm broad, 3-lobed, the lobes obtuse-ovate; petioles 5–15 cm long, pubescent. Flowers 15–25 mm across, varying in color from white or pinkish to bluish purple; perianth with 6–9 segments, with 3 bract-like stem leaves appearing to form a calyx. Achenes hirsute. Very rare; dry to moist woods; eastern Parklands and Boreal forest.

Myosurus mousetail

Small low-growing annual mud plants from fibrous roots. Leaves all basal and linear or thread-like. Flowering scapes bearing 1 flower with 5 sepals and 5 narrow greenish petals. Occasionally 6 or 7 sepals and petals, and in some instances the petals absent. Fruits numerous small achenes crowded on a tall narrow spike, from which the name mousetail was given to the genus.

Myosurus minimus L.

LEAST MOUSETAIL

Small plants, 5–15 cm high. Sepals and petals 3–4 mm long; achenes 1–1.5 mm long, with a very short beak. Not common; in muddy slough margins and marshy areas; throughout Prairies.

Ranunculus buttercup

A large genus, usually with alternate leaves. Flowers (Fig. 7) perfect and regular, with 5 sepals and usually 5 petals, either yellow or white. Fruits numerous achenes borne in a head or a short spike.

1. Plants aquatic; leaves submerged, finely dissected. 2
Plants not aquatic, mud or dryland plants; leaves not finely dissected. 3
2. Flowers white, axillary. *R. aquatilis*
Flowers yellow, terminal or in cymes. *R. gmelinii*
3. Basal leaves elliptic to linear, not deeply divided. 4
Basal leaves ovate to rhomboid, reniform, entire to compound. 6
4. Flowers white; leaves with a petiole almost as wide as the blade. *R. pallasi*
Flowers yellow. 5
5. Basal leaves elliptic to lanceolate, entire; stem leaves lobed. *R. glaberrimus*
Basal leaves linear-lanceolate to filiform. *R. flammula*
6. Achenes pubescent. 7
Achenes glabrous. 9

7. Basal leaves deeply 3-parted or lobed almost to base. *R. uncinatus*
 Basal leaves crenate or lobed to almost halfway. 8
8. Stems and leaves pilose-pubescent or villose. *R. cardiophyllus*
 Stems and leaves subglabrous or puberulent. *R. inamoenus*
9. Basal leaves crenate or lobed to almost halfway. 10
 Basal leaves lobed almost to the base, or divided. 12
10. Plants stoloniferous, creeping; leaves cordate at the base. *R. cymbalaria*
 Plants not stoloniferous, stems upright. 11
11. Leaves deltoid-ovate in outline, acute to rounded at base. *R. rhomboideus*
 Leaves reniform in outline, more or less cordate at base. *R. abortivus*
12. Stem leaves and leaves in the inflorescence sessile or subsessile. 13
 Stem and inflorescence leaves on well-developed petioles. 15
13. Plants small, usually less than 10 cm high; petals as long as or shorter than sepals. *R. pygmaeus*
 Plants usually more than 10 cm high; petals as long as or longer than sepals. 14
14. Segments of stem leaves linear, 1–3 mm wide. *R. pedatifidus* var. *leiocarpus*
 Segments of stem leaves lanceolate, 3–6 mm wide. *R. nivalis*
15. Plants small, rarely more than 5 cm high; leaves mostly 3- or 5-lobed. *R. hyperboreus*
 Plants larger, with the stems usually erect. 16
16. Leaves simple. 17
 Leaves compound. 22
17. Stems bearing a single leaf and a single flower. *R. lapponicus*
 Stems bearing several leaves and flowers. 18
18. Sepals reflexed at the middle or base. 19
 Sepals spreading or curved inward. 20
19. Petals and sepals about the same length, 3–5 mm. *R. uncinatus*
 Petals about 3 times as long as the sepals. *R. occidentalis*
20. Plants villose-pubescent, especially the petioles. *R. acris*

Plants glabrous to subglabrous or puberulent.	21
21. Beak of the achene very short, to 0.1 mm long.	<i>R. sceleratus</i>
Beak of the achene developed, 0.5–1.0 mm long.	<i>R. gmelinii</i>
22. Flowers 6–8 mm across, with the petals shorter than the sepals.	<i>R. pensylvanicus</i>
Flowers larger, with the petals longer than the sepals.	23
23. Leaves divided into many narrow segments.	24
Leaves trifoliate.	25
24. Plants subglabrous; petals about 5 mm long, 4–5 mm wide.	<i>R. gelidus</i>
Plants appressed pubescent; petals about 10 mm long, 5 mm wide.	<i>R. fascicularis</i>
25. Sepals reflexed at base, 3.5–5 mm long; petals 4–7 mm long.	<i>R. macounii</i>
Sepals spreading or incurved; petals 7–15 mm long.	26
26. Plants stoloniferous; beak of the achene about 1 mm long.	<i>R. repens</i>
Plants not stoloniferous; beak of the achene to 3 mm long.	<i>R. septentrionalis</i>

Ranunculus abortivus L.

SMOOTH-LEAVED BUTTERCUP

A biennial plant 15–50 cm high, with a rather fleshy smooth stem. Basal leaves long-stalked, usually cordate-based, and round with wavy margins; stem leaves usually 3-cleft, the upper ones without stalks. Petals usually shorter than the reflexed (turned downward) sepals; flowers yellow, 6–10 mm across. Found in open woodlands, ravines, and moist places. Plants growing on margins of streams or sloughs and wet spots have large and branched inflorescence with numerous small flowers; when growing in less favorable spots, more erect, with fewer flowers and less-branched inflorescence. The two phases may be mistaken for different species. Fairly plentiful throughout Prairies and Parklands; rare in Boreal forest.

Ranunculus acris L.

TALL BUTTERCUP

A tall erect perennial, branched above, with a hairy stem 30–80 cm high. Basal leaves stalked and much cleft and divided; upper leaves divided into 3 lobes, with short stalks. Numerous yellow flowers 20–35 mm across, with petals more than twice as long as sepals. Achenes smooth, flattened, with a short beak. Introduced from Europe. Not common; found occasionally in moist places and along railway grades; throughout the Prairie Provinces. Very plentiful, however, in some irrigated areas, especially in southern Alberta.

Ranunculus aquatilis L.

LARGE-LEAVED WATERCROWFOOT

A fully submersed aquatic plant, with leaves finely dissected, about 10–25 mm long, growing from a sheathing stipule. Flowers either floating on the water or protruding above it, white, about 10–15 mm across. Achenes with a short beak not more than one-third the length of the body of the achene. In slowly moving water and brackish pools that are not saline; found throughout the Prairie Provinces. A very variable species, of which the following varieties occur:

- 1. Beak of the achene more than 0.5 mm long. var. *longirostris* (Godr.) Laws.
 Beak of the achene less than 0.5 mm long. 2
- 2. Leaves stiff, with a grayish green petiole included in the sheath. var. *subrigidus* (Drew) Breit.
 Leaves not stiff, with the green petiole to 1 cm long. 3
- 3. Stem 1.0–1.5 mm thick; plants large. var. *capillaceus* (Thuill.) DC.
 Stem less than 1 mm thick; plants small and reduced. var. *eradicatus* Laest.

Ranunculus cardiophyllus Hook. (Fig. 113)

HEART-LEAVED BUTTERCUP

An erect species 15–50 cm high, with few branches. Basal leaves stalked, round to ovate, sometimes lobed, and usually heart-shaped at the base; stem leaves with narrow, almost linear lobes, very short-stalked. Yellow flowers 10–20 mm across. Not common; in moist places, meadows, and stream banks; in western Parklands.

Ranunculus cymbalaria Pursh

SEASIDE BUTTERCUP

Low perennial plants with runners and predominantly basal leaves on thin stalks. Leaves small, rounded with cordate bases and wavy margins, 5–15 mm long. Flowers with a conical center, yellow, and quite small, usually 5–8 mm across; the petals slightly shorter than the sepals. Achenes numerous, with longitudinal grooves. Common; on margins of sloughs and lakes, on saline wet areas and stream banks; throughout the Prairie Provinces.

Ranunculus fascicularis Muhl.

EARLY BUTTERCUP

Plants with 2–5 stems 10–30 cm high, strigose. Leaves mostly basal, ovate in outline, with the segments lobed. Flowers 15–25 mm across, long-peduncled. Southern Parklands and Boreal forest.

Ranunculus flammula L. var. *ovalis* (Bigel.) Benson

CREEPING SPEARWORT

A low-growing trailing plant, with stems rooting at the nodes and linear or spatulate leaves 10–25 mm long. Bright yellow flowers about 8 mm across borne singly on short stems. Uncommon; on lakeshores and riverbanks; Boreal forest, Parklands. In var. *ovalis* the leaves lanceolate and about 5 mm wide. Plants with filiform leaves, usually 1–2 mm wide, are var. *filiformis* (Michx.) Hooker.



HERBARIUM OF THE RESEARCH STATION
WINDY MOUNTAIN, ALBERTA

ORIGINAL LOCALITY: ...
 DURATION: ...
 DATE: ...
 ELEVATION: ...
 HABITAT: ...



Fig. 113. Heart-leaved buttercup, *Ranunculus cardiophyllus* Hook.

Ranunculus gelidus Kar. & Kir.

DWARF BUTTERCUP

Plants very small, with stems 5–10 cm high, glabrous or nearly so, bearing 1–3 leaves, these 3–5 foliate. Flowers 1–3, long-peduncled. Scree slopes; Rocky Mountains.

Ranunculus glaberrimus Hook.

SHINY-LEAVED BUTTERCUP

A very early low-growing smooth-stemmed buttercup. Basal leaves stalked, some entire and not lobed, others 3-lobed. In the type the lower leaves usually rounded or kidney-shaped and wavy-margined, but the plants usually found are the var. *ellipticus* Greene (called *R. buddii* by Boivin), with entire long-elliptical basal leaves. Flowers 10–20 mm across, yellow, usually with a lavender or purplish tinge on the backs of the sepals. Occasionally large patches of this species with only one floral ring are found (called *R. buddii* f. *monochlamydeus* by Boivin). A very early spring flowering plant, local, and not usually abundant; plentiful in certain parts of the Prairies.

Ranunculus gmelinii DC.

SMALL YELLOW WATERCROWFOOT

A semiaquatic perennial with underwater leaves divided into very narrow, flat segments, the floating leaves having wider lobes. Petals yellow, about 5 mm long; head of fruit globose, about 5 mm thick. Occasionally found in ponds and lakes; throughout the Prairie Provinces except the Prairies.

Ranunculus hyperboreus Rottb.

BOREAL BUTTERCUP

Glabrous aquatic plants, with the stems floating, freely branching. Leaves with simple reniform blades, lobed or 3-parted. Petals 2–4 mm long, the same length as the sepals. Wet areas; Boreal forest. Large plants with cordate leaves and petals 3–4 mm long are var. *intertextus* (Greene) Boiv.; rare; Rocky Mountains.

Ranunculus inamoenus Greene

GRACEFUL BUTTERCUP

A species 15–30 mm high, with stalked basal leaves varying from almost circular to fan-shaped and sometimes divided; stem leaves divided into 3 segments. Petals small, usually less than 6 mm long. Not common; has been found in the Cypress Hills and Rocky Mountains.

Ranunculus lapponicus L.

LAPLAND BUTTERCUP

Plants with slender rootstocks; stems solitary, 10–20 cm high. Basal leaf 1 or rarely 2; cauline leaf solitary or absent. Flower solitary, with 3 sepals and 6–10 petals. Wet woods and muskeg; Boreal forest.

Ranunculus macounii Britt.

MACOUN'S BUTTERCUP

A tall, branching hairy species 30–60 cm high, usually decumbent when partly grown. Leaves in broad divisions, with the leaf segments usually stalked. Yellow flowers 10–15 mm across; achenes having a sharp stout beak, about one-quarter the length of the body. Fairly common; in moist and wet places; throughout the Prairie Provinces.

Ranunculus nivalis L.

SNOW BUTTERCUP

Plants usually unbranched, 10–20 cm high. Basal leaves orbicular-reniform, deeply lobed. Sepals villous; achenes with a beak about as long as the body. Tundra and alpine prairies. In var. *nivalis* the pubescence of the sepals brown to blackish; Hudson Bay; in var. *eschscholtzii* (Schlecht.) Benson the pubescence pale yellow; Rocky Mountains. Syn.: *R. verecundus* Rob.

Ranunculus occidentalis Nutt.

WESTERN BUTTERCUP

Erect or somewhat decumbent plants, with stems 20–70 cm high, branching above. Basal leaves 3-parted, with the lobes cuneate, the petioles pubescent. Flowers large, 15–25 mm across. Meadows and prairies; Rocky Mountains. Several varieties have been described; most plants from Alberta are var. *brevistylis* Greene, with narrowly obovate petals, and the achene 2–2.5 mm long with a straight beak about half as long as the body.

Ranunculus pallassii Schlecht.

PALLAS BUTTERCUP

Plants small, creeping, with stems 10–15 cm high. Leaves obovate-cuneate in outline, deeply 3-lobed. Flowers solitary, 20–25 mm across, white or somewhat violet-tinged; achenes 5–7 mm long. Wet shores; Hudson Bay, Boreal forest.

Ranunculus pedatifidus J. E. Smith var. *leiocarpus* (Trautv.) Fern.

NORTHERN BUTTERCUP

An erect plant 10–30 cm high. Basal leaves deeply dissected into linear divisions. In the typical species, flowers 8–15 mm across with yellow petals and greenish yellow sepals. Fruit has a slender curved beak. The type is apparently a mountain and arctic plant, but the apetalous form has been found on open grassland in Prairies and Parklands. Formerly known as *R. affinis* R. Br. This form has only one floral ring of greenish yellow petal-like sepals.

Ranunculus pensylvanicus L. f.

BRISTLY BUTTERCUP

An erect species, usually annual, very hairy, 30–60 cm high. Leaves divided into 3-stalked segments, again divided into 3 lobes. Yellow flowers 6–8 mm across, with petals shorter than reflexed sepals; achenes with a sharp beak about one-third the length of the body. Not common; occasionally found in wet places; Parklands and Boreal forest.

Ranunculus pygmaeus Wahl.

PYGMEE BUTTERCUP

Small plants with a rootstock, and erect stems 1.5–5 cm high. Leaves about 1 cm long, glabrous. Flowers 5–10 mm across, with petals and sepals about the same length; achenes about 1 mm long. Rare; alpine tundra; Rocky Mountains.

Ranunculus repens L.

CREEPING BUTTERCUP

Plants stoloniferous, with stems 30–45 cm high. Leaves 3-foliolate. Flowers 20–30 mm across, yellow. Introduced species; occasionally found in wet areas.

Ranunculus rhomboideus Goldie

PRAIRIE BUTTERCUP

A short hairy-stemmed low-growing species 15–45 cm high. Rounded or oval wavy-margined basal leaves on long stalks; stem leaves deeply cleft or

divided and without stalks. Flowers yellow, 10–20 mm across; petals rather narrow, appearing to be somewhat widely spaced. Flowering very early in the spring. Fruiting head globular; achenes with a short beak. Perhaps the most common species on the prairies; over all the open plains; the Prairie Provinces. Syn.: *R. ovalis* Raf.

Ranunculus sceleratus L.

CELERY-LEAVED BUTTERCUP

A hollow-stemmed annual plant, with smooth stems 15–60 cm high. Leaves deeply 3- to 5-lobed and thick; basal leaves with long stalks but upper stem leaves stalkless. Yellow flowers small and numerous, 6–8 mm across, with petals about the same length as sepals. Common; along shores of lakes and sloughs and often the dominant plant of the hummocky bed of a drying slough; throughout the Prairie Provinces. Its acrid **poisonous** juice blisters the skin and produces intestinal inflammation if eaten. Sometimes called cursed crowfoot because of its toxic nature.

Ranunculus septentrionalis Poir.

SWAMP BUTTERCUP

Stout-stemmed plants 30–60 cm high, erect or ascending, often rooting at the lower nodes of prostrate stems. Flowers 20–30 mm across; achenes 3–4 mm long, with a straight beak about half the length of the body. Wet areas; southeastern Parklands and Boreal forest.

Ranunculus uncinatus D. Don

HAIRY BUTTERCUP

Plants with erect hirsute stems 30–60 cm high. Leaves uniform, cordate at the base, 3-parted. Flowers about 5 mm across, with the petals 2–3 mm long, the sepals reflexed; achenes 1.5–2 mm long, flattened, with a beak as long as the body, strongly recurved. Moist areas; Rocky Mountains, Peace River.

Thalictrum meadow-rue

Perennial plants from rootstocks, with alternate leaves divided into 3 leaflets, each 3-lobed. The segments and the leaflets are usually stalked. Flowers, either perfect or unisexual, with greenish white sepals and no petals, borne in rather open panicles. The yellow stamens of the male flowers make the inflorescence very attractive. Fruits achenes, usually ribbed, and borne in small clusters.

1. Flowers perfect; achenes flat and half obovoid with a straight back. *T. sparsiflorum* var. *richardsonii*
Flowers unisexual. 2
2. Stem leaves barely stalked; leaves oblong, usually longer than wide, often very finely hairy beneath; stems purplish. *T. dasycarpum*
All leaves with stalks; leaflets almost round, pale, and smooth beneath. 3
3. Leaves thin, not distinctly veined; achenes similarly pointed at both ends. *T. occidentale* var. *palousense*
Leaves thick and prominently veined; achenes obovoid and sharper at the upper end than at the lower. *T. venulosum*

Thalictrum dasycarpum Fisch. & Lall. (Fig. 114)

TALL MEADOW-RUE

A tall, erect purplish-stemmed plant 50–150 cm high. Leaves compound, the long leaflets usually longer than wide, with 3 pointed lobes, dark green above, and pale and prominently veined beneath, often with very fine hairs beneath. Flowers borne in a long open panicle 30 cm long or longer. Flowering in midsummer. A fairly common species; in rich moist woodland; throughout the Prairie Provinces.

Thalictrum occidentale A. Gray var. *palousense* St. John WESTERN MEADOW-RUE

An erect smooth-stemmed perennial 30–60 cm high, with thin pale green leaflets almost round but margins many-lobed. Flowers in an open panicle, appearing fairly early in the spring. In shady moist places, woodlands, and coulees; Cypress Hills, Rocky Mountains, Peace River.

Thalictrum sparsiflorum Turcz. var. *richardsonii* (Gray) Boiv.

FEW-FLOWERED MEADOW-RUE

A species with perfect flowers. Leaflets glandular below and scented. Rare; listed from several localities.

Thalictrum venulosum Trel.

VEINY MEADOW-RUE

Plants 15–200 cm high, with the leaflets rounded, strongly veined, bluish green, pale below. Panicle of flowers narrow and dense. Achenes ovoid. Fairly common; in woodlands; throughout most of the Prairie Provinces. Three varieties can be recognized:

1. Plants 15–40 cm high; filaments 3–4 mm long; fruit 3–4 mm long. var. *venulosum*
Plants larger. 2
2. Plants 60–80 cm high; filaments 4–5 mm long; fruit 4–4.5 mm long. var. *turneri* Boiv.
Plants 1–2 m high, lax; filaments 5–6 mm long; fruit 5–6.5 mm long. var. *lunellii* (Greene) Boiv.

Trollius globeflower

Trollius laxus Salisb.

AMERICAN GLOBEFLOWER

Plants with a thick caudex, and erect stems 20–40 cm high, glabrous, 2- to 4-leaved. Flowers usually solitary, 2–4 cm across, whitish. Follicles about 1 cm long. Marshy areas and alpine meadows; Rocky Mountains.

BERBERIDACEAE—barberry family

Plants shrubby, the leaves pinnate with spinulose toothed leaflets.

Berberis

Plants herbaceous, with a single compound leaf, the leaflets 5–8 cm long.

Caulophyllum



Fig. 114. Tall meadow-rue, *Thalictum dasycarpum* Fisch. & Lall.

Berberis barberry

Berberis aquifolium Pursh

MAHONIA

Shrubby plants 10–100 cm high, the leaves alternate with 5–7 leathery leaflets. Inflorescence a raceme with yellow flowers; fruit a blue berry with a few large seeds. In mountain woods; southern Rocky Mountains. Plants with trailing stems 10–30 cm high are forma *repens* (Lindl.) Boiv.

Caulophyllum blue cohosh

Caulophyllum thalictroides (L.) Michx.

BLUE COHOSH

Plants erect, 30–80 cm high. The leaf inserted at about the middle of the stem, consisting of 3 long-petioled leaflets each of which is twice divided into 3 segments; the ultimate leaflets obovate-oblong. A smaller leaf subtends the inflorescence, consisting of a panicle of yellowish green or purplish flowers, about 1 cm across; seeds drupe-like, blue. Rare; moist woods; southeastern Parklands and Boreal forest.

MENISPERMACEAE—moonseed family

Menispermum moonseed

Menispermum canadense L.

YELLOW PARILLA

Plants woody climbers 2–3 m high; leaves 10–15 cm long, suborbicular, peltate near the margin. Inflorescence a panicle; flowers 3–5 mm across; fruit a blackish drupe. Woods; southeastern Parklands and Boreal forest.

PAPAVERACEAE—poppy family

Petals usually 4; stems leafy; capsule opening

by a circle of pores under the broad stigma. *Papaver*

Petals 8–12; leaf solitary, basal; capsule 2-

valved. *Sanguinaria*

Papaver poppy

1. Plants perennial; the leaves all basal. 2

Plants annual; cauline leaves present. 3

2. Flowers sulfur yellow; leaves hirsute to subglabrous. *P. pygmaeum*

Flowers red or reddish orange; leaves glabrous and glaucous. *P. nudicaule*

3. Culine leaves cordate-clasping, glabrous, not divided. *P. somniferum*

Culine leaves not clasping, pubescent, pinnately divided. *P. rhoeas*

Papaver nudicaule L.

ICELAND POPPY

Plants with a rosette of basal leaves, and scapose stems to 50 cm high bearing solitary flowers to 6 cm across. Introduced ornamental; in various locations escaped and established.

Papaver pygmaeum Rydb.

ALPINE POPPY

Plants with a rosette of basal leaves 2–3 cm long, subglabrous to pubescent. Scape 4–10 cm high, hirsute; flower 2–3 cm across; capsule about 1 cm long, densely bristly. Alpine areas; southern Rocky Mountains.

Papaver rhoeas L.

CORN POPPY

Plants 30–60 cm high, hirsute; leaves pinnately divided, with the pinnae lobed or toothed. Flowers 5–7 cm across, scarlet. Introduced; occasionally escaped and established.

Papaver somniferum L.

COMMON POPPY

Plants 60–100 cm high, usually blue green, glabrous. Flowers 6–10 cm across, white to purplish or red, with the petals black at the base; capsule 5–8 cm long, to 5 cm across. Introduced as ornamental and for poppy seed; occasionally escaped and established.

Sanguinaria bloodroot

Sanguinaria canadensis L.

RED PUCCOON

Plants with a stout rootstock, a single basal leaf, and scape with a solitary flower. Leaf orbicular in outline, 3- to 9-lobed, with the lobes undulate. Flowers white or pinkish, 2–5 cm across. Capsule 3–5 cm long. The name derives from the brilliant red sap, exuding when the plant is wounded. Rare; more or less open woods; southeastern Parklands and Boreal forest.

FUMARIACEAE—fumitory family

- 1. Plants climbers; corolla persistent, the two outer petals saccate. *Adlumia*
- Plants not climbing; corolla deciduous, only one of the petals saccate. 2
- 2. Fruit an elongate capsule with more than one crested seed. *Corydalis*
- Fruit globose, with a single noncrested seed. *Fumaria*

Adlumia fumitory

Adlumia fungosa (Ait.) Greene

CLIMBING FUMITORY

A biennial vine, stemless in the first year, growing to a climbing vine 2–3 m long. Leaves pinnate-ternate, with the rachis of the uppermost leaflets elongating and twining around supporting vegetation. Flowers pink, with the

corolla becoming spongy and enclosing the capsule. Very rare; moist woods; southeastern Boreal forest.

Corydalis corydalis

Flowers yellow; plants usually somewhat prostrate. *C. aurea*

Flowers pink with yellow tips; plants fairly erect. *C. sempervirens*

Corydalis aurea Willd.

GOLDEN CORYDALIS

A much-branched low, sometimes prostrate annual or biennial, with pale green much-dissected leaves. Plants sometimes spread about 45 cm. Racemes of narrow golden yellow flowers, each 10–15 mm long, with a spur at the base about half as long as the body of the corolla. Fruits narrow pods 15–25 mm long containing many small shiny seeds. Common; in edges of woodlands, moist spots, railway grades; throughout the Prairie Provinces. Reported to be somewhat **poisonous** to sheep.

Corydalis sempervirens (L.) Pers. (Fig. 115)

PINK CORYDALIS

A slender-stemmed annual or biennial plant 30–75 cm high, with alternate twice-divided leaves. Flowers 10–20 mm long, in a raceme, pinkish with yellow tips. Fruits long, narrow cylindric capsules, turning brown at maturity, and containing many seeds. Often found; in rocky woodlands and disturbed areas; Boreal forest.

Fumaria fumitory

Fumaria officinalis L.

EARTH-SMOKE

Annual plants 20–80 cm high, with the stems lax, diffusely branched. Inflorescence a many-flowered raceme 2–4 cm long; flowers 7–9 mm long, reddish purple; fruit about 2.5 mm in diam. Introduced; occasionally a weed.

CAPPARIDACEAE—caper family

Pods borne on a stalk about 10 mm long above the calyx; stamens 6. *Cleome*

Pods on a very short stalk above the calyx, clammy to touch; stamens 8–32. *Polanisia*

Cleome spiderflower


Cleome serrulata Pursh (Fig. 116)

SPIDERFLOWER, PINK CLEOME

An erect, branching annual plant 30–100 cm high. Stem smooth and hairless. Leaves trifoliolate with lanceolate leaflets 2–7 cm long. Flowers usually pale pink, sometimes white, stamens protruding above the flower. Pods 2–5 cm long, usually slightly curved, and containing one compartment with a single row of large seeds. Plant strong and unpleasant smelling. Very common; along roadsides and on light soil; Prairies. Syn.: *Peritoma serrulatum* (Pursh) DC.



Fig. 115. Pink corydalis, *Corydalis sempervirens* (L.) Pers.



A.C. Budd

Fig. 116. Spiderflower, *Cleome serrulata* Pursh.

Polanisia clammyweed

Polanisia dodecandra (L.) DC.

CLAMMYWEED

A sticky, glandular annual, branched, 15–50 cm high, with trifoliolate leaves, leaflets 10–25 mm long. Flowers yellowish white or pink, about 6 mm long, with pods 20–35 mm long, glandular. Not common; found on sandy soils; throughout Prairies and southeastern Parklands. Syn.: *P. graveolens* Raf. Large-flowered plants are distinguished as var. *trachysperma* (T. & G.) Iltis, large-flowered clammyweed, with flowers 8–12 mm long, pale yellow, the stamens protruding. Not common; on rocky banks, hillsides, and light soils; same locations as the species.

CRUCIFERAE—mustard family

Annual, winter annual, biennial, or perennial plants with various-shaped alternate leaves. Flowers perfect and regular, with 4 sepals, 4 petals, and 6 stamens divided into 2 groups, 4 long and 2 short stamens. Fruit pods (siliques) (Fig. 117) divided into 2 compartments by a thin partition and usually containing many seeds (Fig. 117A). This family contains many troublesome weeds and also many edible plants.

1. Pods compressed at right angles to the central partition. 2
Pods not compressed at right angles to the central partition. 9
2. Plants with basal leaves only; aquatic, usually submerged. *Subularia*
Plants with leafy stems; mostly terrestrial. 3
3. Pods strongly inflated. *Physaria*
Pods not strongly inflated. 4
4. One seed in each compartment of pod. 5
More than one seed in each compartment of pod. 6
5. Pods not winged or notched at top. *Cardaria*
Pods notched at apex, usually winged above. *Lepidium*
6. Pods strongly flattened. 7
Pods not strongly flattened. 8
7. Pods more or less winged, orbicular. *Thlaspi*
Pods not winged, triangular. *Capsella*
8. Stigma sessile; plants small, not fleshy. *Hymenolobus*
Stigma not sessile; plants with fleshy leaves. *Cochlearia*
9. Pods usually not more than 3 times as long as wide. 10




Fig. 117. Pods of the mustard family (Cruciferae): *A*, a dehiscent pod; *B*, small alyssum, *Alyssum alyssoides* L.; *C*, tower mustard, *Arabis glabra* (L.) Bernh.; *D*, false flax, *Camelina sativa* (L.) Crantz; *E*, shepherd's-purse, *Capsella bursa-pastoris* (L.) Medic.; *F*, hare's-ear mustard, *Conringia orientalis* (L.) Dum.; *G*, yellow whitlow-grass, *Draba nemorosa* L. var. *leiocarpa* Lindbl.; *H*, sand bladderpod, *Lesquerella ludoviciana* (Nutt.) Wats. var. *arenosa* (Rich.) Wats.; *I*, ball mustard, *Neslia paniculata* (L.) Desv.; *J*, wild radish, *Raphanus raphanistrum* L.; *K*, wild mustard, *Brassica kaber* (DC.) L. C. Wheeler; *L*, stinkweed, *Thlaspi arvense* L.

Pods usually at least 4 times as long as wide	20
10. Flowers white.	11
Flowers yellow.	14
11. Large plants; basal leaves 40 cm long or longer; pods subglobose.	<i>Armoracia</i>
Plants not with very large basal leaves; pods flattened.	12
12. Seeds many in each cell of pod.	<i>Draba</i>
Seeds few in each cell of pod.	13
13. Pods almost circular, with flat margins.	<i>Alyssum</i>
Pods short, oblong.	<i>Berteroa</i>
14. Pods not opening or breaking off at maturity.	15
Pods opening lengthwise.	16
15. Pods globular, remaining on seed; seeds 1 or rarely 2 per pod.	<i>Neslia</i>
Pods elongate, breaking in two at the nar- rowed middle.	<i>Rapistrum</i>
16. Pods much flattened, at least twice as wide as thick.	<i>Draba</i>
Pods not much flattened, more or less round in cross section.	17
17. Stem leaves bipinnately divided to bipinnatifid.	<i>Descurainia</i>
Stem leaves entire to pinnately lobed or pinnatifid.	18
18. Valves of pods with no nerves.	19
Valves of pods with one nerve.	<i>Camelina</i>
19. Hairs on plant branched or starry; seeds flat.	<i>Lesquerella</i>
Hairs on plant simple, not starry; seeds round.	<i>Rorippa</i>
20. Pods with a long, distinct beak.	21
Pods scarcely beaked.	26
21. Pods constricted between seeds, breaking up.	<i>Raphanus</i>
Pods opening by valves, not constricted between seeds.	22
22. Leaves entire, clasping stem with a cor- date base.	<i>Conringia</i>
Leaves dentate or divided, not entire.	23
23. Beak of pod flattened, the body almost round.	<i>Eruca</i>
Beak of pod and the body equal in cross section.	24

24. Pods flattened; seeds more or less in 2 rows.	<i>Diplotaxis</i>	
Pods circular in cross section; seeds in a single row.		25
25. Beak joint of capsule containing a seed.	<i>Erucastrum</i>	
Beak joint of capsule seedless.	<i>Brassica</i>	
26. Flowers yellow.		27
Flowers white to pinkish to purple, not yellow.		34
27. Stem leaves entire to dentate, not lobed.		28
Stem leaves deeply lobed, lyrate to tripinnate.		31
28. Pubescence consisting of hairs, attached at the middle.	<i>Erysimum</i>	
Pubescence consisting of simple or several-forked hairs.		29
29. Stem leaves narrowly linear, long attenuate at base.	<i>Sisymbrium</i>	
Stem leaves broad, more or less auriculate at base.		30
30. Stem leaves clasping, with the base cordate; plants glabrous; flowers yellowish white.	<i>Conringia</i>	
Stem leaves with a sagittate base; plants pubescent below; flowers greenish to yellowish white.	<i>Arabis</i>	
31. Stem leaves bi- to tri-pinnately divided; pubescence of forked hairs.	<i>Descurainia</i>	
Stem leaves lyrate to pinnatifid; plants with simple hairs or glabrous.		32
32. Pods 4-angled in cross section, with valves of pod nerveless; seeds globose.	<i>Rorippa</i>	
Pods circular or nearly so in cross section.		33
33. Leaf blades pinnatifid; seeds flat.	<i>Barbarea</i>	
Leaf blades entire or merely toothed; seeds plump.	<i>Erysimum</i>	
34. Leaves deeply lobed to dissected.		35
Leaves entire to more or less deeply toothed.		38
35. Plants densely grayish to silvery pubescent.	<i>Smelowskia</i>	
Plants glabrous to more or less pubescent.		36
36. Pods flattened; leaves and stem usually with forked or starry hairs; leaves usually simple and not deeply pinnately lobed.	<i>Arabis</i>	

- Pods cylindric or 4-angled in cross section. 37
37. Pods straight, linear; leaves and stem hairless or with simple hairs; leaves deeply pinnately lobed. *Cardamine*
 Pods more or less falcate, oblong to globular; leaves glabrous or somewhat pubescent. *Rorippa*
38. Flowers 20–25 mm across, purple; plants to 1.5 m high. *Hesperis*
 Flowers much smaller. 39
39. Central partition with an opening sometimes almost reduced to the margin. *Eutrema*
 Central partition entire. 40
40. Plants glabrous, or the pubescence of simple hairs. 41
 Plants with pubescence mostly or entirely of branched hairs. 43
41. Stem leaves cuneate to petiolate at the base. *Cardamine*
 Stem leaves sagittate or cordate at the base. 42
42. Stem leaves cordate at the base, clasping the stem; pods 6–10 cm long. *Conringia*
 Stem leaves sagittate at the base; pods 10–15 mm long. *Thellungiella*
43. Pods strongly flattened. *Arabis*
 Pods round in cross section. 44
44. Flowers purple, 10–20 mm across; pods 6–10 cm long. *Erysimum*
 Flowers white, less than 10 mm across. 45
45. Pods cylindric, swollen at intervals. *Braya*
 Pods not swollen at intervals, the same width over entire length. *Halimolobos*

Alyssum alyssum

Alyssum alyssoides L.

SMALL ALYSSUM

A low annual 10–30 cm high, covered with starry hairs and bearing linear-oblong to spatulate leaves 6–12 mm long. Creamy white flowers about 5 mm across, in narrow racemes. Pods (Fig. 117*B*) yellowish, almost circular, about 3 mm across, having a narrow flat margin around the edges. An introduced weed, not common; in sandy areas and disturbed places; Prairies, southeastern Parklands, southern Rocky Mountains.

Arabis rock cress

Biennial or perennial plants with straight erect stems. Basal leaves usually in a rosette, with stem leaves stalkless and clasping. Flowers usually small and in racemes; fruits long narrow pods containing many seeds.

1. Pods erect or ascending. 2
Pods spreading or reflexed. 7
2. Pods almost circular in cross section. *A. glabra*
Pods flattened parallel to the partition. 3
3. Plant tufted; basal leaves lyrate lobed;
stem leaves not eared or clasping. *A. lyrata* var. *kamchatica*
Plant slightly tufted; basal leaves not
lyrate lobed; stem leaves often clasping
or eared. 4
4. Stem leaves sessile, not auricled or
sagittate. 5
Stem leaves with the base auricled or
sagittate. 6
5. Pods 0.5–1.5 mm wide, 1–3 cm long;
flowers white. *A. nuttallii*
Pods 2–3 mm wide, 2–5 cm long; flowers
pinkish to purple. *A. lyallii*
6. Leaves coarsely hairy; stem leaves eared
at base. *A. hirsuta*
Leaves not coarsely hairy, the hairs, if
any, 2- or 3-forked. *A. drummondii*
7. Pods spreading to reflexed and
pendulous. 8
Pods spreading, never reflexed or
pendulous. 10
8. Pods always sharply reflexed and
pendulous. *A. retrofracta*
Pods in part spreading, in part
descending. 9
9. Pods 2–5 cm long, 2–3.5 mm wide. *A. lemmonii*
Pods 5–10 cm long, 1–2 mm wide. *A. divaricarpa*
10. Plants 10–20 cm high; pods 1–3 cm long,
1.5–2.5 mm wide. *A. arenicola*
Plants 20–80 cm high; pods 5–10 cm long,
1–2 mm wide. *A. divaricarpa*

Arabis arenicola (Rich.) Gelert

Perennial plants with dentate basal leaves; stem leaves oblong, cuneate at the base, slightly fleshy. Arctic and subarctic gravelly or sandy areas. Glabrous plants are var. *arenicola*, those with lower stem leaves and basal leaves pubescent are var. *pubescens* (Wats.) Gelert. Boreal forest.

Arabis divaricarpa A. Nels.

PURPLE ROCK CRESS

A biennial species, usually with a purplish stem 30–50 cm high. Basal leaves 20–40 mm long with starry white hairs; stem leaves stalkless and narrow-lanceolate. Flowers white or pinkish purple, about 6 mm across. Pods spreading or curved slightly downward, 35–60 mm long. Dry slopes, sandy areas; Prairies and Parklands. A form with the pods sickle-shaped and spreading at various angles, var. *dacotica* (Greene) Boiv., more common than the typical variety.

Arabis drummondii A. Gray

DRUMMOND'S ROCK CRESS

A biennial species with a tall, smooth, erect stem 20–70 cm high. Leaves generally smooth, but basal ones sometimes having a few 2-forked hairs. Stem leaves have arrow-shaped clasping bases. Pink or white flowers borne in a terminal raceme. Pods usually fairly erect, 5–8 cm long. Fairly plentiful; on dry hillsides; throughout most of the Prairie Provinces.

Arabis glabra (L.) Bernh.

TOWER MUSTARD

A tall biennial species with erect stems 20–60 cm high; stems quite smooth above but slightly hairy near the base. Basal leaves have short stalks and are 5–15 cm long and slightly hairy. Stem leaves stalkless with an arrow-shaped base. Flowers usually greenish white or yellowish white and quite small. Pods (Fig. 117C) erect and closely pressed against the stem, 5–10 cm long. Not particularly plentiful; in waste places and fields; throughout the Prairie Provinces and quite common in the Cypress Hills.

Arabis hirsuta (L.) Scop.

HIRSUTE ROCK CRESS

An erect simple-stemmed species 30–60 cm high, either smooth or hairy-stemmed, with leaves coarsely hairy, the basal ones forming a rosette, the stem leaves stalkless and clasping. Flowers white or greenish white. The fairly erect pods are 2–5 cm long. Not plentiful; on dry, rocky places; throughout the Prairies and Parklands. A form with larger flowers, about 2 cm across, and only the lower part of the plant very coarsely pubescent, is var. *glabrata* T. & G. Southern Rocky Mountains.

Arabis lemmonii Wats.

Perennial plants with branching caudex, 5–30 cm high; pubescent, with branching hairs. Basal leaves spatulate; stem leaves oblong-lanceolate. The common form, var. *drepanoloba* (Greene) Rollins, usually 20–30 cm high, and with pods 3–5 cm long, 2.5–3.5 mm wide. Not common; alpine areas; southwestern Rocky Mountains.

Arabis lyallii Wats.

Perennial plants with branching caudex, few to several glabrous stems, 5–25 cm high. Basal leaves oblanceolate, glabrous or pubescent. Alpine areas; Rocky Mountains.

Arabis lyrata L. var. *kamchatica* Fisch.

LYRE-LEAVED ROCK CRESS

A native, tufted perennial or biennial plant 10–30 cm high, with lyrate lobed basal leaves and linear to spatulate stem leaves. Flowers usually white

but occasionally pinkish. Found sparingly; on sandy soils in lightly forested areas; Boreal forest.

Arabis nuttallii Robins.

Perennial plants with several to many stems from a branching caudex, 5–30 cm high, glabrous above, usually hirsute below. Alpine areas; southern Rocky Mountains.

Arabis retrofracta Graham

REFLEXED ROCK CRESS

A biennial or perennial 10–60 cm high, with one or several stems from the base, usually with appressed hairs. Leaves usually covered with fine hairs, the lower leaves forming a rosette and the stem leaves often clasping the stem. Flowers purplish pink to white and borne in a terminal raceme. Pods usually straight, 2–6 cm long. Common; dry slopes, open areas; Prairies and Parklands. The var. *collinsii* (Fern.) Boiv. is similar, but has long spreading hairs at the base of the stem instead of appressed hairs as in the typical variety; similar locations. The var. *multicaulis* Boiv. is many-stemmed and has basal leaves about twice as long as the stem leaves; pods somewhat longer and wider. Western Parklands, Rocky Mountains.

Armoracia horse-radish

Armoracia rusticana P. Gaertner, B. Meyer & Scherb.

HORSE-RADISH

Perennial plants with stems 20–100 cm high, glabrous, much-branched. Basal leaves 10–40 cm long, crenate. Flowers white, 10–15 mm across. Introduced; cultivated for its roots for making table relish; occasionally escaped and established along roadsides and in moist areas.

Barbarea winter cress

Barbarea orthoceras Ledeb.

AMERICAN WINTER CRESS

A smooth-stemmed biennial 20–50 cm high. Leaves alternate and usually lyrate with one or two pairs of narrow lobes and a large terminal lobe on each leaf. Streams and wet places; in the Cypress Hills, Rocky Mountains, Boreal forest. Plants with large flowers 10–15 mm across and the beak of the pod 2–3 mm long are *B. vulgaris* R. Br., an introduced species.

Berteroa hoary-alyssum

Berteroa incana (L.) DC.

HOARY-ALYSSUM

A starry-haired plant 30–60 cm high, with numerous lanceolate leaves 10–25 mm long. Flowers about 3 mm across with white deeply notched petals. Pods starry-hairy, broadly oval, about 6 mm long. An introduced weed reported at various localities across the Prairies and Parklands.

Brassica mustard

1. Pods with a large flat or angled beak, often containing a seed. 2

- Pods with a slender round or conical beak
without seeds. 3
2. Pods densely pubescent; the beak usually
longer than the body. *B. hirta*
Pods glabrous; the beak half as long as
the body. *B. kaber*
3. Upper stem leaves cordate and clasping. *B. campestris*
Upper stem leaves cuneate at the base,
not clasping. *B. juncea*

Brassica campestris L.

BIRD-RAPE

Annual or biennial plants 40–80 cm high. Inflorescence not elongating, the opened flowers overtopping the buds; pods 3–7 cm long, with seeds reddish brown to blackish. Introduced; occasionally grown for seed; weedy in several areas. Plants in which the inflorescence elongates have been distinguished as *B. napus* L.

Brassica hirta Moench

WHITE MUSTARD

Annual plants 30–70 cm high, more or less pubescent. Flowers about 2 cm across; pods about 3 cm long, spreading, hirsute, with seeds yellowish to pale brown. Occasionally grown for seed; rare as a weed.

Brassica juncea (L.) Cosson

INDIAN MUSTARD

An annual erect, weedy plant 30–100 cm high, with almost hairless smooth stems and leaves. Lower leaves lyrate-pinnatifid with a large end lobe, stalked, 10–15 cm long; upper leaves slightly stalked and often not lobed, much smaller. Flowers 10–15 mm across, yellow; pods 2–5 cm long, with a conic beak. Introduced, common; in grainfields and waste places; throughout the Prairies and Parklands.

Brassica kaber (DC.) L. C. Wheeler

WILD MUSTARD

A more or less hirsute annual 20–80 cm high; the lower leaves usually lyrate-pinnatifid, with the terminal lobe large, rounded. Flowers 10–15 mm across; pods (Fig. 117K) 2–5 cm long, with the beak half as long as the body and seeds black or brown. Introduced weed; in waste places and disturbed areas.

Braya

Braya humilis (C. A. Mey.) Robins.

Perennial plants with several stems arising from the caudex, 5–30 cm high, more or less pubescent. Basal leaves linear oblanceolate, 1–3 cm long, thickish, entire or somewhat toothed; stem leaves smaller, distant. Flowers white or purplish-tinged, 6–10 mm across; pods 1–3 cm long, cylindrical, pubescent, swollen at intervals. Gravelly or sandy areas; Boreal forest, Rocky Mountains. A very variable species, in which several varieties have been described that overlap in characters.

Camelina false flax

Introduced annual or winter-annual weeds, with small yellow flowers and long-stalked obovoid (or pear-shaped) pods containing numerous seeds, which have a very short dormant period.

1. Stems and leaves hairy; lower leaves not stalked. *C. microcarpa*
Stems and leaves smooth or slightly hairy; lower leaves with stalks. 2
2. Lower leaves entire. *C. sativa*
Lower leaves dentate or lobed. *C. dentata*

Camelina dentata Pers.

FLAT-SEEDED FALSE FLAX

Stem erect and smooth, 30–50 cm high. Lower leaves lobed, with winged stalks, 35–70 mm long; upper leaves smaller and entire. Flowers yellow; pod obovoid, 5–10 mm long. An introduced weed; sometimes found in grainfields and waste places.

Camelina microcarpa Andrz.

SMALL-SEEDED FALSE FLAX

A densely hairy plant 30–50 cm high. Leaves stalkless, with simple starry hairs. Flowers yellow; pods ovoid, to 5 mm long. An introduced weed, fairly common; in grainfields and waste places.

Camelina sativa (L.) Crantz

FALSE FLAX

An introduced annual or winter annual 30–100 cm high, with a few hairs on the lower leaves and the lower part of the stem. Upper leaves stalkless and clasping; lower ones tapering to a winged stalk. Flowers yellow, about 3 mm across; pods (Fig. 117D) 5–10 mm long, ovoid, containing about 10 seeds. Introduced from Europe, found occasionally; a weed in grainfields and waste places.

Capsella shepherd's-purse

Capsella bursa-pastoris (L.) Medic. (Fig. 118)

SHEPHERD'S-PURSE

An introduced annual or winter-annual plant, with branched stems 15–50 cm high. Basal leaves forming a rosette and often deeply cut and lobed; stem leaves usually clasping with ears at the base. Flowers in terminal racemes, small and white. The seedpods (Fig. 117E) are characteristic of this species; they resemble an inverted triangle, the flat base uppermost and notched, each pod containing about 20 seeds. An introduced weed, common; in gardens and waste places; throughout the Prairie Provinces.

Cardamine bitter cress

1. Stem leaves entire or dentate, not lobed or divided. 2
Stem leaves deeply divided to pinnate. 3
2. Stem leaves entire. *C. bellidifolia*
Stem leaves coarsely dentate. *C. bulbosa*



Fig. 118. Shepherd's-purse, *Capsella bursa-pastoris* (L.) Medic.

3. Flowers 15–25 mm across. *C. pratensis*
 Flowers 3–10 mm across. 4
4. Plants perennial, with rootstocks;
 inflorescence corymbose, appearing
 umbellate; flowers 5–10 mm across. *C. umbellata*
 Plants annual or perennial, without root-
 stocks; inflorescence elongate; flowers
 3–5 mm across. 5
5. Stems mostly pubescent at the base; ter-
 minal leaflets obovate. *C. pensylvanica*
 Stems glabrous throughout; terminal
 leaflets linear-oblongate. *C. parviflora*

Cardamine bellidifolia L.

ALPINE BITTER CRESS

Perennial plants with branched caudex; stems 3–10 cm high, glabrous. Flowers 1–5, white. Rare; alpine areas; southern Rocky Mountains.

Cardamine bulbosa (Schreb.) BSP.

SPRING CRESS

Perennial plants with a fleshy bulb; stem solitary, 15–30 cm high; leaves mostly coarsely toothed. Rare; wet areas; southeastern Boreal forest.

Cardamine parviflora L.

SMALL BITTER CRESS

Annual or biennial plants; stems usually solitary, 10–30 cm high, simple to branched, glabrous. Flowers about 4 mm across. Rare; dry, usually rocky areas; southeastern Boreal forest, Rocky Mountains.

Cardamine pensylvanica Muhl.

BITTER CRESS

A biennial or short-lived perennial plant 10–50 cm high. Leaves very deeply pinnately lobed with 2–8 pairs of lateral segments and a larger terminal segment. Stem usually bearing a few scattered hairs. Flowers small and white. Fruits linear pods 10–30 mm long. Uncommon; in wet or very moist places; Boreal forest, Rocky Mountains, Cypress Hills. Syn.: *C. scutata* Thunb.

Cardamine pratensis L.

MEADOW BITTER CRESS

A hairless perennial 20–50 cm high, with deeply pinnately lobed leaves and white or pink flowers 15–25 mm across. Rare; a plant of bogs and swampy places; Boreal forest, Peace River. Mostly represented by var. *angustifolia* Hook., with stem leaves narrow, not decurrent.

Cardamine umbellata Greene

MOUNTAIN CRESS

Perennial with rootstocks; stems erect, 20–50 cm high, glabrous or nearly so. Flowers 5–10 mm across; pods crowded, 2–2.5 cm long. Rare; wet areas; southern Rocky Mountains.

Cardaria hoary cress

Introduced perennial weeds spreading by seeds and by running roots penetrating deeply into the soil. Stem leaves alternate and clasping; basal ones

with a short stalk. Flowers white in a close raceme. Seedpods varying in shape, a character useful in distinguishing the species.

1. Pods with downy hairs, globular and inflated. *C. pubescens* var. *elongata*
Pods smooth, flattened, not hairy. 2
2. Pods heart-shaped at base, not inflated. *C. draba*
Pods somewhat inflated, not heart-shaped at base. *C. draba* var. *repens*

Cardaria draba (L.) Desv.

HEART-PODDED HOARY CRESS

Perennial from deep running roots, with dense heads of white flowers and heart-shaped smooth seedpods. Introduced from Europe, found occasionally; in gardens and shelterbelts; throughout the Prairie Provinces.

The var. *repens* (Schrenk) O. E. Schulz, lens-podded hoary cress, is similar to the species, but with seedpods slightly inflated and almost round instead of heart-shaped. Introduced, probably from Afghanistan; found in the same locations as the species.

Cardaria pubescens (Meyer) Rollins var. *elongata* Rollins

GLOBE-PODDED HOARY CRESS

Plants similar to *C. draba*, but with leaves more hoary in appearance and seedpods globular, inflated with fine, downy hairs. Apparently introduced from Asia, becoming quite common; in gardens and shelterbelts and sometimes in fields and roadsides; southern part of the Prairie Provinces.

Cochlearia scurvy-grass

Cochlearia officinalis L.

SCURVY-GRASS

Annual plants of very variable size, 1–40 cm high; basal leaves reniform, usually fleshy, glabrous. Flowers white, with petals 3–7 mm long; silicle ovoid to globose, 4–7 mm long. Shores and beaches. Rare; Hudson Bay.

Conringia hare's-ear mustard

Conringia orientalis (L.) Dum.

HARE'S-EAR MUSTARD

An introduced annual or winter-annual weed with taproots and a perfectly smooth stem 15–60 cm high. Seedling leaves round, on stalks, but subsequent leaves clasping and elliptical, entire, eared at the base, quite smooth, and the whole plant covered with a bluish bloom. Flowers creamy white, about 6 mm across. Pods (Fig. 117F) narrow, 4-angled, erect, 7–10 cm long. Common; roadside and field weed.

Descurainia tansy mustard

Annuals or biennials from taproots, usually with much-dissected leaves, yellow flowers, and long narrow seedpods.

1. Both upper and lower leaves 2 or 3 times divided. 3
Upper leaves simply pinnate. 2
2. Seeds in 1 row in each compartment of pod; pods erect and closely pressed to the stem; plant grayish. *D. richardsonii*
Seeds in 2 rows in each compartment of pod; pods divergent from stem. *D. pinnata* var. *brachycarpa*
3. Pubescence of simple hairs; plants glandular. *D. sophioides*
Pubescence of mostly stellate hairs; plants not glandular. *D. sophia*

Descurainia pinnata (Walt.) Britt. var. *brachycarpa* (Richards.) Fern.

SHORT-FRUITED TANSY MUSTARD

Annual plants 20–60 cm high, erect, with a hairy and glandular stem. Leaves dark green; the upper ones pinnate and the lower ones often 2 or 3 times divided, glandular. Flowers yellow. Pods somewhat club-shaped, 5–12 mm long, on stalks 8–15 mm long. Not very common; has been found in southern part of the Prairie Provinces. Syn.: *Sophia brachycarpa* (Richards.) Rydb.

Descurainia richardsonii (Sweet) O. E. Schulz

GRAY TANSY MUSTARD

Tall, erect biennial plants 30–90 cm high, with pinnate leaves, sometimes doubly pinnate. Whole plant covered with very fine hairs, giving it a grayish appearance. Flowers small, pale yellow. Pods small, 5–10 mm long, tightly compressed against stem, and crowded. Common; in fields and waste places; throughout the Prairie Provinces. Syn.: *Sophia richardsonii* (Sweet) Rydb.

Descurainia sophia (L.) Webb.

FLIXWEED

An annual or biennial, branched, with all leaves 2 or 3 times dissected, and with some star-like hairs. Flowers yellow. Pods linear, 10–30 mm long, on stalks 6–12 mm long, and borne at an angle from the stem. An introduced weed, which has become very common; in fields, waste places, and even on prairie land; one of the commonest weeds in Western Canada. Syn.: *Sophia multifida* Gilib.

Descurainia sophioides (Fisch.) O. E. Schulz

NORTHERN FLIXWEED

Much like *D. sophia*, but brighter green and glandular. Inflorescence tardily elongating. Pods sickle-shaped, irregularly spreading to erect. Dry gravel beds; Hudson Bay. Syn.: *Sisymbrium sophioides* Fisch.

Diplotaxis sand-rocket

Diplotaxis muralis (L.) DC.

SAND-ROCKET

Annual plants, branched from base, 30–60 cm high. Stems leafy only near the base, with oblanceolate lobed leaves 5–10 cm long, usually with a slender stalk. Flowers yellow. Pods 20–25 mm long with a short beak, on stalks 10–15 mm long. Introduced from Europe, but occasionally found; in waste places; throughout the Prairies and Parklands.

Draba whitlow-grass

Low tufted plants, usually with a rosette of basal leaves, scapose or with a few stem leaves. Flowers small, white or yellow, usually in racemes. Pods (Fig. 117G) either oval or linear and flat, with seeds in 2 rows in each cell.

1. Flowering stems with 2 or more leaves. 2
Flowering stems leafless, or at most with a reduced leaf. 12
2. Flowers white. 3
Flowers yellow. 10
3. Stem leaves all or mostly opposite. *D. reptans* var. *micrantha*
Stem leaves alternate. 4
4. Stem leaves numerous, usually 6–25. *D. incana*
Stem leaves usually not more than 3–8. 5
5. Pods glabrous. 6
Pods pubescent. 7
6. Basal leaves 1–2 cm long, narrowly lanceolate; pedicels ascending. *D. hirta*
Basal leaves 1–4 cm long, obovate to oblanceolate; pedicels spreading. *D. stenoloba*
7. Stems often with only 1 leaf; leaves ciliate toward the base. *D. cinerea*
Stems usually with 3–5 leaves; leaves not ciliate. 8
8. Pubescence, including that of the pods, of stellate hairs. *D. lanceolata*
Pubescence of the pods of simple or 2-forked hairs. 9
9. Pods mostly 10–15 mm long, linear, about 2 mm wide. *D. praealta*
Pods mostly 7–10 mm long, lanceolate, about 3 mm wide. *D. mccallae*
10. Stem leaves usually numerous, often 10–15. *D. aurea*
Stem leaves usually 1–5. 11
11. Plants annual; pedicels 1.5–4 times as long as the pod. *D. nemorosa*
Plants biennial or perennial; pedicels half as long as the pod. *D. stenoloba*
12. Plants annual; inflorescence short, with stems flexuous; pods 10–15 mm long. *D. reptans* var. *micrantha*
Plants biennial or perennial; inflorescence elongate; pods usually less than 10 mm long. 13
13. Flowers white. 14
Flowers yellow. 17

14. Leaves ciliate, at least toward the base. 15
 Leaves densely stellate throughout, not
 ciliate. *D. nivalis*
15. Plants densely stellate, pubescent
 throughout. *D. cinerea*
 Plants not densely pubescent. 16
16. Basal leaves rather thick, oblong-spatu-
 late, subglabrous. *D. crassifolia*
 Basal leaves oblong-obovate, usually
 sparsely pubescent, not thick. *D. fladnizensis*
17. Basal leaves glabrous, but ciliate toward
 the tip. *D. crassifolia*
 Basal leaves more or less pubescent, at
 least below. 18
18. Stems glabrous; leaves very narrow,
 mostly less than 1 mm,
 stellate-pubescent. *D. oligosperma*
 Stems pubescent; leaves mostly 1 mm
 wide or wider. 19
19. Leaf margins conspicuously long-ciliate,
 pubescent with long tangled hairs. *D. stenopetala*
 Leaf margins not long-ciliate. 20
20. Pubescence of simple or forked hairs. *D. alpina*
 Pubescence mostly of stellate hairs. *D. incerta*

Draba alpina L.

ALPINE WHITLOW-GRASS

Plants with a dense rosette, the leaves elliptic-lanceolate, densely pubescent, entire. Scape to 20 cm high, slightly elongating in fruit. Flowers 6–10 mm across, bright yellow; pods glabrous. Open places in rocky arctic tundra; Boreal forest, Manitoba.

Draba aurea Vahl

GOLDEN WHITLOW-GRASS

Plants with basal rosettes and erect or decumbent leafy stems. Stem leaves entire or somewhat dentate. Flowers 8–12 mm across, yellow; pods 8–16 mm long, puberulent, often twisted. Open forest and alpine meadows; Rocky Mountains. Plants with glabrous pods are var. *leiocarpa* (Payson & St. John) C. L. Hitchc.

Draba cinerea Adams

Plants densely stellate-pubescent in all parts; stems 5–25 cm high, with 1–4 stem leaves. Flowers 6–10 mm across; pods 4–8 mm long, inflated. Sandy or gravelly areas; southern Rocky Mountains.

Draba crassifolia R. C. Graham

THICK-LEAVED WHITLOW-GRASS

Dwarf perennial plants with dense rosettes; leaves rather thick, sparsely pubescent. Stems usually leafless, 5–15 cm high, glabrous. Flowers 4–6 mm across, with the petals little larger than the sepals; pods 4–7 mm long, glabrous. Alpine meadows and scree fields; southern Rocky Mountains.

Draba fladnizensis Wulf.

Small, often dwarf perennials, with stems rarely more than 10 cm high. Stem leaves 0–2; basal leaves oblong-obovate, glabrous except the ciliate margins. Inflorescence 2- to 10-flowered; flowers 5–8 mm across; pods 5–7 mm long. Arctic and alpine, usually stony areas; Boreal forest, southern Rocky Mountains.

Draba hirta L.

HAIRY WHITLOW-GRASS

Plants with stems to 25 cm high, and 1–4 stem leaves, usually unbranched. Basal leaves to 2 cm long, narrowly lanceolate, subentire, ciliate, and more or less densely stellate-pubescent. Flowers 6–10 mm across, creamy white or light yellow; pods 6–12 mm long, glabrous, often somewhat twisted. Gravelly shores and riverbanks; Boreal forest, Rocky Mountains.

Draba incana L.

Plants biennial or short-lived perennial; stems to 35 cm high, erect, simple or branched, with 6–25 stem leaves. Basal leaves 1–2.5 cm long, pubescent with simple or branched hairs. Stem, stem leaves, and inflorescence stellate-pubescent. Flowers 6–10 mm across, white; pods 6–10 mm long, often somewhat twisted. Dry gravel beds; Boreal forest, Manitoba.

Draba incerta Payson

Loosely tufted perennial; scapes 2–10 cm high, often with a reduced leaf below the first flower, pubescent with branched hairs. Leaves linear-oblong-ovate, stellate-pubescent below. Flowers 8–10 mm across, yellow; pods 5–10 mm long, pubescent. Alpine meadows and slopes; Rocky Mountains.

Draba lanceolata Royle

Plants perennial, with leafy simple or branched stems 5–25 cm high, pubescent; basal leaves 1–3 cm long, stem leaves 5–25 mm long, dentate, pubescent with stellate or bushy hairs. Flowers 6–10 mm across, white; pods 4–12 mm long, often twisted, pubescent. Meadows and slopes; Rocky Mountains.

Draba mcallae Rydb.

Plants to 25 cm high, with 3–5 or sometimes more stem leaves. Stem long pilose below; leaves stellate pubescent. Flowers 6–10 mm across; pods 7–10 mm long, densely puberulent. Meadows, slopes, and gravel banks; southern Rocky Mountains.

Draba nemorosa L.

YELLOW WHITLOW-GRASS

A low plant, 10–30 cm high, branching from the rosette of basal leaves. Flowers pale yellow, occasionally fading to white; pods 6–8 mm long, on oval stalks 6–20 mm long. Fairly common; throughout southern part of the Prairie Provinces. The var. *leiocarpa* Lindbl. differing from the type by having smooth pods (Fig. 117G) and being probably more plentiful, especially in the northern and eastern parts. Both the species and the variety flowering quite early in the spring. Syn.: *D. lutea* Gilib.

Draba nivalis Liljeb.

Perennial plants with rosettes on the caudex branches; leaves densely stellate-pubescent, with branched hairs mixed in. Stems 2–10 cm high, leafless or with a single reduced leaf. Flowers 5–10 mm across, white. In var. *nivalis*, the pods less than 1 cm long, flat to slightly twisted; rock outcrops; Boreal forest and southern Rocky Mountains; in var. *elongata* Wats., the pods 10–15 mm long, and conspicuously twisted; alpine areas; southern Rocky Mountains.

Draba oligosperma Hook.

FEW-SEEDED WHITLOW-GRASS

Small perennials with rosettes on the caudex branches. Scapes 1–10 cm high; flowers 6–10 mm across, yellow; pods 3–5 mm long, ovate, pubescent to glabrous. Rocky alpine areas; southern Rocky Mountains.

Draba praealta Greene

Biennial or perennial plants with stems 10–30 cm high, and compact rosettes. Basal leaves 1–3 cm long, entire or somewhat dentate, densely pubescent with branched hairs. Stem leaves 2–6, densely pubescent. Flowers 4–8 mm across, white; pods 8–15 mm long, soft pubescent. Slopes and rocky areas; Rocky Mountains.

Draba reptans (Lam.) Fern. var. *micrantha* (Nutt.) Fern.

CREeping WHITLOW-GRASS

A small plant 2–15 cm high, with a leafy stem. Leaves usually 8–20 mm long. White flowers, 4–6 mm across, but often without petals; pods linear, 8–12 mm long, clustered near the top of the stem, usually covered with fine, stiff hairs. Flowering in early June. Not common; has been found in disturbed sandy areas; Prairies.

Draba stenoloba Ledeb.

Biennial or perennial plants 5–30 cm high, with stems arising from a simple caudex, glabrous above, somewhat pubescent below, with 1–4 pubescent leaves. Basal leaves 1–4 cm long, denticulate, stiffly pubescent. Flowers 6–10 mm across, yellow; pods 8–20 mm long. Alpine slopes; southern Rocky Mountains. Plants with the pubescence mostly of simple hairs and the stem coarsely pilose are var. *nana* (Schulz) C. L. Hitchc.; southern Rocky Mountains.

Draba stenopetala Trautv.

Perennial mat-forming plants with leafless stems 1–6 cm high; leaves crowded, linear to linear-oblongate, the margins ciliate with simple or branched hairs, pubescent below with simple or branched hairs. Flowers 5–9 mm across, light yellow; pods 3–5 mm long, ovate, pubescent to subglabrous. Alpine slopes; southern Rocky Mountains. Syn.: *D. paysonii* Macbr. var. *treleasii* (Schulz) C. L. Hitchc.

Eruca rocket

Eruca sativa L.

GARDEN-ROCKET

Annual plants, with stems 20–100 cm high; leaves lyrate-pinnatifid, with the terminal lobe large, and 2–5 pairs of lateral lobes. Flowers 30–40 mm

across, whitish or yellowish, purplish-veined; pods 12–25 mm long. Introduced; occasionally escaped from cultivation.

Erucastrum dog mustard

Erucastrum gallicum (Willd.) O. E. Schulz DOG MUSTARD

An annual plant 20–50 cm high, erect, with stem hairs pointing downward. Leaves deeply cut, often to the midrib, making them appear pinnate, and often deeply lobed. Leaves varying to 25 cm long. Flowers pale yellow; pods 25–40 mm long, linear, circular in cross section, on a short stalk tipped with a slender style, with one row of seeds in each compartment. An introduced weed; in grainfields; throughout Prairies and Parklands.

Erysimum treacle mustard

Annual or biennial plants with leafy stems and 2-branched hairs, yellow flowers and linear 4-angled seedpods with seeds in one row in each compartment.

- 1. Flowers large, 2–4 cm across, purple. *E. pallassii*
Flowers smaller, yellow. 2
- 2. Petals longer than 10 mm. *E. asperum*
Petals shorter than 10 mm. 2
- 3. Stalks of seedpods almost half as long as
pods. *E. cheiranthoides*
Stalks of seedpods not one-quarter as
long as pods. 4
- 4. Pubescence mostly or entirely of 2-
branched hairs. *E. inconspicuum*
Pubescence mostly or entirely of stellate
hairs. *E. hieracifolium*

Erysimum asperum (Nutt.) DC. (Fig. 119) WESTERN WALLFLOWER

A rough, usually erect plant 20–60 cm high, often much branched and coarse, with appressed small white hairs. Flowers pale yellow, 10–15 mm across; pods 3–10 cm long, narrow, rough, and 4-angled, with a short, thick style at the end, and a very short, stout stalk. Spreading out considerably from the stem in all directions. Common; on light and sandy soil; in Prairies, rare in Parklands. Syn.: *Cheirinia aspera* (Nutt.) Rydb.

Erysimum cheiranthoides L. WORMSEED MUSTARD

An erect plant 20–60 cm high, with lanceolate or oblong-lanceolate dark green leaves 2–10 cm long. Flowers yellow and small, about 5 mm across, in dense terminal clusters 2–3 cm across. Seedpods linear, 10–25 mm long, on slender stalks 6–12 mm long. Not common; a weedy plant found in moist places and fields; throughout the Prairie Provinces. Syn.: *Cheirinia cheiranthoides* (L.) Link.



HERBARIUM OF THE RESEARCH STATION
SWEET CURENT, I.A.K.

SPECIES: *Erysimum asperum* (Nutt.) DC.

LOCATION: Near Medakow

COLL: J. Looma:

DET:

NUMBER: 24043

DATE: 8 June 1970

ELEVATION: 660 m

HABITAT: Sandstone



Fig. 119. Western wallflower, *Erysimum asperum* (Nutt.) DC.

Erysimum hieracifolium L.

GRAY ROCKET

Biennial or perennial plants 30–80 cm high; leaves linear or oblong, wavy-dentate, greenish gray. Flowers 15–20 mm across; pods 30–55 mm long, 4-angled. Rare, introduced weed; southeastern Parklands.

Erysimum inconspicuum (S. Wats.) MacM. SMALL-FLOWERED PRAIRIE-ROCKET

An erect grayish green perennial 20–50 cm high, and not much branched. Leaves narrow, 2–8 cm long, the lower ones having stalks but the upper ones stalkless. Yellow flowers about 6 mm across, on the top of the stem. Linear seedpods 2–6 cm long on a very short stalk and borne erect on the stem. Common; on dry and sandy prairie; throughout Prairies and Parklands. Syn.: *E. parviflorum* Nutt.

Erysimum pallassii (Pursh) Fern.

PURPLE ROCKET

Biennial plants 10–20 cm high, with numerous linear leaves. Pods 6–10 cm long, ascending. Shale and gravel slides; southern Rocky Mountains.

Eutrema

Eutrema edwardsii R. Br.

Rhizomatous perennial to 40 cm high, with rather fleshy ovate to orbicular leaves, truncate to cordate at the base. Flowers 5–8 mm across; pods 6–20 mm long, linear to oblong, erect or appressed. Arctic and alpine tundra; Boreal forest, southern Rocky Mountains.

Halimolobos

Halimolobos virgata (Nutt.) O. E. Schulz

Perennial plants 15–40 cm high, much resembling *Arabis* ssp., densely pubescent with branched and stellate hairs. Flowers 4–8 mm across, white; pods 1–3 cm long, round, slightly constricted between the seeds, strongly ascending. Dry grasslands; western Prairies.

Hesperis rocket

Hesperis matronalis L.

DAME'S-ROCKET

A tall perennial species with erect stems 30–100 cm high, simple or branched above. Leaves simple oblong- to ovate-lanceolate, dentate, pubescent on both sides. Flowers 3–5 cm across, purple, fragrant; pods 5–10 cm long, linear, ascending or spreading. Introduced ornamental, occasionally escaped and established; in tree groves or shrubbery.

Hymenolobus

Hymenolobus procumbens (L.) Nutt.

Small annual or biennial plants 3–30 cm high, with stems procumbent or erect. Lower leaves entire to deeply lyrate-pinnatifid; stem leaves entire, pubescent throughout with sparse simple hairs. Inflorescence many-flowered,

the flowers 3–6 mm across; petals white, spatulate, the same length as or somewhat exceeding the sepals; pods 1.5–5 mm long, elliptic to obovate, with the valves translucent. Very rare; wet, often saline areas; Hudson Bay, Prairies. Syn.: *Hutchinsia procumbens* (L.) Desv.

Lepidium pepper-grass

Annual or perennial herbs, glabrous or pubescent with simple hairs. Inflorescence in dense, bractless racemes; flowers small, usually with whitish petals, or apetalous. Fruit a silicle, strongly flattened at right angles to the septum, orbicular to ovate or obovate; seeds one in each locule. Weedy plants of roadsides, waste places, and disturbed areas.

1. Upper leaves with a sagittate or cordate-clasping base. 2
 Upper leaves tapering at the base. 3
2. Auricles of the leaf base acute, much shorter than the leaf blade; lower leaves similar to the upper leaves. *L. campestre*
 Auricles of the leaf base rounded, about the same length as the leaf blade; lower leaves pinnate. *L. perfoliatum*
3. Silicle not notched at the apex; perennial plant. *L. latifolium*
 Silicle notched at the apex; annual plant. 4
4. Silicle 5–7 mm long, distinctly winged; stamens 6; petals 2 mm long. *L. sativum*
 Silicle 2–4 mm long, not winged; stamens 2 or 4; petals minute or none. 5
5. Plants with short axillary racemes as well as terminal ones; petals linear, shorter than the sepals. *L. ramosissimum*
 Plants with elongated terminal racemes; petals none or rudimentary. 6
6. Silicles oval or elliptic-ovate, gradually narrowed toward the apical teeth; basal leaves bipinnatifid. *L. ruderale*
 Silicles round-cordate or round-obovate, rounded at the apex; basal leaves at most pinnatifid. *L. densiflorum*

Lepidium campestre (L.) R. Br.

PEPPERWORT

An annual or biennial species, with simple or branched stems 20–50 cm high, grayish green, densely pubescent. Basal leaves entire or lyrate; lower stem leaves petiolate, the upper ones clasping the stem with sagittately lobed leaves. Flowers about 3 mm across, white; silicle about 5 mm long. Rare; introduced weed, roadsides, railway grades; western Prairies.

Lepidium densiflorum Schrad.

COMMON PEPPER-GRASS

An annual or winter annual with erect stem, much branched above, 15–60 cm high. Stem leaves lanceolate, with a few coarse teeth; basal leaves often deeply incised and divided. Flowers minute, with petals missing or very rudimentary. Seedpods borne on short stalks and very numerous on the stem, the spike of pods often 10–15 cm long; each pod about 2–3 mm wide, heart-shaped with a notch at the top, and containing a single seed in each compartment. Common; in fields, waste places, and roadsides; throughout the Prairie Provinces. Syn.: *L. apetalum* A. Gray. Plants with flattened pedicels and some axillary racemes are var. *bourgeauanum* (Thell.) C. L. Hitchc.

Lepidium latifolium L.

BROAD-LEAVED PEPPER-GRASS

Tall plants 50–100 cm high, with a thick, branching rootstock, glabrous or nearly so. Basal leaves petiolate, to 30 cm long; stem leaves sessile, much reduced. Racemes numerous in a large panicle; flowers about 3 cm across, with the sepals white-margined. Very rare; introduced weed; southwestern Prairies.

Lepidium perfoliatum L.

PERFOLIATE PEPPER-GRASS

An annual weed with a stem much branched and bushy above, 20–50 cm high. Basal leaves pale green, very finely dissected and divided; subsequent leaves more entire, the upper ones entire with a pointed end and a cordate to entirely clasping base. Flower terminal in racemes and very small with yellow petals. Numerous seedpods about 3 mm long, oval, and containing a seed in each compartment. When growing, the plants have a reddish tinge to the leaves, which, with the yellow petals, makes them easy to notice. An introduced weed, common in dry areas of western USA but rare in Canada; western Prairies.

Lepidium ramosissimum A. Nels.

BRANCHED PEPPER-GRASS

An annual or biennial, very similar to *L. densiflorum*, but branched diffusely from the base, and with white petals shorter than the sepals. Seedpods rather pointed at the notched apex, whereas in *L. densiflorum* the pods are more rounded. Probably fairly common throughout southern part of area, but distinction between the two species is obscure.

Lepidium ruderales L.

ROADSIDE PEPPER-GRASS

Annual or biennial with a single erect or ascending stem, 10–30 cm high, sparsely pubescent. Basal leaves 5–7 cm long, long-petioled, pinnatifid to bipinnatifid. Flowers about 1.5 mm across; silicle 2–2.5 mm long, deeply notched. Introduced weed, rare; southeastern Parklands.

Lepidium sativum L.

GARDEN CRESS

Annual with a single erect stem, glabrous. Basal leaves long-petioled, lyrate, with the lobes toothed; upper stem leaves linear, entire. Flowers 3–5 mm across; silicle 5–6 mm long, deeply notched. Introduced, occasionally escaped from cultivation.

Lesquerella bladderpod

Low tufted annual or perennial plants with clustered basal leaves and a few stem leaves. Plants with starry hairs. Flowers perfect, in racemes, yellow or purple. Seedpods (Fig. 117H) inflated, globose or oval, with 2 to many seeds in each compartment.

1. Pods ovate; plants densely tufted; flowers bright yellow. *L. alpina* var. *spathulata*
Pods globose; plants not densely tufted; flowers reddish yellow. 2
2. Pedicels ascending in fruit; pods glabrous or nearly so. *L. arctica*
Pedicels recurved in fruit; pods densely pubescent. *L. ludoviciana* var. *arenosa*

Lesquerella alpina (Nutt.) S. Wats. var. *spathulata* (Rydb.) Payson

SPATULATE BLADDERPOD

A deep-rooted tufted perennial 5–15 cm high, with several stems. Numerous basal leaves 1–3 cm long; very few linear stem leaves. Flowers yellow, about 6 mm across; pods ovoid, very slightly compressed. Not common; dry hillsides and badlands; Prairies, southern Rocky Mountains.

Lesquerella arctica (Wormsk.) Wats.

NORTHERN BLADDERPOD

Small plants with spatulate or oblanceolate basal leaves. Flowers long-pedicellate, yellow; pods on straight pedicels, 1–2 cm long. Gravelly areas; Hudson Bay. Plants with the pods scurfy and few white stellate hairs are var. *purshii* Wats.; southern Rocky Mountains.

Lesquerella ludoviciana (Nutt.) Wats. var. *arenosa* (Rich.) Wats. (Fig. 120)

SAND BLADDERPOD

A very slender stemmed spreading and decumbent plant 5–30 cm across, with mostly basal linear-oblanceolate leaves. Flowers dull yellow or reddish-tinged, about 6 mm across. Pods globose (Fig. 117H), 3–5 mm across, on curved stalks. Common; on dry hillsides and prairie; throughout the southern part of the Prairie Provinces, not necessarily on sandy soil, specimens having been collected on Regina Plains.

Neslia ball mustard

Neslia paniculata (L.) Desv.

BALL MUSTARD

A tall erect annual or winter annual 30–60 cm high, on much-branched stems. The whole plant is pale yellowish green, with starry hairs. Lower leaves somewhat stalked and lanceolate; stem leaves arrow-shaped and clasping the stems at the base. Flowers small, orange yellow, about 3 mm across, and clustered on the ends of the stems. Fruit pods (Fig. 117I) in long racemes, small, round, about 2 mm long. Pods usually contain only 1 seed and remain on the seed when ripe. Introduced from Europe, a weed of crop lands and waste places; throughout the Prairie Provinces.



Fig. 120. Sand bladderpod, *Lesquerella ludoviciana* (Nutt.) Wats. var. *arenosa* (Rich.) Wats.

Physaria bladderpod

Physaria didymocarpa (Hook.) Gray

TWIN BLADDERPOD

A tufted perennial with dense rosettes and several stems 5–15 cm long. Rosette leaves broadly obovate or spatulate, 1–4 cm wide, densely silvery stellate; stem leaves few and small. Flowers 1–2 cm across, bright yellow; pods 2-lobed, strongly inflated, often 2 cm or more across; seeds 4 in each locule. Slopes and rocky areas; Rocky Mountains.

Raphanus radish

Raphanus raphanistrum L.

WILD RADISH

An erect annual or winter-annual weed 30–70 cm high. Basal and lower leaves 10–20 cm long, deeply lobed, with a large terminal lobe; upper leaves smaller. Flowers yellow with purplish veins, 10–20 mm across. Seedpods (Fig. 117J) 3–4 cm long, constricted between each seed, and breaking rapidly into 1-seeded sections when handled. Introduced from Europe, becoming increasingly common; a weed of abandoned fields and waste places; in eastern Prairies and Parklands.

Rapistrum

Rapistrum perenne (L.) All.

PERENNIAL MUSTARD

A biennial or perennial, with stems 30–80 cm high, densely hispid below, glabrous above. Lower leaves pinnate, coarsely serrate. Flowers 10–15 mm across, bright yellow; pods 7–10 mm long, the lower part cylindrical, the upper part ovoid, strongly ribbed. Introduced, occasional weed; southeastern Parklands.

Rorippa yellow cress

- 1. Creeping-rooted perennials. 2
 Fibrous-rooted plants without creeping
 roots; winter annual or biennial. 4
- 2. Pods not more than 3 times as long as
 wide; leaves entire or merely toothed. *R. austriaca*
 Pods 4 or more times as long as wide;
 leaves dissected or lobed. 3
- 3. Leaves with sharply toothed or incised
 divisions. *R. sylvestris*
 Leaves with blunt lobes. *R. sinuata*
- 4. Stems erect, branched above; petals 1.5–2
 mm long; sepals 2 mm. *R. islandica*
 Stems branched from the base; petals
 1.0–1.4 mm long; sepals 1.5 mm. *R. tenerrima*

Rorippa austriaca (Crantz) Besser

AUSTRIAN CRESS

A tall-growing introduced perennial 40–90 cm high, with running roots and smooth stem. Leaves alternate, almost entire or merely toothed, not dis-

sected; the lower leaves with stalks and the upper ones without. Flowers yellow and small, in racemes. Pods small, almost globular, 2–3 mm long, on stalks 5–15 mm long. An introduced weed found occasionally; in grainfields in Prairies and Parklands, but uncommon at present. However, it should be watched to prevent its spread.

Rorippa islandica (Oeder) Borbas

MARSH YELLOW CRESS

Native annual or biennial branching plants 20–60 cm high, with deeply dissected leaves. The lower leaves stalked, 6–15 cm long, and the upper ones less dissected or lobed and without stalks. Yellow flowers, small, usually less than 6 mm across. Pods either globose or linear, with two rows of seeds in each compartment. Authorities have split this species into several varieties, two of which are common in the Prairie Provinces. The var. *fernaldiana* Butt. & Abbe with almost or entirely hairless stems and linear to linear-oblong pods 6–9 mm long. It was formerly called *R. palustris* (L.) Besser. The var. *hispida* (Desv.) Butt. & Abbe has the stem covered with bristly hairs and the pods globose or oval, not more than 6 mm long. This was formerly known as *R. hispida* (Desv.) Britt. Both varieties common on lakeshores, sloughs, and wet places; throughout the Prairie Provinces.

Rorippa sinuata (Nutt.) Hitchc.

SPREADING YELLOW CRESS

A native perennial with smooth stems 10–40 cm high, and cleft or divided leaves with rather obtuse lobes. Leaves 3–8 cm long. Yellow flowers about 4 mm across. Pods 8–12 mm long, linear-oblong. Not particularly common; but has been found in several locations throughout the Prairie Provinces as far north as northern Alberta.

Rorippa sylvestris (L.) Besser

CREEPING YELLOW CRESS

An introduced perennial with creeping roots, smooth stem, and pinnately-divided leaves with sharp-toothed divisions. Yellow flowers 6–8 mm across. Pods linear, 8–12 mm long, on slender stalks, with 1 row of seeds in each compartment. Found in a few isolated places in the Prairie Provinces. Another potential pest that should be watched to prevent its spread.

Rorippa tenerrima Greene

SLENDER CRESS

An annual species 10–30 cm high; stems numerous from the root crown, slender, prostrate or erect. Leaves deeply pinnatifid, with the lobes rounded, sinuately toothed. Wet places; southern Rocky Mountains.

Sisymbrium *sisymbrium*

1. Pods 1–2 cm long, conical, closely appressed to the stem. *S. officinale*
 Pods 2–10 cm long, linear, spreading or ascending. 2
2. Leaves all or mostly entire, very narrow. *S. linifolium*
 Leaves all pinnatifid to pinnate. 3
3. Stem with spreading hairs; upper leaves with narrow, linear divisions; pods 5–10 cm long. *S. altissimum*

Stem with reflexed hairs; upper leaves with lanceolate divisions; pods about 4 cm long. *S. loeselii*

Sisymbrium altissimum L.

TUMBLING MUSTARD

An introduced annual or winter-annual weed 30–100 cm high. Basal leaves a rosette of pale green soft hairy divided leaves; the stem leaves of various shapes from deeply lobed or pinnate to entire. Flowers pale yellow, about 8 mm across. Very thin pods 5–10 cm long, each containing over 100 small light brown seeds. At maturity the plant dries and breaks off, acting as a tumbling weed and rolling with the wind, leaving seeds in its wake. A very common weed; on cultivated land; throughout Prairie Provinces, but more prevalent on open plains where it can travel with the wind.

Sisymbrium linifolium Nutt.

NARROW-LEAVED MUSTARD

Perennial plants, with much-branched glabrous stems and narrow mostly entire leaves. Pods spreading, 3–6 cm long. Rare weed, introduced from the west; western Parklands.

Sisymbrium loeselii L.

TALL HEDGE MUSTARD

A tall branched annual weed 40–100 cm high. Lower parts with downward-pointed hairs; upper parts usually smooth. Leaves deeply lobed. Flowers bright yellow; seedpods linear, 2–4 cm long. Introduced, fortunately not common; found in disturbed areas; Prairies and Parklands.

Sisymbrium officinale (L.) Scop.

HEDGE MUSTARD

Annual or biennial plants 5–80 cm high, erect, branched above. Lower leaves petiolate, pinnatifid, with the terminal lobe rounded. Flowers yellow, 4–6 mm across; pods 10–20 mm long, pubescent, or glabrous as in var. *leiocarpum* DC. Introduced, not common; a weed of waste areas.

Smelowskia rockcress

Smelowskia calycina (Stephan) C. A. Mey.

var. *americana* (Regel & Herder) Drury & Rollins

SILVER ROCKCRESS

Densely tufted perennial, with branched caudex. Stems 5–20 cm high, more or less pubescent; leaves pinnately divided, silvery gray pubescent, with the hairs mostly branched; basal leaf bases ciliate. Flowers white or cream-colored, about 1 cm across; pods 5–10 mm long, linear or narrowly oblong. Rock slides, stony areas; southern Rocky Mountains.

Subularia awl-wort

Subularia aquatica L.

WATER AWL-WORT

A glabrous annual, with the leaves all basal, 2–7 cm long, terete, awl-shaped, entire. Flowers 2–12, white; silicle 2–5 mm long. In shallow waters; eastern Boreal forest.

Thellungiella

Thellungiella salsuginea (Pall.) O. E. Schulz

MOUSE-EAR CRESS

An erect divaricately branched annual 10–35 cm high. Stem leaves oblong-ovate, with rounded auricles. Flowers 6–8 mm across, white; pods linear, 12–16 mm long, ascending or erect. Saline areas; Prairies and Parklands.

Thlaspi pennycress

Thlaspi arvense L.

STINKWEED

An introduced annual or winter-annual weed 2–40 cm high, with hairless stems and smooth leaves. Basal leaves stalked and oblanceolate, soon withering and falling off; stem leaves oblong to lanceolate, eared at the base and clasping the stems. Flowers small and white, about 3 mm across, in a cluster at the head of the stem. Pods (Fig. 117L) flat, oval, and broadly winged, 6–12 mm across, deeply notched at the top, on stalks 1–2 cm long. They form large racemose clusters, at first bright green but turning yellow or orange when mature. Seeds purplish chocolate brown, about 1 mm across, and bearing concentric grooves like a fingerprint on each side. A very common unpleasant-smelling weed, which when eaten by cattle taints their milk, butter, and meat. Often flowering and producing seed when only 3–6 cm high in early spring. Because of its early spring growth, it depletes soil of moisture. In fields and waste places; throughout the Prairie Provinces.

RESEDACEAE—mignonette family

Reseda mignonette

Introduced perennials with lobed or pinnatifid alternate leaves bearing small glandular stipules. Flowers in narrow racemes, with the stamens on one side of the flower. Seeds contained in numerous globose capsules.

Petals greenish yellow, 3 or 4 of them divided. *R. lutea*

Petals white, all cleft or divided. *R. alba*

Reseda alba L.

WHITE CUT-LEAVED MIGNONETTE

An erect hairless plant 30–50 cm high, with deeply cut or pinnate leaves, and spikes of white flowers. Rarely found, but has been introduced in seeds in some places.

Reseda lutea L.

YELLOW CUT-LEAVED MIGNONETTE

An erect plant, sometimes slightly hairy, 20–60 cm high, with deeply cut leaves, which have spatulate lobes. Flowers yellowish green in a long spike-like raceme. Occasionally found as a weed in imported lawn grass seed.

DROSERACEAE—sundew family

Perennial bog herbs with basal leaves bearing glandular sticky hairs that close over insects and entrap them. Flowers in a 1-sided raceme on an elongated stem, and numerous seeds enclosed in a capsule.

Drosera sundew

1. Leaf blades almost round, broader than long. *D. rotundifolia*
Leaf blades longer than broad. 2
2. Leaf blades spatulate or oblanceolate. *D. anglica*
Leaf blades linear. *D. linearis*

Drosera anglica Huds. OBLONG-LEAVED SUNDEW

Leaves sticky-haired, spatulate, 15–25 mm long. Flowers white, on a stem 4–10 cm high. Found in cold bogs; Boreal forest.

Drosera linearis Goldie SLENDER-LEAVED SUNDEW

Leaves linear, 1–6 cm long. Flowers white, solitary or a few on a stem 4–10 cm high. Rare; in bogs; Boreal forest.

Drosera rotundifolia L. ROUND-LEAVED SUNDEW

Leaves roundish, 6–10 mm long, broader than long, with the upper surface covered with fine glandular insect-catching hairs. Flowers white, about 4 mm wide, on a stem 10–20 cm high. The commonest of the sundews; in moist swamps and bogs; Boreal forest.

SARRACENIACEAE—pitcherplant family

Sarracenia pitcherplant

Sarracenia purpurea L. PITCHERPLANT

A peculiar and conspicuous bog plant, with pitcher-shaped leaves 10–30 cm long, erect, greenish yellow with purple veins, and hood-like top containing downward-pointing bristly hairs that trap insects. The hollow lower part of the leaf usually containing water in which insects drown and are absorbed by the plant. Flower large, about 5 cm across, nodding, yellow and purple, and borne on a long stalk 20–50 cm high. Not common; occasionally in cold bogs; in Boreal forest.

CRASSULACEAE—orpine family

Sedum stonecrop

Succulent or fleshy-leaved perennials with perfect flowers borne in cymes. They have 4 or 5 sepals and 4 or 5 petals and their numerous seeds are con-

tained in follicle-like capsules. Plants of rocky places, usually associated with mountainous locations.

1. Leaves very thick, less than 3 mm wide. 2
 Leaves flat, mostly more than 5 mm
 wide. 4
2. Stem leaves less than 5 mm long, persist-
 ent in drying. *S. acre*
 Stem leaves about 10 mm long, deciduous
 in drying. 3
3. Leaves narrowed at the base, not
 bulbiferous. *S. lanceolatum*
 Leaves broadened at the base, bulbiferous
 in upper part of the stem. *S. stenopetalum*
4. Flower parts mostly in 4's; flowers dark
 purplish. *S. rosea*
 Flower parts mostly in 5's; flowers yellow
 or pinkish. 5
5. Flowers pinkish red. *S. telephium*
 Flowers yellow. 6
6. Leaves lanceolate, with the margins
 serrate. *S. aizoon*
 Leaves spatulate, with the margins den-
 tate toward the apex. *S. hybridum*

Sedum acre L.

MOUNTAIN MOSS

Glabrous, tufted, mat-forming plants with numerous short, sterile shoots, and flowering stems 5–12 cm high. Leaves 3–6 mm long, elliptic in cross section. Flowers about 15 mm across, yellow, in small cymes. Introduced, rarely escaped from cultivation; dry rocky places.

Sedum aizoon L.

Plants with erect stems 30–40 cm high; leaves 5–8 cm long, lanceolate. Flowers 15–20 mm across, yellow, in dense compound cymes. Introduced ornamental, occasionally escaped from cultivation.

Sedum hybridum L.

Plants with creeping woody stems, short sterile shoots, and flowering stems 15–20 cm high. Leaves 2–3 cm long, oblong-cuneate. Flowers 10–20 mm across, golden yellow, numerous in terminal corymbs. Introduced ornamental, and rarely escaped; dry rocky places.

Sedum lanceolatum Torr.

LANCE-LEAVED STONECROP

Plants mat-forming, with numerous sterile shoots, and flowering stems 5–15 cm high. Leaves 6–15 mm long, linear, round in cross section. Flowers about 1 cm across, yellow, borne in a terminal cyme. Not common; dry stony slopes and hillcrests; western Prairies, Cypress Hills.

Sedum rosea (L.) Scop. var. *integrifolium* (Raf.) Berger.

ROSE-ROOT

Plants with a thick, fleshy rootstock, fragrant when cut. Flowering stems 5–35 cm high, erect; leaves linear-oblong to orbicular-ovate, entire. Flowers 6–10 mm across, dark purple to dull reddish yellow, borne in a terminal cyme. Alpine meadows and scree fields; Rocky Mountains.

Sedum stenopetalum Pursh

NARROW-PETALED STONECROP

A low fleshy-leaved tufted perennial, branched, 5–15 cm high. Leaves round in cross section (terete), 6–15 mm long, usually crowded around the base, and overlapping. Yellow flowers borne at the head of a stem 5–15 cm long, fruiting follicles erect and close to the stem. On gravelly slopes and rocky places; southern Rocky Mountains.

Sedum telephium L.

STONECROP

Plants with tuberous roots, usually simple stems 15–60 cm high, with leaves 2–7 cm long, suborbicular to oblong. Flowers 6–10 mm across, pinkish to reddish, borne in large terminal corymbs. Introduced, rarely escaped from cultivation.

SAXIFRAGACEAE—saxifrage family

Shrubs and herbs with opposite or alternate leaves with no stipules. Usually 5 sepals and 5 petals, and flowers perfect and regular. Stamens 5 or 10. The fruit usually, but not always, a capsule.

- 1. Shrubs. 2
- Herbs. 3
- 2. Fruit a berry; stamens 5. *Ribes*
- Fruit a capsule; stamens 20–40. *Philadelphus*
- 3. Petals missing; low plants. 4
- Petals present. 5
- 4. Leaves reniform, crenate; leaves and flowers clustered at tip of stem. *Chrysosplenium*
- Leaves lanceolate, serrate; flowers in terminal cymes. *Penthorum*
- 5. Bundles of sterile stamens (staminodia) alternating with stamens; flowers singly on long stems. *Parnassia*
- Sterile stamens not present. 6
- 6. Petals divided. 7
- Petals entire. 8
- 7. Petals 3- to 5-lobed; leaves deeply lobed or divided. *Lithophragma*
- Petals finely lobed, filament-like; leaves shallowly to more or less deeply lobed. *Mitella*
- 8. Inflorescence a simple raceme. *Conimitella*
- Inflorescence not a raceme. 9

9. Stamens 5. 10
 Stamens 10. 11
10. Inflorescence a spike-like panicle; root-
 stock elongate. *Heuchera*
 Inflorescence a loose panicle; rootstock
 corm-like. *Suksdorfia*
11. Petals linear, filiform. *Tiarella*
 Petals spatulate or wider. 12
12. Leaves leathery, 3–8 cm long, glossy
 green above. *Leptarrhena*
 Leaves not leathery, smaller, not glossy
 green. 13
13. Flowers pink to purple; leaves reniform;
 styles slender, elongate, partly fused. *Telesonix*
 Flowers white or yellowish; leaves vari-
 ous; styles short, not fused. *Saxifraga*

Chrysosplenium golden saxifrage

Chrysosplenium tetrandrum (Lund) Fries. GOLDEN SAXIFRAGE

Stems erect, 5–15 cm high, with alternate round leaves. Flowers greenish, mostly at the end of the stems, with no petals, but the sepals usually orange yellow inside. Rare; found in wet coniferous forests; Boreal forest. Plants with outer sepals larger and wider, and 4–8 stamens are often distinguished as *C. iowense* Rydb.

Conimitella

Conimitella williamsii (Eat.) Rydb.

Plants with basal leaves only, these reniform, glandular puberulent. Scapes 15–30 cm high, bearing 5–10 flowers, with white petals 4–5 mm long. Montane forests; southern Rocky Mountains.

Heuchera alumroot

Perennial plants from scaly rootstocks, with broad, rounded cordate or reniform, stalked, all-basal leaves; the flowers in a narrow panicle on a long stem. Fruits are capsules opening at top, with spreading beaks.

1. Flowers small; the calyx 2–4 mm long. 2
 Flowers large; the calyx 5–12 mm long. 3
2. Panicle open; teeth of the leaves acute. *H. glabra*
 Panicle very narrow; teeth of the leaves
 rounded. *H. parvifolia* var. *dissecta*
3. Stamens exerted; the calyx strongly
 oblique. *H. richardsonii*
 Stamens included; the calyx not strongly
 oblique. *H. cylindrica*

Heuchera cylindrica Dougl.

STICKY ALUMROOT

Plants 20–50 cm high; leaves 2–5 cm long, to 4 cm wide, deeply cordate at the base, glabrous, 5- to 7-lobed. Inflorescence a narrow panicle; petals included and inconspicuous or absent; base of the calyx glandular pubescent. Plants with the leaves crenate, with broad, somewhat mucronate teeth are var. *glabella* (T. & C.) Wheelock. Not common; rocky slopes, ledges, and ridges; southern Rocky Mountains.

Heuchera glabra Willd.

ALPINE ALUMROOT

Leaves ovate to rounded in outline, deeply 5- to 7-lobed, cordate at the base, sparingly pubescent when young, soon glabrous, 3–10 cm long, almost as wide as long. Stems 10–60 cm high, with the panicle 5–20 cm long, lax, glandular puberulent, open. Rare; riverbanks, rock slides; southern Rocky Mountains.

Heuchera parvifolia Nutt. var. *dissecta* M. E. Jones SMALL-LEAVED ALUMROOT

Leaves 15–30 mm broad, rounded cordate, but deeply cleft one-third to half their length in 7–9 wedge-shaped lobes. The space between the lower lobes of the leaves is very narrow and the lobes frequently overlap. Flowering stalk 20–30 cm high; flowers greenish. Found only in southern Rocky Mountains and in Cypress Hills.

Heuchera richardsonii R. Br. (Fig. 121)

ALUMROOT

Leaves coarse, basal, rounded cordate with broadly ovate teeth, 2–6 cm across, dark green. Flowering stalks 30–50 cm high with a few glandular hairs, especially near the top. Petals purplish, a trifle longer than sepals. Indians and early settlers chewed the scaly rootstocks of this plant as a cure for diarrhea. Common; on the prairie, especially in lower or moister places; throughout Prairies, Parklands, and southern fringe of Boreal forest.

Leptarrhena leather-leaved saxifrage

Leptarrhena pyrifolia (D. Don) R. Br.

LEATHER-LEAVED SAXIFRAGE

Plants with creeping or ascending rootstocks. Stems 10–40 cm high, glandular pubescent above, with 1 or 2 clasping leaves. Leaves alternate and rather crowded on the caudex, 3–8 cm long, glabrous, deep green above, pale green or brownish below; stem leaves few and distant. Flowers arranged in dense cymules in a short panicle, whitish or pinkish, 4–5 mm across. Not common; wet areas; southern Rocky Mountains.

Lithophragma starflower

Upper leaf axils usually bearing bulblets. *L. bulbifera*

Upper leaf axils not bearing bulblets. *L. parviflora*

Lithophragma bulbifera Rydb.

ROCKSTAR

A small plant from bulblet-bearing fibrous rootstocks. Stems erect, 10–20 cm high, glandular hairy. Leaves small, thrice-divided, stalked, 5–10 mm




Fig. 121. Alumroot, *Heuchera richardsonii* R. Br.

across; stem leaves usually having several very tiny bulblets in the axils. Flowers white or rose, with petals 3- to 5-cleft, 3–6 mm long. Very uncommon; on dry hills; southern Rocky Mountains and Cypress Hills.

Lithophragma parviflora (Hook.) Nutt.

SMALL-FLOWERED ROCKSTAR

Plants with more or less densely glandular hairy stems 10–30 cm high. Basal leaves white pubescent, 3- to 5-lobed to the base, with the divisions ternately divided into entire or toothed segments. Raceme 3- to 9-flowered; the calyx densely glandular hairy; petals 4–10 mm long, 3- to 5-cleft. Eroded, gravelly slopes; southern Rocky Mountains.

Mitella miterwort

1. Petals 3-cleft at the tip, white or purplish. *M. trifida*
 Petals pinnatifid or fringed, greenish or yellow. 2
2. Stamens 10; rootstocks creeping, very slender. *M. nuda*
 Stamens 5; rootstocks short, thick. 3
3. Stamens opposite sepals; leaves reniform, obscurely lobed. *M. breweri*
 Stamens opposite petals; leaves broadly cordate, shallowly lobed. *M. pentandra*

Mitella breweri Gray

BREWER'S MITERWORT

Plants with brown scaly rootstocks, and scapes 10–25 cm high. Leaves 4–8 cm wide, orbicular to reniform, with a deep basal sinus; petioles long pubescent. Petals usually with 3 pairs of filiform divisions. Moist woods; southern Rocky Mountains.

Mitella nuda L.

BISHOP'S-CAP

A low-growing perennial with scaly rootstocks and long-stalked, mostly basal, rounded cordate leaves 2–5 cm across. Flowers borne in a raceme on a stalk 5–20 cm high, 5–6 mm across, with 4 greenish sepals and 5 greenish white petals, very finely divided and branched, pinnatifid. Fairly common; in cold, wet woodlands; throughout Rocky Mountains, Boreal forest, and also in the Cypress Hills.

Mitella pentandra Hook.

Much like *M. breweri*, but with the petioles glabrous to subglabrous and the calyx often purplish inside. Mountain forests; Rocky Mountains.

Mitella trifida Graham

Perennials with scaly rootstocks, 1 to several scapes, 10–40 cm high. Leaves round-reniform to cordate, 2–4 cm wide, 7- to 9-lobed, with the lobes shallow, rounded, obscurely glandular-toothed, pubescent with white hairs. Inflorescence a one-sided raceme; the flowers subtended by small, fringed bracts; petals white, wedge-shaped, 3-cleft. Mountain woods; southern Rocky Mountains.

Parnassia grass-of-Parnassus

Perennials with rootstocks and stalked, entire, basal leaves. Flowers single on a long stalk, with sometimes a leaf on the stalk, and 5 sepals and 5 white petals with green veins. There are 5 fertile stamens, alternating with 5 clusters of gland-tipped infertile stamens termed staminodia. Fruit a 1-celled capsule opening at the top.

- 1. Petals fringed on margins, at least below;
leaves reniform to reniform-cordate. *P. fimbriata*
Petals not fringed; leaves ovate to
cordate. 2
- 2. Petals shorter than, to as long as, the sepals;
stem leafless or with a leaf at base. *P. kotzebuei*
Petals much larger than sepals; stem
leafless or with a leaf near the middle. 3
- 3. Petals 3 or more times longer than sepals;
stem leafless. *P. glauca*
Petals 1.5 times longer than sepals; stem
bearing 1 leaf near the middle. *P. palustris* var. *tenuis*

Parnassia fimbriata Konig

FRINGED GRASS-OF-PARNASSUS

Plants with leaves 2–4 cm wide, reniform, with a broad sinus. Stems 10–30 cm high, with a single ovate leaf near the middle. Flowers about 20 mm across; petals 5-nerved. Moist, springy areas and banks; Rocky Mountains.

Parnassia glauca Raf.

GLAUCOUS GRASS-OF-PARNASSUS

Low plants with broadly ovate basal leaves 2–5 cm long. Flowers white, strongly veined, 15–30 mm across, and borne singly on stems 20–40 cm high; 3–5 staminodia in each cluster. In cold bogs; Boreal forest. Syn.: *P. americana* Muhl.

Parnassia kotzebuei Cham. & Schl.

SMALL GRASS-OF-PARNASSUS

Plants about 10 cm high, with leaves ovate or oval and the base rounded or subcordate. Stem leafless or sometimes with an oval leaf near the base. Flowers 8–12 mm across; petals 3-nerved. Wet areas, bogs; Boreal forest and Rocky Mountains.

Parnassia palustris L. var. *tenuis* Wahl. (Fig. 122)

NORTHERN GRASS-OF-PARNASSUS

Cordate leaves 10–25 mm wide. The flowering stalk 10–30 cm high with a single clasping stem leaf and at the top the white flower 15–25 mm across. Petals much longer than sepals, distinguishing this variety from var. *montanensis*. The most common species of the genus; found in wet shady places; throughout Boreal forest and less plentiful in favorable sites in Parklands. In var. *montanensis* (Fern. & Rydb.) C. L. Hitchc., the petals slightly longer than the sepals, the plants generally smaller; wet areas; Rocky Mountains. Still smaller is var. *parviflora* (DC.) Boiv., usually less than 20 cm high, and with most of the flowers less than 15 mm across; wet areas; Boreal forest and Rocky Mountains.



Fig. 122. Northern grass-of-Parnassus, *Parnassia palustris* L. var. *tenuis* Wahl.

Penthorum stonecrop

Penthorum sedoides L.

DITCH STONECROP

Perennial plants with stems 20–70 cm high arising from a creeping base. Leaves 5–10 cm long, lanceolate, serrate. Inflorescence terminal, glandular; flowers 4–6 mm across; petals none, in cymes 2–8 cm long. Rare; shores and ditches; southeastern Boreal forest.

Philadelphus mock orange

Philadelphus lewisii Pursh

MOCK ORANGE

Shrubs with freely branching stems 1–3 m high; young twigs reddish, glabrous; leaves 2–6 cm long, ovate-lanceolate, opposite. Flowers showy, fragrant, in racemose cymes; 4 or 5 white petals 1–2 cm long; sepals usually 4, rarely 5, pubescent at the inside of the tips; stamens 20–40. Rare; mountain slopes; southern Rocky Mountains.

Ribes currant

Shrubs, with palmately divided and veined leaves, and regular perfect flowers. Fruit a globose or ovoid pulpy berry. Some plants armed with prickles, especially at the nodes of the stem. Fruit edible, but sometimes not very tasty.

- 1. Inflorescence a reduced raceme, 1- to 3-flowered; most of the stems spiny. *R. oxyacanthoides* var. *oxyacanthoides*
 Inflorescence an elongated raceme, flowers numerous. 2
- 2. Stems densely spiny along the internodes. *R. lacustre*
 Stems not spiny, or spines only at the nodes. 3
- 3. Flowers golden yellow to somewhat reddish; leaves rolled in buds. *R. aureum*
 Flowers not golden yellow; leaves folded in buds. 4
- 4. Ovary and fruit more or less densely glandular pubescent. 5
 Ovary and fruit glabrous, or with a few sessile glands. 7
- 5. Leaves and young stems glandular puberulent. *R. viscosissimum*
 Leaves and young stems not glandular. 6
- 6. Fruit red, glandular only. *R. glandulosum*
 Fruit purplish black, pubescent and glandular. *R. laxiflorum*
- 7. Leaves abundantly glandular dotted on the underside. 8
 Leaves without glands. 9

8. Pedicels much longer than the bracts. *R. hudsonianum*
 Pedicels much shorter than the bracts. *R. americanum*
9. Lobes of leaves closely and uniformly serrate from base to tip. *R. rubrum* var. *propinquum*
 Lobes of leaves coarsely dentate above the middle. *R. diacanthum*

Ribes americanum Mill.

WILD BLACK CURRANT

A low shrub 1–2 m high, with unarmed stems and 3- to 5-lobed leaves 2–7 cm across and somewhat hairy and resinous-dotted on the underside. Flowers greenish white, with a tubular calyx 3–5 mm long, and borne in drooping racemes. Fruit black, 6–10 mm across, and smooth. Fairly common; in moist woodlands; throughout the Prairie Provinces. Syn.: *R. floridum* L'Her.

Ribes aureum Pursh

GOLDEN CURRANT

An erect shrub 1–2 m high, with 3-lobed leaves 2–5 cm across. Flowers bright yellow tipped with red, with a long cylindric calyx tube 6–10 mm long, clove-scented, in small racemes. Fruit, about 6 mm in diam, varying from pale yellow through shades of red to black. Often grown as an ornamental garden shrub. Only found in the Prairies; plentiful in some localities on the southern slope of Cypress Hills. Syn.: *Chrysobotrya aurea* (Pursh) Rydb.

Ribes diacanthum Pall.

RED CURRANT

Shrubs 1–2 m high; leaves 2–6 cm long, rather deeply 3-lobed. Flowers in racemes, functionally dioecious, with the organs of one sex rudimentary; each flower subtended by a conspicuous bract; corolla greenish; fruit scarlet. Introduced and cultivated, occasionally escaped.

Ribes glandulosum Grauer

SKUNKBERRY

A low spreading or reclining shrub about 1 m long, with no thorns or prickles. Leaf blades 5- to 7-lobed, cordate at base, hairy on the veins below, 2–7 cm across, and having a somewhat skunk-like odor. Flowers in a short raceme. Fruit a red, glandular, bristly berry about 6 mm across. Common; in damp woodlands; throughout Parklands and Boreal forest.

Ribes hudsonianum Richards.

NORTHERN BLACK CURRANT

An erect shrub 1–1.5 m high, with unarmed stems and 3-lobed (occasionally 5-lobed) leaves, wider than long, 2–10 cm wide, more or less hairy, and dotted with resinous glands on the undersides. Flowers white, in racemes. Fruit black, 5–10 mm in diam. Common in shady woodland; in the northern parts of the Prairie Provinces and in the Cypress Hills.

Ribes lacustre (Pers.) Poir.

SWAMP GOOSEBERRY

Shrubs 1–2 m high, with clusters of slender spines; stems and branches densely bristly. Leaves 2–5 cm long, deeply 5- to 7-lobed. Flowers in racemes, green or purplish. Fruit a densely hairy, glandular reddish berry about 6 mm in diam. Found occasionally; in swamps; in Boreal forest and in the Riding Mountain. Syn.: *Limnobotrya lacustris* (Pers.) Rydb.

Shrubs with ascending or spreading stems; leaves more or less orbicular in outline, 5- to 7-lobed, cordate at base. Flowers in ascending racemes, with the pedicels glandular, the corolla red, the petals 3–4 mm long, pubescent on the outside; berry purplish black to blue black. Swamps and woods; southern Rocky Mountains.

Ribes oxycanthoides L. var. *oxycanthoides* (Fig. 123) NORTHERN GOOSEBERRY

A low, bristly shrub or bush to 1 m high, with lobed leaves 1–4 cm broad and greenish purple or white sepals and petals; bracts glandular ciliate. Calyx tube not longer than lobes. Fruit a globose berry 10–15 mm across, turning reddish purple when ripe. Common in woodlands and shrubbery, especially in Boreal forest. Syn.: *R. setosum* Lindl. The var. *saxosum* (Hook.) Cov. having the bracts long-ciliate but not glandular, generally less spiny, particularly at the nodes and on older stems. Woodlands; in northern Parklands and Boreal forest. Syn.: *R. hirtellum* Michx.

Ribes rubrum L. var. *propinquum* Trautv. & Mey. SWAMP RED CURRANT

A shrub about 1 m high, unarmed; leaves paler beneath, usually 3-lobed, sometimes 5-lobed, 3–8 cm across. Flowers usually purplish; fruit a smooth red berry, very similar to the garden red currant, about 6 mm across. Found occasionally; in rich poplar woods; especially in Parklands, Boreal forest, and Riding Mountain. Syn.: *R. triste* Pall.

Ribes viscosissimum Pursh STICKY CURRANT

A rather stout shrub 50–100 cm high, with spreading branches. Leaves 4–7 cm across, orbicular or reniform in outline, 3- or sometimes 5-lobed, cordate, sticky. Racemes 3- to 7-flowered, glandular puberulent; flowers greenish white tinged with pink; fruit black, glandular hairy, about 1 cm long. Mountain forests and slopes; southern Rocky Mountains.

Saxifraga saxifrage

- 1. Leaves opposite, densely crowded, 4-ranked on matted stems; petals 6–8 mm long, purple. *S. oppositifolia*
- Leaves alternate or basal. 2
- 2. Stems leafless, except for reduced leaves subtending branches of inflorescence. 3
- Stems with several well-developed leaves. 9
- 3. Basal leaves 10–20 cm long; scapes 30–80 cm high. *S. pensylvanica*
- Basal leaves much smaller; scapes lower. 4
- 4. Leaves subcordate to deeply cordate at base. 5
- Leaves cuneate to acute at the base. 6
- 5. Flowers in part replaced by clusters of bulblets. *S. mertensiana*



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Ribes oxycanthoides* L.
var. *oxycanthoides*

LOCATION: Montrose Plains

COLLECTOR: C. D. Loope

HT:

COLLECTOR: 1946

DATE: 12 June 1946

ELEVATION: 1900'

TERMINAL: none



Fig. 123. Northern gooseberry, *Ribes oxycanthoides* L. var. *oxycanthoides*.

- Flowers normal, bulblets absent. *S. punctata* var. *porcildiana*
6. Sepals sharply reflexed, pendent. 7
 Sepals spreading or ascending. 8
7. Plants glabrous or somewhat puberulent
 above. *S. lyallii*
 Plants densely glandular pubescent
 throughout. *S. ferruginea*
8. Petals 2–4 mm long, obovate to oblong. *S. occidentalis*
 Petals 4–5 mm long, oblanceolate to
 lanceolate. *S. virginiensis*
9. Leaves 3-parted to palmately lobed. 10
 Leaves shallowly 3-lobed to entire. 12
10. Plants with bulblets in the upper axils. *S. cernua*
 Plants not bulbiferous. 11
11. Lobes of leaves linear to
 linear-lanceolate. *S. cespitosa*
 Lobes of leaves ovate to rounded. *S. rivularis*
12. Flowers white, the petals often
 purplish-dotted. 13
 Flowers yellow, the petals not dotted. 15
13. Leaves soft, not spine-tipped. *S. adscendens* ssp. *oregonensis*
 Leaves stiff, thick, spine-tipped. 14
14. Leaves entire, spinulose on the margins. *S. bronchialis* var. *austromontana*
 Leaves 3-toothed at the apex, the teeth
 spine-tipped. *S. tricuspidata*
15. Plants long-stoloniferous, the stolons
 whip-like. *S. flagellaris*
 Plants not stoloniferous. 16
16. Leaves all sessile. *S. aizoides*
 Basal leaves long-petioled, stem leaves
 subsessile. *S. hirculus*

Saxifraga adscendens L. ssp. *oregonensis* (Raf.) Bacig.

WEDGE-LEAVED SAXIFRAGE

A small perennial, with stems 2–8 cm high; several stem leaves. Basal leaves in compact rosettes 5–15 mm long, oblong-spatulate or cuneate, entire or 3-toothed; stem leaves cuneate, 3-toothed. Petals 3–5 mm long, white. Moist slopes and ledges; southern Rocky Mountains.

Saxifraga aizoides L.

YELLOW MOUNTAIN SAXIFRAGE

Perennial plants with stems from a multiple caudex; flowering stems 5–20 cm high, with several stem leaves. Lower leaves 1–2 cm long, linear-oblong, thick, often ciliate. Cymes usually several-flowered; flowers erect, 5–8 mm across, yellow, predominantly orange-dotted. Dry, gravelly slopes; southern Rocky Mountains.

Saxifraga bronchialis L. var. *austromontana* (Wieg.) G. N. Jones

Plants forming dense mats or cushions; the flowering stem 5–15 cm high. Leaves 8–15 mm long, loosely imbricate, linear to narrowly lanceolate, with the margin spinulose. Inflorescence a corymbose cyme; flowers 6–12 mm across, pale yellow or whitish, often red-spotted. Usually on rock outcrops; Rocky Mountains.

Saxifraga cernua L.

Plants with usually simple stems 10–20 cm high, more or less glandular puberulent. Basal leaves reniform, 15–25 mm wide, 5- to 7-lobed; the cauline leaves similar but progressively smaller, all with reddish bulblets in the axils. Usually a solitary terminal flower 15–20 mm across, white. Alpine rockslides; southern Rocky Mountains.

Saxifraga cespitosa L.

Plants with short leafy shoots, more or less erect, forming lax to rather dense cushions. Stems 4–10 cm high. Leaves 5–15 mm long, mostly 3-lobed, cuneate, glandular pubescent. Flowers 8–10 mm across, dull white or slightly greenish. Arctic and alpine gravel slopes; northeastern Boreal forest, southern Rocky Mountains.

Saxifraga ferruginea Graham

Plants with stems 10–30 cm high, more or less glandular pubescent. Basal leaves crowded, 2–6 cm long, spatulate to oblanceolate, densely pubescent. Flowering stems with branches ascending, flowers white, the 3 upper petals with 2 yellow spots at the base. In f. *vreelandii* (Small) St. John & Thayer, some flowers replaced by bulblets. Moist ledges and banks; southern Rocky Mountains.

Saxifraga flagellaris Stemb. & Willd.

Plants with erect stems 5–20 cm high, with whip-like stolons to 15 cm long from the base, each terminating in a rosette. Leaves mainly in a basal rosette, more or less glandular pubescent. Flowers few, in a lax cyme 15–20 mm across, bright yellow. High alpine soils; southern Rocky Mountains.

Saxifraga hirculus L.

Plants caespitose or with short stolons; stems 15–25 cm high; leafy below. Leaves 10–25 mm long, lanceolate, pubescent with reddish brown hairs. Flowers solitary or 2–4 in a loose corymb 25–30 mm across, bright yellow, sometimes red-spotted. Wet tundra; northeastern Boreal forest.

Saxifraga lyallii Engler

Plants with a prominent caudex; leaves fan-shaped to cuneate-obovate, 1–4 cm long, usually coarsely toothed above the middle. Scapes 5–10 cm high, glabrous or sparsely glandular pubescent above. Inflorescence open; flowers 5–8 mm across, white, with deep red sepals. In var. *hultenii* Calder & Savile, plants 10–30 cm high, with larger basal leaves and paniculate inflorescence. In

var. *laxa* Engler, plants to 40 cm high, with the basal leaves orbicular, cuneate to subtruncate at base, and paniculate inflorescence. Along alpine creeks and rivulets; southern Rocky Mountains.

Saxifraga mertensiana Bong.

Plants with stout, erect, often bulb-forming rootstocks. Leaves basal, glandular pubescent, with the blades orbicular or ovate-reniform, cordate at the base, shallowly lobed; the lobes 3-toothed. Flowering stems 10–30 cm high, minutely glandular pubescent, with 1 or 2 leaves at the base, bracteose above. Inflorescence few-flowered; often all but the terminal flower replaced by bulblets; flowers white, with the filaments conspicuously club-shaped. Moist cliffs; southern Rocky Mountains.

Saxifraga occidentalis Wats.

RHOMBOID-LEAVED SAXIFRAGE

A low plant with fleshy basal leaves, usually ovate, 1–4 cm long, wavy-margined, and smooth. Flowers borne in a branched inflorescence at the head of a glandular haired stalk 8–20 cm high. Flowers about 5 mm across, with white petals. Fruits are follicle-like capsules. Moist meadows and slopes; southern Rocky Mountains, Cypress Hills. Syn.: *Micranthes rhomboidea* (Greene) Small; *S. rhomboidea* Greene.

Saxifraga oppositifolia L.

RED-FLOWERED SAXIFRAGE

Mat-forming plants, with stems procumbent or ascending. Leaves 2–6 mm long, suborbicular to obovate-lanceolate, dull bluish green, thick, keeled below. Flowering stems very short, leafy, glandular pubescent; flowers solitary, pale pink to deep purple. Rocky and gravelly areas, arctic–alpine; northeastern Boreal forest, Rocky Mountains.

Saxifraga pensylvanica L. (Fig. 124)

MARSH SAXIFRAGE

Stout plants with erect stems 30–80 cm high. Basal leaves 10–20 cm long, oblong to lanceolate or oblong-lanceolate, entire to sparsely glandular serrulate, sparingly pubescent. Inflorescence at first compact, becoming elongate, lax; flowers 5–8 mm across, greenish white. Rare; bogs and wet meadows; southeastern Boreal forest.

Saxifraga punctata L. var. *porsildiana* (Calder & Savile) Boiv.

SPOTTED SAXIFRAGE

Plants with scapes 8–15 cm high. Basal leaves long-petioled, with the blades nearly orbicular, deeply cordate at base, 2–5 cm across, crenate-dentate, glabrous or nearly so. Scapes solitary or few tufted; inflorescence closely corymbose, glandular puberulent; flowers 6–10 mm across, white; sepals reflexed, ciliate. Moist areas; Rocky Mountains.

Saxifraga rivularis L.

ALPINE BROOK SAXIFRAGE

Plants with short creeping rootstocks; erect stems 6–15 cm high, glabrous to glandular puberulent. Basal leaves 10–15 mm wide, 3- to 5-lobed; cauline leaves soft pubescent. Flowers 6–10 mm across, white or pale pink. Moist, springy areas; Rocky Mountains, northeastern Boreal forest.



Fig. 124. Marsh saxifrage, *Saxifraga pensylvanica* L.

Saxifraga tricuspidata Rottb.

THREE-TOOTHED SAXIFRAGE

Plants with branched caudex, the branches mat-forming; leaves closely crowded, 1–2 cm long, narrowly oblong, ciliate, the apex with 3 sharp erect teeth. Stems 10–20 cm high, with several stem leaves 3–10 mm long; flowers in a cyme 12–15 mm across, white with light purplish dots. Rock outcrops and rocky slopes; Boreal forest, Rocky Mountains.

Saxifraga virginensis Michx.

EARLY SAXIFRAGE

Plants with basal leaves 2–5 cm long, oblong to ovate, entire or serrate, the base narrowed to a petiole, glandular pubescent. Scapes 10–20 cm high, glandular pubescent; inflorescence branched, at first compact, later elongating and open; flowers 8–12 mm across; sepals spreading to ascending. Moist or dry open woods; southeastern Boreal forest.

Suksdorfia suksdorfia

Suksdorfia violacea Gray

BLUE SUKSDORFIA

A slender perennial, with more or less glandular pubescent stems 10–20 mm high. Basal leaves kidney-shaped, 1–3 cm wide, with 5–7 rounded lobes, on petioles 2–8 cm long, glandular puberulent; upper leaves sessile. Inflorescence a few-flowered panicle; flowers long-pedicel, 10–15 mm across, violet. Wet rocks along streams; southern Rocky Mountains.

Telesonix

Telesonix jamesii (Torr.) Raf. var. *heucheriformis* (Rydb.) Bacig.

Low perennial plants with a thick scaly rootstock; stems 5–15 cm high, glandular pubescent. Basal leaves reniform to orbicular, 2–4 cm wide, deeply crenate-dentate, with sessile glands; upper cauline leaves fan-shaped to cuneate, short-petioled or sessile. Inflorescence a short, leafy panicle; flowers 3–5 mm long, deep violet. Rock crevices; southern Rocky Mountains.

Tiarella false miterwort

Leaves compound, usually trifoliolate. *T. trifoliata*
Leaves simple, shallowly 3- to 5-lobed. *T. unifoliata*

Tiarella trifoliata L.

LACEFLOWER

Perennial plants with slender, sparsely hirsute stems 20–60 cm high, 2- to 4-leafed, glandular hirsute above. Leaves with 3 leaflets, on petioles 5–15 cm long, glabrous to villous-hirsute; middle leaflet more or less 3-lobed, 2–8 cm long. Inflorescence an elongated, narrow panicle; sepals about 1.5 mm long, white; petals white, linear, about 5 mm long. Moist coniferous forests; Rocky Mountains.

Tiarella unifoliata Hook.

SUGARSCOOP

Slender perennial, with stems 10–50 cm high, glabrous or hirsute with white hairs, glandular puberulent above. Basal leaves cordate in outline, 4–10

cm wide, 3- to 5-lobed, with the lobes broadly ovate, acute or obtuse, crenate-dentate, and the petioles 5–15 cm long. Stem leaves 1–4, similar to but smaller than the basal leaves. Inflorescence a narrow panicle; sepals white to pinkish, 1.5–2.5 mm long; petals white, linear, about 6 mm long. Moist woods; Rocky Mountains and northwestern Boreal forest.

ROSACEAE—rose family

An extremely variable family, including herbs, shrubs, and small trees, all having alternate leaves with stipules. Flowers in racemes or cymes, perfect and regular, with 5 petals and 5 sepals, often with 5 bracts below, alternating with the sepals. Many stamens and from one to many pistils. Fruits varying: dry achenes, follicles, fleshy receptacles (strawberry), fleshy drupes or drupelets (raspberry), and berry-like pomes (saskatoon).

1. Shrubs or trees.	2
Herbs or half-shrubs.	14
2. Leaves simple.	3
Leaves compound or lobed.	9
3. Flowers solitary on a long peduncle; creeping dwarf shrubs.	<i>Dryas</i>
Flowers usually more numerous; plants taller.	4
4. Spiny shrubs or small trees.	5
Shrubs or trees not spiny.	6
5. Spines never leafy.	<i>Crataegus</i>
Spines leafy in the first year, leafless thereafter.	<i>Prunus</i>
6. Leaves entire; ovary inferior.	<i>Cotoneaster</i>
Leaves serrate or toothed.	7
7. Fruit a leathery follicle.	<i>Spiraea</i>
Fruit fleshy.	8
8. Fruit a red or black drupe with a single stone.	<i>Prunus</i>
Fruit a berry-like pome with several seeds.	<i>Amelanchier</i>
9. Leaves lobed, not distinctly divided.	10
Leaves compound, distinctly divided into leaflets.	11
10. Shrubs tall; flowers in dense racemose clusters.	<i>Physocarpus</i>
Shrubs usually low; flowers in loose racemes or solitary.	<i>Rubus</i>
11. Flowers yellow; fruit dry; leaves with 3–7 small leaflets.	<i>Potentilla</i>

- Flowers white or pink; fruit more or less fleshy. 12
12. Small trees; leaflets 11–17; flowers white in flat-topped racemes. *Sorbus*
 Shrubs; leaflets 3–11. 13
13. Leaves with 3–5 leaflets; flowers white; fruit a cluster of many fleshy drupelets. *Rubus*
 Leaves with 5–11 leaflets; flowers pink or white; fruit a bottle-shaped “hip.” *Rosa*
14. Leaves much dissected into linear lobes. 15
 Leaves compound. 16
15. Plant a half-shrub, with trailing woody stems. *Luetkea*
 Plant with dense rosettes, entirely herbaceous. *Chamaerhodos*
16. Calyx simple, of 5 lobes. 17
 Calyx double, with 2 rings of lobes, the outer ones smaller than the inner. 18
17. Flowers yellow, in a spike-like raceme; fruit with hooked prickles. *Agrimonia*
 Flowers white, in a terminal panicle; fruit not prickly. *Sorbaria*
18. Fruit fleshy; leaves with 3 leaflets; plants stoloniferous. *Fragaria*
 Fruit not fleshy; leaves various. 19
19. Style much longer than the achene, hooked or plumose. *Geum*
 Style much shorter than the achene, often deciduous. *Potentilla*

Agrimonia agrimony

Agrimonia striata Michx.

AGRIMONY

An erect-stemmed perennial herb 30–80 cm high, sometimes branched above, with soft, fine brownish hairs. Leaves pinnate, with 7–9 leaflets, deeply indented, 2–7 cm long, smooth above, and softly hairy below. Small yellow flowers borne in a spike-like raceme. The fruiting calyx reflexed (pointing downward), with short hooked bristles at the top, and containing 2 seeds. Found occasionally; in small poplar woods and along roadsides; throughout the Prairie Provinces, but very rare in Prairies.

Amelanchier juneberry

Small trees or bushes without spines or prickles, with simple alternate leaves, and with white flowers in racemes. Fruit a berry-like, sweet-flavored, several-seeded pome.

- 1. Pedicels short, mostly 6–8 mm. *A. alnifolia*
 Pedicels long, the lower ones 15–20 mm
 long. 2
- 2. Leaves glabrous or nearly so, rounded to
 cordate. *A. florida*
 Leaves white pubescent below, obtuse to
 rounded. *A. sanguinea*

Amelanchier alnifolia Nutt. (Fig. 125)

SASKATOON

A tall shrub or small tree 1–4 m high, with smooth chocolate brown stems and branches. Leaves simple, rounded at the ends, 1–5 cm long, with a few serrate teeth at the apex. Flowers white, about 8–12 mm across, in racemes quite early in the season. Fruits purple and berry-like, very sweet when ripe, globular, 6–10 mm across, and used for preserves. Used by the Indians and early settlers as a constituent of pemmican. Very common; in coulees, bluffs, and open woodlands; throughout the Prairie Provinces. The most commonly found *Amelanchier* in the area.

Amelanchier florida Lindl.

SERVICEBERRY

A shrub or small tree 2–5 m high, with reddish brown branchlets. Leaves broadly oblong to suborbicular, 3–4 cm long, subcordate to rounded at base, coarsely toothed above the middle, more or less tomentose below. Flowers 20–30 mm across; fruit dark purple. Slopes of valleys and along rivers; Rocky Mountains, Peace River, and Cypress Hills.

Amelanchier sanguinea (Pursh) DC.

EASTERN SERVICEBERRY

An erect or straggling shrub or small tree 1–3 m high, usually in clumps of several stems. Leaves densely tomentose at flowering time, only half-grown; later becoming glabrous, oblong to subrotund, to 7 cm long, finely to coarsely toothed above the middle. Racemes loose; flowers 20–30 mm across; fruit purple. Forest margins and openings; southeastern Boreal forest.

Chamaerhodos chamaerhodos

Chamaerhodos erecta (L.) Bunge var. *parviflora* (Nutt.) C. L. Hitchc.

CHAMAERHODOS

Low-growing perennial or biennial glandular herbs 10–30 cm high, from a single stem, but sometimes branching above; the usual form, however, being a very narrow pyramid. Basal leaves, from a stout root, usually forming a rosette, and later becoming a small clump; leaves small, with 3 leaflets, each leaflet again divided into 3 lobes, and sometimes these lobes again 3 times cleft. Stem leaves similar to the basal ones, but less divided. Inflorescence much-branched, of many small white flowers about 3 mm across. Very common; on dry hillsides and light soil; throughout Prairies and Parklands. Syn.: *C. nuttallii* Pickering.

Cotoneaster cotoneaster

Young shoots and leaves yellowish tomentose

on underside, later pubescent. *C. acutifolia*




Fig. 125. Saskatoon, *Amelanchier alnifolia* Nutt.

Young shoots and leaves white tomentose on underside, remaining so. *C. melanocarpa*

Cotoneaster acutifolia Turcz. COTONEASTER

Shrubs with branches and leaves 2-ranked, forming flattish sprays. Leaves 3–4 cm long, ovate, dark green above, pale green below; tomentum of young leaves bright to brownish yellow. Inflorescence a small corymb; flowers yellow; fruit black with 2 nutlets. Introduced and cultivated as an ornamental or hedge; occasionally spreading into native woodlands or shrubbery.

Cotoneaster melanocarpa Lodd. COTONEASTER

Similar to the previous species, but smaller, with the tomentum denser, pure white, and persistent; the fruit with 4 nutlets. A cultivated introduction, occasionally escaped.

Crataegus hawthorn

Shrubs or small trees with sharp thorns or spines on the stem, and alternate, sometimes lobed, leaves. Flowers white, in clusters. Fruit a round berry-like pome. A difficult genus to separate into species because of apparent hybridizing between species, thereby obscuring the characters.

1. Sepals short, triangular, acute but not attenuate; fruit black or very dark purple. *C. douglasii*
Sepals attenuate, long-pointed; fruit purple or red. 2
2. Teeth of leaves not gland-tipped; nutlets deeply pitted on faces. *C. succulenta*
Teeth of leaves gland-tipped; nutlets not pitted. *C. rotundifolia*

Crataegus douglasii Lindl. DOUGLAS HAWTHORN

A tall shrub or small tree 3–6 m high with short spines 6–25 mm long. Leaves ovate to obovate, 15–50 mm long; fruit black or very dark purple. Found occasionally; in western part of the Prairie Provinces and in Cypress Hills.

Crataegus rotundifolia Moench (Fig. 126) ROUND-LEAVED HAWTHORN

A round-topped shrub 1–3 m high with stout thorns, usually about 3 cm long, on the stems and branches. Leaves usually almost round, 2–5 cm across, sometimes lobed, but usually double-toothed. Flowers white, 10–15 mm across, borne in clusters at the ends of the branches, and followed by red berry-like fruits about 1 cm across containing several bony carpels. Fairly common; in coulees, stream banks, and open woods; throughout the Prairie Provinces. Syn.: *C. chrysoarpa* Ashe; *C. columbiana* Howell.

Crataegus succulenta Link LONG-SPINED HAWTHORN

A shrub very similar to *C. rotundifolia* with no dark glandular tips on the leaf teeth and with deeply pitted nutlets. Plentiful in eastern parts but appears to be replaced by *C. rotundifolia* in the west.



HERBARIUM OF THE RESEARCH CENTER
WEST VIRGINIA, 1954

CRATAEGUS ROTUNDFOLIA MOENCH

TYPE AND COLLECTOR

C. DC.

1798

FRANCE

HERBARIUM

PARIS




Fig. 126. Round-leaved hawthorn, *Crataegus rotundifolia* Moench.

Dryas mountain-avens

- 1. Leaves entire or nearly so. *D. integrifolia*
 Leaves crenate to the tip. 2
- 2. Flowers yellow; leaves tapering to base. *D. drummondii*
 Flowers white; leaf base blunt. *D. octopetala* var. *hookeriana*

Dryas drummondii Richardson YELLOW MOUNTAIN-AVENS

Depressed, mat-forming shrubs; the young branches white tomentose. Leaves elliptic to obovate, 1–3 cm long, densely white tomentose below, dark green and rugose above. Scape 5–20 cm long, tomentose; calyx black glandular pubescent; petals yellow, about 1 cm long; styles plumose in fruit, 3–5 cm long. Gravel banks, scree fields, and slopes; Rocky Mountains, northern Boreal forest.

Dryas integrifolia Vahl WHITE MOUNTAIN-AVENS

Depressed, densely mat-forming shrub, very leafy. Leaves 8–15 mm long, oblong-lanceolate, entire, often cordate at base. Scape 3–10 cm high; calyx tomentose; petals about 1 cm long, white; styles 15–25 mm long in fruit. Gravel banks along rivers and rocky slopes; northeastern Boreal forest, Rocky Mountains.

Dryas octopetala L. var. *hookeriana* (Juz.) Breit. MOUNTAIN-AVENS

Similar to *D. drummondii*, but forming small mats. Leaves strongly rugose above, truncate to subcordate at base, and black glandular as well as tomentose; calyx white tomentose and black pubescent; petals white. Alpine gravels and rock outcrops; Rocky Mountains.

Fragaria strawberry

Perennial low-growing herbs with 3-foliolate leaves, and running stems rooting at tips and producing new plants. Flowers white, with 5 petals, 5 sepals, and 5 sepal-like bracts. Fruit is the enlarged fleshy receptacle of the flower, bearing very juicy and sweet achenes on the surface.

Hairs on the stems ascending and closely pressed to stems. *F. virginiana*

Hairs on the stems spreading in various directions. *F. vesca* var. *americana*

Fragaria vesca L. var. *americana* Porter AMERICAN WILD STRAWBERRY

Similar to the following species, but with leaves paler, flowers smaller, and leaflets stalkless (leaflets of *F. virginiana* having short stalks). Fruit ovoid or conical, with seeds borne on the surface. Rocky woodlands; in Boreal forest. Syn.: *F. americana* (Porter) Britt.

Fragaria virginiana Dcne. SMOOTH WILD STRAWBERRY

A low-growing herb with coarsely toothed, broadly ovate leaflets, sometimes silky beneath. Flowers white, 15–20 mm across, appearing fairly early in the season. Fruit almost round, 10–15 mm in diam, with seeds sunk in shallow

pits. The commonest strawberry; in low spots on prairie, open woodlands, and moist areas; throughout the Prairie Provinces. Syn.: *F. virginiana* Dcne. var. *glauca* Wats.; *F. glauca* (S. Wats.) Rydb.

Geum avens

Perennials from stout rootstocks, with lyrate or pinnate leaves, perfect regular flowers, and many achenes. The long styles, remaining on the achenes, either jointed or feathery.

- 1. Styles feathery and not jointed; leaves pinnate. *G. triflorum*
- Styles not feathery, but jointed; leaves lyrate-pinnate. 2
- 2. Sepals erect or spreading; flowers purple or flesh-colored; upper part of style at least half as long as lower part. *G. rivale*
- Sepals reflexed; flowers yellow; upper portion of style less than one-third the length of lower part. 3
- 3. Upper portion of style hairy. *G. aleppicum*
- Upper portion of style not hairy, or with only a few stiff hairs. *G. perincisum*

Geum aleppicum Jacq.

YELLOW AVENS

A hairy erect plant 40–120 cm high, with broad leafy stipules and basal leaves lyrate-pinnate (terminal lobe longer than others) with 5- to 7-toothed or divided leaflets and often a few very small interspersed leaflets. Stem leaves with 3–5 leaflets, and either with or without very short stalks. Flowers bright yellow, 10–25 mm across, and followed by a fruiting head about 15 mm in diam, on which the characteristic hooked or bent styles are located. Common; in wet or moist locations on prairies and in meadows and open woods; throughout the Prairie Provinces. Syn.: *G. strictum* Ait.

Geum perincisum Rydb.

LARGE-LEAVED AVENS

Very similar to *G. aleppicum*, but the terminal segment of the basal leaves large, often 3-lobed, and having 5–15 leaflets. Flowers yellow and the fruiting head similar to that of the preceding species except that the upper portion of the style on the fruit is entirely or almost hairless. Not so common as yellow avens; found in moist meadows and open woods; throughout the Prairie Provinces.

Geum rivale L.

PURPLE AVENS

An erect, simple, and little-branched herb, more or less hairy, 30–60 cm high, with lyrate-pinnate basal leaves and thrice-divided stem leaves. Sepals not bent abruptly downward as those of the two preceding species. Flowers 15–20 mm across, flesh-colored or purple, often with a yellowish tinge. Fruiting heads very similar to those of the preceding species, but the upper portion of the style at least half as long as the lower part. Found occasionally; in moist places and wet, swampy ground; Boreal forest and the Cypress Hills.

Erect perennial herbs 15–40 cm high from a thick, coarse, almost black rootstock, with coarse black roots. The many basal leaves pinnate, with many wedge-shaped lobed leaflets, often with smaller leaflets mixed with the larger ones. Flowering stem usually with a tuft of small leaves halfway up, and some thin leafy bracts at summit. Flowers usually 3, 12–20 mm across; sepals purplish pink; petals pink, yellowish, or flesh-colored. Fruiting head bearing long, persistent, feathery styles 2–5 cm long. A very common spring flower on the prairies, the bright green leaves being among the first new foliage to show in spring and the flowers among the very early spring blooms. Plentiful; everywhere on open prairie; Prairies, Parkland, and Rocky Mountains. Syn.: *Sieversia triflora* (Pursh) R. Br.

Luetkea luetkea

Luetkea pectinata (Pursh) Ktze.

LUETKEA

A caespitose plant, semishrubby, with prostrate, stoloniferous branches. Leaves dissected into linear divisions, 10–15 mm long, acute, grooved above. Raceme narrow, 1–5 cm long; petals 3–3.5 mm long, white; sepals 2 mm long, ovate, acute. Moist slopes and mountain meadows; southwestern Rocky Mountains.

Physocarpus ninebark

Physocarpus malvaceus (Greene) Ktze.

MALLOW-LEAVED NINEBARK

A shrub 1–2 m high, with brown branches, these glabrous or stellate pubescent. Leaves 2–6 cm long, round-ovate, 3- to 5-lobed, more or less doubly crenate, mostly cordate at base, and stellate pubescent on both sides. Inflorescence of half-round corymbs, 3–5 cm wide; flowers white, about 1 cm across. Rare; thickets and forest openings; southern Rocky Mountains.

Potentilla cinquefoil

Generally perennial, but some species annual or biennial, with alternate leaves, either palmately or pinnately compound. Flowers perfect, with 5 sepals, 5 petals, and many stamens. Fruit a head with many achenes.

- 1. Flowers reddish purple; leaves with 5–7 leaflets. *P. palustris*
- Flowers white or yellow. 2
- 2. Plant shrubby or with woody bases; achenes hairy. 3
- Plant herb-like, not woody based; achenes generally smooth. 4
- 3. Shrub 15–150 cm high; flowers yellow; pinnate leaves. *P. fruticosa* ssp. *floribunda*
- Low plant, woody at base; flowers white; palmate leaves. *P. tridentata*



Fig. 127. Three-flowered avens, *Geum triflorum* Pursh.

4. Low creeping plant with runners; leaves basal and pinnate; yellow flowers singly on long stalks from base of plant.	<i>P. anserina</i>
Plant without runners; flowers borne in terminal cymes and not singly.	5
5. Annual or biennial plant without perennial rootstocks; inflorescence many-flowered and leafy.	6
Perennial plant with stout rootstocks; often showing old bases of leaves.	8
6. Lower leaves pinnate, with 5–11 leaflets.	7
Lower leaves digitate, with 3 leaflets.	<i>P. norvegica</i>
7. Lower leaves with 7–11 leaflets; achenes ribbed with a corky enlargement on one side.	<i>P. paradoxa</i>
Lower leaves with 5–7 leaflets; achenes smooth.	<i>P. rivalis</i>
8. Plants glandular pubescent; petals white or yellowish; stamens 25–30.	9
Plants not usually glandular; petals yellow; stamens 5–20.	10
9. Cymes short, crowded; petals white, slightly longer than the sepals.	<i>P. arguta</i>
Cymes open; petals white to yellowish, as long as the sepals.	<i>P. glandulosa</i> var. <i>intermedia</i>
10. Leaves all or mostly pinnate, with the upper stem leaves sometimes trifoliate.	11
Leaves all trifoliate or digitate with 5–7 leaflets.	21
11. Leaflets serrate to shallowly lobed.	12
Leaflets dissected or incised more than halfway to midrib.	15
12. Leaflets green on both sides.	13
Leaflets gray or white tomentose below.	14
13. Plants with 0–2 stem leaves.	<i>P. drummondii</i>
Plants with 4–7 stem leaves.	<i>P. paradoxa</i>
14. Leaves white tomentose below.	<i>P. hippiana</i>
Leaves grayish pilose to hirsute below.	<i>P. pensylvanica</i>
15. Leaflets green on both sides.	<i>P. plattensis</i>
Leaflets white or a different shade of green below.	16
16. Leaflets pale green or grayish pilose to glandular below.	<i>P. pensylvanica</i>
Leaflets white tomentose below.	17
17. Leaflets pectinately toothed or lobed; leaf margins revolute.	18

- Leaflets not pectinately incised; leaf margins flat. 19
18. Stipules of the upper stem leaves ovate, more or less lacinate. *P. bipinnatifida*
 Stipules of the upper stem leaves linear to lanceolate, entire. *P. multifida*
19. Leaflets 5; flowers solitary; plants cushion-forming. *P. concinna*
 Leaflets 3–5; plants usually several-flowered. 20
20. Plants low, tufted; basal leaves usually digitate. *P. nivea*
 Plants tufted; stems to 20 cm high; basal leaves pinnate. *P. saximontana*
21. Leaves all or mostly trifoliolate. 22
 Leaves all or mostly digitate. 24
22. Leaves wedge-shaped, 3-toothed at apex. *P. sibbaldii*
 Leaves not wedge-shaped, with more than 3 teeth or lobes. 23
23. Leaflets densely and more or less white tomentose below. *P. nivea*
 Leaflets green below; stem leaves one or none. *P. flabellifolia*
24. Leaflets densely grayish or white tomentose below. 25
 Leaflets green below, and less densely pubescent. 29
25. Stems leafy above and in the inflorescence. *P. argentea*
 Stems with 0–3 leaves. 26
26. Stems usually less than 10 cm high; plants densely tufted. 27
 Stems usually more than 10 cm high; plants not densely tufted. 28
27. Flowers solitary; basal leaves usually digitate. *P. concinna*
 Flowers 1 to several; basal leaves usually 3-foliolate. *P. nivea*
28. Leaflets commonly 3–5; plants usually less than 25 cm high. *P. quinquefolia*
 Leaflets commonly 5–9; plants usually more than 25 cm high. *P. gracilis*
29. Petals minute; flowers in a leafy cyme. *P. rivalis*
 Petals large; inflorescence leafy only at base. 30
30. Stem leaves 4 or more. *P. recta*
 Stem leaves 1–3 below the inflorescence. 31

31. Leaflets entire below the middle, dentate
 above. *P. diversifolia* var. *glaucophylla*
 Leaflets dentate along the margins. *P. gracilis*

Potentilla anserina L.

SILVERWEED

A low tufted perennial plant, spreading by runners. Leaves pinnate, 10–45 cm long, with 7–25 leaflets, often with some smaller interspersed leaflets. In the typical form, the leaflets are green above and silky white woolly beneath, but in forma *sericea* (Hayne) Hayek the leaves are silvery white on both sides. Both the type and the form have been found in the same clump of plants. Flowers bright yellow, 20–25 mm across, borne singly on a long stalk, blooming early in season until fall. Very common; in low, wet places and slough margins; throughout the Prairie Provinces.

Potentilla argentea L.

SILVERY CINQUEFOIL

Perennial with white woolly freely branching stems 10–50 cm high, ascending or erect. Lower leaves mostly 5-digitate, with the leaflets white tomentose on underside, toothed to pinnatifid above the middle and the margins revolute. Cyme leafy, many-flowered; petals yellow, about the same length as the sepals. Roadsides and waste areas; eastern Prairies, Parklands, and Boreal forest.

Potentilla arguta Pursh

WHITE CINQUEFOIL

Perennial glandular erect herbs 30–90 cm high, with pinnate leaves; the lower leaflets smaller than the upper ones. Leaflets 7–11 on the stalked basal leaves, fewer on the stem leaves. Flowers in a rather dense inflorescence, 12–20 mm in diam; petals white. Common; in moist places, slough margins on prairie; throughout Prairies and Parklands. Syn.: *Drymocallis agrimonioides* (Pursh) Rydb.

Potentilla bipinnatifida Dougl.

PLAINS CINQUEFOIL

A many-stemmed perennial 20–50 cm high, with pinnate leaves of 3–7 leaflets. Leaflets silky and green above, snowy white woolly beneath, deeply cut into narrow lobes. Fairly well distributed; on dry prairies; across Prairies and Parklands. Syn.: *P. pensylvanica* L. var. *bipinnatifida* (Dougl.) T. & G.

Potentilla concinna Richardson

EARLY CINQUEFOIL

A low-growing perennial from a very coarse, woody rootstock, with 5 (rarely 7) oblong or obovate leaflets, either palmately or pinnately arranged. Leaflets 12–30 mm long and toothed, greenish silky above and whitish below. Flowers yellow, 6–12 mm across, opening early in the season. Common; on dry hillsides and prairie; throughout Prairies, Parklands, and southern Rocky Mountains.

Potentilla diversifolia Lehm. var. *glaucophylla* Lehm.

SMOOTH-LEAVED CINQUEFOIL

A perennial plant 15–50 cm high, with palmately lobed leaves; the leaflets green and almost hairless on both sides. Leaflets usually coarsely toothed along the upper two-thirds of their margins, with forward-pointing teeth. A

mountain species found in the southern Rocky Mountains and occasionally in the Cypress Hills.

Potentilla drummondii Lehm.

DRUMMOND'S CINQUEFOIL

Plants with a short caudex; the stems 30–60 cm high, more or less hirsute. Basal leaves 5–10 cm long, more or less hirsute; leaflets 2–5 cm long, deeply incised. Flowers in a cyme, long-pedicelled; petals 6–10 mm long. Mountain meadows, riverbanks; southern Rocky Mountains.

Potentilla flabellifolia Hook.

MOUNTAIN CINQUEFOIL

Plants with branching, scaly rootstocks; the stems slender, 15–30 cm high, minutely puberulent. Basal leaves 3-foliolate, very thin; leaflets fan-shaped, deeply incised to serrate, short-pubescent. Cymes few-flowered; petals 8–10 mm long. The var. *emarginata* (Pursh) Boiv. includes more densely pubescent plants, about 10 cm high, and the marginal teeth with tufts of hairs. Moist mountain meadows; southern Rocky Mountains.

Potentilla fruticosa L. ssp. *floribunda* (Nutt.) Elk. (Fig. 128)

SHRUBBY CINQUEFOIL

A much-branched shrub from branching rootstocks, 15–150 cm high. Leaves pinnate, leathery, with 5–7 leaflets, 12–25 mm long, linear-oblong and pointed at both ends. Flowers yellow, 15–25 mm across. Achenes densely hairy. Much used as a garden ornamental. Very abundant on low moist ground; especially on Cypress Hills – Wood Mountain bench and in the Rocky Mountains, but also found across the Prairie Provinces. Syn.: *Dasiphora fruticosa* (L.) Rydb.; *P. fruticosa* Auct.; *P. floribunda* Pursh.

Potentilla glandulosa Lindl. var. *intermedia* (Rydb.) C. L. Hitchc.

STICKY CINQUEFOIL


Plants with slender, viscid, and glandular hirsute stems 15–30 cm high. Basal leaves with 5–9 leaflets, glabrous above, or nearly so, sparingly glandular hirsute below; leaflets 1–3 cm long, obovate, serrate; stem leaves reduced. Cymes many-flowered; sepals 6–7 mm long, glandular hirsute; petals about the same length as the sepals. Alpine slopes and mountain meadows; southern Rocky Mountains.

Potentilla gracilis Dougl.

GRACEFUL CINQUEFOIL

A somewhat tufted species with root crowns bearing brownish remains of old stipules, and several stems 30–60 cm high. Basal leaves long-stalked and digitate, of 5–7 narrowly oblanceolate leaflets. Flowers yellow, 15–20 mm across. A very variable plant, some of its varieties have been considered separate species. Two varieties are distinguished here.

Leaflets oblanceolate, serrate to pinnatifid,
green silky to white tomentose below. var. *gracilis*
Leaflets fan-shaped, deeply incised, densely
tomentose below. var. *flabelliformis* (Lehm.) Nutt.



A. C. Budd

Fig. 128. Shrubby cinquefoil, *Potentilla fruticosa* L. ssp. *floribunda* (Nutt.)
Elk.

Both varieties occur throughout the Prairie Provinces, north to the southern parts of the Boreal forest; var. *flabelliformis* is usually found in swampy or somewhat alkali meadows; var. *gracilis* in moist grasslands, slopes, and thickets.

Potentilla hippiana Lehm.

WOOLLY CINQUEFOIL

A perennial from a stout rootstock, with several woolly stems 10–25 cm high, and mostly basal pinnate leaves with 7–11 leaflets 1–5 cm long, toothed, densely silky above and white woolly below. Stem leaves smaller, with fewer leaflets. Flowers fairly numerous, in a terminal cyme 6–10 mm across. Found fairly often on prairie land and in valleys in southwestern parts of the Prairie Provinces. The var. *argyrea* (Rydb.) Boivin, silvery cinquefoil, has leaflets white woolly on both upper and lower sides; found occasionally on dry hill-sides; in the Prairies and in the Cypress Hills.

Potentilla multifida L.

BRANCHED CINQUEFOIL

Plants with strigose stems 20–40 cm high. Basal leaves with 7 leaflets; stem leaves with 5–7 leaflets; the leaflets deeply incised, green below, with revolute margins. Cymes few- to many-flowered; the petals 6–10 mm long. Open areas and gravel banks; Boreal forest.

Potentilla nivea L.

SNOW CINQUEFOIL

A caespitose perennial with a stout caudex; the flowering stems 10–20 cm high. Leaves mostly basal, 3-foliolate or rarely 5-digitate; leaflets 1–3 cm, oblong or obovate, densely white tomentose beneath, deeply 3- to 5-dentate on each side. Flowers 10–15 mm across. Alpine, subarctic, and arctic meadows; Boreal forests, Rocky Mountains. Besides the typical variety described, two other varieties occur: var. *villosa* (Pall.) Regel & Tiling, more coarsely and densely villous throughout, including the upper surface of the thick, rugose leaflets; and var. *pulchella* (R. Br.) Durand, with the leaves mostly short-pinnate with 5 coarse leaflets, and rather large, brown stipules.

Potentilla norvegica L.

ROUGH CINQUEFOIL

A very coarse hairy annual or biennial plant, with erect branched stems 15–60 cm high. Leaves digitately 3-foliolate, with leaflets 2–10 cm long, obovate or elliptic, and much-toothed. Flowers numerous, 6–12 mm across, yellow, in a fairly dense cyme; the sepals a little longer than the petals. A common plant; moist meadows and waste places, a bad weed in gardens; throughout the Prairie Provinces. Syn.: *P. monspeliensis* L.

Potentilla palustris (L.) Scop.

MARSH CINQUEFOIL

A somewhat decumbent herb 20–50 cm high, from long, creeping, weedy rootstocks. Leaves pinnate, often purple; the lower ones long-stalked, with 5–7 oblong or oval leaflets 2–7 cm long, tapering to the base and toothed. Upper leaves short-stalked, with 3–5 leaflets. Flowers conspicuous, 15–35 mm across, purple or maroon; the sepals much longer than the petals and also purplish. Found in shallow water and bogs; in Boreal forest and rarely in Parklands. Syn.: *Comarum palustre* L.

Potentilla paradoxa Nutt.

BUSHY CINQUEFOIL

An annual, biennial, or short-lived perennial species, with spreading or somewhat ascending stems 20–50 cm high. The pinnate leaves of 5–11 almost smooth leaflets. Flowers about 6 mm across, in a leafy cluster. Achenes or fruits ribbed lengthwise and having a corky enlargement on one side. Found occasionally in low moist places; throughout the Prairie Provinces.

Potentilla pensylvanica L. (Fig. 129)

PRAIRIE CINQUEFOIL

A low-growing tufted species, 10–40 cm high, with the crown usually bearing the brown remains of previous leaf stipules. Two varieties of this species occur throughout the Prairies and Parklands.

Hairs on stem very short and appressed.

Leaflets grayish and hairy on underside. var. *pensylvanica*

Hairs on stem obvious and spreading. Leaflets

lobed to about halfway to midrib. var. *atrovirens* (Rydb.) T. Wolf

The var. *atrovirens* common in dry grasslands in the Prairies; var. *pensylvanica* common in moister grasslands of the Prairies, Parklands, and Cypress Hills.

Potentilla plattensis Nutt.

LOW CINQUEFOIL

A low decumbent or spreading perennial 10–20 cm high, with many basal pinnate leaves of 9–17 leaflets. Leaflets light green, deeply divided into oblong to nearly linear divisions. Flowers few and in rather open terminal cymes; petals yellow. Found occasionally in valleys; Prairies.

Potentilla quinquefolia Rydb.

FIVE-FINGERED CINQUEFOIL

A low tufted perennial 10–20 cm high, with small leaves, palmately divided into 3–5 leaflets. Leaflets green above and white woolly beneath, deeply divided into narrow lobes. Flowers about 1 cm across, in a fairly dense cluster. Quite rare; but has been found in Parklands and Boreal forest.

Potentilla recta L.

ROUGH-FRUITED CINQUEFOIL

A leafy-stemmed erect plant 15–50 cm high, with loosely hairy stems and leaf stalks. The deeply toothed leaves are digitate, usually of 5–7 leaflets, more or less hairy, but paler on the underside. Flowers on erect stalks 15–25 mm across, pale yellow or sulfur. This introduced plant has been found at various places throughout the Prairie Provinces.

Potentilla rivalis Nutt.

BROOK CINQUEFOIL

An erect annual or biennial species, branching above, and with finely hairy, sometimes sticky, stems 20–40 cm high. The lower leaves have 5 obovate coarsely toothed leaflets 2–5 cm long; upper stem leaves have 3 leaflets. Yellow flowers less than 5 mm across, with petals shorter than sepals. Many authorities consider this species and *P. millegrana* as forms of the same species, differing mainly in the number of leaflets on the basal leaves. Found occasionally; in river valleys; Prairies.



Fig. 129. Prairie cinquefoil, *Potentilla pensylvanica* L.

A low perennial with stout caudex; the stems 10–20 cm high, glabrous to pubescent. Leaves mostly basal, pinnate, usually with 5 leaflets 1–5 cm long, deeply cleft, the divisions narrow, white tomentose or strigose below, silky hairy above. Flowers few to several, 10–15 mm across. Alpine slopes; southern Rocky Mountains.

Potentilla sibbaldii Hall. f.

A low densely tufted herb with short decumbent or creeping woody stems. Leaves 3-foliolate, with slender petioles; leaflets 3- to 5-toothed at the apex, obovate, cuneate at the base, sparsely pubescent on both surfaces. Flowers on axillary peduncles; sepals 2–3 mm long; petals minute, yellow. Alpine slopes and scree fields; Rocky Mountains.

Potentilla tridentata Ait.

A tufted, low-growing, woody-based subshrub, much-branched, 10–20 cm high. Stems have silky, appressed hairs. Leaves mostly stalked with 3 narrow wedge-shaped leaflets 12–25 mm long, dark green and shiny on the upper side and paler beneath, having 3 teeth at their apex. Flowers 1–6 in a cyme, white, about 6–8 mm across. Fairly common; in dry sandy places, in pine forests; Boreal forest. Syn.: *Sibbaldiopsis tridentata* (Ait.) Rydb.

Prunus plum

Shrubs or trees with bark almost black; alternate leaves; perfect regular flowers with 5 petals and 5 sepals; fruit a fleshy drupe with a smooth bony stone.

- 1. Flowers in long racemes at the ends of the branches. 2
- Flowers in small umbels or corymbs. 3
- 2. Fruit red or purplish, very astringent; leaves thin. *P. virginiana*
- Fruit black and sweetish; leaves thick. *P. virginiana* var. *melanocarpa*
- 3. Stone more or less flattened, with a groove on the end. 4
- Stone not flattened or grooved. 5
- 4. Teeth of leaves not very deep and tipped with a small gland; lobes of sepals toothed and glandular; fruit oval. *P. nigra*
- Teeth of leaves deeper and not gland-tipped; lobes of sepals not toothed or glandular; fruit round. *P. americana*
- 5. Tall shrub or small tree; flowers 12–25 mm across, in corymbs. *P. pensylvanica*
- Low shrub; flowers 6–12 mm across, in umbels. *P. pumila*

Prunus americana Marsh.

AMERICAN PLUM

A tree 3–8 m high, with more or less thorny branches. Leaves narrowly obovate, 3–10 cm long, with double teeth and a pointed apex. Flowers white, 15–25 mm across, appearing before the leaves. Fruit a red or yellow plum, almost round, 15–25 mm long. Found occasionally; in moist woods and along riverbanks; southeastern Parklands and Boreal forest.

Prunus nigra Ait.

CANADA PLUM

A tree or shrub 2–8 m high, with oval or obovate leaves 7–12 cm long, pointed at the apex, dark green with gland-tipped teeth. Flowers white, turning pink, about 15–30 mm across, opening before the leaves. Fruit a yellow to orange plum, oval, 25–35 mm long. Found occasionally in woodlands and bluffs; southeastern Parklands and Boreal forest.

Prunus pensylvanica L. f.

PIN CHERRY

A small tree 4–8 m high, with lanceolate finely toothed leaves 3–10 cm long. Flowers small, white, on long stalks in corymbose clusters 6–10 mm across, opening about the same time as the leaves. Fruit a small, sour, bright red cherry, 5–8 mm in diam. Fairly common; in bluffs, ravines, and hillsides; throughout the Prairie Provinces.

Prunus pumila L.

LOW SAND CHERRY

A low, spreading bush with oblanceolate to spatulate leaves, dark green above and pale beneath. Flowers in clusters of 2–4, white, about 1 cm across. Fruit a dark brown to purplish cherry, 6–12 mm across. Found on dry prairie and in sandhills; southeastern Parklands and Boreal forest.

Prunus virginiana L.

RED-FRUITED CHOKE CHERRY

A shrub 1–3 m high, with grayish stems and thin ovate leaves 5–10 cm long. Flowers white, about 12 mm across, in loose racemes. Fruit a red, very astringent cherry 8–10 mm in diam. Quite common; on riverbanks and open woodlands; in the southeastern Parklands. It is replaced further west by the var. *melanocarpa*.

Prunus virginiana L. var. *melanocarpa* (A. Nels.) Sarg. (Fig. 130)

BLACK-FRUITED CHOKE CHERRY

A small tree or large shrub 2–6 m high, with smooth reddish brown stems. Leaves obovate or oval, rather thick, 2–8 cm long, smooth on both sides. Flowers numerous, white or pale cream, about 1 cm across, in dense racemes. Fruit a black cherry, slightly astringent, round, 6–8 mm in diam. Very common; in bluffs, ravines, sandhills, and open woodlands; throughout the Prairie Provinces. Leaves of choke cherry injured by frost or extreme drought are reported to be **poisonous** to sheep and cattle.

Rosa rose

From low shrubs or herbaceous plants to large bushes, usually with prickly stems. Leaves pinnate, with odd number of small leaflets. Flowers large and showy, generally pinkish and fragrant. Fruit a hip, or berry-like enlarged




Fig. 130. Black-fruited choke cherry, *Prunus virginiana* L. var. *melanocarpa* (A. Nels.) Sarg.

calyx tube containing numerous achenes. A difficult genus to identify, local variations in form being so common that authorities do not agree on nomenclature, and some specialists have separated them into many species. Probably only about 4 species, with many slight variations, occur in the Prairie Provinces.

1. Plant with few, if any, bristles. *R. blanda*
 Plant definitely bristly. 2
2. Thorns present below the stipules; bristles
 broad and usually flattened at the base;
 fruit globose without constricted neck. *R. woodsii*
 Thorns not present; bristles usually not
 broad or flattened at base. 3
3. Plant shrubby; leaflets usually 5–9 (9 on
 young shoots); flowers usually borne
 singly; fruit oval with distinct neck. *R. acicularis*
 Plant partly shrubby, often dying back
 close to the ground in winter; leaflets
 9–11. *R. arkansana*

Rosa acicularis Lindl.

PRICKLY ROSE

A low bushy plant 30–120 cm high, with stems densely covered with straight weak bristles. Leaves usually of 5–7 hairy leaflets 12–50 mm long, elliptic or oval. Stipules broad, usually both hairy and glandular. Flowers usually borne singly, 5–7 cm across. Fruit usually ovoid with a constricted neck, about 1.5 cm long. Common; in bluffs, around woods, in fields, and along roadsides; throughout the Prairie Provinces.

Rosa arkansana Porter

LOW PRAIRIE ROSE

Low shrubs 20–30 cm high with densely prickly stems, which usually die off annually close to the root. Leaflets 9–11, usually without hairs, smooth and shiny, 2–5 cm long. Flowers in corymbs of 2 or 3 flowers. Fruit almost globular, about 12 mm across. Common; on hills and sandy open prairie; throughout the Prairies.

Rosa blanda Ait.

SMOOTH ROSE

A low bush 60–120 cm high, with an unarmed stem or with a very few straight prickles. Stipules rather broad; leaves of 5–7 leaflets 25–40 mm long. Flowers pink, usually borne singly, may be up to 7 cm across. Not common; but has been found in eastern Parklands.

Rosa woodsii Lindl.

WOOD'S ROSE

A bush 50–200 cm high, with the stems armed with straight or slightly curved prickles, often broad and flattened at the base. Spines below the stipules usually well-defined. Leaves of 5–9 oval or obovate leaflets 15–35 mm long. Flowers 2–5 cm across. Fruit globular without a constricted neck, about 1 cm across. In bluffs, on ravines, and in sandhills; throughout the Prairies and Parklands.

Rubus raspberry

Perennial or biennial, shrubby or herbaceous, from rootstocks, often with prickly stems. Leaves various, from simply lobed to 3–5 times pinnately or palmately compound. Flowers regular, unisexual or perfect, usually with 5 sepals and 5 petals, either pink or white. Fruit a berry composed of many fleshy drupelets.

1. Stems more or less woody; leaves 3–5 foliolate, usually pinnate or lobed. 2
Stems herbaceous, dying down annually, not prickly; leaves with 3 leaflets or merely lobed. 3
2. Stems prickly as well as bristly; leaves compound. *R. idaeus* var. *aculeatissimus*
Stems not prickly; leaves lobed, 10–20 cm wide. *R. parviflorus*
3. Leaves merely lobed; petals spreading; flowers unisexual. *R. chamaemorus*
Leaves compound; petals erect. 4
4. Leaves 5-digitate. *R. pedatus*
Leaves 3-foliolate. 5
5. Central leaflet with rounded tip. *R. acaulis*
Central leaflet with pointed tip. *R. pubescens*

Rubus acaulis Michx.

STEMLESS RASPBERRY

A low, herbaceous, unarmed perennial 5–20 cm high, with pinnately divided leaves of 3 broadly ovate leaflets 15–35 mm long. Flowers usually borne singly, pink; fruit red. Only found in bogs and meadows; in Boreal forest.

Rubus chamaemorus L.

CLOUDBERRY

A herbaceous perennial 5–20 cm high, with 2 or 3 stalked round leaves 2–7 cm across, each with 5–9 rounded lobes. Flowers borne singly, terminal, white, 10–25 mm across; fruit reddish, turning golden yellow when ripe. Found in bogs and swamps on Boreal forest.

Rubus idaeus L. var. *aculeatissimus* Regel & Tiling (Fig. 131)

WILD RED RASPBERRY

A large bush 1–2 m high, with brownish more or less bristly stems and pinnate leaves with 5 leaflets (the floral branches with 3 leaflets). Leaflets ovate, the terminal one often being 3-lobed, 5–10 cm long, dark green above and white woolly beneath. New shoots fairly bristly, but rarely glandular. Flowers white, 8–12 mm across; fruit round, light red, about 1 cm across. Probably the most common raspberry; in shady, wooded places, on burned-over woodlands, bluffs, riverbanks; throughout the Prairie Provinces. Syn.: *R. melanolasius* Focke; *R. strigosus* Michx.




Fig. 131. Wild red raspberry, *Rubus idaeus* L. var. *aculeatissimus* Regel & Tiling.

Rubus parviflorus Nutt.

THIMBLEBERRY

An erect shrub 50–200 cm high, unarmed, with the young growth more or less glandular hairy. Leaves 10–20 cm wide, simple, round to reniform in outline, few-lobed, with the lobes triangular, glabrous or pubescent below. Flowers 2–4 cm across, white; fruit red, 15–20 mm across. Forest margins and openings; common in southern Rocky Mountains, rarer in northern Rocky Mountains and Cypress Hills.

Rubus pedatus J. E. Smith

DWARF BRAMBLE

A low or creeping unarmed herbaceous species, with glabrous stems rooting at the nodes, to 1 m long. Branches short; leaves 1–4, digitately 5-foliolate, or rarely 3-foliolate; leaflets 1–5 cm long, obovate, irregularly incised, glabrous or very sparsely pubescent. Flowers solitary on slender peduncles 4–10 cm long; sepals green, 8–10 mm long; petals white, the same length as the sepals; fruit reddish, 8–10 mm long. Moist woods; Rocky Mountains, western Parklands, and Boreal forest.

Rubus pubescens Raf.

DEWBERRY

A trailing or climbing herbaceous plant, with pinnate leaves usually of 3 leaflets (rarely 5), ovate or rhombic, 2–10 cm long, sharply toothed, and green on both sides. Flowers borne in groups of 2 or 3 with white or pink petals, 8–12 mm across, with reflexed sepals. Fruit reddish purple, about 1 cm across. Common; in rich woods and bluffs; Parklands, Boreal forest, and Cypress Hills.

Sorbaria false spiraea

Sorbaria sorbifolia (L.) A. Braun

ASH-SPIRAEA

Shrubs to 2 m high; young branches tomentose with stellate hairs. Leaves pinnate; leaflets 3–7 cm long, acuminate, sharply double serrate. Inflorescence a panicle to 30 cm long, flowers white, 5–6 mm across. Introduced ornamental, occasionally escaped and persisting.

Sorbus rowan

- 1. Trees 6–10 m high. *S. aucuparia*
- Shrubs or small trees less than 5 m high. 2
- 2. Leaflets usually 11–13, serrate to close to the base; inflorescence broad, flat-topped. *S. americana*
- Leaflets usually 7–11, entire toward the base; inflorescence small, rounded. *S. sitchensis*

Sorbus americana Marsh.

WESTERN MOUNTAIN ASH

A small tree 1–4 m high, with pinnate leaves of 11–13 elliptic-lanceolate leaflets 3–6 cm long, bright green on both sides. Flowers white, in dense terminal clusters, 6–10 mm across, with the flower clusters up to 10 cm broad; fruit

globose, red, about 1 cm across. Not common; but found in Parklands, Boreal forests, Riding Mountain, and Cypress Hills. Syn.: *Sorbus scopulina* Greene.

Sorbus aucuparia L.

ROWAN TREE

A tree to 10 m high, round-topped, with young branches more or less pubescent; winter buds white villose. Leaflets 9–15, oblong, 3–5 cm long, serrate, usually long-pubescent below. Inflorescence to 20 cm broad, flat-topped; fruit bright orange red. Introduced from Europe, extensively planted, and occasionally escaped.

Sorbus sitchensis Roemer

MOUNTAIN ASH

A tall shrub, usually 1–3 m high, with stems dull brown, sparsely branched; twigs pubescent with red brown hairs. Leaves commonly with 7–11 leaflets, 2–4 cm long, serrate above the middle. Inflorescence small, round-topped; fruit orange, red, or purplish. Woods and openings in mountains; southern Rocky Mountains.

Spiraea meadowsweet

Shrubs with simple leaves and no stipules. Flowers perfect, with 5 sepals and 5 petals; fruit a follicle opening along one side.

1. Inflorescence flat-topped. 2
 Inflorescence elongate. *S. alba*
2. Petals pink or roseate; inflorescence
 small, dense. *S. densiflora*
 Petals white; inflorescence broad, open. *S. lucida*

Spiraea alba Du Roi

NARROW-LEAVED MEADOWSWEET

An erect shrub 50–100 cm high, with brown twigs and narrowly oblanceolate leaves, pointed at both ends, 3–6 cm long, sharply toothed, and sometimes with fine hairs beneath on the veins. Inflorescence finely hairy in dense terminal panicles; flowers small and white. Fairly common; Parklands and Boreal forest.

Spiraea densiflora Nutt.

PINK MEADOWSWEET

A slender shrub 50–150 cm high, with branches ascending, reddish brown. Leaves oval to elliptic, 15–30 mm long, rounded at base and tip, crenate or serrate above the middle, glabrous or nearly so. Inflorescence 2–5 cm broad, dense; flowers showy, very sweet-scented, with petals about 1.5 mm long. Moist thickets and meadows; southern Rocky Mountains.

Spiraea lucida Dougl.

SHINING-LEAVED MEADOWSWEET

A low shrub 30–100 cm high; stems and branches erect, usually dying down to close to the base annually. Leaves 2–5 cm long, obovate to oval, on short stalks, shiny green above, paler below. Flowers white, small, in a flat-topped panicle. Common; in the Cypress Hills and Rocky Mountains.

LEGUMINOSAE—pea family

Shrubs or herbs with alternate compound leaves with stipules. Flowers (see Fig. 7) perfect and irregular, having 5 more or less united sepals and 5 petals. The upper petal, larger than the others, is called the standard; the two side petals are called wings, the two lower ones, united, form the keel. The pistil and stamens are contained in the keel. The stamens, usually 10, are arranged in various groupings, their arrangement being one of the characters used in separating the genera of this large family. They can all be distinct and separate, or all joined in one group, or diadelphous (separated into two groups of 9 and 1). The fruit is either a 1- or 2-compartment pod (legume) or a pod constricted between the seeds (loment).

- | | |
|--|-------------------------------|
| 1. Shrubs. | 2 |
| Herbs. | 3 |
| 2. Flowers yellow; leaves pinnate, leaflets even-numbered, the terminal leaflet lacking. | <i>Caragana</i> |
| Flowers purple; leaves pinnate, leaflets odd-numbered, the terminal leaflet present. | <i>Amorpha</i> |
| 3. The terminal leaflet replaced by a tendril. | 4 |
| The terminal leaflet normal. | 5 |
| 4. Styles filiform, bearded at the tip; wings of flower coherent with the keel. | <i>Vicia</i> |
| Styles flattened, bearded along the inside; wings free from the keel. | <i>Lathyrus</i> |
| 5. Leaves palmately divided. | 6 |
| Leaves pinnately divided. | 16 |
| 6. Plants very low cushion plants. | 7 |
| Plants not cushion plants. | 8 |
| 7. Flowers white, 2 cm long. | <i>Astragalus gilviflorus</i> |
| Flowers purple, 6–8 mm long. | <i>Astragalus spatulatus</i> |
| 8. Leaves glandular-dotted. | <i>Psoralea</i> |
| Leaves not glandular-dotted. | 9 |
| 9. Leaflets clearly toothed. | 10 |
| Leaflets entire. | 12 |
| 10. Pods strongly curved or coiled. | <i>Medicago</i> |
| Pods straight or almost straight. | 11 |
| 11. Flowers in long spike-like racemes; the deciduous petals free from the stamen tube. | <i>Melilotus</i> |
| Flowers in short loose or dense heads; the petals more or less adhering to the stamen tube. | <i>Trifolium</i> |
| 12. Stems twining on vegetation; upper flowers with petals, the lower ones apetalous. | <i>Amphicarpa</i> |

- Stems not twining; flowers all alike. 13
13. Pods usually with 3–5 one-seeded segments,
 which separate at maturity. *Desmodium*
 Pods not segmented. 14
14. Flowers about 4 mm long, usually solitary
 in the leaf axils. *Lotus*
 Flowers 10–20 mm long, in spikes or racemes. 15
15. Plants with rootstocks; flowers yellow;
 pods 4–7 cm long, sickle-shaped. *Thermopsis*
 Plants tufted; flowers purple or yellow;
 pods 2–4 cm long, straight. *Lupinus*
16. Pods segmented or jointed, breaking up at maturity. 17
 Pods not segmented or jointed. 18
17. Inflorescence a loose few-flowered head-like umbel. *Coronilla*
 Inflorescence an elongated raceme. *Hedysarum*
18. Leaves glandular-dotted. 19
 Leaves not glandular. 20
19. Pods with hooked prickles; flowers 10–15 mm long. *Glycyrrhiza*
 Pods not prickly; flowers small
 to 5 mm; stamens 5. *Petalostemon*
 Pods not prickly; flowers 10–12 mm
 long, purple with yellowish base; stamens 10. *Oxytropis viscida*
20. Keel tipped with a point. *Oxytropis*
 Keel not tipped with a point. *Astragalus*

Amorpha false indigo

Shrubs with pinnate leaves, bearing odd number of glandular-dotted leaflets; midrib projecting from end of leaflets. Flowers having the standard petal only, other petals missing. Flowers borne in long narrow spikes. Fruit a short, oblong, curved, 1- or 2-seeded pod.

1. Plants densely pubescent. *A. canescens*
 Plants sparsely pubescent to glabrous. 2
2. Low shrubs; leaflets to 1 cm long. *A. nana*
 Tall shrubs; leaflets to 3 cm long. *A. fruticosa*

Amorpha canescens Pursh

LEADPLANT

A bushy shrub, densely white hairy, 30–100 cm high. Leaves very dense, 5–10 cm long, with 21–51 oval leaflets, each 8–12 mm long. Flowers in dense, clustered, spike-like racemes 5–15 cm long, small, with a bluish purple petal. Pods single-seeded, about 3–5 mm long. Found occasionally on dry prairies; southeastern Parklands.

Amorpha fruticosa L.

FALSE INDIGO

A fairly tall shrub 2–5 m high, with stalked leaves 15–40 cm long and 11–25 oval leaflets 10–30 mm long. The spike-like racemes of flowers either

clustered or borne singly, 7–15 cm long, with a violet purple petal. Found occasionally on riverbanks; southeastern Parklands.

Amorpha nana Nutt.

DWARF FALSE INDIGO

A low shrub, usually less than 30 cm high, almost hairless, with numerous leaves 2–8 cm long, each having 13–31 stiff elliptic leaflets 6–12 mm long. Spike-like racemes of flowers borne singly, 2–7 cm long, fragrant with a purplish petal. Not common; moist prairie and waste places; southeastern Parklands.

Amphicarpa hog-peanut

Amphicarpa bracteata (L.) Fern.

HOG-PEANUT

Twining plants with stems to 1 m long, pubescent with retrorse appressed hairs. Leaflets thin, somewhat pubescent, with the terminal one 2–6 cm long. Racemes with 2–10 flowers, white to pale purplish; pods sparsely pubescent, mainly at maturity, those of the upper flowers 3-seeded, those of the basal ones 1-seeded and often underground. Not common; woods along rivers and in ravines; southeastern Parklands and Boreal forest.

Astragalus milk-vetch

Perennial herbs with pinnate leaves of 3 to many leaflets, and perfect irregular flowers having a blunt keel. Stamens 10, diadelphous (9 being united in a bundle and 1 separate). Flowers in spikes or racemes. Fruit a pod, sometimes only 1-celled and often appearing 2-celled because of the ingrowth of one or both of the joints between the halves of the pods. *Astragalus* is of considerable importance in the Prairie Provinces, most of the species occurring in the native grasslands and woodlands, where they are accessible to grazing livestock. Several species are **poisonous** or potentially poisonous to livestock, and although poisoning by these species seems to be rare, in periods of drought or under other unfavorable conditions they can be dangerous. In the Rocky Mountains, *A. miser*, and possibly some other species, produce an alkaloid that causes partial paralysis, which has led to the name “locoweed,” because of the erratic movements of affected animals. Other species, such as *A. pectinatus* and *A. bisulcatus*, can accumulate the element selenium, which can cause sickness or death when eaten.

Some authors have divided the genus *Astragalus* into several small genera, or sections, by using the characters of the legumes. The legumes (or pods) vary from short to long, compressed to inflated, papery to woody, 1-celled to 2-celled, and few to many seeds. The inflorescence is a raceme with few to many flowers more or less long peduncled, and axillary in the leaf axils.

- 1. Plants low, cushion-like, matted or straggling 2
- Plants with erect or decumbent stems. 12
- 2. Leaflets spine-tipped, plants prickly. *A. kentrophyta*
- Leaflets not spine-tipped. 3

3. Flowers white, borne in the crown of the plant; leaves with 3 leaflets, densely villous. *A. gilviflorus*
 Flowers purplish or yellowish, borne on more or less well developed peduncles. 4
4. Leaves usually with a single leaflet, occasionally with 3 or 5 leaflets; flowers purple. *A. spatulatus*
 Leaves with at least 7 leaflets. 5
5. Flowers yellowish, with or without purplish tips. 6
 Flowers purple or blue, occasionally whitish. 8
6. Plants cushion-like; leaves silky villous, tomentose; flowers about 2 cm long. *A. purshii*
 Plants not cushion-like; stems spreading. 7
7. Plants large; flowers about 2 cm long; leaves with 17–25 leaflets; pods glabrous. *A. crassicarpus*
 Plants small; flowers about 1 cm long; leaves with 7–15 leaflets; pods densely pubescent. *A. lotiflorus*
8. Plants with taproots and caudex. 9
 Plants with rootstalks. 11
9. Flowers about 2 cm long; leaves silvery pubescent. *A. missouriensis*
 Flowers about 1 cm long; leaves not silvery pubescent. 10
10. Stems long, few-leafed; leaflets 15–21; inflorescence elongating at maturity; pods cylindrical. *A. flexuosus*
 Stems short, densely leafed; leaflets 7–11; inflorescence not elongating; pods flattened. *A. vexilliflexus*
11. Inflorescence a dense, head-like raceme; flowers and pods erect; pods densely pubescent. *A. danicus*
 Inflorescence a short, loosely flowered raceme; flowers spreading; pods reflexed, black hairy. *A. alpinus*
12. Flowers white, yellow, or greenish. 13
 Flowers at least partly reddish, purplish, or bluish. 21
13. Leaflets very narrowly linear. *A. pectinatus*
 Leaflets not narrowly linear. 14
14. Plants densely hairy. *A. drummondii*
 Plants not densely hairy. 15

15. Inflorescence densely flowered.	16
Inflorescence loosely flowered.	19
16. Flowers ascending, greenish white; pods woody.	<i>A. canadensis</i>
Flowers ascending to spreading, creamy white or yellowish; pods not woody.	17
17. Flowers creamy white, spreading; pods triangular in cross section.	<i>A. racemosus</i>
Flowers greenish yellow or yellow; pods not triangular in cross section.	18
18. Flowers yellow; pods inflated, globular to ovoid, black pubescent.	<i>A. cicer</i>
Flowers greenish yellow; pods inflated, ellipsoid, glabrous.	<i>A. frigidus</i>
19. Leaflets glabrous or nearly so; pods oblong, pointed at both ends, drooping.	<i>A. tenellus</i>
Leaflets strigose, at least beneath; pods not pointed at both ends.	20
20. Pods sessile, linear-oblong, straight on both sutures, drooping.	<i>A. miser</i>
Pods long-stipitate, elliptic, with the lower suture arched, spreading, arched downward.	<i>A. aboriginum</i>
21. Inflorescence a short, dense, head-like raceme, about as long as wide; flowers and pods all erect.	22
Inflorescence not head-like, if densely flowered an elongated raceme; some flowers pendent.	23
22. Plants with slender rootstalks; pubes- cence of the leaflets composed of basi- fixed hairs.	<i>A. danicus</i>
Plants with taproots and caudex; pubes- cence of the leaflets composed of mal- pighian hairs.	<i>A. striatus</i>
23. Flowers brick red to purplish; inflorescence a loose raceme; pods strongly inflated.	<i>A. iochrous</i>
Flowers purplish to whitish purple; pods not inflated.	24
24. Plants low; stems erect or ascending, 10–30 cm high.	25
Plants up to 60 cm high; stems erect.	27
25. Plants with slender rootstalks.	<i>A. alpinus</i>
Plants with taproots or caudex.	26
26. Pods erect, 5–7 mm long; leaflets 7–15.	<i>A. yukonis</i>

- Pods drooping, 10–15 mm long; leaflets 13–23. *A. bourgovii*
27. Flowers in a dense raceme, somewhat reflexed; pods 2-grooved on the upper side. *A. bisulcatus*
- Flowers in loose racemes; pods not 2-grooved. 28
28. Pods stipitate, 1.5–2 cm long, spreading or arched downward. *A. robbinsii*
- Pods sessile, 1 cm long, drooping. *A. eucosmus*

Astragalus aboriginum Richardson

INDIAN MILK-VETCH

Plants with a stout taproot, few- to many-stemmed from a branched caudex. Stems 20–40 cm long, ascending or erect, densely and finely pubescent with spreading hairs. Leaves with 7–15 leaflets 1–3 cm long, elliptic to linear-lanceolate, pubescent on both sides. Inflorescence 2–3 cm long in flower, elongating to 10–15 cm in fruit; flowers 6–10 mm long, whitish, bluish purple tipped; calyx 3–4 mm long, black pubescent. Legumes 1.5–2.5 cm long, straight to somewhat falcate, flattened, glabrous to white pubescent, drooping, borne on a stipe 5–8 mm long. An almost glabrous form has been named var. *major* Gray. Gravel banks along rivers and on slopes; throughout the Prairie Provinces. Syn.: *A. aboriginorum* Richardson; *Atelophragma aboriginum* (Richardson) Rydb.

Astragalus alpinus L.

ALPINE MILK-VETCH

Plants with extensively creeping roots; stems 10–40 cm high, solitary or in small tufts, slender, decumbent or ascending, glabrous or nearly so. Leaves with 11–25 leaflets 1–2 cm long, oblong to oval, with the apex retuse or obtuse, glabrous above or somewhat pubescent on both sides. Inflorescence 3–4 cm long, at first dense, soon elongating to 5–8 cm in fruit; flowers 7–13 mm long, light purplish blue to almost white; calyx 3–4 mm long, black pubescent. Legume 8–12 mm long, straight or falcate, compressed, black pubescent, reflexed, stipitate. The var. *brunetianus* Fern. is distinguishable by having a short glabrescent calyx, 2–2.5 mm long, and a glabrescent pod. Forest margins, open and disturbed areas; Boreal forest and Rocky Mountains. Syn.: *Atelophragma alpinum* (L.) Rydb.

Astragalus bisulcatus (Hook.) Gray (Fig. 132A)

TWO-GROOVED MILK-VETCH

Plants densely tufted, often many-stemmed, erect; stems stout, 30–80 cm high, usually reddish to purplish, finely pubescent to glabrous. Leaves with 13–29 leaflets 1–3.5 cm long, oblong-elliptic, somewhat pubescent to glabrous. Inflorescence 10–18 cm long, dense; flowers 10–15 mm long, reddish purple, usually somewhat pendent; calyx about 4 mm long, inflated at base. Legume 18–22 mm long, pendent, linear oblong, the upper side deeply 2-grooved. Plant with an unpleasant odor. Prairies and Parklands. Syn.: *Diholcos bisulcatus* (Hook.) Rydb.



Fig. 132. *A*, Two-grooved milk-vetch, *Astragalus bisulcatus* (Hook.) Gray; *B*, narrow-leaved milk-vetch, *Astragalus pectinatus* Dougl.

Astragalus bourgovii Gray

Plants densely tufted from a stout caudex. Stems erect or ascending, 10–30 cm high. Leaves with 13–25 leaflets 5–12 mm long, lanceolate to oblong-elliptic, pubescent on both sides. Inflorescence lax, loosely flowered; flowers 8–11 mm long, bluish purple or violet; calyx 4–5 mm long, black pubescent. Legume 1–1.5 cm long, oblong elliptic, flattened, black pubescent. Montane and alpine meadows; Rocky Mountains.

Astragalus canadensis L.

CANADIAN MILK-VETCH

Plants with creeping roots; stems single or a few together. Stems stout, 40–100 cm high, glabrous or thinly pubescent. Leaves with 13–27 leaflets 2–4.5 cm long, elliptic to oblong, glabrous to thinly pubescent. Inflorescence dense, 8–10 cm long; flowers greenish white to white, 1.0–1.5 cm long, ascending to spreading; calyx 4–5 mm long, strigose. Legume 1.2–1.6 cm long, woody, glabrous or nearly so, terete, sessile. Moist areas, woodland; throughout the Prairie Provinces.

Astragalus cicer L.

CICER MILK-VETCH

Plants with creeping roots; stems single, 40–60 cm high. Leaves with 23–33 leaflets 5–20 mm long, pubescent on both sides. Inflorescence dense, 4–6 cm long, black pubescent; flowers yellow, 10–15 mm long; calyx 6–8 mm long, black pubescent. Legume 10–15 mm long, ovoid to globose, inflated, thin-walled, black pubescent. An introduced species, rare as a weed; Manitoba, Alberta.

Astragalus crassicaarpus Nutt.

GROUND-PLUM

Plants with a stout caudex; many-stemmed; stems procumbent, straggling, 20–35 cm long, finely pubescent. Leaves with 13–27 leaflets 8–20 mm long, oblong to linear, strigose below, glabrous above. Inflorescence 4–5 cm long, few-flowered; flowers 1.5–2.0 cm long, whitish with purplish tip or bluish purple; calyx 5–8 mm long, pubescent with black and white hairs. Legume subglobose, 1.5–2.5 cm in diam, deep reddish purple when ripe. Grasslands; Prairies and Parklands. Syn.: *A. caryocarpus* Ker; *Geoprimum crassicaarpum* (Nutt.) Rydb.

Astragalus danicus Retz.

PURPLE MILK-VETCH

Plants with slender creeping roots; stems single or tufted, slender, 10–30 cm high, sparsely pubescent to glabrous. Leaves with 11–21 leaflets 1–2 cm long, lanceolate to linear-oblong, sparsely pubescent on both sides. Inflorescence dense, 2–4 cm long; flowers 14–17 mm long, purplish, erect; calyx black pubescent, 5–6 mm long. Legume about 1 cm long, densely pubescent. A Eurasian steppe species. North American plants are considered to be var. *dasyglottis* (Fisch.) Boiv. A white-flowered form is f. *virgultulus* (Sheld.) Boiv. Grasslands; Prairies and Parklands. Syn.: *A. goniatus* Nutt.; *A. hypoglottis* Richardson.

Astragalus drummondii Dougl.

DRUMMOND'S MILK-VETCH

Plants with well-developed caudex; many-stemmed; stems 30–60 cm high, more or less densely pubescent with long, spreading hairs. Leaves with 23–33

leaflets 10–15 mm long, linear-oblong to elliptic, glabrous or nearly so above, densely soft pubescent below. Inflorescence 5–15 cm long, dense at first, soon elongating; flowers 1.5–2.0 cm long, white, pendent; calyx 5–6 mm long, pubescent with mixed white and black hairs. Legumes pendulous, 2–3 cm long, linear, glabrous, with the stipe 6–8 mm long. Grassland; western Prairies and Parklands. Syn.: *Tium drummondii* (Dougl.) Rydb.

Astragalus eucosmus Robinson

Plants with creeping roots; stems solitary or in small tufts, 30–60 cm high, glabrous or nearly so. Leaves with 11–17 leaflets 8–25 mm long, elliptic to oblong, glabrous or nearly so above, sparsely pubescent on underside. Inflorescence at first dense, soon elongating to 10–15 cm long; flowers 7–10 mm long, purple to whitish purple or almost white; calyx 3–4 mm long, black pubescent. Legume 8–10 mm long, obliquely elliptic, reflexed, sessile, densely pubescent. The var. *eucosmus* has black hairy pods; plants with white hairy pods are f. *leucocarpus* Lepage. Open woods, riverbanks; Boreal forest and Rocky Mountains.

Astragalus flexuosus Dougl.

SLENDER MILK-VETCH

Plants with stout creeping roots; stems solitary or few together, 30–50 cm long, straggling, thinly pubescent. Leaves with 13–23 leaflets 5–15 mm long, linear to oblong, glabrous above, somewhat pubescent below. Inflorescence 5–10 cm long in flower, elongating to 15 cm in fruit; flowers white, tipped purplish to reddish purple; calyx 3–3.5 mm long, pubescent. Legume 1–2 cm long, almost terete, linear, spreading or reflexed, with the stipe shorter than the calyx tube. Prairies and Parklands throughout the Prairie Provinces. Syn.: *Pisophaca flexuosa* (Dougl.) Rydb.

Astragalus frigidus (L.) Gray

AMERICAN MILK-VETCH

Plants with a stout caudex; stems usually several, erect, seldom branching, glabrous or nearly so. Leaves with 9–15 leaflets 2–5 cm long, oblong-elliptic to elliptic, glabrous above, somewhat pubescent below. Inflorescence 5–7 cm long in flower, elongating to 10 cm in fruit; flowers 13–15 mm long, yellowish white, at first ascending, later reflexed; calyx oblique, about 5 mm long, with very short teeth. Legume 1.5–2.0 cm long, ellipsoid, inflated, pendent, with the stipe about 5 mm long. In the var. *frigidus* of Eurasia, the legume is at first black pubescent, and later glabrescent. In Canada, the legume is glabrous, or very sparsely pubescent; the plants are distinguished as var. *americanus* (Hook.) Wats. (Fig. 133). Moist woods, riverbanks, and openings; Boreal forest, Rocky Mountains. Syn.: *A. americanus* (Hook.) M. E. Jones; *Phaca americana* (Hook.) Rydb.

Astragalus gilviflorus Sheld.

CUSHION MILK-VETCH

Plants with a caudex and deep taproot, stemless, forming a dense cushion. Leaves with 3 leaflets 1–2.5 cm long, densely silvery silky villous on both sides. Inflorescence 1 or 2 flowers, subsessile at base of leaves; flowers white, with purplish-spotted keel, 1.5–3.0 cm long; calyx 15 mm long, densely pubescent. Legume about 15 mm long, oblong-ovate, terete, silky pubescent. Dry eroded hills, slopes, and disturbed areas; western Prairies. Syn.: *A. triphyllus* Pursh.; *Orophaca caespitosa* (Nutt.) Britt.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Astragalus frigidus* (L.) Gray
var. *americanus* (Hook.) Wats.

LOCATION: DelaRonde Creek

COLL: J. Looman

DET:

COLL. NO: 17508

DATE: 15 July 1972

ELEVATION: 500 m

HABITAT: Forest opening




Fig. 133. American milk-vetch, *Astragalus frigidus* (L.) Gray var. *americanus* (Hook.) Wats.

Astragalus iochrous Barneby (Fig. 134)

AUSTRIAN FIELD-PEA

Plants with long creeping roots; stems solitary or few together, 40–90 cm high. Leaves with 11–17 leaflets 5–25 mm long, oblong to elliptic, glaucous green. Inflorescence 5–10 cm long, loosely flowered; flowers 10–15 mm long, brick red; calyx 3–5 mm long. Legume 15–25 mm long, ovoid, glabrous, with the stipe about twice as long as the calyx. An introduced species, rare; known from Maple Creek, Sask. Syn.: *Swainsona salsula* (Pall.) Taub.

Astragalus kentrophyta Gray

PRICKLY MILK-VETCH

Plants with a deep taproot; stems pubescent, straggling. Leaves 2–3 cm long; leaflets 1–2 cm long, tipped with a sharp spine. Inflorescence 1–2 cm long, almost hidden among the leaves; flowers 4–5 mm long, white with purplish tinge; calyx 2–3 mm long, pubescent. Legume about 5 mm long, ovoid, somewhat compressed. Rare; sand dunes and other sandy areas; southwestern Prairies. Syn.: *Kentrophyta montana* Nutt.

Astragalus lotiflorus Hook.

LOW MILK-VETCH

Plants with a deep taproot; stems erect, pubescent, 10–15 cm high. Leaves with 9–13 leaflets 8–15 mm long, elliptic to oblong, thinly pubescent above, densely pubescent below. Inflorescence 2–3 cm long; flowers 8–10 mm long, yellow white, often purplish-tipped; calyx 3–3.5 mm long, densely pubescent. Legumes 1.5–2.0 cm long, long-pointed, densely pubescent, usually sessile amongst the leaf bases, sometimes on stems about 5 cm long. Not common; grassland and openings on light soils; Prairies and Parklands. Syn.: *Batidophaca lotiflora* (Hook.) Rydb.

Astragalus miser Dougl.

TIMBER MILK-VETCH

Plants with several stems arising from caudex. Stems slender, 5–20 cm high, ascending, somewhat pubescent or glabrous. Leaves with 9–15 leaflets 3–10 mm long, linear to linear-oblong, pubescent on both sides. Inflorescence 6–10 cm long; flowers 8–10 mm long, white to yellowish, purple-tipped; calyx 3–4 mm long, pubescent with mixed white and black hairs. Legume 15–20 mm long, linear, subsessile, drooping, somewhat pubescent. Rare; in Rocky Mountains. The var. *serotinus* (Gray) Barneby, with the calyx 2–2.5 mm long, flowers 6–8 mm long, purple. More common; open woods, slopes; Rocky Mountains. Syn.: *Homalobus serotinus* (A. Gray) Rydb.; *A. serotinus* A. Gray.

Astragalus missouriensis Nutt.

MISSOURI MILK-VETCH

Plants with a deep stout taproot and caudex. Stems tufted, to 20 cm long, usually ascending, more or less densely gray or silvery pubescent. Leaves with 9–21 leaflets 5–15 mm long, elliptic to obovate, densely silvery pubescent. Inflorescence 1.5–5 cm long, dense at first, and elongating in fruit; flowers 15–20 mm long, deep reddish purple to bluish purple; calyx 7–8 mm long, silvery pubescent. Legume 2.5–3 cm long, oblong, pubescent, erect. Slopes and eroded hillsides; Prairies. Syn.: *Xylophacos missouriensis* (Nutt.) Rydb.

Astragalus pectinatus Dougl. (Fig. 132B)

NARROW-LEAVED MILK-VETCH

Plants with a deep taproot, and few-stemmed caudex. Stems 20–50 cm long, decumbent to ascending, glabrous, often reddish at base. Leaves with




Fig. 134. Austrian field-pea, *Astragalus iochrous* Barneby.

9–17 leaflets 2–6 cm long, narrowly linear, 1–2 mm wide, sparingly pubescent. Inflorescence 5–8 cm long; flowers 1.5–2.5 cm long, white to yellowish; calyx 7–8 mm long. Legume 10–20 mm long, oblong, terete, glabrous, fleshy at first, becoming woody when dry. Grasslands and slopes; Prairies. Syn.: *Cnemidophacos pectinatus* (Dougl.) Rydb.

Astragalus purshii Dougl.

PURSH'S MILK-VETCH

Cushion-like plants with a deep taproot and caudex. Stems 5–8 cm long, densely pubescent. Leaves with 9–15 leaflets 8–12 mm long, oblanceolate to ovate, densely long pubescent. Inflorescence 2–3 cm long; flowers 2–2.5 cm long, yellowish white, purple-tipped; calyx to 10 mm long, densely pubescent. Legume 15–20 mm long, ovoid, densely long pubescent. Rare; dry prairie slopes and eroded hills; western Prairies. Syn.: *Xylophacos purshii* (Dougl.) Rydb.

Astragalus racemosus Pursh

RACEMOSE MILK-VETCH

Plant several-stemmed from a stout taproot and caudex. Stems 40–80 cm high, sparingly pubescent. Leaves with 15–27 leaflets 1–2 cm long, glabrous above, somewhat pubescent below, linear to linear-oblong. Inflorescence 5–7 cm in flower, elongating to 10 cm in fruit; flowers 15–20 mm long, yellowish white; calyx 2.5–3 mm long. Legume 1.5–2.5 cm long, glabrous, triangular in cross section, drooping, with the stipe 5–7 mm long. Rare; dry prairie slopes and hillsides; Saskatchewan. Syn.: *Tium racemosum* (Pursh) Rydb.

Astragalus robbinsii (Oakes) Gray

Plants with several stems arising from caudex. Stems 30–50 cm high, erect, pubescent. Leaves with 9–17 leaflets 1.5–2.5 cm long, elliptic to oval or oblong, glabrous above, pubescent below. Inflorescence 5–10 cm long in flower, elongating to 15 cm in fruit; flowers 8–10 mm long, bluish purple to almost white; calyx 3–4 mm long, black hairy. Legume 15–20 mm long, flattened, black pubescent, drooping, stipitate. Riverbanks, lakeshores; Rocky Mountains.

Astragalus spatulatus Sheld.

TUFTED MILK-VETCH

Cushion-like plant with a deep taproot and caudex. Stems 5–7 cm long, silvery gray pubescent. Leaves usually with a single leaflet, occasionally 3 leaflets 1–4 cm long, oblong to linear oblong, densely silvery gray pubescent. Inflorescence 2–3 cm long in flower, the peduncle elongating to 7–8 cm in fruit; flowers 6–8 mm long, bluish purple; calyx 2–2.5 mm long, pubescent. Legume 6–12 mm long, flattened, elliptic, glabrous, erect. Locally moderately common; eroded hillsides and slopes; western Prairies. Syn.: *A. caespitosus* (Nutt.) Gray; *Homalobus caespitosus* Nutt.

Astragalus striatus Nutt.

ASCENDING PURPLE MILK-VETCH

Plants with several to many stems arising from a stout taproot. Stems decumbent or ascending, 15–40 cm high, pubescent. Leaves with 9–19 leaflets 1–2 cm long, elliptic to oblong, pubescent, with the hairs attached in the middle. Inflorescence 4–5 cm long, dense; flowers 12–15 mm long, purplish, rarely white; calyx 5 mm long, pubescent with black and white hairs. Legume 7–15

mm long, ovoid, pubescent. Open grasslands and slopes; Prairies and Parklands. Syn.: *A. adsurgens* Hook.

Astragalus tenellus Pursh

LOOSE-FLOWERED MILK-VETCH

Plants with few to several stems from a stout caudex. Stems 20–50 cm high, somewhat pubescent, erect. Leaves with 11–21 leaflets 8–12 mm long, linear to linear-oblong, glabrous above, somewhat pubescent below. Inflorescence 4–5 cm long in flower, elongating to 7 cm in fruit; flowers 8–10 mm long, whitish; calyx 2–2.5 mm long, pubescent. Legume 8–12 mm long, flattened, glabrous, drooping, stipitate. Coulees, forest margins, lakeshores; Prairies and Parklands. Syn.: *Homalobus tenellus* (Pursh) Britt.

Astragalus vexilliflexus Sheld.

FEW-FLOWERED MILK-VETCH

Plants matted, many-stemmed from a deep taproot and caudex. Stems 10–30 cm long, straggling, pubescent. Leaves with 7–11 leaflets 5–18 mm long, oblong to oblong-lanceolate, pubescent. Inflorescence 1.5–3.0 cm long; flowers 6–8 mm long, purplish, occasionally whitish; calyx 2 mm long, pubescent. Legume 7–10 mm long, somewhat pubescent, flattened, sessile. Rare, locally moderately common; eroded slopes and hills; western Prairies, Rocky Mountains. Syn.: *Homalobus vexilliflexus* (Sheld.) Rydb.

Astragalus yukonis M. E. Jones

Plants with several to many stems arising from a stout taproot and caudex. Stems 5–30 cm high, slender, decumbent to ascending, glabrous. Leaves with 7–15 leaflets 4–12 mm long, oblong, apex often retuse, glabrous above, pubescent below. Inflorescence 2–4 cm long on a long peduncle; flowers 7–10 mm long, bluish- or purple-tipped; calyx 2–2.5 mm long, black pubescent. Legume 5–10 mm long, ellipsoid, black pubescent, erect. Grassy openings; Boreal forest.

Caragana Siberian peatree

Caragana arborescens Lam.

COMMON CARAGANA

A bush 3–4 m high; leaves pinnate with 8–12 pale green leaflets 10–25 mm long; the leaf stem ending in a short spine. Flowers bright yellow, 15–25 mm long, followed by dark brown linear pods 3–5 cm long. Introduced as a hedge and ornamental plant from Siberia and now established in many places.

Coronilla crown-vetch

Coronilla varia L.

FIELD CROWN-VETCH

Perennial plants with stems 20–100 cm high. Leaves with 6–10 pairs of oblong or elliptic leaflets 6–20 mm long. Heads mostly 10- to 20-flowered; corolla 10–15 mm long, white to purplish; pods 20–60 mm long, with 3–7 joints, 4-angled. Introduced and occasionally spreading from cultivation; southeastern Parklands.

Desmodium tick-trefoil

Desmodium canadense (L.) DC.

BEGGAR'S-LICE

Plants perennial, with erect pubescent stems, mostly about 1 m, occasionally to 2 m high. Leaves trifoliate, with the leaflets oblong to oblong-lanceolate, the terminal one petioled, 5–9 cm long; racemes densely flowered; flowers 10–15 mm long, purplish; legume 2–3 cm long, with 3–5 joints. Rare; moist open areas; southeastern Parklands and Boreal forest.

Glycyrrhiza wild licorice

Glycyrrhiza lepidota (Nutt.) Pursh (Fig. 135)

WILD LICORICE

A coarse erect branching herb 30–100 cm high, from a thick sweet-tasting rootstock having a slight licorice flavor. Leaves of 11–19 lanceolate or oblong leaflets, pale green, glandular-dotted, 20–35 mm long, and pointed at both ends. Flowers yellowish white, 10–15 mm long, and borne in rather dense racemes 2–6 cm long, arising from the axils of the leaves. Fruit in clusters of oblong reddish brown pods 10–15 cm long, densely covered with long hooked prickles, and containing several large seeds. Rootstocks were chewed by Indians. Palatable as hay, but seldom grazed. Very common in southern portion, but becoming scarcer toward the north; found in low spots on the prairie, slough margins, riverbanks, and coulees; throughout the Prairies and Parklands.

Hedysarum sweet-broom

Perennial herbs with pinnate leaves; the flowers usually reflexed in long, spike-like racemes, and the pods flat and jointed or constricted between the seeds. The plants make good forage.

- 1. Calyx teeth very unequal, the upper ones nearly triangular, shorter than the calyx tube; veins of leaflets conspicuous. 2
- Calyx teeth almost equal, linear, about as long as the calyx tube; veins of leaves not conspicuous. *H. boreale*
- 2. Flowers violet or pale pink. *H. alpinum* var. *americanum*
- Flowers sulfur yellow. *H. sulphurescens*

Hedysarum alpinum L. var. *americanum* Michx. (Fig. 136)

AMERICAN HEDYSARUM

An erect plant 15–80 cm high, usually with few branches. Leaves of 11–21 oblong leaflets 10–30 mm long. Flowers on a long raceme, pinkish or violet, 10–15 mm long, usually pointing downward. Pods usually have 3–5 internodes or enlargements, hairless except perhaps on the margins. Common; in semi-open prairie and open woods; especially in the Boreal forest and the Cypress Hills. The most common species of the genus in the Prairie Provinces, and readily eaten by livestock. A variety with hairy pods has been found in the southwestern part of the Prairie Provinces. Syn.: *H. americanum* (Michx.) Britt.




Fig. 135. Wild licorice, *Glycyrrhiza lepidota* (Nutt.) Pursh.




Fig. 136. American hedysarum, *Hedysarum alpinum* L. var. *americanum* Michx.

Somewhat similar to *H. alpinum* but the lobes of the calyx longer and very narrow and the veins on the leaflets not distinct. Leaflets almost smooth and flowers reddish purple and somewhat reflexed.

Three forms of this species occur:

- 1. Inflorescence short and dense; flowers to 20 mm long; stipules whitish. var. *mackenzii*
 Inflorescence elongate and loose; flowers about 15 mm long; stipules brownish. 2
- 2. Leaflets glabrous or sparsely pubescent; pods rather smooth. var. *boreale*
 Leaflets densely silky hairy; pods rugose. var. *cinerascens*

Both var. *boreale* and var. *cinerascens* (Rydb.) Rollins occur on slopes of ravines and coulees in Prairies; var. *cinerascens* in drier areas, var. *boreale* in moister situations and also occasionally in Parklands. The var. *mackenzii* (Richardson) C. L. Hitchc., Mackenzie's hedysarum, is fairly common in meadows and on slopes in Parklands and Boreal forest.

An erect species 30–60 cm high; leaves with 11–15 oblong to oval leaflets 10–85 mm long. Flowers 12–15 mm long, sulfur yellow, in a long loosely-flowered raceme. A species readily distinguishable by the flower color. Open woods, slopes, and meadows; southern Rocky Mountains, rare farther north.

Perennial twining vines, dying to the ground each year and bearing pinnate leaves with a tendril taking the place of the terminal leaflet. The flowers perfect, the stamens diadelphous. Style hairy along its inner side, and somewhat flattened, distinguishing this genus from *Vicia* (the true vetch), with merely a tuft of hairs at the end of an unflattened style. Pods somewhat flattened containing several seeds. Very palatable, a valuable native fodder in many localities. Grazed out readily, but making good growth again when protected from livestock. The sweet pea, *Lathyrus odoratus* L., occurs occasionally as an escape, but is not persistent.

- 1. Leaves with only 2 leaflets. 2
 Leaves with more than 2 leaflets. 3
- 2. Stem clearly winged. *L. sativus*
 Stem wingless. *L. tuberosus*
- 3. Flowers creamy white or yellowish. *L. ochroleucus*
 Flowers purplish. 4
- 4. Racemes with 15–20 or more flowers. *L. venosus*
 Racemes with fewer than 15 flowers. 5
- 5. Leaflets linear to linear-lanceolate; stem narrowly winged, at least above. *L. palustris*
 Leaflets oblong to ovate; stem wingless. *L. japonicus*

A somewhat glaucous glabrous or pubescent perennial, usually straggling with stems to 1 m long. Leaves with 4–10 leaflets, often without a tendril. Racemes 2- to 12-flowered; corolla 15–25 mm long, purplish; legume 3–5 cm long, with 4–10 seeds. Not common; lakeshores and beaches; Hudson Bay, Lake Winnipeg, Lake Manitoba. The typical form has 2- to 7-flowered racemes, and a pubescent calyx; plants with 5- to 12-flowered racemes and glabrous calyx distinguishable as subsp. *maritimus* (L.) R. W. Ball.

Lathyrus ochroleucus Hook. (Fig. 137)

CREAM-COLORED VETCHLING

A slender smooth climber up to 1 m long, with a somewhat angled stem. Stipules large and almost cordate. Leaves of 6–10 broad oval leaflets 2–5 cm long. Flowers cream-colored, about 15 mm long, in racemes of 5–10 flowers. Pods about 4 cm long. Very common; in bluffs, open woodlands, and among bushes; throughout the Prairie Provinces.

Lathyrus palustris L.

MARSH VETCHLING

A smooth climber 30–100 cm long, with a somewhat winged stem and leaves of 4–8 linear or linear-oblong leaflets 10–65 mm long. Stipules small and almost linear. Flowers purple, 10–15 mm long, 2–8 in each raceme. Pods 3–5 cm long. Fairly common; in moist places and damp woodlands; Boreal forest.

Lathyrus sativus L.

CHICKEN VETCH

Annual plants with winged stems to 1 m high; leaflets 2–15 cm long, linear-lanceolate. Flowers solitary, white, pink, or bluish; legume 2–4 cm long, 1–2 cm wide. Introduced, occasionally sown for fodder, and rarely escaped.

Lathyrus tuberosus L.

TUBEROUS VETCHLING

Perennial with a tuberous root; stems 30–100 cm high, glabrous or subglabrous. Leaflets 15–45 mm long, elliptic to oblong. Racemes 2- to 7-flowered; corolla 12–20 mm long, bright reddish purple; legume 2–4 cm long. Introduced, occasionally sown for fodder, and spreading from cultivation.

Lathyrus venosus Muhl.

WILD PEAVINE

A climbing plant 50–100 cm long, sometimes somewhat finely hairy. Stems strongly 4-angled. Leaves having 8–12 oblong-ovate blunt-tipped leaflets 2–5 cm long. Flowers purple, 10–18 mm long, 15–20 in each raceme. Pods 3–5 cm long and veiny. This wild peavine provided a valuable source of forage and hay in the earlier days of settlement of northern bushlands. Common; around bushes and woodlands; eastern Parklands and Boreal forest. The pubescent form is var. *intonsus* Butt. & St. John.

Lotus trefoil

Annual or perennial herbs; leaves pinnate, with odd-numbered leaflets. Flowers solitary or in heads; legume cylindrical. Mostly Eurasian; several species useful as forage crops, two of these introduced in Canada.




Fig. 137. Cream-colored vetchling, *Lathyrus ochroleucus* Hook.

- 1. Plants annual; flowers solitary in upper leaf axils. *L. americanus*
Plants perennial; flowers in umbellate heads. 2
- 2. Stems solid; calyx lobes incurved in bud, 1.5–2.0 mm long; leaflets 5–15 mm long. *L. corniculatus*
Stems hollow; calyx lobes recurved in bud, 2–4 mm long; leaflets 10–25 mm long. *L. pedunculatus*

Lotus americanus (Nutt.) Bisch. SPANISH CLOVER

Plants 10–40 cm high, with stems erect, branched or simple, pubescent. Leaves subsessile, 3-foliolate, 1–2 cm long, with the terminal one stalked. Flowers pinkish, 5–7 mm long; pods 2–3 cm long, deflexed. Not common; dry to moist grasslands; southeastern Parklands. Syn.: *Trigonella americana* Nutt.; *Lotus purshianus* (Benth.) Clem. & Clem.

Lotus corniculatus L. BIRD'S-FOOT TREFOIL

Stems prostrate, erect or ascending, to 60 cm long, solid or nearly so. Inflorescence 1- to 5-flowered; flowers yellow, often tinged with red; pods 2–4 cm long. Introduced as forage crop and locally established.

Lotus pedunculatus Cav. TREFOIL

Resembling *L. corniculatus*, but the stem with a large cavity, the inflorescence 5- to 10-flowered, and the leaflets usually larger. Legume 2–4 cm long. Probably introduced with *L. corniculatus*; rare as an escape.

Lupinus lupine

Annual or perennial showy herbs with alternate palmate leaves, each bearing 5–12 leaflets. Flowers perfect and borne in terminal racemes. Ten stamens united into one bundle with the anthers alternately elongated and short. Pods flattened, consisting of 2 cells, and containing 1–6 seeds. Some species of lupines **poisonous** to stock, particularly sheep; the seedpods apparently being the most dangerous part of the plants.

- 1. Plants annual; stem and leaves densely pilose to hirsute; flowers pale blue to white. *L. pusillus*
Plants perennial. 2
- 2. Standard (or banner) pubescent on the back over most of its surface. *L. sericeus*
Standard glabrous or with a few cilia. 3
- 3. Leaflets sericeous above; plants silky-sericeous, up to 20 cm high; the leaves mostly basal. *L. minimus*
Leaflets glabrous above, or occasionally sparsely puberulent or strigose; plants more than 20 cm high. 4

4. Keel densely ciliate along most of its upper edge; petioles usually less than twice the length of the leaflets. *L. nootkatensis*
 Keel ciliate only toward the tip, or glabrous; petioles of various lengths. 5
5. Leaves mostly basal, long-petioled; leaflets 10–17, glabrous above, sparsely strigose below, the largest to 12 cm long, to 25 mm wide. *L. polyphyllus*
 Leaves mostly cauline, short-petioled; leaflets 7–9, glabrous above, sericeous below, the largest to 4.5 cm long, to 5 mm wide. *L. argenteus*

Lupinus argenteus Pursh (Fig. 138)

SILVERY LUPINE

A rather shrubby much-branched herb 30–60 cm high, with stems covered with appressed silky hairs. Leaves of 6–9 narrowly oblanceolate leaflets 2–5 cm long, sometimes silvery hairy or smooth above. Flowers varying from light violet or purplish to almost white, in long terminal racemes. Pods densely silky hairy, 15–25 mm long, containing up to 5 seeds. Plentiful; on submontane prairie; southern Rocky Mountains, Cypress Hills, and Wood Mountain.

Lupinus minimus Dougl.

ALPINE LUPINE

Stems erect to decumbent, unbranched, 15–20 cm high, silky pubescent, with 1 or 2 stem leaves. Basal leaves long-petioled, with 5–9 leaflets 2–3 cm long. Very rare; alpine slopes; southern Rocky Mountains.

Lupinus nootkatensis Donn

NOOTKA LUPINE

Plants 20–60 cm high, with stems erect, densely villose pubescent. Leaves with 7–15 leaflets; the petioles 1.5–2 times as long as the leaflets. Racemes to 20 cm long; flowers 12–16 mm long. Pods loosely pubescent, 3–4 cm long. Not common; moist alpine meadows; southern Rocky Mountains.

Lupinus polyphyllus Lindl.

LARGE-LEAFED LUPINE

Plants 50–120 cm high, stout-stemmed, appressed to spreading pubescent. Leaves long-petioled, with up to 17 leaflets. Inflorescence 15–30 cm long, rather loosely flowered; petals blue to reddish or yellowish. Pods pubescent, 3–5 cm long. Not common; grassland, open woods, and meadows; southern Rocky Mountains.

Lupinus pusillus Pursh

SMALL LUPINE

A low-growing annual plant 10–25 cm high, with decumbent branches and very hairy. Leaves usually of 5 oblong leaflets, smooth above but with long hairs beneath, usually rounded at the ends, 20–35 mm long. Flowers tinged with purple or rose, sometimes almost white, in short dense racemes on very short stalks. Pods about 2 cm long with 1 or 2 seeds and somewhat constricted between the seeds. Found locally in sandhills and among sand dunes, but not common.




Fig. 138. Silvery lupine, *Lupinus argenteus* Pursh.

A branching plant 40–80 cm high, with grayish appressed silky hairs. Leaves of 6–10 narrowly oblanceolate leaflets 2–5 cm long, densely appressed silky hairy on both sides. Dark blue flowers in dense terminal spikes. Densely white hairy pods 20–35 mm long, containing 2–5 seeds. A very showy species. In grasslands and open woods; southern Rocky Mountains. Syn.: *L. flexuosus* Lindl.

Medicago medick

Annual or perennial plants, not native to the prairie, with trifoliolate leaves, toothed only beyond the center. Flowers perfect, borne in racemes or spikes; stamens diadelphous. Pods curved or spirally twisted.

- 1. Plants annual or biennial; flowers yellow; leaflets about 10 mm long, obovate. 2
- Plants perennial; flowers yellow or some shade of purple; leaflets about 15 mm long, linear. 3
- 2. Flowers in dense head-like racemes, 10- to 30-flowered; legume kidney-shaped, to 3 mm across, black. *M. lupulina*
- Flowers in a raceme with 1–5 flowers; legume spirally coiled, to 8 mm across, spiny. *M. polymorpha*
- 3. Flowers some shade of purple, sometimes whitish; legume coiled; corolla 7–11 mm. *M. sativa* ssp. *sativa*
- Flowers yellow; legume falcate to almost straight; corolla 5–8 mm. *M. sativa* ssp. *falcata*

Medicago lupulina L.


A prostrate branched annual weed 10–60 cm across, with trifoliolate leaves. Leaflets obovate, toothed above the middle, 3–10 mm long. Flowers yellow, about 3 mm long, in dense head-like racemes, less than 1 cm long. Pods small and black, containing a single seed. Introduced; common in disturbed areas in Parklands and Boreal forest; not common in Prairie.

Medicago polymorpha L.

Annual plants 15–40 cm high, with glabrous or sparingly pubescent stems. Racemes of 1–8 yellow flowers 3–5 cm long. Pods 4–8 mm in diam, spirally coiled, shiny. Introduced, rarely weedy.

Medicago sativa L. (Fig. 139)

A fairly erect perennial 30–80 cm high, much-branched with trifoliolate leaves. Leaflets 10–35 mm long, obovate, and sharply toothed toward the apex. Flowers 5–11 mm long, in a dense oblong raceme 10–45 mm long. Introduced as a fodder crop from Europe and now very common; along roadsides and waste places; throughout most of the Prairie Provinces.



A. C. Budd

Fig. 139. Alfalfa, *Medicago sativa* L. ssp. *sativa*.

This species consists of several entities that hybridize quite freely and are often considered as separate species. Two of these are commonly used as forage, along with many cultivated varieties (cultivars) resulting from the breeding program.

- Corolla blue to purple, 7–11 mm; pods coiled in 1.5–3.5 turns. ssp. *sativa*
- Corolla yellow, 5–8 mm; pods straight to sickle-shaped. ssp. *falcata* (L.) Arcangeli

Cultivars of the two subspecies have flower colors and pod shapes between those of the parents.

Melilotus sweet-clover

Annual or biennial introduced legumes with trifoliolate leaves; the leaflets toothed almost to the base. Flowers perfect, small, and in elongate, spike-like racemes. Pods short, thick, and straight with 1 or few seeds. Excellent for honey production and also as pasture for stock. Very common roadside weeds.

- 1. Flowers white. 2
- Flowers yellow. 3
- 2. Corolla 3–3.5 mm long, with pedicels 2–4 mm long; racemes 5–10 cm long. *M. wolgica*
- Corolla 4–5 mm long, with pedicels 1–1.5 mm long; racemes to 20 cm long. *M. alba*
- 3. Corolla 2–3 mm long; legume 1.5–3 mm long. *M. indica*
- Corolla 4–7 mm long; legume 3–5 mm long. *M. officinalis*

Melilotus alba Medic. (Fig. 140) WHITE SWEET-CLOVER

An erect plant 50–250 cm high, with palmately or pinnately trifoliolate leaves; the leaflets 10–25 mm long, toothed almost to the base. Flowers white, in long narrow spike-like racemes. Introduced as a forage plant from Europe and Asia.

Melilotus indica (L.) All. YELLOW SWEET-CLOVER

Erect plants to about 50 cm high. Racemes dense; flowers pale yellow. Occasionally cultivated, rare as a weed; southeastern Parklands.

Melilotus officinalis (L.) Pall. YELLOW SWEET-CLOVER

Similar to *M. alba*, but with yellow flowers. Introduced as a forage plant; common along roadsides and in disturbed areas.

Melilotus wolgica Poir. WHITE SWEET-CLOVER

Also similar to *M. alba*, but with longer pedicels, 2–4 mm, and smaller flowers. Introduced as a forage plant, and rarely escaped.



Fig. 140. White sweet-clover, *Melilotus alba* Medic.

Oxytropis locoweed

Perennial herbs, usually with no apparent stem (except in *O. deflexa*), pinnate leaves with odd-numbered leaflets, and perfect flowers borne in a spike or a raceme. Stamens diadelphous; keel extended to a protruding point, distinguishing this genus from *Astragalus*. Several species of this genus **poisonous** to livestock, causing well-known loco disease, affecting the nervous system.

1. Plant caulescent; stipules adnate to the base of the petiole only; pods pendulous. *O. deflexa*
 Plants acaulescent; stipules adnate for half their length; pods erect. 2
2. Leaflets whorled on the rachis or appearing so; few leaflets in pairs. 3
 Leaflets all paired or scattered on the rachis. 5
3. Inflorescence capitate, few-flowered. *O. bellii*
 Inflorescence elongate, many-flowered. 4
4. Corolla purple; plants silky pubescent. *O. splendens*
 Corolla yellow; plants appressed-pubescent. *O. campestris*
5. Corolla purple to bluish. 6
 Corolla yellow to whitish. 11
6. Pubescence in part composed of malpighian hairs. *O. lambertii*
 Pubescence entirely composed of basifixed hairs. 7
7. Plants more or less glandular-viscid. *O. viscida*
 Plants not glandular-viscid. 8
8. Pods ovoid, stipitate, strongly inflated. *O. podocarpa*
 Pods not ovoid, not strongly inflated. 9
9. Corolla small, the keel to 13 mm long. *O. campestris*
 Corolla large, the keel 15–25 mm long. 10
10. Plants very densely silky and spreading pubescent; bracts of the inflorescence linear-lanceolate, membranous with involute blades. *O. lagopus*
 Plants hispid-hirsute to subappressed-pilose; bracts of the inflorescence rhombic-lanceolate, herbaceous with flat blades. *O. besseyi*
11. Keel of the corolla to 13 mm long; leaves with 17–33 leaflets; pods not rigid at maturity. *O. campestris*
 Keel of the corolla to 20 mm long; leaves with 11–19 leaflets; pods rigid at maturity. *O. sericea*

Oxytropis bellii (Britt.) Palibine

Plants 10–15 cm high. Leaves with 17–35 leaflets 5–10 mm long, linear-lanceolate, acute, silky pubescent, verticillate, mostly 3 or 4 leaflets per verticil. Inflorescence with 4–7 flowers in a dense raceme; flowers 15–20 mm long, large, dark blue purple; calyx 4–5 mm long, villous. Legume 8–10 mm long, black pubescent, ovoid, erect. Very rare; Hudson Bay.

Oxytropis besseyi (Rydb.) Blank. (Fig. 141)

BESSEY'S LOCOWEED

A tufted silvery hairy plant 10–20 cm high. Leaflets 5–20 mm long. Inflorescence a short dense spike 2–5 cm high; flowers reddish purple. The plant is easily mistaken for Missouri milk-vetch. Pods about 2 cm long, covered with long silky hairs. This is a southern species, which has been found, but very rarely; on dry hillsides; Wood Mountain.

Oxytropis campestris (L.) DC.

LATE YELLOW LOCOWEED

Plants with a deep taproot and branching caudex, 15–40 cm high. Leaves with 7–33 leaflets 1–2 cm long, oblong-lanceolate, silky pubescent. Inflorescence 5–10 cm long; flowers 12–15 mm long, variable in color; calyx 5–7 mm long, pubescent with both white and black hairs. Legume 16–20 mm long, oblong-ovate, pubescent with both white and black hairs, semimembranous.

A circumboreal species, very variable. It has been divided into a large number of "small" species by various authors. North American plants may be considered as varieties of ssp. *gracilis* (Nelson) Boiv., in which the legumes lack a fully developed septum, whereas in the Eurasian ssp. *campestris* the septum is present. Some authorities consider all these forms as varieties of *O. campestris*.

- 1. Flowers bluish or purplish. 2
 Flowers yellow or creamy white. 3
- 2. Leaves of two sizes, the long ones about
 twice as long as the short ones; leaflets
 often in part verticillate. var. *dispar* (Nels.) Barneby
 Leaves all about the same length; leaflets
 all paired. var. *johannensis* Fern.
- 3. Stipules villous, often with club-shaped
 processes on margins. var. *varians* (Rydb.) Barneby
 Stipules glabrous or glabrate, without
 club-shaped processes. 4
- 4. Leaves with 17–33 leaflets; scapes 15–30
 cm; inflorescence many-flowered. var. *gracilis*
 Leaves with 7–17 leaflets; scapes 1.5–15
 cm; inflorescence few-flowered. var. *cusickii* (Greenm.) Barneby

Of these forms, var. *gracilis* is common in moist grassland, open woods, and openings throughout the Prairie Provinces; var. *cusickii* occurs in alpine and subalpine meadows and rockslides in the Rocky Mountains; var. *dispar*, rare, on slopes of ravines and margins of groves, in southeastern Parklands; var. *johannensis*, gravel bars, Hudson Bay; var. *varians*, rocky hillsides and meadows, Hudson Bay.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Oxytropis besseyi* (Rydb.) Blank.

LOCATION: East of Rock Glen

COLL: J. Looman

DET:

COLL. NO: 14338

DATE: 24 June 1970

ELEVATION: 930 m

HABITAT: Eroded crest
of steep hill



Fig. 141. Bessey's locoweed, *Oxytropis besseyi* (Rydb.) Blank.

Plants with a deep taproot and caudex. Stems well-developed, 10–40 cm long, decumbent to ascending, or acaulescent with the scape 10–20 cm long, loosely pubescent. Leaves appearing flattened, with 17–41 leaflets 5–15 mm long, lanceolate, pubescent on both sides. Inflorescence loosely flowered, elongating in fruit; flowers 6–10 mm long, bluish or creamy white with purple tip; calyx 3–4 mm long, black pubescent, reflexed.

This species has been divided into the following varieties:

- 1. Lateral sinuses of the calyx similar to the ventral pair, broad and obtuse; petals ample, purple, with the banner broadly obcordate, about twice as long as broad. 2
- Lateral sinuses similar to the dorsal pair, narrow and acute, rarely a little broader and rounded; petals narrower, mostly blue- or purplish-tinged, with the banner oblanceolate, about 3 times as long as broad. var. *sericea* T. & G.
- 2. Plants copiously and loosely villous-pilose; stems commonly developed, with at least 1 apparent internode at maturity; racemes 5–10 to 20-flowered, usually elongating in fruit. var. *deflexa*
- Plants sparingly pilose, green, with the hairs appressed; stems usually none, occasionally with 1 or 2 internodes; racemes 2–7 to 10-flowered, almost always compact in fruit. var. *foliolosa* (Hook.) Barneby

Of these forms, var. *sericea* is found in grasslands and openings throughout Parklands and Boreal forest; var. *foliolosa* in openings and on slopes in Rocky Mountains; var. *deflexa* does not occur in the Prairie Provinces.

Oxytropis lagopus Nutt.

Plants with a stout taproot, densely pubescent. Leaves with 9–15 leaflets 5–15 mm long, elliptic to lanceolate. Scape 15–20 cm high; inflorescence 3–5 cm long; flowers 15–20 mm long, dark bluish purple; calyx 5–6 mm long. Legume 12–15 mm long, oblong ovoid, densely long villous. Very rare; dry hills; southwestern Prairies.

Oxytropis lambertii Pursh

Plants with a stout taproot, 15–25 cm high. Leaves with 11–17 leaflets 15–30 mm long, lanceolate, somewhat pubescent on both sides. Scape to 25 cm high; inflorescence 7–12 cm long; flowers 15–20 mm long, dark bluish purple; calyx about 7 mm long. Legume 15–20 mm long, lanceolate, pubescent. Some of the pubescence of this species is malpighian, which makes it distinguishable from the other purple-flowered species. Grasslands; southeastern Parkland.

Oxytropis podocarpa Gray

Plants with a deep taproot and branched caudex, somewhat cushion-like. Leaves with 11–25 leaflets 3–10 mm long, linear to oblong, silky pubescent. Scape 3–6 cm high; inflorescence with 1–3 dark bluish purple flowers; calyx 6–7 mm long, villose pubescent. Legume 15–20 mm long, ovoid-ellipsoid, inflated, black pubescent, stipitate. Arctic–alpine; Rocky Mountains, Alberta.

Oxytropis sericea Nutt.

EARLY YELLOW LOCOWEED

Plants with a stout taproot and branched caudex. Leaves with 7–15 leaflets 10–30 mm long, elliptic or oblong, silky pubescent on both sides. Scape 10–20 cm high; inflorescence 5–7 cm long; flowers 18–20 mm long, yellowish; calyx 6–7 mm long, silky pubescent, black hairy on the calyx lobes. Legume 20 mm long, oblong, pubescent with both black and white hairs, rigid. The var. *spicata* (Hook.) Barneby (Fig. 142) is common in grasslands in Prairies, less common in Parklands; var. *sericea* with flowers light purple to whitish is more southern.

Oxytropis splendens Dougl.

SHOWY LOCOWEED

Plants with a deep taproot and branched caudex. Leaves with up to 60 leaflets arranged in verticils of 3–6; leaflets 10–25 mm long, linear-lanceolate, long silky pubescent. Scape 20–30 cm high; inflorescence 4–10 cm long; flowers 12–15 mm long, dark blue; calyx 6–7 mm long, densely long silky pubescent. Legume 8–12 mm long, ovoid, densely long pubescent. Grasslands and open woods; Boreal forest, Parklands, Rocky Mountains. The var. *richardsonii* Hook., wooly locoweed, differs from the species in the type of hairiness: the variety has hairs appressed (lying flat) and silky, whereas the typical plant has long, soft, spreading hairs. Flowers dark purple; leaflets lanceolate. Common in southern Rocky Mountains, decreasing in abundance toward the east.

Oxytropis viscida Nutt.

VISCID LOCOWEED

Plants with a deep taproot and branched caudex. Leaves with 21–35 leaflets 5–12 mm long, lanceolate, somewhat pubescent, and glandular. Scape 5–20 cm high; inflorescence 3–8 cm long; flowers 10–12 mm long, purple with yellowish base; calyx 4–6 mm long, densely glandular pubescent. Legume 10–15 mm long, finely black pubescent, ovate. Southwestern Alberta, Prairies, and Parklands.

Petalostemon prairie-clover

Rather low-growing, often prostrate perennial plants, with glandular-dotted pinnate leaves having odd-numbered leaflets. Flowers perfect, borne in dense spikes at head of flowering stalks. Five stamens united into one bundle. Fruits short; pods containing 1 or 2 seeds. Many authorities call this genus *Petalostemum*; both spellings are acceptable.

- 1. Leaves with 7–17 leaflets; plants densely hairy throughout. *P. villosum*
- Leaves with 3–5 (sometimes 7) leaflets; plants glabrous or sparsely pubescent. 2




Fig. 142. Early yellow locoweed, *Oxytropis sericea* Nutt. var. *spicata* (Hook.) Barneby.

- 2. Flowers purple; leaflets narrowly linear, about 15–25 mm long and 1–1.5 mm wide. *P. purpureum*
- Flowers white; leaflets linear-oblong, about 15–25 mm long and 2–3 mm wide. *P. candidum*

Petalostemon candidum (Willd.) Michx. WHITE PRAIRIE-CLOVER

Stems 20–50 cm high, usually erect. Leaves of 7–9 linear-oblong leaflets 1–2 cm long. Flowers in a compact spike 2–8 cm long, white, and a little under 6 mm long. Common; on dry prairie and hillsides; throughout Prairies and Parklands.

Petalostemon purpureum (Vent.) Rydb. (Fig. 143) PURPLE PRAIRIE-CLOVER

A several-stemmed plant, erect or decumbent, 20–50 cm high, but usually prostrate. Leaves of 3–7 linear leaflets 5–20 mm long, sparingly hairy or glabrous. Flower spikes dense and cylindric, 1–5 cm long, with red or purple flowers. Common; on hillsides, dry banks, and prairie; throughout Prairies. A densely hairy form, var. *pubescens* (Gray) Fassett, is sometimes found in the southwest.

Petalostemon villosum Nutt. HAIRY PRAIRIE-CLOVER

A densely hairy plant 20–50 cm high, branching from the base; leaves bearing 7–17 closely packed leaflets 5–15 mm long, covered with silky hairs. Flower spikes either single or clustered, 2–10 cm long, flowers reddish purple or pink. Found occasionally; in sandhills; southeastern Parklands.

Psoralea breadroot

Perennial herbs with glandular-dotted foliage and leaves. Leaves palmately compound, of 3–7 leaflets. Flowers in spikes or racemes, perfect, with stamens in 1 or 2 bundles. Pods ovoid, short, 1-seeded, not splitting open at maturity, but opening very irregularly. Roots, especially those of *P. esculenta*, were used by Indians for food.

- 1. Plants densely long-hairy; flowers in a short, dense spike. *P. esculenta*
- Plants not densely long-hairy; flowers in racemes or loose spikes. 2
- 2. Plants silvery hairy; leaves with 3–5 leaflets; inflorescence loose. *P. argophylla*
- Plants not silvery hairy; leaves and stem glandular-dotted; leaves with 3 leaflets; inflorescence dense. *P. lanceolata*

Psoralea argophylla Pursh SILVERLEAF PSORALEA

An erect much-branched plant 30–60 cm high with silvery whitish hairiness throughout. Leaves of 3–5 obovate silvery-haired leaflets 10–35 mm long. Flowers borne on interrupted spikes in clusters of 2 or 4, about 6 mm long, blue fading during drying. Common; on dry to moist grassland; throughout Prairies and Parklands. Syn.: *Psoralidium argophyllum* (Pursh) Rydb.




Fig. 143. Purple prairie-clover, *Petalostemon purpureus* (Vent.) Rydb.

A low, stout short-stemmed plant 10–50 cm high, densely covered with loose white hairs, growing from a large tuberous starchy root or cluster of roots. Leaves of 5 leaflets 2–5 cm long. Flowers in a dense oblong spike 3–8 cm long, blue, a little longer than the greenish sepals. Fairly common on prairie and in sheltered places, or sandy banks throughout Prairies. Roots edible, raw or cooked. Syn.: *Pedimelum esculentum* (Pursh) Rydb.

A low-growing semiprostrate or erect plant 20–50 cm high, with glandular-dotted stems; the whole plant pale yellowish green. Very long stringy roots with ramifications extending for many meters. Leaves of 3 linear-lanceolate leaflets 10–35 mm long. Flowers about 6 mm long, pale bluish white, in short dense spikes. Fruit a globular dotted lemon-shaped pod 5–8 mm in diam containing a single seed. A species of sandhills and sandy land, often the dominant plant in some areas, especially on partly stabilized dunes. The rough roots often exposed for long distances, bridging the gap where sand has been blown out between dunes. Very common in suitable sandy sites throughout the southwestern portion of the area, but not found on heavier soils. Not palatable to livestock. Syn.: *Psoralidium lanceolatum* (Pursh) Rydb.

Thermopsis golden-bean

An erect, branched perennial 15–50 cm high, usually in large patches, from running rootstocks. Leaves of 3 obovate leaflets 2–4 cm long, with appressed silky gray hairs, leaf stalks large and leaf-like, stipules at junction of stem. Very bright golden yellow flowers, 1–2 cm long, in rather dense racemes; stamens separate. Pods 3–7 cm long, curved, grayish hairy, and containing 10–13 seeds. An early blooming plant and one of the most striking and colorful early spring flowers. The milk from cows that have eaten the flowers of this species is said to have a peculiar odor and flavor. The fruit of golden-bean has caused **severe sickness** in children. Very common; in great masses along roadsides, on edges of buffalo wallows, and on hillsides; throughout Prairies, also in sandy areas in Parklands.

Trifolium clover

Perennial or biennial herbs, with leaves of 3 leaflets and flowers in short very dense head-like racemes. Flowers perfect; stamens diadelphous. European plants much used for forage and lawns, but escaped from cultivation.

- 1. Flowers yellow or white. 2
- Flowers purple or pinkish. 4
- 2. Flowers white; perennial, with extensively creeping stems, rooting at the nodes. *T. repens*
- Flowers yellow; annual, with erect or ascending, not creeping, stems. 3



Fig. 144. Indian breadroot, *Psoralea esculenta* Pursh.



Fig. 145. Lance-leaved psoralea, *Psoralea lanceolata* Pursh.




Fig. 146. Golden-bean, *Thermopsis rhombifolia* (Nutt.) Richardson.

3. Leaflets to 15 mm long, with the terminal one nearly sessile; stipules not dilated. *T. aureum*
 Leaflets to 10 mm long, with the terminal one petiolulate; stipules dilated below. *T. campestre*
4. Plants pubescent; flowers sessile in the heads. *T. pratense*
 Plants glabrous or nearly so; flowers pedicelled in the heads. *T. hybridum*

Trifolium aureum Poll.

YELLOW CLOVER

Plants biennial or annual; stems 15–30 cm high, erect, branched. Leaflets to 15 mm long, oblong-lanceolate; petiole of terminal leaflet as long as that of the lateral ones. Flowers 5–8 mm long, golden yellow; pods 1-seeded. Introduced, cultivated, and occasionally escaped.

Trifolium campestre Schreb.

YELLOW FIELD CLOVER

Similar to *T. aureum*, but the petiole of the terminal leaflet distinctly longer than that of the lateral ones, and the stipules larger. Flowers 4–5 mm long, lemon yellow. Introduced for cultivation, rarely escaped.

Trifolium hybridum L.

ALSIKE CLOVER

An erect species 30–60 cm high, with long-stalked leaves of 3 obovate leaflets 10–25 mm long, smooth. Flowers pink, in globose head-like racemes. Common; in waste places and roadsides, where it has escaped from cultivation; Parklands and Boreal forest.

Trifolium pratense L.

RED CLOVER

An erect somewhat hairy biennial or perennial species. Leaflets large, 1–5 cm long, often with a reddish inverted V on the upper surface of the leaflets. Flowers red, in globose heads. Rarely found in Prairies, but fairly common in waste places in Parklands and Boreal forest.

Trifolium repens L.

WHITE CLOVER

A creeping perennial with smooth hairless leaflets 5–20 mm long, often having a whitish or pale inverted V on the upper surface of the leaflets. Flowers white or somewhat pinkish-tinged, in round head-like racemes. Often used for lawns and occasionally found along roadsides, in meadows, and throughout forest areas, where it has escaped from cultivation.

Vicia vetch

Annual or perennial herbaceous vines, with pinnate leaves and even-numbered leaflets, the terminal leaflet being replaced by tendrils. Flowers perfect, in spikes or racemes; stamens diadelphous. Vetches distinguishable from vetchlings (*Lathyrus*) by having the style or female organ not flattened and merely a tuft of hairs at its summit instead of down one side. Good forage for livestock.

1. Flowers 1 or 2 (seldom 4) in the upper leaf axils. *V. sativa* ssp. *nigra*
 Flowers many in a more or less one-sided raceme. 2
2. Plants annual; villous with more or less long-spreading pubescence. *V. villosus*
 Plants perennial; glabrous or sparsely appressed pubescent. 3
3. Inflorescence with 10–40 flowers in a dense raceme. *V. cracca*
 Inflorescence with 3–9 flowers in a loose raceme. *V. americana*

Vicia americana Muhl. (Fig. 147)

AMERICAN VETCH

A smooth trailing or climbing plant 40–80 cm long. Leaves of 8–14 ovate or elliptic leaflets 15–35 mm long, very strongly veined. Flowers 15–20 mm long, bluish purple, in loose 3- to 9-flowered racemes. Pods smooth, 2–4 cm long. Very common; around bluffs and shady parts of prairie; throughout the Prairie Provinces.

The var. *truncata* (Nutt.) Brewer, Oregon vetch, differs from the species by being somewhat hairy below, especially when young, and having the leaflets abruptly flattened at the apex and sometimes toothed. Sometimes found in southeastern Parkland. Syn.: *V. oregana* Nutt.

The var. *minor* Hook., narrow-leaved vetch, a prostrate trailing plant with 8–12 narrowly linear strongly-veined leaflets 10–35 mm long and very narrow. Racemes 2- to 6-flowered; flowers almost 2 cm long, bluish purple. Pods a little over 25 mm long. Common; on open prairie and dry soil; throughout the Prairies, often persisting after cultivation. Syn.: *V. americana* Muhl. var. *angustifolia* Nees.

Vicia cracca L.

TUFTED VETCH

A tufted weak-stemmed vetch 50–125 cm long, with leaves formed of 8–24 linear-oblong leaflets. Flowers in a dense 1-sided raceme. Rarely found, but has escaped from cultivation; in some localities; Parklands and Boreal forest.

Vicia sativa L. ssp. *nigra* (L.) Erhr.

VETCH

A pubescent annual 50–80 cm high. Leaflets 3–8 pairs, linear, 6–25 mm long. Flowers 1 or 2 together (or rarely 4), 10–30 mm long, purple; pods 25–70 mm long. Introduced, occasionally escaped.

Vicia villosus Roth

HAIRY VETCH

An annual with stems 30–150 cm high, villose. Leaves with 4–12 pairs of linear to elliptic leaflets. Flowers 10–20 mm long, purple to violet, occasionally the wing white or yellowish; the calyx strongly swollen at the base; pods 2–4 cm long. Introduced, occasionally escaped from cultivation.




Fig. 147. American vetch, *Vicia americana* Muhl.

GERANIACEAE—geranium family

Herbs with opposite stipulate leaves. Flowers perfect and regular, with 5 petals and 5 sepals. Stamens either 5 or 10. Style extending into a long beak-like column with short appendages at tip, and splitting from bottom upward into 5 sections, each with a single-seeded capsule bearing a long tail. Plants of this family not palatable to livestock.

- Leaves pinnately lobed or dissected; tails of capsules twisted at maturity; capsule spindle-shaped. *Erodium*
- Leaves palmately lobed or divided; tails of capsules merely curved; capsule round. *Geranium*

Erodium stork's-bill

Erodium cicutarium (L.) L'Hér. STORK'S-BILL

A low prostrate annual with pinnately divided leaves. Sepals 5, somewhat awn-tipped. Flowers about 1 cm across, pink or purplish, and borne in clusters of 2–12 on long flower stalks. Stamens 5 fertile and 5 sterile. Style column very long, 2–4 cm, and splitting into 5 segments, each with a long spirally twisted tail. Introduced from Europe, and occasionally found; around towns; in southern and eastern Prairie Provinces.

Geranium geranium

1. Flowers not over 12 mm across; petals scarcely longer than sepals; annuals or biennials. 2
- Flowers over 12 mm across; petals much longer than sepals; perennials. 4
2. Sepals without bristle tips; seeds smooth. *G. pusillum*
- Sepals bristle-tipped; seeds rough. 3
3. Stalks of individual flowers more than twice as long as calyx; inflorescence loose. *G. bicknellii*
- Stalks of individual flowers not more than twice as long as calyx; inflorescence compact. *G. carolinianum*
4. Petals white. *G. richardsonii*
- Petals rose or purple. *G. viscosissimum*

Geranium bicknellii Britt. BICKNELL'S GERANIUM

A fairly erect annual or biennial plant 15–50 cm high, with a loosely hairy stem. Leaves very deeply dissected into narrow oblong segments 2–6 cm broad. Stalks of leaves and flowers hairy and glandular; petals rose-colored, about the same length as sepals. Mature style column about 25 mm long, with narrow beak about 6 mm long. Inflorescence loose. Fairly common; in Parklands and Boreal forest and on pathways and roads in Cypress Hills.

An erect annual species 15–40 cm high, with stems loosely hairy and somewhat glandular. Leaves deeply cut into wedge-shaped lobed segments 2–6 cm across. Flowers pale pink or whitish, about the same length as sepals, and borne in rather compact clusters. Style column about 25 mm long with a very short narrow beak. Not so plentiful as *G. bicknellii*; found in meadows and waste places; throughout the Prairie Provinces.

A weak-stemmed annual, spreading and hairy, 10–50 cm long. Leaves 1–5 cm long, 5- to 7-lobed, the lobes either entire or 3-toothed. Sepals not bristle-tipped. Flowers pale purple, 6–10 mm across, with often only 5 fertile stamens. Introduced from Europe and occasionally found; in waste places near towns; in eastern Parklands and Boreal forest.

An erect perennial species 30–80 cm high, often with spreading hairs. Leaves 3–10 cm across, deeply 3- to 5-lobed, with cut and toothed segments. Flowers 2–4 cm across, white with pink veins; style column 20–35 mm long. Plentiful; in meadows and open forest; Rocky Mountains and in Cypress Hills.

An erect branching perennial species 30–60 cm high, with the stem and leaf stalks sticky glandular. Leaves 4–10 cm across, hairy, 3–5 times cleft into sharply toothed segments. Sepals hairy and awn-tipped; flowers pink purple, 3–4 cm across, and very showy. Common; in open woodlands; in southern Rocky Mountains and also on south slope of Cypress Hills, but rare elsewhere in the area. *G. strigosum* Rydb., a very similar plant, having stems and leaf stalks with reflexed hairs and not glandular, occasionally found in the north-western Boreal forest and in the Peace River District.

OXALIDACEAE—wood-sorrel family

Low herbs with rootstocks, and palmately divided leaves with 3-leaflets; the leaflets broadly inverted heart-shaped, indented at the apex. Flowers perfect and regular, with 5 petals, 5 sepals, and 10 stamens. Fruit a capsule.

Oxalis wood-sorrel

- 1. Stem creeping, often rooting at nodes. *O. corniculata*
 Stem erect or decumbent in age, not
 rooting. 2
- 2. Stem with appressed hairs; capsule hairy. *O. stricta*
 Stem with loose spreading hairs or nearly
 smooth; capsule smooth. *O. europaea*



HERBARIUM OF THE UNIVERSITY OF ALBERTA
 ST. JOHN'S, ALBERTA

PROCESSED BY: _____

DATE: _____

NO. _____

BY: _____

BY: _____



Fig. 148. Sticky purple geranium, *Geranium viscosissimum* Fisch. & Mey.

Oxalis corniculata L.

CREeping WOOD-SORREL

A decumbent creeping plant, branched at base, with sparse loose hairiness. Leaves trifoliolate, with leaflets 3–10 mm wide. Petals pale yellow, 5–10 mm long. Escaped from cultivation and occasionally found around buildings. Syn.: *Xanthoxalis corniculata* (L.) Small.

Oxalis europaea Jordan

BUSH'S YELLOW WOOD-SORREL

A slender-stemmed erect plant 10–20 cm high, loosely hairy. Leaves trifoliolate, with leaflets 5–20 mm long, bright green. Flowers bright yellow, 10–15 mm across. Capsule smooth, 10–15 mm long. Found in dry soil; has been reported, though seldom, from Boreal forest.

Oxalis stricta L.

YELLOW WOOD-SORREL

A low pale green plant, somewhat decumbent, usually branched from the base. Leaves trifoliolate, with leaflets 1–2 cm wide. Flowers pale yellow, in umbel-like cymes; petals almost 12 mm long. Capsule hairy, 15–25 mm long, rather abruptly pointed at the tip, and borne erect on reflexed stems. Fairly common locally; along roadsides, gardens, and waste places; in Prairies. Syn.: *Xanthoxalis stricta* (L.) Small.

LINACEAE—flax family

Annual or perennial plants having simple leaves without stalks and perfect regular flowers with 5 sepals, 5 petals, and 5 stamens. Fruit a round capsule divided into 4 or 5 cells, each containing 2 flat seeds. Petals fall very readily from the plant, usually not lasting more than a day.

Linum flax

- 1. Flowers blue; sepals without glands. 2
Flowers yellow; inner sepals with marginal glands. 3
- 2. Sepals more than 5 mm long when mature, more than half as long as capsule. *L. lewisii*
Sepals less than 5 mm long when mature, less than half as long as capsule. *L. pratense*
- 3. Sepals persistent; capsule not thickened at base. *L. sulcatum*
Sepals falling off; capsule with firm thickenings at base. *L. rigidum*

Linum lewisii Pursh

LEWIS WILD FLAX

An erect perennial plant, sometimes branched from base, growing 20–60 cm high from a woody root, hairless throughout. Leaves linear, 10–25 mm long, somewhat crowded on stem. Flowers blue, 20–35 mm across; the petals soon falling off. Capsules round, 5 mm in diam. Common; on dry prairie; throughout Prairies and Parklands.

Linum pratense (Norton) Small

MEADOW WILD FLAX

Very similar to *L. lewisii*, but the sepals are less than 5 mm long and less than half the length of the capsule, and the flowers are smaller, 15–25 mm across. Occasionally found but not common on dry prairie; Prairies and Parklands.

Linum rigidum Pursh (Fig. 149)

LARGE-FLOWERED YELLOW FLAX

An erect pale green species 15–40 cm high, with the stem simple below and somewhat branched above. Leaves few, linear, 10–25 mm long, and very easily knocked off. Flowers yellow, 20–25 mm across. Common locally; in sand hills and sandy or very light soils; throughout Prairies and Parklands. Not generally common, but very plentiful where found. Syn.: *Cathartolinum rigidum* (Pursh) Small; *L. compactum* Nelson.

Linum sulcatum Riddell

GROOVED YELLOW FLAX

An annual 5–50 cm high, with the stem angled and somewhat winged, simple below but much-branched above. Leaves linear, 10–25 mm long, soon falling off. Flowers yellow, 8–12 mm across. Not common, but found occasionally; in dry sandy soils; eastern Parklands. Syn.: *Cathartolinum sulcatum* (Riddell) Small.

BALSAMINACEAE—touch-me-not family

Somewhat succulent herbs having simple leaves without stipules. Flowers perfect, but irregular; 3 sepals, 2 of them small and green, the other large, petal-like, and extended back into a bag-like pouch terminating in a nectar-filled spur. Only 3 apparent petals, 2 pairs of petals being united; 5 stamens. Fruit a 5-celled capsule, springing open at maturity and forcibly expelling the seeds. Plants found in moist places.

Impatiens touch-me-not

Flowers orange, spotted, and sharply contracted to spur. *I. biflora*

Flowers light yellow, not spotted, gradually tapering to spur. *I. noli-tangere*

Impatiens biflora Walt.

SPOTTED TOUCH-ME-NOT

An annual, branched, rather succulent plant 30–150 cm high, with stems often tinged with red, and leaves ovate, 2–10 cm long, green or purplish. Petals and the large pouch-like sepals orange, copiously dotted with reddish brown or purplish spots; flower 20–25 mm long. Found on banks of rivers and lakes; throughout Parklands and Boreal forest and also in Cypress Hills. Syn.: *I. capensis* Meerb.




Fig. 149. Large-flowered yellow flax, *Linum rigidum* Pursh.

An annual very similar to *I. biflora*, but with light green or straw-colored stems and pale green leaves. Flowers pale yellow, unspotted, tapering to a spur about 10 mm long. Rare in the Prairie Provinces, but has been found in wet places in woodlands; Boreal forest. Syn.: *I. occidentalis* Rydb.

POLYGALACEAE—milkwort family

Herbs with simple leaves and no stipules. Irregular flowers with 5 sepals, 2 of which are large, colored, and petal-like; either 3 or 5 more or less united petals; keel of petals with a fringed crest.

Polygala milkwort

- 1. Leaves whorled; plants annual. *P. verticillata*
 Leaves alternate; plants perennial. 2
- 2. Flowers, showy, rose purple. *P. paucifolia*
 Flowers white. 3
- 3. Leaves linear or linear-oblong; inflorescence conic. *P. alba*
 Leaves lanceolate to ovate; inflorescence oblong, cylindrical. *P. senega*

Polygala alba Nutt.

WHITE MILKWORT

An erect plant with several stems growing from the base, 15–40 cm high, with alternate linear leaves 5–25 mm long. Flowers borne in spike-like racemes 2–5 cm long, white or somewhat green-tinged. Found occasionally; in dry ground; along extreme southern border of the Prairie Provinces, but its natural habitat is farther south.

Polygala paucifolia Willd.

FRINGED MILKWORT

An erect plant 5–20 cm high, usually branched, with small leaves near the lower part of the stem and larger ones above; upper leaves oval, 1–3 cm long. Flowers few, 3 or 4, rose purple or pink, 1–2 cm long; keel of corolla with a fringed crest. Found occasionally in Boreal forest.

Polygala senega L.

SENECA SNAKEROOT

An erect plant with several stems growing from a thick rootstock to 10–50 cm high. Leaves numerous on the stems, lanceolate, 2–5 cm long. Flowers greenish white, borne in terminal spike-like racemes 2–6 cm long. Roots used for medicinal purposes. Fairly common; around edges of bluffs and in semi-wooded prairie; throughout Parklands.

Polygala verticillata L.

WHORLED MILKWORT

Erect plants with stems 10–40 cm high, divergently branched. Leaves 1–2 cm long, linear to linear-oblong, with the lower ones mostly in whorls of 2–5.

Inflorescence with the lower branches opposite or whorled; racemes 5–15 mm long; flowers 2–3 mm long, whitish to pinkish. Moist grasslands; southeastern Parklands.

EUPHORBIACEAE—spurge family

Erect or prostrate herbs with an acrid milky juice. Annual or perennial with simple, entire opposite or alternate leaves. Flowers unisexual (but both sexes on the same plant), with sepals reduced to a minute scale. Involucre resembling a calyx, with numerous male flowers consisting of a single stamen and a minute bract and one female flower consisting of a 3-lobed ovary, which, when fertilized, extends upward on a thin stalk and bears the capsule containing the seeds. In some cases stalks arising from the involucre bear still more involucre and flowers, thus forming a large branched inflorescence.

Euphorbia spurge

1. Prostrate mat-like annual plants;
inflorescence in leaf axils. 2
- Erect plants; inflorescence in terminal
umbels. 3
2. Leaves entire; seeds not compressed,
3-angled. *E. geyeri*
- Leaves denticulate; seeds compressed,
4-angled. *E. serpyllifolia*
3. Upper leaves with a conspicuous white
margin. *E. marginata*
- Upper leaves not with a white margin. 4
4. Stem leaves finely serrulate; inflorescence
very leafy. *E. helioscopia*
- Stem leaves entire; inflorescence not very
leafy. 5
5. Leaves obovate, rounded or blunt at end. *E. pepus*
- Leaves linear or lanceolate, pointed at
end. 6
6. Leaves less than 3 mm wide, crowded on
stem. *E. cyparissias*
- Leaves more than 3 mm wide, not very
crowded. 7
7. Leaves usually not over 1 cm wide. *E. esula*
- Leaves usually 1–3 cm wide. *E. lucida*

Euphorbia cyparissias L.

CYPRESS SPURGE

A tufted introduced perennial plant growing from a rootstock to 15–60 cm high, with numerous linear pale green leaves 1–2 cm long and less than 3 mm wide densely crowded toward the upper part of the stems. Flowers in a terminal umbel-like inflorescence and also in the upper leaf axils. Escaped from cultivation and becoming weedy in several localities.

An erect bluish green perennial growing from running roots to 15–75 cm high. A few scattered linear or oblong stem leaves 10–35 mm long. A whorl of narrow leaves is located below the inflorescence. Inflorescence somewhat umbel-like; flowers borne on a pair of pale yellowish green leaf-like bracts, with many inconspicuous male flowers and one female flower for each pair of bracts. Female flowers extend upward on a short stalk, forming 3-seeded capsules. The capsules bursting and expelling the seeds, sometimes for a long distance. It is extremely difficult to eradicate this pernicious weed, which spreads by creeping underground rootstocks and also by seed. Becoming very plentiful in many localities in the Prairie Provinces. An introduced plant, native of Europe and Asia.

Euphorbia geyeri Engelm.

PROSTRATE SPURGE

An annual plant, prostrate in a mat on the ground, 5–25 cm across, with small oblong leaves 5–10 mm long, opposite and pale green. Inflorescence minute, in axils of leaves; seeds smooth, reddish. Rare; sandy soils; southeastern Parklands.

Euphorbia helioscopia L.

SUN SPURGE

An erect introduced annual, often branching from the base, 15–60 cm high. Leaves obovate or spatulate, 2–5 cm long, bluntly rounded at tip and narrowed at base. Flowers borne in leafy inflorescence at head of stem. Very scarce; reported as a weed from several locations in the Prairie Provinces.

Euphorbia lucida Waldst. & Kit.

SHINING SPURGE

An introduced perennial 20–70 cm high, from a thick rootstock. Leaves 15–30 mm wide and 5–10 cm long. Very similar to *E. esula*, but differing in the wider leaves and thicker rootstocks. At present very scarce; but has been reported from east central Alberta.

Euphorbia marginata Pursh

SNOW-ON-THE-MOUNTAIN

Erect annual plants 30–80 cm high, softly pubescent. Stem leaves 4–10 cm long, broadly ovate to elliptic. Inflorescence subtended by whorled white-margined leaves. Introduced as an ornamental plant; occasionally escaped and reseeding itself.

Euphorbia peplus L.

PETTY SPURGE

Annuals with erect stems 10–30 cm high; stem leaves 1–2 cm long, obovate to suborbicular, rounded at the tip. Inflorescence with 3–5 rays; capsule winged; seeds pitted. Introduced, and locally weedy.

Euphorbia serpyllifolia Pers.

THYME-LEAVED SPURGE

A native annual plant, usually prostrate and forming mats 5–50 cm across, usually smooth with somewhat reddish stems. Leaves opposite, small, 5–15 mm long, dark green, usually with a conspicuous red line down the center. Flowers inconspicuous in the leaf axils; seeds pitted and wrinkled. The var. *hirtella* (Engelm.) L. C. Wheeler is more or less hairy, but the typical variety is quite smooth. Common; on dry soil and in yards and waste places; throughout the Prairie Provinces. Includes *E. glyptosperma* Engelm.



Fig. 150. Leafy spurge, *Euphorbia esula* L.

CALLITRICHACEAE—water-starwort family

A small slender-stemmed aquatic perennial with opposite entire leaves. Leaves linear but some floating leaves obovate or spatulate. Flowers unisexual (but both sexes on the same plant), growing either singly or in groups of 2 or 3 in the axils of leaves. Male flower having 1 stamen and female flower a 4-celled ovary with 2 stigmas or style branches. Fruit small, nut-like, pendulous, and divided into 4 single-seeded sections.

Callitriche water-starwort

All leaves submersed, linear, and 1-nerved. *C. hermaphroditica*

Upper floating leaves obovate or spatulate, 3-nerved, but submersed leaves linear and 1-nerved. *C. palustris*

Callitriche hermaphroditica L. NORTHERN WATER-STARWORT

A completely submersed aquatic perennial, with stems 10–40 cm long. Leaves opposite, linear, 5–15 mm long. Usually rather crowded on stem and sometimes slightly indented at apex. Found occasionally; in flowing water; throughout the Prairie Provinces, but more frequently in the northern portion. Syn.: *C. autumnalis* L.

Callitriche palustris L. VERNAL WATER-STARWORT

An aquatic perennial, usually with floating stems 2–30 cm long. Submersed leaves stalkless, up to 2 cm long, with a single nerve. Floating leaves obovate. Some plants with all leaves submersed and linear. Found occasionally; in ponds and ditches; throughout the Prairie Provinces.

EMPETRACEAE—crowberry family

Empetrum crowberry

Empetrum nigrum L. BLACK CROWBERRY

A bushy-branched prostrate species, with stems to 40 cm long. Leaves numerous, crowded, 4–8 mm long, widely spreading or reflexed. Flowers small, purple; berries black. Bogs, tundra, and rocky areas; Boreal forest, Rocky Mountains.

ANACARDIACEAE—sumach family

Small trees, shrubs, or very low herbs, with a somewhat acrid sap. Leaves consisting of from 3 to many leaflets. Both perfect and imperfect flowers, with 5 sepals, 5 petals, and 5 stamens. Fruit a drupe.

Rhus sumach

1. Leaflets 13–31 *R. glabra*
Leaflets 3. 2

- 2. Low single-stemmed shrubs; leafing before flowering; leaflets 3–10 cm long. *R. radicans* var. *rydbergii*
- Large shrubs; flowering before leaves appear; leaflets not over 3 cm long. *R. aromatica* var. *trilobata*

Rhus aromatica Ait. var. *trilobata* (Nutt.) Gray

SKUNKBUSH

A shrub 1–2 m high, much-branched, and usually flowering before the leaves appear. Leaves 3-foliolate, with dark green leaflets, somewhat paler beneath, often 3-cleft, 1–3 cm long. Flowers yellowish green, minute, in clusters. Fruit a red globular drupe about 6 mm in diam. Found occasionally; in coulees, thickets, and open-wooded places; throughout Prairies. The unpleasant-smelling bush is not known to be poisonous.

Rhus glabra L.

SMOOTH SUMACH

A shrub 2–4 m high, with pinnate leaves having 13–31 lanceolate leaflets, dark green above, paler beneath. Flowers in terminal clusters, bright green. Bright red fruits borne in dense clusters. Occasionally found on hillsides and dry soils; southeastern Parkland and Boreal forest.

Rhus radicans L. var. *rydbergii* (Small) Rehder (Fig. 151)

POISON-IVY


A single-stemmed erect shrub from a creeping rootstock, 10–30 cm high, with a rather woody stem. Leaves with 3 large bright green strongly veined leaflets 3–10 cm long. Flowers whitish yellow, in dense panicles from axils of leaves. Fruits somewhat globose berries, dull whitish color, about 6 mm in diam. Cattle appear to eat it with impunity, but pollen, sap, and even exhalations from the plant affect susceptible persons and cause severe skin eruptions and other troubles. Found in ravines, shady woodlands; Prairies and Parklands, but rare in Boreal forest. Syn.: *Toxicodendron rydbergii* (Small) Greene.

ACERACEAE—maple family

Trees with a sweet sap and opposite, lobed, or pinnately compound leaves. Flowers either perfect or unisexual, sometimes appearing before the leaves. Fruit (a samara) consisting of 2 carpels joined at the base.

Acer maple

- 1. Leaves pinnate with 3–5 leaflets; flowers unisexual, with male and female flowers on separate trees. *A. negundo* var. *interius*
- Leaves lobed, not pinnate. 2
- 2. Leaves with lobes rhomboid, the base cordate. *A. saccharinum*
- Leaves with lobes deltoid, the base not cordate. 3
- 3. Inflorescence a many-flowered racemose panicle. *A. spicatum*
- Inflorescence a few-flowered corymb. *A. glabrum* var. *douglasii*



A.C. Budd

Fig. 151. Poison-ivy, *Rhus radicans* L. var. *rydbergii* (Small) Rehder.

Acer glabrum Torr. var. *douglasii* (Hook.) Dippel

MOUNTAIN MAPLE

A small tree or tall shrub, with leaves palmately 3- to 5-lobed, coarsely and irregularly serrate. Panicle with fascicles of 2-4 flowers; samaras 15-25 mm long. Found occasionally; in deciduous and mixed forests; southern Rocky Mountains.

Acer negundo L. var. *interius* (Britt.) Sarg.

MANITOBA MAPLE, BOX ELDER

A tree, occasionally 6-7 m high, with rough grayish bark. Leaves of 3-5 lanceolate or ovate toothed leaflets, pale green, 5-12 cm long, each with a short stalk. Flowers appearing slightly before the leaves; unisexual, with male and female flowers on separate trees. Female flowers in small racemes, very small, and greenish. Male flowers consisting of 4 or 5 very small sepals and 4 or 5 stamens, reddish, borne in drooping clusters. Fruit a samara consisting of 2 oval carpels about 10 mm long, joined at the base and each terminating in a broad membranous wing about 25 mm long and 10 mm wide. Found along streams, in ravines, and wooded valleys; on Prairies and Parklands. Syn.: *Negundo interius* (Britt.) Rydb.

Acer saccharinum L.

SILVER MAPLE

Trees 10-20 m high, with leaves deeply 5-lobed, the base distinctly cordate. Flowers in clusters from lateral buds, either staminate or pistillate. Samaras 3.5-5 cm long. Planted as shade trees, and occasionally escaped.

Acer spicatum Lam.

WHITE MAPLE

Small trees 3-10 m high; the leaves clearly 3-lobed with 3 obscure basal lobes. Flowers in 3- to 7-flowered corymbs; samaras 3-4 cm long. Mixed woods and copses; eastern Parklands and Boreal forest.

CELASTRACEAE—staff-tree family

Leaves alternate. *Celastrus*

Leaves opposite. *Pachystima*

Celastrus staff-tree

Celastrus scandens L.

BITTERSWEET

Plants climbing to several meters high, with the stems becoming woody and eventually strangling the supporting trees. Leaves elliptic to ovate, acute, serrulate, 5-10 cm long. Panicles terminal, 3-8 cm long; fruits orange, clustered, subglobose, to 1 cm long. Woods, especially in sandy areas, river valleys, and ravines; southeastern Parklands and Boreal forest.

Pachystima boxwood

Pachystima myrsinites (Pursh) Raf.

MOUNTAIN BOXWOOD

Small much-branched leafy shrubs 30-100 cm high; sometimes spreading to prostrate. Leaves ovate to oblanceolate, 15-30 mm long, serrulate, evergreen. Flowers axillary, solitary or in clusters of 2 or 3; petals reddish brown; capsule 4-5 mm long. Coniferous or mixed woods; Rocky Mountains.

RHAMNACEAE—buckthorn family

- Flowers numerous, in terminal umbels. *Ceanothus*
 Flowers in few-flowered axillary umbels or
 solitary. *Rhamnus*

Ceanothus

- Leaves lanceolate; panicles hemispheric or
 short-ovate. *C. ovatus*
 Leaves ovate or elliptic, glandular serrate;
 panicles compound with numerous umbels. *C. velutinus*

Ceanothus ovatus Desf.

NEW JERSEY TEA

A bushy shrub about 1 m high; leaves 2–6 cm long, obtuse to subacute. Panicles on peduncles 2–5 cm long, with the umbels crowded. Rare; semiopen woods on light soils; southeastern Boreal forest.

Ceanothus velutinus Dougl.

STICKY LAUREL

Shrubs to 2 m high, with stout branches. Leaves evergreen, 2–8 cm long, obtuse or rounded at the tip, with the base subcordate, closely glandular denticulate. Panicles usually compound, with the umbels not crowded. Rare; open woods and slopes; southern Rocky Mountains.

Rhamnus buckthorn

Trees or shrubs, with leaves simple, usually alternate; flowers small, greenish or white; fruit a capsule or berry-like drupe.

1. Some branches ending in a short thorn. *R. catharticus*
 Branches not thorny. 2
2. Flowers unisexual, apetalous; leaves
 serrate. *R. alnifolius*
 Flowers perfect, with petals present;
 leaves entire. *R. frangula*

Rhamnus alnifolius L'Hér.

ALDER-LEAVED BUCKTHORN

A small shrub 1–2 m high, with grayish, very finely hairy branches. Leaves ovate to elliptical, strongly veined, 2–6 cm long. Flowers small and greenish, either single or in umbels of 2 or 3 on short stalks in axils of leaves. Fruit berry-like, black, 6–8 mm in diam, and **poisonous**. Fairly common; in moist woodlands and swamps; in northern and eastern fringes of the Prairie Provinces.

Rhamnus catharticus L.

BUCKTHORN

Shrubs or small trees to 3 m high; lateral branches usually ending in a thorn. Leaves 3–6 cm long, broadly elliptic. Flowers having 4 parts; drupe black, 5–6 mm in diam. Introduced; occasionally establishing in shrubbery.

Tall shrubs, with leaves 5–8 cm long, usually obovate or oblong. Flowers in axillary umbels; drupe at first red, becoming black. Introduced; occasionally establishing in shrubbery.

VITACEAE—grape family

Woody vines climbing by means of tendrils. Leaves alternate, either lobed or palmately divided. Flowers in panicles, usually unisexual, sometimes both perfect and unisexual. Fruit a several-seeded berry.

- Leaves simple, lobed. *Vitis*
- Leaves palmately compound, with 5–7 leaflets. *Parthenocissus*

Parthenocissus Virginia creeper

- Leaves dull above, paler beneath; tendrils adhesive with disks; berry about 6 mm in diam. *P. quinquefolia*
- Leaves glossy above, slightly paler beneath; tendrils not adhesive, without disks; berry 8–10 mm in diam. *P. inserta*

Parthenocissus inserta (Kerner) Fritsch LARGE-TOOTHED VIRGINIA CREEPER

A straggling vine with smooth bark and long tendrils. Leaves of 5 or 6 lanceolate leaflets 4–10 cm long, large-toothed, glossy on upper side, and turning red in the fall. Tendrils branched with few, if any, adhesive disks at ends. Berries bluish black. Not common; but found in moist woods and shady banks; in eastern Parklands and Boreal forest as far north as Riding Mountain. Syn.: *Psedera vitacea* (Knerr) Greene.

Parthenocissus quinquefolia (L.) Planch. VIRGINIA CREEPER

A tall climbing vine, usually with rather warty branches. Tendrils bearing an adhesive disk at end. Leaves usually of 5 ovate dull green leaflets 5–12 cm long, and coarsely toothed. Fruit blue with a slight bloom. Found occasionally; in moist woodlands; in southeastern Boreal forest. Syn.: *Psedera quinquefolia* (L.) Greene.

Vitis grape

Vitis riparia Michx. RIVERBANK GRAPE

A climbing or trailing vine, with greenish, somewhat angled branches. Leaves not lobed or sometimes 2-lobed near the apex and cordate at the base. The inflorescence with a fairly compact panicle; the berries blue, 6–12 mm in diam, and covered with a heavy bloom. In moist woods; in the eastern Parklands and Boreal forest, as far north and west as Riding Mountain.

TILIACEAE—linden family

Tilia basswood

Tilia americana L.

BASSWOOD

Trees to 40 m high. Leaves broadly ovate to suborbicular, sharply serrate, cordate to truncate at the base, glabrate to stellate pubescent above, and with conspicuous tufts of hairs in the vein axils below. Flowers in axillary cyme-like clusters; the petiole with a large, adnate foliaceous bract. Woods in ravines and along rivers; southeastern Parklands and Boreal forest.

MALVACEAE—mallow family

Herbs with alternate, lobed or dissected, nearly round leaves. Flowers perfect, regular, and either single in clusters or in racemes in axils of leaves. Five partly united sepals and 5 petals united at base. Numerous stamens; fruit a many-segmented capsule. Velvetleaf, *Abutilon theophrasti* Medic., hollyhock, *Althea rosea* Cav., and gay mallows, *Lavatera thuringiaca* L., are used as ornamentals, and occasionally escape but are not long persistent.

1. Leaves palmately divided into 5 lobes;
these entire or divided. *Malvastrum*
Leaves more or less lobed, but not
divided. 2
2. Flowers in axillary racemes. *Iliamna*
Flowers in axillary clusters or solitary. *Malva*

Iliamna wild hollyhock

Iliamna rivularis (Dougl.) Greene

MOUNTAIN HOLLYHOCK

A stout perennial 50–200 cm high, sparsely stellate-pubescent. Leaves 5–15 cm across, cordate or reniform in outline, irregularly 4- to 7-lobed, the lobes toothed. Flowers about 5 cm across, pink to purplish or sometimes white; staminal column stellate-pubescent at the base; fruit segments 2- or 3-seeded. Mountain slopes and meadows; southern Rocky Mountains.

Malva mallow

1. Petals 15–35 mm long; erect perennial
plants. 2
Petals 4–15 mm long; erect or prostrate
annuals. 4
2. Flowers mostly in a terminal corymb. *M. moschata*
Flowers in the leaf axils. 3
3. Lower flowers solitary in the leaf axils. *M. alcea*
Lower flowers 2 or more in the leaf axils. *M. sylvestris*

4. Petals about twice as long as sepals. 5
 Petals scarcely longer than sepals. 6
5. Plants erect; carpels net-veined. *M. crispa*
 Plants decumbent, prostrate; carpels
 smooth. *M. neglecta*
6. Lower part of petals hairless; 8–12 car-
 pels, with net-like venation on backs;
 calyx reflexed at maturity. *M. parviflora*
 Lower parts of petals hairy-margined;
 12–15 carpels with smooth backs; calyx
 incurved at maturity. *M. pusilla*

Malva alcea L.

PINK MALLOW

Plants 50–150 cm high, with leaves 3- to 5-lobed, 5–10 cm long. Flowers 5–8 cm across, pale pink or mauve. Introduced as an ornamental; occasionally escaped and established.

Malva crispa L.

CRISP MALLOW

An erect annual 50–150 cm high, with almost circular or kidney-shaped leaves 5–20 cm across, wavy and lobed at the edges. Flowers purplish or white, 10–15 mm across, without stalks, and crowded in leaf axils. Introduced from Europe; occasionally escaped.

Malva moschata L.

MUSK MALLOW

Plants 50–150 cm high, with leaves palmately 5- to 7-lobed, the lobes pin-
 natifid. Flowers in a terminal corymb, the lower ones solitary in the leaf axils,
 4–5 cm across, white to pink. Introduced as an ornamental; occasionally
 escaped and established.

Malva neglecta Wallr.

COMMON MALLOW

A prostrate annual weed with lobed, wavy-margined leaves, roughly kid-
 ney-shaped, 2–7 cm across. Flowers blue, lilac, or whitish, 10–15 mm across,
 petals about twice the length of sepals. Fruit a circular series of 12–15 smooth
 carpels. Introduced from Europe; becoming an increasingly abundant weed of
 gardens and roadsides.

Malva parviflora L.

SMALL-FLOWERED MALLOW

A prostrate mat-forming annual weed with many branches. Leaves 2–6
 cm across, roughly kidney-shaped in outline, with about 7 wavy-margined
 lobes. Flowers pink or lilac, about 10 mm across, with sepals almost as long as
 petals. Fruit consisting of a series of 8–11 carpels or small capsules arranged in
 a circle. Introduced from Europe, but becoming a common weed; of roadsides
 and waste places; throughout most of the Prairie Provinces.

Malva pusilla Sm.

ROUND-LEAVED MALLOW

A much-branched prostrate annual or biennial plant similar to *M.*
parviflora, with pale lilac flowers and wavy-margined, somewhat kidney-
 shaped leaves. An introduced weed, becoming common; on wasteland and
 roadsides.

Plants 50–150 cm high; leaves reniform to orbicular in outline, 5–15 cm across, usually with 5 acute lobes. Flowers 25–60 mm across, dark pink to purple. Introduced as an ornamental; occasionally escaped and established.

Malvastrum false mallow

Malvastrum coccineum (Pursh) Gray (Fig. 153)

A native perennial, with a woody base and running rootstocks. Grows 5–20 cm high, often forming large patches along roadsides and in disturbed prairie. Leaves roughly round in outline and divided to the base into wedge-shaped, lobed, and cleft leaflets. Leaves covered with fine white star-shaped hairs, giving the plant a grayish appearance. Flowers borne in a dense, short, raceme-like inflorescence, brick red (a shade rarely found in prairie flowers), 10–25 mm across. Common on dry prairie in lighter soils throughout Prairies and Parkland. Though common on virgin prairie, it is not generally noticeable because it does not flower profusely or form large clumps, but on disturbed soil or roadside cuts it often takes full possession of large areas and flowers freely.

HYPERICACEAE—St. John’s-wort family

Hypericum St. John’s-wort

- 1. Leaves narrowly lanceolate. 2
 Leaves ovate or elliptic. 3
- 2. Leaves 1- to 3-nerved, linear to oblanceolate; the margins of opposite leaves not meeting. *H. canadense*
 Leaves 5- to 7-nerved, lanceolate; the margins of opposite leaves meeting around the stem. *H. majus*
- 3. Petals yellow, at least twice as long as the sepals; leaves, sepals, and petals black glandular. *H. formosum* var. *nortoniae*
 Petals pinkish to greenish; plants with translucent glands. *H. virginicum* var. *fraseri*

Hypericum canadense L.

Slender-stemmed annuals 10–40 cm high. Leaves 1–3 cm long, usually less than 3 mm wide, 1- or 3-nerved. Inflorescence few-flowered, with yellow petals the same length as or shorter than the sepals. Rare; lakeshores, wet meadows; eastern Boreal forest.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Malva sylvestris* L.

LOCATION: Saskatchewan

COLL: J. L. L.

HT:

YILL: 1960

DATE: 14 September 1960

ELEVATION: 500'

LOCALITY: Anderson farm
O'Brien




Fig. 152. Purple mallow, *Malva sylvestris* L.



A. C. Budd

Fig. 153. Scarlet mallow, *Malvastrum coccineum* (Pursh) Gray.

Hypericum formosum HBK. var. *nortoniae* (M. E. Jones) C. L. Hitchc.

WESTERN ST. JOHN'S-WORT

Perennials with rootstocks; stems 15–60 cm high, few or solitary, simple or branched at the summit. Leaves ovate to oval-lanceolate, sessile, the base more or less clasping. Flowers 15–20 mm across, yellow, in more or less paniculate cymes. Wet mountain meadows and shores; southern Rocky Mountains.

Hypericum majus (Gray) Britt.

LARGE CANADA ST. JOHN'S-WORT

Similar to *H. canadense*, but with leaves to 4 cm long, 3–9 mm wide, 5- to 7-nerved, and somewhat clasping the stem; the lower leaf margins meeting or overlapping. Not common; wet meadows and lakeshores; Boreal forest.

Hypericum virginicum L. var. *fraseri* (Spach) Fern.

MARSH ST. JOHN'S-WORT

Perennial plants with erect stems 30–60 cm high. Leaves ovate-oblong to elliptic, 3–6 cm long, cordate to subcordate at the base. Flowers 15–20 mm across, with petals the same length as or somewhat longer than the sepals. Lakeshores and marshy or boggy areas; eastern Boreal forest.

ELATINACEAE—waterwort family

Elatine waterwort

Elatine triandra Schkuhr var. *americana* (Pursh) Fassett

MUD-PURSLANE

A tufted annual plant growing in either mud or water, rooting along the stems, 10–35 mm long. Leaves obovate, 2–6 mm long. Minute flowers occurring singly in the leaf axils. Neither leaves nor flowers stalked, but attached directly to the stem. Found occasionally along margins of ponds or slow-moving streams; in Boreal forest.

CISTACEAE—rock-rose family

- 1. Petals 3, dark red, minute. *Lechea*
- Petals 5, yellow, conspicuous. 2
- 2. Leaves lanceolate or oblong, the main stem leaves 20–30 mm long. *Helianthemum*
- Leaves narrowly linear or scale-like, 1–4 mm long. *Hudsonia*

Helianthemum rock-rose

Helianthemum bicknellii Fern.

FROSTWEED

Perennials with stems solitary or few together from rootstocks, erect or nearly so. Early flowers 5–12 in a loose terminal raceme, 15–25 mm across; later flowers apetalous, crowded on short axillary branches. Not common; open sandy or stony soils; southeastern Boreal forest.

Hudsonia false heather

Hudsonia tomentosa Nutt.

SAND-HEATHER

A low, densely tufted, shrubby plant 10–20 cm high. Leaves 1–4 mm long, oval, densely imbricated (overlapping like shingles), and almost scale-like, pale grayish hoary. Flowers small, yellow, about 1 cm across, and borne near ends of branches. Found occasionally; on sandy shores and pine lands; Boreal forest.

Lechea pinweed

Underside of leaves glabrous except on the midrib and margins; seeds 4–6, shaped like a section of orange. *L. intermedia*

Underside of leaves pubescent over entire surface; seeds 3 or 4, 2-sided or obscurely 3-sided. *L. stricta*

Lechea intermedia Leggett

PINWEED

Perennial plants with pubescent stems 20–60 cm high. Leaves of the basal shoots lanceolate, 3–7 mm long; stem leaves linear-oblong, sparsely pubescent. Panicle slender, cylindric about half the height of the plants, branches to about 5 cm long. Flowers about 3 mm long, deep red. Rare; sandy forest openings; southeastern Boreal forest.

Lechea stricta Leggett

HAIRY PINWEED

Resembling *L. intermedia*, but coarser, with larger leaves, and more pubescent. Rare; sandy forest openings; southeastern Boreal forest.

VIOLACEAE—violet family

Viola violet

Perennial (rarely annual) herbs with either basal or alternate simple leaves with stipules. Flowers of two kinds: early spring flowers, showy and cross-fertilized; summer flowers, remaining closed and self-fertilized (cleistogamous). The spring flowers perfect, but irregular in shape, having 5 sepals and 5 petals, the lowest petal prolonged into a spur. Fruit an ovoid or cylindric capsule containing 20–60 obovate seeds.

- 1. Plants with well-developed leafy stems. 2
- Plants stemless. 9
- 2. Stipules large, about the same length as the leaf blades; plants annual. 3
- Stipules much smaller than the leaf blades; plants perennial. 4
- 3. Corolla the same length as or shorter than the calyx, 10–15 mm across. *V. arvensis*

Corolla distinctly longer than the calyx, 10–35 mm across.	<i>V. tricolor</i>
4. Flowers yellow.	5
Flowers blue, mauve, or white.	8
5. Leaves lanceolate; cuneate to rounded at base.	<i>V. nuttallii</i>
Leaves orbicular or deltoid in outline; cordate at base.	6
6. Stem leaves few, much smaller than the basal leaves.	<i>V. orbiculata</i>
Stem leaves numerous, the same size as the basal leaves.	7
7. Plants glabrous or nearly so; stipules 2–10 mm long.	<i>V. glabella</i>
Plants softly and densely pubescent; sti- pules 8–20 mm long.	<i>V. pubescens</i>
8. Flowers purple to whitish blue; stipules toothed.	<i>V. adunca</i>
Flowers white to mauve; stipules entire.	<i>V. rugulosa</i>
9. Leaves deeply divided into linear or lan- ceolate lobes.	<i>V. pedatifida</i>
Leaves not deeply divided, entire to crenate.	10
10. Flowers yellow.	<i>V. nuttallii</i>
Flowers white to purple.	11
11. Rhizome thick, fleshy; lateral petals bearded at base.	<i>V. cucullata</i>
Rhizome thin, elongate; petals not bearded.	12
12. Flowers generally purplish or blue, some- times white-centered or with white lines.	13
Flowers generally white, with blue or pur- ple spots or lines.	14
13. Spur about 2 mm long; plants stoloniferous.	<i>V. palustris</i>
Spur about 8 mm long; plants not stoloniferous.	<i>V. selkirkii</i>
14. Plants glabrous or nearly so; leaves acute at apex.	<i>V. blanda</i>
Plants pubescent throughout; leaves rounded at apex.	<i>V. renifolia</i>

Viola adunca J. E. Smith (Fig. 154A)

EARLY BLUE VIOLET

A plant 5–30 cm high, from a woody rootstock, usually showing the remains of the previous season's growth of foliage. Leaves ovate with somewhat cordate bases, 10–20 mm wide. Flowers violet or purple, 10–20 mm long,



Fig. 154. Violets: *A*, early blue violet, *Viola adunca* J. E. Smith; *B*, northern bog violet, *Viola cucullata* Ait.; *C*, Nuttall's yellow violet, *Viola nuttallii* Pursh; *D*, Western Canada violet, *Viola rugulosa* Greene.

with somewhat bearded side petals and a spur almost as long as the petals. Flowering early in May. Fairly common; on prairie that is not too dry, and in shady places; throughout Prairies, Parklands, and southern fringe of Boreal forest.

Viola arvensis Murr.

WILD PANSY

An irregularly branched annual with leafy stems 10–30 cm high. Leaf blades varying from ovate or round lower leaves to oblong or oblanceolate upper ones. Stipules large, toothed, and leaf-like; leaves and center sections of stipules round-toothed at apex. Flowers pale yellow, with upper petals sometimes violet-tipped. Introduced from Europe, has been found as a field weed.

Viola blanda Willd.

SWEET WHITE VIOLET

Plants with slender creeping rhizomes. Leaf blades cordate-ovate, dark green. Flowers fragrant; petals white, with the 3 lower ones veined brownish purple near the base. Rich woods; eastern Boreal forest, southern Rocky Mountains.

Viola cucullata Ait. (Fig. 154B)

NORTHERN BOG VIOLET

A stemless species with all leaves basal. Earliest leaves round, later ones broadly ovate to reniform, heart-shaped at base, wavy-margined, and 3–6 cm across. Flowers fairly large and violet-colored; the side petals bearded and the spurred petal hairy. Flowering from mid-May until late July. Probably the most common of the blue violets; found in moist places on the prairies, woodlands, slough margins, and bogs; throughout the Prairie Provinces.

Viola glabella Nutt.

YELLOW WOOD VIOLET

A species usually with both basal and stem leaves. Plants smooth, with the only hairiness at the upper part of the stem and on the underside of the leaf veins. Flowers yellow, springing from the leaf bases. Found in woodlands; in southern Rocky Mountains.

Viola nuttallii Pursh (Fig. 154C)

NUTTALL'S YELLOW VIOLET

A somewhat hairy species with many basal leaves and a few stem leaves. Leaves mostly lanceolate, tapering to the stalk, and somewhat wavy-margined. Flowers yellow; petals occasionally somewhat bearded. Very early flowering. Common; on open prairie and hillsides; throughout Prairies and Parklands.

Viola orbiculata Geyer

ROUND-LEAVED WOOD VIOLET

Plants small, with 1–3 small stem leaves; the basal leaves 2–5 cm across, orbicular in outline, deeply cordate. Petals pale yellow with the centers purplish-veined and the lateral petals bearded. Moist woods; in Rocky Mountains.

Viola palustris L.

MARSH VIOLET

A stemless species growing from creeping rootstocks. Stalks and leaves smooth and hairless; the leaf blades broadly ovate or heart-shaped, 2–6 cm wide. Flowers pale lilac, occasionally nearly white, with darker veins; the side

petals may be somewhat bearded and the spur short and thick. Quite rare; in cold bogs and along stream banks; Boreal forest and Rocky Mountains.

Viola pedatifida G. Don

CROWFOOT VIOLET

A plant with all leaves basal and leaf blades cleft almost to the base into 3 divisions, each further cleft into 2–4 lobes. Leaves varying to 10 cm wide, and slightly hairy on margins. Flowers violet, very showy. Never abundant; found on prairies and exposed banks; Parklands, Wood Mountain, Cypress Hills.

Viola pubescens Ait.

DOWNY YELLOW VIOLET

A stout-stemmed species 15–30 cm high, covered with soft, downy hairs. Rarely some basal leaves and a few stem leaves near the top of the stem. Leaf blades broadly ovate with heart-shaped bases; stipules fairly large and ovate-lanceolate. Flowers bright yellow, growing on stalks from axils of stem leaves, with bearded side petals and a short spur. Found often; in moist woodlands; eastern Parklands and southeastern Boreal forest.

Viola renifolia Gray

KIDNEY-SHAPED VIOLET

A low-growing plant with running rootstocks and kidney-shaped, wavy-margined leaves with heart-shaped bases. Flowers white, without bearded petals. Cleistogamous flowers purple, on horizontal stalks. Flowering very early in spring. In cold woodlands, forests, and swamps; eastern Parklands, Boreal forest, and Cypress Hills.

Viola rugulosa Greene (Fig. 154D)

WESTERN CANADA VIOLET

A tall woodland species with stems 20–60 cm high growing from numerous stolons. Cordate or heart-shaped leaves pointed at apex, borne on long stalks, and often up to 10 cm across, although becoming small toward the top of stems; often densely hairy beneath. Flowers white with pinkish or purplish veins, although the complete flower is sometimes pale pink. The common white violet of woodlands throughout Western Canada. Transplants readily into gardens and then forms a low, dense mass of plants flowering copiously quite early in the season. Spreads rapidly by its slender, numerous, white roots and may become too aggressive. Very common; in shady woodlands; throughout the Prairie Provinces.

Viola selkirkii Pursh

LONG-SPURRED VIOLET

A delicate species 6–15 cm high, tufted from elongate slender rhizomes. Leaf blades 15–30 mm long, elongating after flowering. Flowers about 15 mm across, pale violet, with the blunt spur 5–8 mm long. Moist or wet woods; Parklands, Boreal forest, and Rocky Mountains.

Viola tricolor L.

PANSY

Annual, biennial, or perennial plants 10–40 cm high, glabrous or somewhat pubescent. Flowers usually 10–25 mm across, sometimes to 35 mm, violet, yellowish, or 3-colored, with a spur 3–6.5 mm long. An introduced species, very variable; occasionally escaped from cultivation, and established.

LOASACEAE—loasa family

Mentzelia sand-lily

Mentzelia decapetala (Pursh) Urban & Gilg. (Fig. 155)

EVENINGSTAR

Stout, erect, biennial plants 15–60 cm high, with rough, pale gray stems. Leaves alternate, oblanceolate to lanceolate, 5–15 cm long; the lower ones with short stalks, but the upper ones stalkless. Blades sharply and coarsely toothed and covered with tiny, white, stiff bristles, making them very rough to touch. Flowers borne singly or in clusters of 2 or 3 at ends of stems or branches; sharp-pointed sepals about 25 mm long; and petals 35–50 mm long, creamy white, narrow, and pointed at apex. Only 5 petals appearing to be 10, the inner ones being petal-like sterile stamens. Flowers opening only in evening. Fruit an oblong capsule about 35 mm long and up to 12 mm thick, opening at the top, and containing many seeds. Occasionally leafy bracts are attached to base or side of covering of capsule. Plentiful locally, but not common; on eroded hillsides and clay banks; throughout Prairies and on badlands. Syn.: *Nuttallia decapetala* (Pursh) Greene.

CACTACEAE—cactus family

Almost leafless perennial plants with stems fleshy, thickened, succulent, and covered with spines. Flowers perfect and regular with many sepals and petals. Fruit a fleshy berry relished by antelope and sheep.

Globose or cushion-like plants; flowers red or

purple. *Mamillaria*

Branching or jointed plants; stems flattened;

flowers large, yellow or orange. *Opuntia*

Mamillaria ball cactus

Mamillaria vivipara (Nutt.) Haw.

PURPLE CACTUS

A cushion-like cactus 3–20 cm high and 3–30 cm across, covered with somewhat cone-shaped tubercles each bearing a cluster of 3–8 reddish brown spines 12–20 mm long. Flowers borne between tubercles, 3–5 cm across, with numerous purple or dark red narrow petals and a yellow center of many stamens. Fruit a pale green fleshy berry 1–2 cm long, turning brown with age, and very sweet and edible when ripe. Very common; on open prairie and hillsides; throughout Prairies and southern fringe of Parklands. Syn.: *Neomamillaria vivipara* (Nutt.) Britt. & Rose.

Opuntia prickly-pear

Cacti with flattened stems jointed and divided into somewhat plate-like sections called internodes. Plant often bearing small, scale-like, reddish leaves, soon falling off. Spines long, with a tuft of barbed bristles at the base. Flowers very showy, with large waxy petals. Fruit quite sweet and edible.



Fig. 155. Eveningstar, *Mentzelia decapetala* (Pursh) Urban & Gilg.

- Internodes very fleshy, often circular in cross section, 3–5 cm long, the terminal one easily breaking loose. *O. fragilis*
- Internodes never circular in cross section, 5–15 cm long, always broader than thick. *O. polyacantha*

Opuntia fragilis (Nutt.) Haw.

BRITTLE PRICKLY-PEAR

A low-growing decumbent cactus, often forming very large mats, red or reddish green. Spines in divaricate groups, 10–25 mm long. Flowers pale yellow, about 5 cm across. Fruit a fleshy berry 15–25 mm long. Common on dry prairie throughout Prairies; rare in western Parklands and Peace River.

Opuntia polyacantha Haw.

PRICKLY-PEAR

A prostrate bright green plant growing in large clumps. The internodes large and much flattened, 5–15 cm long with reddish brown spines 1–5 cm long. Flowers showy, yellow to pinkish orange, 5–8 cm across. Fruit a prickly berry 25–35 mm long containing numerous seeds. Very common; on dry prairies and light soils, increasing greatly through overgrazing and erosion; Prairies.

ELAEAGNACEAE—oleaster family

Shrubs or small trees with silvery scurfy leaves. Flowers either perfect or unisexual, sometimes both kinds of flowers on the same plant and sometimes only one sex on a plant. Flowers with 4 sepals but no petals, and 4 or 8 stamens. Fruit drupe-like. All species in this family have nodules on the root system in which nitrogen compounds are formed.

- Leaves alternate; flowers perfect, with 4 stamens. *Elaeagnus*
- Leaves opposite; flowers unisexual, the sexes being on different plants, with 8 stamens. *Shepherdia*

Elaeagnus oleaster

- Leaves lanceolate; shrubs spiny; twigs white stellate-pubescent. *E. angustifolia*
- Leaves elliptic; shrubs not spiny; twigs brown stellate-pubescent. *E. commutata*

Elaeagnus angustifolia L.

RUSSIAN OLIVE

Shrubs or small trees 4–7 m high. Leaves lanceolate, 3–10 cm long. Flowers inconspicuous, in small clusters; sepals spreading, yellow within, about 4 mm across; fruit about 1 cm long, yellowish with silvery scales. An introduced species, planted as an ornamental and occasionally escaped.

Elaeagnus commutata Bernh.

SILVERBERRY, WOLF-WILLOW

Shrubs or small trees 2–5 m high, with brown scurfy twigs. Alternate leaves silvery, scurfy on both sides, oblong or elliptic, 2–6 cm long. Flowers in

clusters of 2 or 3 in axils of leaves, yellowish, very fragrant. Fruit oval, drupe-like, silvery, about 1 cm long, containing a large stony seed. Very common on lighter soils (where moisture is plentiful); throughout the Prairie Provinces. Spreads rapidly in overgrazed pastures throughout Parklands. In sparse stands, the plant is kept grazed and does not spread.

Shepherdia buffaloberry

Leaves oblong, silvery on both sides; shrub or small tree bearing long thorns. *S. argentea*

Leaves oval, green above, silvery below; low undershrub without thorns. *S. canadensis*

***Shepherdia argentea* Nutt. (Fig. 156)**

BUFFALOBERRY

A thorny shrub 1–5 m high, with whitish branches. Leaves oblong, 2–5 cm long, and densely silvery scurfy on both sides. Flowers unisexual, all flowers on a plant being the same sex, brownish, in small clusters at nodes formed in the preceding season. Fruit rounded, 3–5 mm across, orange and very sour, but after a hard frost being a good jelly fruit. Common around sloughs, in coulees, and on light soils; Prairies.

***Shepherdia canadensis* (L.) Nutt.**

CANADA BUFFALOBERRY

An unarmed undershrub 0.5–3 m high, with brown scurfy branches. Oval or ovate leaves 2–4 cm long, with green upper surface, but with silvery star-shaped hairs on underside. Flowers yellowish, borne at leaf nodes. Female plant producing round or oval fruit 3–5 mm long, reddish or yellowish, and insipid tasting. Fairly common in wooded places and riverbanks in Parklands and Boreal forest, rare in Cypress Hills, along South Saskatchewan River breaks, and adjacent areas.

LYTHRACEAE—loosestrife family

Lythrum loosestrife

***Lythrum salicaria* L.**

PURPLE LOOSESTRIFE

An erect perennial herb with stems 50–100 cm high; foliage glabrous to sparsely pubescent; leaves opposite or whorled, 3–10 cm long. Inflorescence 10–40 cm long, with foliaceous bracts; flowers 15–20 mm across, purple. Introduced, marsh species; Boreal forest.

ONAGRACEAE—evening-primrose family

Annual or perennial herbs with either opposite or alternate leaves. Flowers perfect and generally regular, with usually 4 sepals and 4 petals, and with as many, or twice as many, stamens as sepals. Style slender, either knobbed or



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Shepherdia argentea* Nutt.

LOCATION: Prosser Cr. Sask.

COLL. J. Loomis

1937

CANADIAN

DATE: 7/2/37

ELEVATION: 2000

HABITAT: ...



Fig. 156. Buffaloberry, *Shepherdia argentea* Nutt.

4-lobed at the summit. Fruit either a capsule or a small nut. In some genera seeds silky-tufted.

- 1. Parts of flowers in 2's; fruit covered with hooked bristles; leaves opposite; plants with rootstocks. *Circaea*
 Parts of flowers in 4's; fruit not covered with hooks. 2
- 2. Fruit not splitting open when mature, 1- to 4-seeded; short and ribbed. *Gaura*
 Fruit splitting open either at sides or summit when mature. 3
- 3. Seed with a tuft of silky hairs. *Epilobium*
 Seed without a tuft of silky hairs. 4
- 4. Capsules opening with 2 valves. *Gayophytum*
 Capsules opening with 4 valves. 5
- 5. Petals purple; anthers attached near their base. *Boisduvalia*
 Petals white or yellow; anthers attached near their center. *Oenothera*

Boisduvalia boisduvalia

Boisduvalia glabella (Nutt.) Walp. SMOOTH BOISDUVALIA

Native annual plants 10–30 cm high, with leaves alternate, stalkless, lanceolate-ovate, 10–20 mm long. Flowers small and in leafy terminal spikes, purple or violet, with petals almost 3 mm long. Rare; found in moist, mostly alkaline areas; Prairies.

Circaea enchanter's-nightshade

Leaves ovate, mostly less than twice as long as wide; fruit oblanceolate, not furrowed. *C. alpina*

Leaves oblong-ovate, usually more than twice as long as wide; fruit obovoid, deeply furrowed. *C. quadrisulcata* var. *canadensis*

Circaea alpina L. SMALL ENCHANTER'S-NIGHTSHADE

A slender plant 5–20 cm high, with leaves opposite, stalked, heart-shaped, coarsely toothed, and 2–5 cm long. Flowers borne in a terminal raceme, very small and white, on slender stalks; 2 sepals and 2 notched white petals, with 2 stamens. Fruit an ovoid or club-shaped capsule 2–3 mm long covered with fine hooked hairs. In moist woodlands; Boreal forest, eastern Parklands.

Circaea quadrisulcata (Max.) Franch. & Sav. var. *canadensis* (L.) Hara LARGE ENCHANTER'S-NIGHTSHADE

Plants with erect stems to 60 cm high or more, glabrous below, minutely pubescent above. Leaves 6–12 cm long, rounded or subcordate at the base.

Racemes to 20 cm long; fruit 3.5–5.0 mm long. Rare; moist woods; southeastern Boreal forest.

Epilobium willowherb

Flowers usually in spikes or racemes, with 4 sepals and 4 petals. Leaves opposite or alternate, often with lobed blades. Fruit a long, linear, nearly 4-angled capsule containing many seeds; each seed bearing a tuft of silky hairs at the upper end; capsule splitting lengthwise at maturity.

1. Flowers large, more than 12 mm across. 2
Flowers small, less than 12 mm across. 3
2. Racemes long and narrow, not leafy;
leaves with a vein running parallel to
leaf margin; bracts small. *E. angustifolium*
Racemes very leafy; leaves without a vein
parallel to leaf margin; bracts leaf-like. *E. latifolium*
3. Annual with shreddy, straw-colored bark;
stigma 4-cleft. *E. paniculatum*
Perennial; stigmas entire or merely cleft. 4
4. Leaves linear or narrowly lanceolate, not
toothed, often with rolled margins;
stems not angled. *E. palustre*
Leaves lanceolate and toothed, not rolled
at margins. 5
5. Plants low, often decumbent; leaves ovate
to elliptic. *E. alpinum*
Plants mostly erect; leaves lanceolate to
oblong. *E. ciliatum*

Epilobium alpinum L.

ALPINE WILLOWHERB

Stems usually flexuous, 5–15 cm high. Basal leaves small, 3–10 mm long; stem leaves 10–25 mm long. Flowers solitary or in pairs, 8–12 mm across, pink or rose. Alpine meadows, moist areas; Rocky Mountains.

Epilobium angustifolium L. (Fig. 157)

FIREWEED

An erect, fairly stout perennial plant 60–150 cm high, with alternate, very short-stalked, entire lanceolate leaves 5–15 cm long. Venation of leaves of this species interesting, because of lateral veins running parallel to leaf margins. Leaves slightly paler below than on upper side. Flowers pink to purple, 15–35 mm across, in a long terminal raceme, with a small bract below each flower stalk. Fruit a long, linear, somewhat 4-angled capsule 5–8 cm long, splitting lengthwise to release the numerous tufted seeds. One of the most ubiquitous of plants, being found over most of the northern hemisphere. A good source of honey, and fairly palatable to livestock, especially the upper and more tender tips. Very common; in woodlands, edges of forest, burned-over forests; throughout Boreal forest and Parklands; occasionally along roadsides and in moist places in Prairies. Syn.: *Chamaenerion spicatum* (Lam.) S. F. Gray; *C. angustifolium* (L.) Scop.



Fig. 157. Fireweed, *Epilobium angustifolium* L.

Epilobium ciliatum Raf.

NORTHERN WILLOWHERB

An erect perennial plant 30–100 cm high, sometimes rather sticky. Leaves mostly opposite, lanceolate or ovate-lanceolate, 2–7 cm long. Flowers pink, usually nodding in their early stage, about 6 mm across, with a long ovary tube beneath the flower. Fruit many-seeded, 3–5 cm long; seeds with a white tuft of hairs. Common; in sloughs and wet places; throughout the Prairie Provinces. Syn.: *E. adenocaulon* Hausskn.; *E. glandulosum* Lehm. var. *adenocaulon* (Hausskn.) Fern.

Epilobium latifolium L.

BROAD-LEAVED FIREWEED

An erect or somewhat decumbent perennial plant, usually rather branched, 15–50 cm high. Leaves ovate to ovate-lanceolate, entire, 2–5 cm long, some opposite, some alternate. Flowers large and showy, purple, 2–5 cm across, in short leafy racemes; leaves intermixed with flowers. Found along banks of streams; in southern Rocky Mountains and northeastern Boreal forest. Syn.: *Chamaenerion latifolium* (L.) Sweet.

Epilobium palustre L.

MARSH WILLOWHERB

An erect, branched perennial plant 30–60 cm high. Leaves very narrow and linear, 2–5 cm long, glabrous or covered with curled hairs, too fine to be seen without a small lens. Flowers few, pink or whitish, about 6 mm across. Fruit a capsule, 3–5 cm long; seeds bearing a dingy white tuft of hairs. Found in wet places and swamps; throughout the Prairie Provinces, but nowhere very common. Syn.: *E. lineare* Muhl.; *E. leptophyllum* Raf.

Epilobium paniculatum Nutt.

ANNUAL WILLOWHERB

An annual with a pale, straw-colored, somewhat shreddy stem, erect, branched, 25–50 cm high. Leaves linear or linear-lanceolate, 20–35 mm long. Flowers pink, about 1 cm across, at the end of a tube-like ovary about 1 cm long. Capsule 10–25 mm long, splitting open to release the seeds, each bearing a tuft of white silky hairs. Common; on light and sandy soil; throughout southwestern part of the Prairie Provinces. Syn.: *E. adenocladon* (Hausskn.) Rydb.

Gaura butterflyweed

Perennial, branching herbs with alternate, narrow leaves. Flowers have a long calyx tube, and 4 reflexed sepal lobes, 4 petals, and 8 stamens. Fruit a nut-like capsule, containing 1–4 seeds, without hairy tufts.

Gaura coccinea Pursh

SCARLET GAURA

A perennial, decumbent to erect, 10–30 cm high, almost entirely covered with fine grayish hairs. Leaves numerous, oblong or lanceolate, alternate, without stalks, sometimes wavy-margined or shallowly toothed, 10–30 mm long. Flowers in terminal racemes, white when they emerge, but becoming scarlet in a few hours, about 1 cm across. Capsule about 6 mm long containing 1–4 seeds. Common; on dry prairies and hillsides; throughout Prairies and Parklands. The var. *glabra* (Lehm.) Torr. & Gray very similar to the species, but hairless, except for basal parts of flowers.

Gayophytum willowherb

Gayophytum humile Juss.

LOW WILLOWHERB

Plants 5–15 cm high, glabrous, branched from the base; leaves linear to linear-lanceolate, 1–3 cm long, entire. Flowers axillary, white, 2–3 mm across; capsule 10–15 mm long, flattened. Rare; disturbed areas; southern Rocky Mountains.

Oenothera evening-primrose

A variable genus, some species being tall and erect, others low and stemless. Flowers having the tube of the calyx extended beyond the ovary, with 4 lobes turning downward, 4 petals, and 8 stamens. Flowers very variable in color and size; fruit a capsule opening at the top with 4 lobes and containing many seeds not bearing a hairy tuft.

- 1. Plants distinctly stemmed. 2
 Plants stemless or nearly so. 7
- 2. Much-branched perennials with brown woody stems; flowers yellow; stigma not deeply 4-cleft. *O. serrulata*
 Plants without brown woody stems; stigma deeply 4-cleft. 3
- 3. Flowers white or pinkish; stems white with shreddy bark, usually much-branched. *O. nuttallii*
 Flowers yellow; plant usually not much-branched. 4
- 4. Petals 1–3 mm long. *O. andina*
 Petals more than 5 mm long. 5
- 5. Capsules rounded on the angles, not winged. *O. biennis*
 Capsules 4-angled, winged. 6
- 6. Petals 5–9 mm long, inflorescence nodding. *O. perennis*
 Petals 10–25 mm long, inflorescence erect. *O. fruticosa* var. *linearis*
- 7. Stigma not 4-cleft. *O. breviflora*
 Stigma divided into 4 linear lobes. 8
- 8. Flowers yellow, turning pink with age; petals less than 25 mm long; capsule narrowly winged at angles. *O. flava*
 Flowers white, turning pink with age; petals usually more than 25 mm long; capsule with double crests at angles. *O. caespitosa*

Oenothera andina Nutt.

UPLAND EVENING-PRIMROSE

Low, slender-stemmed annuals 2–15 cm high, with branches spreading from the base, finely pubescent. Leaves alternate, linear to narrowly lanceo-

late. Flowers axillary, yellow; capsule 5–6 mm long. Rare; dry sandy soils; western Prairies.

Oenothera biennis L.

YELLOW EVENING-PRIMROSE

An erect biennial from a taproot, 30–200 cm high. Leaves lanceolate or ovate-lanceolate, 2–15 cm long, stalkless except for short stalks on lower leaves. Flowers yellow, 2–5 cm across, opening in evening, borne in a leafy terminal spike. Capsules 2–3 cm long, finely hairy, and opening at the top when mature. The type species is fairly common in the southeast, where it is an introduced weed. However, the common forms found over most of the Prairie Provinces are var. *canescens* Torr. & Gray and var. *hirsutissima* Gray.

The var. *canescens* Torr. & Gray, western yellow evening-primrose, very similar to the typical form, but usually distinguishable by the length of the free tips or lobed portions of the sepals: 3 mm or more in the typical forms, but only slightly over 1.5 mm in the variety. Very common; on lighter soils; throughout southwestern portion of the Prairie Provinces, and occasionally found in Boreal forest.

Oenothera breviflora (Nutt.) Torr. & Gray

TARAXIA

A stemless perennial growing from a taproot. Leaves finely hairy, deeply incised, 5–12 cm long. Flowers yellow, turning reddish when dried, each 10–15 mm across with 4 sepals, 4 petals, and 8 stamens. Stigma or female portion of flower knobbed at end and not divided into 4 linear lobes. Fruit a 4-winged capsule about 15 mm long. Very rare; has been found on heavy soil on a slough margin; in western Prairies. Syn.: *Taraxia breviflora* Nutt.

Oenothera caespitosa Nutt. (Fig. 158)

GUMBO EVENING-PRIMROSE

A low, stemless perennial from a thick woody root. Leaves oblanceolate to lanceolate, 7–20 cm long, growing from short, winged stalks, sometimes toothed and sometimes wavy-margined. Flowers 3–8 cm across, borne on stalks from the root crown, sweet-scented, white, opening in early morning, but soon fading to a pale pink. Capsules about 3 cm long, without stalks, in cluster on root crown. Found on dry hillsides of gumbo or clay soil and sometimes on gumbo flats; throughout Prairies. Syn.: *Pachylophus caespitosus* (Nutt.) Raim.

The var. *montana* (Nutt.) Durand is similar to the typical form but with smaller flowers. Capsules only about 2 cm long and wavy-margined leaves with fine white hairs around their margins. Flowers very early in the season. One of the sweetest-smelling flowers on the prairie, and a good garden plant. Found more often than the typical form; on clay hillsides; in western Prairies. Syn.: *Pachylophus montanus* (Nutt.) A. Nels.

Oenothera flava (A. Nels.) Garrett

YELLOW LAVAUXIA

A low-growing, stemless perennial from a fleshy taproot. Leaves long and narrow, oblong-lanceolate, very deeply incised, medium green, 10–25 cm long, midrib very prominent on the underside. Flowers yellow, turning pink; petals 10–20 mm long. Capsules winged at the 4 angles, 2–3 cm long. Not common, but found occasionally; in valleys, slough margins, and drainage channels; in western Prairies. Syn.: *Lavauxia flava* A. Nels.



Fig. 158. Gumbo evening-primrose, *Oenothera caespitosa* Nutt.

Oenothera fruticosa L. var. *linearis* (Michx.) Wats.

LARGE SUNDROPS

Stems spreading or erect, pubescent, 30–50 cm high. Stem leaves to 6 cm long, linear to lanceolate. Flowers 2–5 cm across; sepals pubescent; capsules 6–10 mm long. Rare; introduced from the east; southeastern Parklands.

Oenothera nuttallii Sweet

WHITE EVENING-PRIMROSE

An erect, often branched perennial 30–100 cm high from a white, fleshy rootstock. Stems white, somewhat shiny, and with a shreddy bark. Leaves entire, linear, 2–10 cm long, pale green, and with wavy margins. Unpleasantly scented flowers borne from axils of upper leaves, white but turning pinkish as they fade, about 4 cm across. Usually they open in the morning. Capsules narrow, somewhat curved, 4-angled, and about 25 mm long. Very common; on roadsides in light soil areas, and on sandy land, persistent in cultivated fields; Prairies and Parklands. Syn.: *Anogra nuttallii* (Sweet) A. Nels.

Oenothera perennis L.

SUNDROPS

Stems 20–60 cm high, erect, usually simple. Leaves 3–6 cm long, oblanceolate to elliptic. Inflorescence nodding, becoming erect during flowering; flowers 1–2 cm across; capsules 5–10 mm long. Rare; gravelly soils; southeastern Parklands and Boreal forest.

Oenothera serrulata Nutt.

SHRUBBY EVENING-PRIMROSE

An erect or decumbent, woody-crowned, brown, shrubby perennial 10–40 cm high. Leaves alternate, spatulate to linear-oblong or linear, 1–5 cm long, often entire but sometimes with small teeth, pale green. Flowers bright yellow, 10–25 mm across, with an almost disk-like stigma. Capsules 15–20 mm long. Not common; found on dry prairie and hillsides; Prairies and Parklands. Syn.: *Meriolix serrulata* (Nutt.) Walp.

HALORAGACEAE—water-milfoil family

Perennial aquatic or marsh plants with creeping roots. Leaves alternate, or in whorls, with the inconspicuous flowers borne in the axils of the leaves. Petals 4 or absent; fruit a nutlet with 2 or 4 single-seeded sections.

Leaves entire and in whorls; flowers perfect,

with 1 stamen and a pistil. *Hippuris*

Submersed leaves dissected; flowers perfect or

unisexual, with 2–8 stamens. *Myriophyllum*

Hippuris mare's-tail

Hippuris vulgaris L.

MARE'S-TAIL

An aquatic plant with creeping rootstocks, 20–50 cm high, with simple unbranched stems. Leaves in whorls of 6–12 around the stem, linear, 1–3 cm long; those under water are soft and flaccid, the ones above water firm. Flowers lack sepals and petals, but are usually perfect, and borne in the axils of the leaves. Very common; in streams and sloughs, and in water and mud along the banks; throughout the Prairie Provinces.

Myriophyllum water-milfoil

- 1. Most or all of the bracts and flowers alternate. *M. alterniflorum*
- Most or all of the bracts and flowers whorled. 2
- 2. Leaves all in whorls; stamens 8; floral leaves shorter than flowers, and usually entire. *M. spicatum*
- Some leaves alternate; stamens 4; floral leaves longer than flowers, and toothed. *M. pinnatum*

Myriophyllum alterniflorum DC.

WATER-MILFOIL

Plants with very slender stems 15–20 cm long; leaves whorled, usually 5–10 mm long. Spikes emersed, 2–5 cm long; male flowers solitary or in opposite pairs, female flowers whorled at the base of the spike. Shallow water; Boreal forests.

Myriophyllum pinnatum (Walt.) BSP.

PINNATE WATER-MILFOIL

A smaller aquatic perennial 10–20 cm high, with whorled submersed leaves in 3–5 pairs of thread-like divisions. Floral leaves longer than flowers and toothed. Male flowers purplish, with 4 stamens, on spikes 10–20 cm long. A southern species, very uncommon; found in extreme southern part of the Prairie Provinces. Syn.: *M. scabratum* Michx.

Myriophyllum spicatum L.

SPIKED WATER-MILFOIL

An aquatic perennial with branched stems 30–150 cm long. Leaves in whorls, 10–30 cm long, divided into many thread-like divisions. Flowers purplish, small, with 4 petals and 8 stamens, borne in whorls on a spike 2–8 cm long protruding above the water. Common; in streams, sloughs, and lakes; throughout the Prairie Provinces. Syn.: *M. spicatum* L. var. *exalbescens* (Fern.) Jepson; *M. exalbescens* Fern.

ARALIACEAE—ginseng family

- Leaves compound. *Aralia*
- Leaves simple. *Oplopanax*

Aralia wild sarsaparilla

- 1. Plants scapose; leaves and peduncle arising from the rootstock, glabrous. *A. nudicaulis*
- Plants with leafy stems. 2
- 2. Umbels 2–10 in a loose cluster; stems bristly below. *A. hispida*
- Umbels numerous in a compound panicle; stems not bristly. *A. racemosa*

Aralia hispida Vent. (Fig. 159)

BRISTLY SARSAPARILLA

Plants with stout rootstocks; stems up to 1 m high, bristly at the base with slender spines. Leaves few, with the petioles usually shorter than the blades; leaflets 2–6 cm long, oblong to ovate. Inflorescence terminal, with 2–10 umbels; berries globose. Dry woods and rock outcrops; Boreal forests.

Aralia nudicaulis L. (Fig. 160)

WILD SARSAPARILLA

A perennial from a creeping rootstock, usually with a single leaf on a stalk 15–30 cm high. This stalk divides into 3 parts, each of which is again divided into 3–5 leaflets. Leaflets oval, pointed at apex, 5–15 cm long, finely toothed on margins, dark green above, and very pale green below. Flowers greenish, usually in 3 umbels 2–5 cm across, on a flowering stalk 10–40 cm high, growing from the rootstock. Fruit globular, purplish black, about 6 mm long. Common; in shady, rich woodlands and deep wooded ravines; throughout the Prairie Provinces. Occasionally forming the major part of herbaceous vegetation in forest.

Aralia racemosa L.

SPIKENARD

Stout herbs to 2 m high. Leaves few, to 80 cm long, the three primary segments pinnately compound; leaflets up to 15 cm long, serrate. Inflorescence a large panicle with numerous umbels; fruit dark purple. Rare; rich woods; southeastern Boreal forest.

Oplopanax devil's-club

Oplopanax horridum (J. E. Smith) Miq.

DEVIL'S-CLUB

A densely prickly shrub 2–4 m high, with an unpleasant odor. Leaves rounded in outline, 15–50 cm broad, cordate at base, palmately 5- to 7-lobed, petioles and veins spiny. Inflorescence terminal, 10–30 cm long, pubescent and prickly; berry scarlet, 4–5 mm long. Moist woods; Rocky Mountains, western Boreal forest.

UMBELLIFERAE—parsley family

Hollow-stemmed herbs with alternate usually divided leaves. Bases of leaf stalks usually enlarged and sheathing stem. Flowers small, in umbels, simple or compound. Sepals minute, 5 petals, 5 stamens, and 2 styles. Fruit a mericarp consisting of 2 single-seeded carpels united into one capsule, usually ribbed and sometimes winged. A large and widespread family, often difficult to identify positively without the fruit.

(Figs. 159 and 160 overleaf)



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Aralia hispida* Vent.

LOCATION: Mile 103, Hwy 106

COLL: C. Jensen

HT: 1.5 m

C. L. No. 1002

DATE: 14 July 1967

ELEVATION

HABITAT: Shrub




Fig. 159. Bristly sarsaparilla, *Aralia hispida* Vent.



A.C. Budd

Fig. 160. Wild sarsaparilla, *Aralia nudicaulis* L.

1. Foliage spiny; inflorescence bluish; involucre spiny. *Eryngium*
 Foliage not spiny; inflorescence not bluish. 2
2. Fruit with hooked spines; foliage digitately compound. *Sanicula*
 Fruit not spiny; foliage not digitately compound. 3
3. Leaves simple or divided into few well-developed leaflets. 4
 Leaves much divided, with leaflets or segments numerous. 16
4. Leaves simple or with 3 leaflets. 5
 Leaves with more than 3 leaflets. 8
5. Leaves simple, entire. *Bupleurum*
 Leaves serrate to trifoliolate. 6
6. Leaflets very large, 10 cm or more wide. *Heracleum*
 Leaflets much smaller. 7
7. Flowers white; rays of umbel very uneven. *Cryptotaenia*
 Flowers yellow; rays rather even. *Zizia*
8. Leaves pinnately divided. 9
 Leaves ternately divided. 10
9. Leaflets linear- to ovate-lanceolate; flowers white. *Sium*
 Leaflets ovate to oblong; flowers yellow. *Pastinaca*
10. Leaflets entire or with a few lobes. 11
 Leaflets finely to deeply serrate. 12
11. Plants tall, 1–2 m high; leaflets cuneate. *Levisticum*
 Plants low, scapose; leaflets linear or lanceolate. *Lomatium*
12. Fruit strongly flattened, broadly winged. *Angelica*
 Fruit flattened but wingless. 13
13. Leaves divided into leaflets of about equal size, usually 3 segments each with 3 leaflets. *Aegopodium*
 Leaves divided into leaflets of unequal size, often with 5 or more segments. 14
14. Fruit linear, more than 1 cm long, usually pubescent. *Osmorhiza*
 Fruit not linear, less than 1 cm long, glabrous. 15
15. Flowers yellow; the central flower in the umbellets sessile. *Zizia*
 Flowers white; the central flower pedicellate. *Cicuta*

16. Flowers in part replaced by axillary bulblets.	<i>Cicuta</i>
Flowers not replaced by bulblets.	17
17. Involucral bracts large, pinnately dissected.	<i>Daucus</i>
Involucral bracts small, slightly or not dissected.	18
18. Umbel simple, few-flowered.	<i>Scandix</i>
Umbel compound, many-flowered.	19
19. Plants scapose.	20
Plants with stem leaves.	22
20. Flowers white; fruit conspicuously winged on marginal and dorsal nerves.	<i>Cymopterus</i>
Flowers yellow; fruit winged on the mar- ginal nerves only or wingless.	21
21. Fruit wingless.	<i>Musineon</i>
Fruit winged.	<i>Lomatium</i>
22. Lower stem leaves opposite; most leaves basal.	<i>Musineon</i>
Lower stem leaves alternate; stem leaves sometimes opposite in the inflor- escence.	23
23. Leaf segments few, usually 5, long and linear.	<i>Perideridia</i>
Leaf segments more than 5.	24
24. Flowers yellow.	25
Flowers white or pinkish.	26
25. Plants strongly scented; introduced annuals.	<i>Anethum</i>
Plants not scented; native perennials.	<i>Lomatium</i>
26. Stout plants with purplish-spotted stems, to 3 m high.	<i>Conium</i>
Plants with stems not purplish-spotted, to 1 m high.	<i>Carum</i>

Aegopodium goutweed

Aegopodium podagraria L.

GOUTWEED

A perennial herb with long creeping rootstocks; stems erect, 40–90 cm high. Lower leaves long-petioled, mostly with 9 leaflets, very similar in size and shape; leaflets 3–8 cm long, oblong to ovate, sharply serrate. Umbels 6–15 cm wide, dense, with 15–25 primary rays; fruit 3–4 mm long. Introduced; formerly much cultivated for medicinal purposes and as a vegetable; occasionally escaped.

Anethum dill

Anethum graveolens L.

DILLSEED

Glabrous and more or less glaucous herbs, to 1.5 m high, often much branched above. Leaves ovate in outline, pinnately dissected into numerous filiform segments, 5–20 mm long. Umbels to 15 cm across, usually with 30–40 primary rays; fruit 3–5 mm long. Introduced; occasionally escaped from cultivation.

Angelica angelica

- 1. Flowers yellow; the involucrel bracts large. *A. dawsonii*
- Flowers white or pinkish; the involucrel bracts small or none. 2
- 2. Ovaries glabrous; pedicels conspicuously webbed. *A. arguta*
- Ovaries hispid; pedicels glabrous. *A. genuflexa*

Angelica arguta Nutt.

LYALL'S ANGELICA

A tall, stout, smooth-stemmed plant 60–150 cm high, with a woody root. Leaves several times pinnately compound; leaflets ovate to lanceolate, 2–10 cm long, with toothed margins. Flowers white, in large compound umbels. Fruit oblong and smooth, 4–7 mm long, often reddish or purplish, making the fruiting plant very conspicuous. Whole plant with a pleasing odor. Found only in wet meadows and along mountain streams; southern Rocky Mountains. Syn.: *A. lyallii* S. Wats.

Angelica dawsonii Wats.

MOUNTAIN PARSNIP

A stout plant, 40–80 cm high, with a thick, woody, strongly scented root, glabrous. Leaves with 9–15 leaflets, these 3–6 cm long, sharply serrate. Umbels solitary on long peduncles; the involucre of bracts 1–3 cm long, toothed or divided, sometimes rather inconspicuous. Moist woods; southern Rocky Mountains.

Angelica genuflexa Nutt.

KNEELING ANGELICA

Stout plants 40–100 cm high, with the foliage glabrous to somewhat scabrous. Leaves ternate-pinnate or biternate, with the main divisions commonly reflexed, the rachis geniculate; leaflets 4–10 cm long, serrate to incised. Inflorescence hispid to pilose; umbels with 20–40 uneven rays; bractlets linear; fruit 3–4 mm long. Moist forests; western Parkland, Boreal forest, and Peace River.

Bupleurum thorough-wax

Bupleurum americanum Coult. & Rose

THOROUGH-WAX

Perennial plants with a woody caudex; stems 10–30 cm high. Basal leaves entire, linear-lanceolate; stem leaves clasping, oblong to linear. Flowers yellow, in small umbels; bractlets conspicuous. Rare; gravelly and rocky areas; southern Rocky Mountains.

Carum caraway

Carum carvi L.

CARAWAY, ANIS

Biennial plants, with glabrous stems to 1 m high. Leaves ovate, pinnate; the leaflets once or twice pinnately divided into narrow segments, 5–15 mm long. Umbels with 7–15 rays, bractless or with a few linear bracts; fruit 3–4 mm long. Introduced; occasionally escaped from cultivation.

Cicuta water-hemlock

Marsh plants with smooth stems and pinnately compound leaves, growing from stout, often tuberous, rootstocks. Flowers white, in compound umbels; fruit oblong and slightly flattened. All species **very poisonous**. Genus contains some of Canada's most **poisonous** plants, and as yet no antidote is known. Although all parts of the plant contain toxic substances, the heaviest concentration occurs in the root. Humans as well as livestock are affected.

1. Leaflets very narrow; bulblets in axils of upper leaves. *C. bulbifera*
Leaflets lanceolate; no bulblets in axils of leaves. 2
2. Leaflets very narrowly lanceolate, length-to-width ratio 10:1 or more. *C. mackenzieana*
Leaflets lanceolate, length-to-width ratio about 5:1. *C. maculata* var. *angustifolia*

Cicuta bulbifera L.

BULB-BEARING WATER-HEMLOCK

An erect, rather slender plant 30–90 cm high, with a few fleshy tuberous roots. Leaves two or three times pinnate, with leaf segments very narrowly linear, 2–5 cm long, sparsely toothed. Axils of upper leaves bearing bulblets. Fruits few, orbicular, less than 3 mm long. Common; in swamps and wet meadows; in Boreal forest and Parklands; **very poisonous** to all forms of livestock and to humans.

Cicuta mackenzieana Raup.


WATER-HEMLOCK

Stout marsh plants, with stems to 1.5 m high, often more than 1 cm thick; rootstock short, rather thin; tuberous roots hardly developed. Leaflets 5–8 cm long, usually less than 5 mm wide. Umbels several, 3–8 cm across; fruit 2–2.5 mm long. Lakeshores and bogs; Boreal forest. **Very poisonous**.

Cicuta maculata L. var. *angustifolia* Hook. (Fig. 161)

WATER-HEMLOCK

A stout-stemmed plant from a swollen, bulbous rootstock, which is divided horizontally into chambers. Plants 50–200 cm high, often much-branched. Leaves twice-pinnate with lanceolate to linear-lanceolate leaflets 5–8 cm long, 1–2 cm wide, sharply toothed. Base of leaf stalk swollen and sheathing the stem. Flowers small and white, in compound umbels 3–10 cm across. Seeds oval, slightly less than 3 mm long, yellow, with dark brown ribs, and slightly grooved between the two carpels. Common; in wet and marshy



A.C. Budd

Fig. 161. Water-hemlock, *Cicuta maculata* L. var. *angustifolia* Hook.

places, stream banks, and lakeshores; throughout Prairies and Parklands. **Very poisonous** to all stock and humans. Syn.: *C. occidentalis* Greene; *C. douglasii* (DC.) Coult. & Rose. In var. *maculata*, the leaflets are 2–3 cm wide, and the fruit somewhat larger. Marshes and wet places; southeastern Parklands. **Very poisonous.**

Conium hemlock

Conium maculatum L.

POISON HEMLOCK

An erect branching annual or biennial 60–300 cm high; usually glabrous, nauseous smelling when cut or bruised. Leaves large, divided into numerous leaflets. Umbels terminal, with 10–15 rays. Introduced; occasionally established. **Very poisonous.**

Cryptotaenia honewort

Cryptotaenia canadensis (L.) DC.

HONEWORT

An erect branching plant 30–80 cm high, with lower leaves long-stalked, of 3 thin, incised leaflets 3–10 cm long. Small white flowers on unequal stalks in compound umbels, with neither bracts nor bractlets. Fruit narrowly oblong, 5–6 mm long. Scarce; but has been found in woodlands; southeastern Parklands and Boreal forest.

Cymopterus cymopterus

Cymopterus acaulis (Pursh) Raf. (Fig. 162)

PLAINS CYMOPTERUS

A very low plant from a deep, thick root, with a short stem rarely more than 5 cm high. Leaves bright green, usually ascending, from pinnate to twice pinnate, 7–20 cm long. Flowers white, in an umbel up to 3 cm across. Fruit oval, about 6 mm long, with winged ribs on side and back. Fairly common; on dry prairie and hillsides; throughout Prairies.

Daucus carrot

Daucus carota L.

WILD CARROT

Erect annual or biennial with a taproot, 30–90 cm high. Leaves much divided, with the ultimate segments lanceolate or linear. Umbels terminal, with numerous rays; bracts about as long as the rays, with lobes linear. Introduced; occasional weed in disturbed areas and shores.

Eryngium eryngo

Eryngium planum L.

CROSS-THISTLE

Perennial plants, with stems 25–100 cm high. Basal leaves persistent, 5–10 cm long, 3–6 cm wide, ovate-oblong, cordate at the base, the margins serrate, spinulose. Inflorescence usually bluish with several ovoid-globose heads, 1–2 cm long, 10–15 mm wide; bracts 15–25 mm long, linear-lanceolate, with up to 4 pairs of spiny teeth. Introduced ornamental; occasionally escaped.




Fig. 162. Plains cymopterus, *Cymopterus acaulis* (Pursh) Raf.

Heracleum cow-parsnip

Heracleum lanatum Michx. (Fig. 163)

COW-PARSNIP

A tall, coarse plant, with somewhat woolly or hairy stem, 1–2.5 m high. Leaves stalked, 10–30 cm wide, divided into 3 broad segments, very hairy below. Flowers white, in large flat umbels 15–30 cm across. Fruit oval, 8–12 mm long, with very fine hairs, pale tawny when mature. Common; in shady woodland and moist places; throughout the Prairie Provinces. Easily distinguished from *Cicuta* or *Sium* by its broad leaves and unpleasant odor. Known to cause dermatitis in humans. Syn.: *H. maximum* Bartr.

Levisticum lovage

Levisticum officinale Koch

LAVAS

Stout perennials, with stems 1–2 m high. Leaves to 70 cm long, 60 cm wide, triangular in outline; leaflets long-cuneate, entire below, dentate in the upper part. Umbels 3–10 cm across, with 12–20 rays; bracts numerous, deflexed; fruit 5–7 mm long. Introduced; cultivated as an aromatic herb; the root useful in veterinary medicine; occasionally escaped.

Lomatium prairie parsley

Very short-stemmed perennial growing from a thickened tuberous root, with ternately or pinnately dissected somewhat hairy leaves, and yellow, purple, or white flowers. No bracts below head of flowers but usually small bractlets below the separate small umbels making up the head. Plants covered with fine gray hairs.

- 1. Leaf divided into few elongate, linear leaflets. *L. triternatum*
- Leaf divided into numerous short, narrow ultimate segments. 2
- 2. Ovary and fruit densely puberulent. 3
- Ovary and fruit glabrous. 4
- 3. Bractlets fused at the base; plants villous throughout. *L. villosum*
- Bractlets free from the base; plants sparingly pubescent. *L. sandbergii*
- 4. Bractlets oblanceolate; root bulbous. *L. cous*
- Bractlets lanceolate, widest at base; root not bulbous. 5
- 5. Stem glabrous; foliage puberulent. *L. dissectum* var. *multifidum*
- Stem and foliage more or less densely pubescent. 6
- 6. Stems usually with at least one pair of leaves near the base. *L. macrocarpum*
- Stems leafless or with a single leaf at the base. *L. orientale*




Fig. 163. Cow-parsnip, *Heracleum lanatum* Michx.

Plants stemless or with a leaf at the base, 20–25 cm high, from a globose or somewhat elongate tuberous root. Leaves ternate, with divisions 2- or 3-pinnate; the ultimate segments 1–5 mm long, usually glabrous. Umbel with 10–20 unequal rays; flowers yellow; fruit 7–10 mm long, with the wings narrower than the body. Very rare; grassland slopes; Cypress Hills.

Lomatium dissectum (Nutt.) Mathias & Constance

var. *multifidum* (Nutt.) Mathias & Constance

MOUNTAIN WILD PARSNIP

A plant with a thick, knobby spindle-shaped root, 30–90 cm high. Leaves in 2 or 3 sections of several times divided leaflets, the ultimate divisions being linear. Flowers yellow, in a large umbel, followed by flattened fruits 8–12 mm long, with thickened corky side wings. Found occasionally; Foothills region. Syn.: *Leptotaenia multifida* Nutt.

Lomatium macrocarpum (Hook. & Arn.) Coult. & Rose LONG-FRUITED PARSLEY

A semidecumbent plant growing from a thick, deep taproot, usually with very short stems. Leaves and leaf stalks covered with fine grayish hairs. Leaves 3 or 4 times pinnately divided. Flowers white, in compound umbels. Fruit flattened, about 9 mm or more long, with broad creamy white wings almost as wide as body of fruit. Found occasionally; on dry and rocky hillsides; Prairies. Syn.: *Cogswellia macrocarpa* (Nutt.) M. E. Jones.

Lomatium orientale Coult. & Rose

WHITE-FLOWERED PARSLEY

Resembling the preceding species but having white or pinkish flowers, translucent-margined bractlets, and fruit not longer than 6 mm. Rare; but has been reported from dry hillsides; southeastern Parklands. Syn.: *Cogswellia orientalis* (Coult. & Rose) M. E. Jones.

Lomatium sandbergii Coult. & Rose

SANDBERG'S WILD PARSLEY

Plants with a stout taproot; the stems 10–25 cm high, more or less clearly leafy at base. Umbels 2–5 cm across; the bractlets narrowly elongate, sometimes lobed at the apex; flowers yellow; fruit 5–7 mm long, densely puberulent. Rare; open alpine areas; southern Rocky Mountains.

Lomatium triternatum (Pursh) Coult. & Rose

WESTERN WILD PARSLEY

A species 30–60 cm high, with doubly ternate leaves; the leaflets narrow, 5–10 cm long. Yellow flowers, in a compound umbel with no bracts, but usually a few bractlets. Fruit winged and hairless or finely puberulent. Grassland, openings, and slopes; southern Rocky Mountains. Includes *L. simplex* (Nutt.) Macbr.

Lomatium villosum Raf. (Fig. 164)

HAIRY-FRUITED PARSLEY

A low, usually prostrate finely hairy plant, from an enlarged taproot. Leaf stalks usually short; leaves finely dissected, and covered with fine grayish hairs. Flowers yellow, in compound umbels. Very early flowering. Fruit flattened, about 6 mm long, and finely hairy. Common; particularly on heavier soils; Prairies. Syn.: *L. foeniculaceum* (Nutt.) Coult. & Rose.



HERBARIUM OF THE RESEARCH STATION
 SWEET SPRING, S.S.P.R.
 SPECIES: LOMATIUM VILLOSUM
 DATE: 1934
 LOCALITY: KANSAS
 COLLECTOR: [illegible]
 [A small rectangular stamp with illegible text is located to the right of the main text block.]

Fig. 164. Hairy-fruited parsley, *Lomatium villosum* Raf.

Musineon musineon

Low-growing plants with long thickened roots and much-dissected bright green leaves. Flowers yellow, in compound umbels; fruit not flattened as in *Lomatium*. Differing also from *Lomatium* by having no hairs on stem or leaves.

Musineon divaricatum (Pursh) Nutt. LEAFY MUSINEON

A low-growing plant 10–20 cm high, with doubly pinnate bright green leaves and yellow flowers in compound umbels 2–6 cm across. Fruit smooth, 2–3 mm long. Fairly common; on dry ground; throughout Prairies and southern Parklands. Includes *M. trachyspermum* Nutt.

Osmorhiza sweet cicely

Perennial herbs from thick sweet-scented roots. Stems tall and leafy with white or yellow flowers and narrow, linear, bristly-ribbed, long-stalked fruit.

- 1. Flowers yellow or greenish; fruit glabrous. *O. occidentalis*
- Flowers white; fruit pubescent. 2
- 2. Involucral bracts present. *O. aristata*
- Involucral bracts absent. *O. chilensis*


Osmorhiza aristata (Thunb.) Mak. & Yabe SMOOTH SWEET CICELY

A plant 30–90 cm high, from thick, rough, fleshy roots. Stems smooth; leaf stalks twice or three times divided into 3's, ending in lanceolate or ovate leaflets 2–7 cm long, pointed at apex, and coarsely toothed. Flowers small, white, in compound umbels, with green leaf-like bracts at base of each umbel-let or terminal small umbel. Fruit 15–25 mm long, pointed at apex, on a long stalk. Fairly common; in shady moist woodlands. Two varieties occur. In var. *brevistylis* (DC.) Boiv. the foliage pubescent and the style 0.5–2.0 mm long; found in southeastern Parklands and Boreal forest. In var. *longistylis* (Torr.) Boiv. (Fig. 165) the stem glabrous, the foliage glabrous or nearly so, and the style 2.0–3.5 mm long; found throughout Prairies, Parklands, and Boreal forest.

Osmorhiza chilensis Hook. & Arn. BLUNT-FRUITED SWEET CICELY

A plant 30–90 cm high with much-branched stems and fleshy roots. Leaves thin; stalks twice divided into 3's. The compound umbels bearing small white flowers. Fruit bearing a conical beak slightly longer than 1.5 mm. Fairly plentiful in woods. Syn.: *O. divaricata* (Britt.) Suksd. Three varieties are distinguished in this species:

- 1. Fruits 20–25 mm long, longer than the pedicels. var. *chilensis*
- Fruits less than 20 mm long. 2
- 2. Fruits about 15 mm long, mostly shorter than the pedicels. var. *cupressimontana* Boiv.
- Fruits about 10 mm long, shorter than the pedicels, blunt. var. *purpurea* (Coul. & Rose) Boiv.



A.C. Budd

Fig. 165. Smooth sweet cicely, *Osmorhiza aristata* (Thunb.) Mak. & Yabe var. *longistylis* (Torr.) Boiv.

Of these, var. *chilensis* occurs in Cypress Hills and Rocky Mountains; var. *cupressimontana* (*O. obtusa* (Coul. & Rose) Fern.) throughout Parklands; and var. *purpurea* (*O. purpurea* (Coul. & Rose) Suksd.) in southern Rocky Mountains.

Osmorhiza occidentalis (Nutt.) Torr.

WESTERN SWEET CICELY

Plants 30–100 cm high, villous at the nodes; leaves 10–20 cm long, ovate or oblong, 1- to 3-ternate, with the ultimate divisions 2–10 cm long, lanceolate to ovate, serrate to lobed. Umbels 3–7 cm across, with 5–12 rays, stiffly ascending to spreading; fruit linear-fusiform 12–20 mm long; styles 1 mm long or less, glabrous or rarely bristly at base. Open woods and meadows; southern Rocky Mountains.

Pastinaca parsnip

Pastinaca sativa L.

WILD PARSNIP

Plants biennial, to 1.5 m high. Leaflets 5–15, usually ovate or oblong, 5–10 cm long, serrate or lobed. Umbel 10–20 cm across, with 15–25 rays; fruit 5–7 mm long. Introduced; occasionally escaped from cultivation.

Perideridia squawroot

Perideridia gairdneri (Hook. & Arn.) Mathias

SQUAWROOT

A narrow, erect plant 30–80 cm high. Fleshy tuberous roots, often with a fascicle or cluster of tubers. Leaves pinnate, of very narrow leaflets 2–15 cm long. Flowers very small, white, in compound umbels 3–8 cm across. Fruit somewhat flattened, brown, slightly longer than 1.5 mm. The roots were used by Indians for food. Found in meadows, woodlands, and ravines; in southern Rocky Mountains and Cypress Hills. Syn.: *Atenia gairdneri* Hook. & Arn.

Sanicula snakeroot

Sanicula marilandica L. (Fig. 166)

SNAKEROOT

An erect plant 30–100 cm high, with long-stalked basal leaves and stalkless upper leaves. Leaves of 5 or 7 palmately arranged leaflets, oblanceolate, sharply toothed, 4–20 cm long. Flowers greenish white, in compound umbels of several almost globular umbellets, each 6–15 mm in diam. Fruit about 6 mm long, densely covered with fine hooked bristles. Common; in moist, rich woodlands; throughout the Prairie Provinces.

Scandix shepherd's-needle

Scandix pecten-veneris L.

VENUS'-COMB

An annual plant 20–40 cm high, branching from the base, and hispidulous throughout. Leaves pinnately decomposed, with the ultimate segments 2–5 mm long, linear. Umbels 2–4 cm across, with 1–3 primary rays; fruit 5–12 mm long, with a flat straight beak, usually about 4 cm long. An introduced weed, rare.




Fig. 166. Snakeroot, *Sanicula marilandica* L.

Sium water-parsnip

Sium suave Walt. (Fig. 167)

WATER-PARSNIP

Tall, erect marsh plants from stout rootstocks, 60–200 cm high, with smooth, hollow stems. Two kinds of leaves: early underwater leaves twice or three times pinnate with thread-like leaflets; later growing abovewater leaves singly pinnate with linear or narrowly lanceolate sharply toothed leaflets 2–10 cm long. Flowers white, in compound umbels 5–8 cm across. Fruit about 3 mm long, ovate and somewhat compressed, bearing prominent ribs. Very common; in sloughs and wet places; throughout the Prairie Provinces. Thought to have slightly **poisonous** properties, but not so dangerous as the water-hemlock, which it closely resembles. Eaten readily by all classes of livestock without bad effect. Water-parsnip has singly pinnate leaves, with narrow leaflets and many small bracts at base of compound flower umbel; water-hemlock has compound pinnate leaves, with lanceolate leaflets and usually no bracts at the base of the umbel. Both species have bracts at the base of each separate umbellet or single portion of the compound umbel. Syn.: *S. cicutaefolium* Schrank.

Zizia alexanders

Smooth, shiny plants growing to 60 cm high. Leaves simple or divided into broad serrated leaflets. Flowers yellow, with bracts only at base of umbellets.

Basal leaves simple and cordate. *Z. aptera*

All leaves divided into 3 leaflets. *Z. aurea*

Zizia aptera (Gray) Fern.

HEART-LEAVED ALEXANDERS

An erect plant 30–60 cm high, with long-stalked cordate basal leaves and thrice-divided stem leaves having ovate leaflets. Flowers bright yellow, in compound umbels, early flowering. Very common; in moist places; throughout the Prairie Provinces. Syn.: *Z. cordata* (Walt.) Koch.

Zizia aurea (L.) Koch

GOLDEN ALEXANDERS

Very similar to *Z. aptera*, but all leaves twice or three times divided into leaflets. Fairly common; in meadows and woodlands; eastern Prairies and Parkland.

CORNACEAE—dogwood family

Cornus dogwood

Herbs, shrubs, or small trees with simple leaves and perfect flowers; the floral parts in 4's. Fruit a small drupe with a 2-seeded stone.



Fig. 167. Water-parsnip, *Sium suave* Walt.

1. Low herbs, with 4 large petal-like bracts surrounding minute heads of flowers. *C. canadensis*
 Shrubs or small trees, with cymes of flowers not surrounded by bracts. 2
2. Leaves alternate; fruit bluish. *C. alternifolia*
 Leaves opposite; fruit white or blue. 3
3. Twigs pale green, mottled with purple; fruit blue. *C. rugosa*
 Twigs not purplish-mottled; fruit white. 4
4. Leaves mostly lanceolate; branches and inflorescence practically hairless; pith of young branches tawny; bark grayish. *C. racemosa*
 Leaves mostly ovate; branches and inflorescence with appressed hairs and often woolly; pith of young branches white; bark reddish or reddish brown. *C. alba*

Cornus alba L.

RED-OSIER DOGWOOD

A shrub 1–2 m high, with bright reddish-colored twigs and opposite leaves. Leaves generally ovate, with rounded base and pointed apex, 2–8 cm long, paler beneath, and with a few short appressed hairs. Small white flowers borne in flat-topped clusters 2–5 cm across. Flowering in early June, producing globular white fruit about 6 mm in diam. Very conspicuous in winter because of its reddish branches. Common; in woodlands and coulees; throughout the Prairie Provinces. Syn.: *C. stolonifera* Michx.; *Svida stolonifera* (Michx.) Rydb. Besides the typical form, two varieties are recognized. In var. *interior* (Rydb.) Boiv. the lower leaf surface, young twigs, and inflorescence are densely pubescent; found mainly in shrubbery in Prairies. The var. *baileyi* (Coul. & Evans) Boiv. differs from the species by undersides of leaves becoming woolly at maturity; common in southeastern Parklands.

Cornus alternifolia L. f.

ALTERNATE-LEAVED DOGWOOD

A shrub or small tree 2–6 m high, with green branches striped with white. Leaves borne alternately on the stems, ovate, long, 5–12 cm long, pale beneath. Flowers white, followed by bluish fruit. Very rare; but has been found in southeastern Parklands and Boreal forest.

Cornus canadensis L. (Fig. 168)

BUNCHBERRY

Low-growing herbs from a horizontal slender rootstock, with woody-based stems 5–15 cm high. One pair of small leaves about the middle of stem and an apparent whorl of ovate leaves 2–8 cm long near the head of the stem. Four petal-like white involucre bracts 10–25 mm long, at head of stem, surrounding a cluster of tiny greenish flowers. Fruit a bright red drupe about 6 mm across, borne in a cluster. Very common; in shady woodlands; throughout the Prairie Provinces. Syn.: *Chamaepericlymenum canadense* (L.) Aschers. & Graebn.



Fig. 168. Bunchberry, *Cornus canadensis* L.

Cornus racemosa Lam.

PANICLED DOGWOOD

A shrub 2–3 m high, with smooth gray bark. Leaves usually lanceolate, opposite, 4–10 cm long. Flowers and fruit white, stone of fruit slightly grooved. Not common; but has been found in woodlands and along streams; southeastern Boreal forest.

Cornus rugosa Lam.

SPOTTED DOGWOOD

A shrub 1–3 m high, with the young branches yellow green, mostly mottled with purple or red. Leaves ovate to rotund, 7–12 cm long, softly white pubescent below. Inflorescence flat-topped or slightly convex, 3–6 cm across; fruit about 6 mm across, light blue. Shady woods; southeastern Parklands and Boreal forest.

PYROLACEAE—wintergreen family

Perennial, usually evergreen herbs with long rootstocks. Leaves thick and leathery. Flowers perfect, in racemes or corymbs, with 4 or 5 sepals, 4 or 5 petals, and 8 or 10 stamens. Fruit a capsule.

- 1. Plants with leafy stems and corymbose inflorescence. *Chimaphila*
- Plants with leaves in a basal rosette; flowers singly or in a raceme. 2
- 2. Flowers borne singly. *Moneses*
- Flowers in a raceme. *Pyrola*

Chimaphila prince's-pine

Chimaphila umbellata (L.) Bart. var. *occidentalis* (Rydb.) Blake PIPSISSEWA

A low perennial herb with decumbent stems 10–20 cm high, with whorled oblanceolate leaves 2–8 cm long, dark green and shiny above, paler beneath. Flowers pinkish white, borne 4–7 in a cyme. Not common; in forested areas on dry soil; Parklands, Boreal forest, and Cypress Hills.

Moneses one-flowered wintergreen

Moneses uniflora (L.) Gray (Fig. 169A) ONE-FLOWERED WINTERGREEN

A small herb 5–15 cm high, from a slender rootstock. Leaves round to ovate, 10–25 mm long, and borne in 1 or 2 pairs or in whorls near base of stem. Flowers (Fig. 7) solitary and nodding at head of short stem, each flower usually with 5 waxy white petals, fragrant, 10–15 mm across. Two stamens for each petal and a straight style with a knobbed summit. Found in cool, moist woodlands; Parklands, Boreal forest, and Cypress Hills. Syn.: *Pyrola uniflora* L.




Fig. 169. Wintergreens: *A*, one-flowered wintergreen, *Moneses uniflora* (L.) Gray; *B*, pink wintergreen, *Pyrola asarifolia* Michx. var. *incarnata* Fern.; *C*, one-sided wintergreen, *Pyrola secunda* L.

Pyrola wintergreen

Low-growing evergreen herbs from slender creeping rootstocks, with round or oval stalked basal leaves. Inflorescence a long narrow raceme of perfect, regular, nodding flowers, usually with a protruding style. Fruit a small round capsule.

- 1. Style straight, not protruding conspicuously from flower. *P. minor*
 Style conspicuously protruding from flower. 2
- 2. Flowers crowded on one side of the stem (secund); style straight. *P. secunda*
 Flowers not all on one side of stem; style curved. 3
- 3. Petals pink or purplish. 4
 Petals white or greenish. 5
- 4. Leaves subentire; the apex rounded, dull above, thin. *P. asarifolia*
 Leaves denticulate-margined; the apex acute, shiny above, firm. *P. bracteata*
- 5. Leaves ovate-lanceolate, acute; the main veins whitish. *P. picta*
 Leaves rotund to elliptic; the veins not whitish. 6
- 6. Sepals oblong to elliptic, much longer than wide; leaves subrotund. *P. rotundifolia*
 Sepals more or less triangular, as wide as long; leaves elliptic or rotund. 7
- 7. Leaves broadly elliptic to oblong, 3–7 cm long, mostly longer than the petioles. *P. elliptica*
 Leaves obovate to subrotund, 1–3 cm long, often shorter than the petioles. *P. virens*

Pyrola asarifolia Michx.

PINK WINTERGREEN

A plant 10–30 cm high with leathery, shiny basal leaves, cordate at base, 2–5 cm wide. Flowers 7–15, pinkish, 8–12 mm across when fully opened, in an open raceme and usually nodding; with 5 sepals, 5 petals, and a protruding style. Fairly common; in moist woods; throughout the Prairie Provinces. The var. *incarnata* Fern. (Fig. 169B) differs from the typical form by having leaf blades rounded or tapering at base instead of being cordate. Found in similar locations as the typical form.

Pyrola bracteata Hook.

LARGE WINTERGREEN

Plants with stems 20–30 cm high, with 1 or 2 scarious bracts on the stem below the inflorescence. Leaves with petioles about the same length as the blades, 3–8 cm long, ovate to subrotund, dark green above, pale or reddish brown below. Flowers rose purple or dull red; petals 6–8 mm long; floral bracts conspicuous, to twice as long as the pedicel. Rare; damp coniferous woods; southern Rocky Mountains.

Pyrola elliptica Nutt.

COMMON SHINLEAF

Leaves broadly elliptic or oblong, with the petioles often very short. Stems 10–30 cm high; flowers 8–12 mm across, white, often greenish- or pinkish-veined. Not common; damp coniferous woods; Boreal forest.

Pyrola minor L.

LESSER WINTERGREEN

A small species with thin dark green oval or rounded leaves 1–3 cm long, on fairly long basal stalks. Flowers small, about 6 mm across, white or faintly pinkish, in a rather crowded raceme on a stem 5–20 cm high. Found occasionally; in woodlands; Boreal forest, Riding Mountains, and Cypress Hills. Syn.: *Braxilia minor* (L.) House.

Pyrola picta Sm.

WHITE-VEINED SHINLEAF

Stems 10–20 cm high; leaves 2–7 cm long, entire or denticulate, with the petioles about the same length as, or shorter than, the blade, white-mottled along the principal veins. Flowers yellowish white; petals 7–8 mm long; floral bracts shorter than the pedicel. Rare; coniferous forests; southern Rocky Mountains.

Pyrola rotundifolia L.

COMMON WINTERGREEN

Stems 10–30 cm high, with 1 or 2 bracts; leaves 3–8 cm long, subrotund or somewhat ovate or obovate, entire to crenate. Flowers white; petals 6–10 mm long; floral bracts the same length as, to longer than, the pedicel. Coniferous woods; Boreal forest.

Pyrola secunda L. (Fig. 169C)

ONE-SIDED WINTERGREEN

A rather small species, usually growing in colonies from a much-branched rootstock. Leaf blades thin, oval to lanceolate, pointed at both ends, 2–6 cm long. Flowers small, about 6 mm across, and crowded on one side of the short stem, 5–20 cm high. Fairly common; in woodlands and bluffs; throughout the Prairie Provinces. Syn.: *Orthilia secunda* (L.) House.

Pyrola virens Schweigg.

GREENISH-FLOWERED WINTERGREEN

Basal leaves round or broadly oval, rounded at the apex. Leaves thick and dull, on rather long stalks; blades 1–3 cm wide. Flowers greenish white, 8–12 mm across when opened, and borne racemosely on a stem 10–20 cm high; 3–10 flowers in an inflorescence. Found in moist coniferous forest; throughout the Prairie Provinces. Syn.: *P. chlorantha* Swartz.

MONOTROPACEAE—Indian-pipe family

Plants parasitic on the roots of other plants or feeding on dead organic matter (saprophytic). No green coloring in leaves, which are reduced to scales. Flowers perfect, usually drooping, with 6–12 stamens. Fruit a single-celled capsule with numerous seeds.

Petals united; stem sticky with an enlarged

bulb-like base. *Pterospora*

Petals distinct; stem not enlarged and bulb-

like at base, not sticky. *Monotropa*

Monotropa pinesap

Flowers solitary at summit of stem. *U. uniflora*

Flowers several in a raceme. *M. hypopithys* var. *latisquama*

Monotropa hypopithys L. var. *latisquama* (Rydb.) Kear. & Peebles PINESAP

A pinkish- or yellowish-stemmed plant 15–30 cm high, with drooping flowers 10–15 mm long. No leaves, but merely stalkless scales on stem. Very rare; found in rich soil in forests; southern Rocky Mountains and Cypress Hills. Syn.: *Hypopithys latisquama* Rydb.

Monotropa uniflora L. INDIAN-PIPE

Stems 10–30 cm high, scaly, white, with a single drooping flower. Plants turning black during drying. Rare; found in damp woodlands; Boreal forest.

Pterospora pinedrops

Pterospora andromedea Nutt. PINEDROPS

Tall sticky-stemmed plants with enlarged bases, growing from a rounded mass of roots. Stems 15–90 cm high, often more than 2 cm in diam, usually pinkish or purplish. Leaves reduced to numerous narrow scales. Flowers borne in an open raceme, whitish, 6–10 mm across. Fruit a capsule containing numerous small seeds. Rare; found in rich coniferous woods; southern Rocky Mountains and Cypress Hills.

ERICACEAE—heath family

Perennial plants, usually shrubby, with perfect flowers. Fruit usually a capsule, but sometimes a berry or drupe, with the ovary superior.

- 1. Stems erect. 2
- Stems prostrate to decumbent or creeping. 7
- 2. Leaves scaly resinous or rusty tomentose below. *Ledum*
- Leaves not scaly resinous or rusty tomentose below. 3
- 3. Leaves white glaucous below. 4
- Leaves not white glaucous below. 5
- 4. Leaves opposite; corolla rotate, 6–12 mm across. *Kalmia*
- Leaves alternate; corolla urn-shaped, 5–7 mm long. *Andromeda*
- 5. Leaves brownish scurfy; inflorescence a leafy secund raceme. *Chamaedaphne*
- Leaves green, not scurfy; inflorescence not as above. 6

6. Flowers large, about 2 cm across; leaves narrowly oblanceolate. *Rhododendron*
 Flowers smaller, about 1 cm long; leaves obovate. *Menziesia*
7. Leaves 2–6 cm long, oval or spatulate-obovate. 8
 Leaves smaller, usually less than 2 cm long. 10
8. Leaves and petioles pilose; leaf base rounded or cordate. *Epigaea*
 Leaves and petioles glabrous; leaf base tapered. 9
9. Leaves crenate-serrate, conspicuously reticulate-veined below. *Arctostaphylos*
 Leaves very shallowly crenate, with teeth bristle-tipped, not reticulate-veined below. *Gaultheria*
10. Leaves and branches densely brown scaly and scurfy; flowers 1–2 cm across. *Rhododendron*
 Leaves and branches not scaly or scurfy; flowers smaller. 11
11. Leaves leathery, elliptic or obovate, 6–20 mm long. *Arctostaphylos*
 Leaves not as above. 12
12. Leaves linear, closely crowded on the stem. 13
 Leaves not as above. 14
13. Leaves thick, blunt, 4-ranked; flowers solitary. *Cassiope*
 Leaves flat, grooved below, alternate; flowers in terminal clusters. *Phyllodoce*
14. Plants shrubby, with decumbent woody stems, diffusely branched. *Loiseleuria*
 Plants not shrubby, with stems creeping, very slender. *Gaultheria*

Andromeda bog-rosemary

Andromeda polifolia L. BOG-ROSEMARY

A shrub 10–40 cm high; linear-oblong leaves, with margins usually rolled toward underside, dark green above and white below, 2–5 cm long. Inflorescence a few-flowered umbel of pinkish white urn-shaped flowers, each about 6 mm long. Found in swamps and muskeg; Boreal forest.

Arctostaphylos bearberry

Leaves 2–5 cm long, strongly reticulate below, serrulate, not evergreen. *A. alpina*

Leaves 1–3 cm long, not strongly reticulate
below, entire, evergreen. *A. uva-ursi*

Arctostaphylos alpina (L.) Spreng. ALPINE BEARBERRY

Stems tufted or prostrate, with branches 10–20 cm long. Leaves thin, oblanceolate to spatulate-obovate. Flowers 4–5 mm long; berry black, or red in var. *rubra* (Rehder & Wilson) Bean, 6–10 mm long. Swamps, bogs, and muskeg; Boreal forest.

Arctostaphylos uva-ursi (L.) Spreng. BEARBERRY

A prostrate, trailing shrub forming large mats on ground. Leaves spatulate, entire-margined, evergreen and shiny dark green, 1–3 cm long, usually with a number of brown or reddish leaves intermixed. Flowers pinkish white, urn-shaped, about 5 mm long, in short few-flowered racemes. Fruit a bright red berry 6–10 mm in diam. Common; in woodlands on sandy hills and eroded slopes; throughout the Prairie Provinces.

Cassiope mountain-heather

Leaves distinctly grooved on the back. *C. tetragona*

Leaves not grooved, but keeled on the back. *C. mertensiana*

Cassiope mertensiana (Bong.) G. Don WESTERN MOUNTAIN-HEATHER

Low, creeping alpine shrub, with branches 10–30 cm high, ascending. Leaves 3–6 mm long, rounded or keeled on the back. Flowers solitary, 4–6 mm long, white, on pedicels 5–20 mm long, puberulent. Rocky alpine slopes and forests; southern Rocky Mountains.

Cassiope tetragona (L.) D. Don WHITE MOUNTAIN-HEATHER

Very similar to *C. mertensiana*, but the leaves with a conspicuous groove down the back, densely ciliate margins, and the pedicels to 3 cm long and finely glandular. Alpine and subalpine slopes and forests. In the southern Rocky Mountains var. *saximontana* (Small) C. L. Hitchc. occurs, with shorter pedicels and smaller flowers than var. *tetragona*, occurring in the northern Rocky Mountains.

Chamaedaphne leatherleaf

Chamaedaphne calyculata (L.) Moench. (Fig. 170) LEATHERLEAF

Branching shrubs 20–100 cm high, with oblong or obovate, slightly toothed scurfy leaves 1–5 cm long. Flowers white, somewhat urn-shaped, about 6 mm long, and borne in 1-sided racemes. Fruit an angular round capsule about 5 mm across. Found in swamps, bogs, and muskeg; Boreal forest.

Epigaea mayflower

Epigaea repens L. MAYFLOWER

Trailing plants with stems 20–40 cm long, branched, hirsute. Leaves ovate to oblong, 2–10 cm long, obtuse to acute, rounded to cordate at the base, more or less pilose. Inflorescence 2–5 cm long; flowers 10–15 mm long. Rare; coniferous forest; southeastern Boreal forest.



HERBARIUM OF THE RESEARCH STATION
SKIPTOCHT, SASK.

PLANTS OF: CANADA, Sask., and U.S.A., Minn.
SPECIES: *Chamaedaphne calyculata* (L.) Moench.

LOCATION: near by

COLL: J. Looman 1957

COLL. NO: 1242 DATE: 17 July 1957 ELEV: 300 m

HABITAT: open along highway

Fig. 170. Leatherleaf, *Chamaedaphne calyculata* (L.) Moench.

Gaultheria wintergreen

- 1. Leafy stems prostrate; branches less than 10 cm high. 2
- Leafy stems upright; branches 10–20 cm high. *G. procumbens*
- 2. Leaves 5–10 mm long, entire; fruit white. *G. hispidula*
- Leaves 10–15 mm long, serrulate; fruit scarlet. *G. humifusa*

Gaultheria hispidula (L.) Muhl. CREEPING SNOWBERRY

Plants with prostrate stems 20–40 cm long, leafy, bristly, especially when young. Leaves short-petioled, broadly elliptic to rotund, bristly beneath. Flowers few, on pedicels about 1 mm long; fruit white, 5–10 mm long. Damp woods, muskeg, and bogs; Boreal forest. Syn.: *Chiogenes hispidula* (L.) Torr. & Gray.

Gaultheria humifusa (Graham) Rydb. ALPINE WINTERGREEN

Creeping shrub with branches usually less than 10 cm high, slender, glabrous or puberulent. Leaves oval to round-oval, rarely over 15 mm long. Flowers solitary on short-bracted peduncles, axillary; fruit 5–7 mm long, scarlet. Mountain slopes; southern Rocky Mountains.

Gaultheria procumbens L. CHECKERBERRY

Plants with leafy stems from a creeping rootstock, erect, 10–20 cm high; leaves 2–5 cm long, crowded at summit of stem. Flowers 8–10 mm long, on nodding pedicels 5–10 mm long; fruit 7–10 mm long, bright red. Coniferous forests; southeastern Boreal forest.

Kalmia sheep-laurel

Kalmia polifolia Wang. PALE LAUREL

A shrub growing 30–60 cm high, with sharply 2-edged twigs. Leaves opposite, almost stalkless, 1–3 cm long, linear-lanceolate, green above but white and finely hairy below, edges often rolled. Flowers in clusters from upper leaf axils, 10–15 mm across, deep pink to red. Fruits ovoid to spherical capsules about 6 mm long. Found in bogs and swampy places; Boreal forest.

Ledum Labrador-tea

Branching evergreen shrubs with leaves alternate, entire-margined, rolled edges, and covered on underside with rusty-colored woolly hairiness or glandular. Flowers with 5 white petals, borne in umbels or short corymbs. Fruit an oblong or ovate many-seeded capsule.

- Leaves densely woolly or felty tomentose below. *L. palustre*
- Leaves glaucous, glandular-dotted below. *L. glandulosum*

Ledum glandulosum Nutt.

TRAPPER'S-TEA

Stout erect shrub to 2 m high, with twigs puberulent and glandular. Leaves pale green, rugose above, with margins little revolute, oblong to broadly elliptic, glaucous, and glandular-dotted below. Flowers white; petals 5–8 mm long. Moist woods; southern Rocky Mountains.

Ledum palustre L.

LABRADOR-TEA

Erect shrubs to 1.5 m high, with twigs tomentose. Leaves with strongly rolled margins, densely white or rusty red tomentose below, rugose, dark green above. Flowers white, 5–8 mm long. Muskeg, bogs, and wet woods; in alpine and boreal to arctic regions. Two varieties occur:

Leaves lanceolate, length-to-width ratio about

6:1 var. *latifolium* (Jacq.) Michx.

Leaves linear, length-to-width ratio about

12:1. var. *decumbens* Ait.

The var. *latifolium* (*L. groenlandicum* Oeder) is common in northern fringes of Parklands, Boreal forest, and Rocky Mountains; var. *decumbens* occurs in the northern parts of Boreal forest.

Loiseleuria alpine azalea

Loiseleuria procumbens (L.) Desv.

ALPINE AZALEA

Stems woody; branches to 30 cm long, about 5 or 6 cm high. Leaves opposite, narrowly elliptic, 5–8 mm long, evergreen, strongly revolute. Flowers pink or white; petals 3–4 mm long. Rocky tundra; Boreal forest.

Menziesia menziesia

Menziesia ferruginea Sm. var. *glabella* (Gray) Peck

WESTERN MINNIEBUSH

An erect or straggling shrub 1–2 m high, with shredding bark; young twigs finely glandular pubescent. Leaves oblong to obovate, 3–6 cm long, somewhat serrate and ciliate, appressed pubescent above, pubescent on the veins below. Flowers 7–8 mm long, whitish or yellowish to purplish pink, on pedicels 15–20 mm long, glandular pilose. Moist woods and slopes; Rocky Mountains.

Phyllodoce mountain-heather

1. Flowers yellow; peduncles and calyx densely glandular.

P. glanduliflora

Flowers pink to purple. 2

2. Peduncles, calyx, and twigs densely glandular.

P. caerulea

Peduncles glandular; calyx and twigs glabrous.

P. empetriformis

Phyllodoce caerulea (L.) Bab.

PURPLE MOUNTAIN-HEATHER

Shrubby plants with a stout, woody root; stems decumbent, 10–20 cm high. Flowers terminal, 1–4 together on peduncles 1–2 cm long, elongating in fruit. Tundra; northeastern Boreal forest.

Phyllodoce empetrifomis (Sm.) D. Don

BLUE MOUNTAIN-HEATHER

Similar to *P. caerulea*, but flowers few to many together, calyx glabrous, and plants usually less glandular. Mountain meadows, forest openings; Rocky Mountains.

Phyllodoce glanduliflora (Hook.) Cov.

YELLOW MOUNTAIN-HEATHER

Similar to *P. empetrifomis*, but flowers yellow, peduncles conspicuously long glandular pubescent, hairs as long as the diameter of the peduncle, and plants usually very glandular. Mountain meadows, ridges, and forest openings; Rocky Mountains.

Rhododendron rose-bay

Flowers borne in 1- to 3-flowered lateral clusters, axillary. *R. albiflorum*

Flowers borne in a terminal umbelliform cluster. *R. lapponicum*

Rhododendron albiflorum Hook.

WHITE ROSE-BAY

Erect shrubs to 2 m high, with slender branches and shredding bark. Leaves 2–7 cm long, thin, deciduous. Flowers creamy white, about 2 cm across. Mountain forests; southern Rocky Mountains.

Rhododendron lapponicum (L.) Wahl.

LAPLAND ROSE-BAY

Dwarf shrubs 10–30 cm high, freely branched. Leaves leathery, evergreen, 10–15 mm long. Flowers 1–5 in a cluster, with the corolla about 15 mm wide, bright purple. Arctic and alpine tundra; southeastern Boreal forest, southern Rocky Mountains.

VACCINIACEAE—huckleberry family

Mostly shrubs or shrubby plants with alternate leaves. The members of this family are sometimes considered genera within Ericaceae, but have the ovary wholly inferior.

Petals appearing separate, turned backward; dwarf plants with very slender creeping stems. *Oxycoccus*

Petals united into an urn-shaped or campanulate tube. *Vaccinium*

Oxycoccus cranberry

Oxycoccus palustris Pers.

SWAMP CRANBERRY

A dwarf creeping plant with slender trailing stem 10–40 cm long, bearing small ovate leaves 3–8 mm long, green above and whitish below. Flowers in terminal umbels, pink, about 8 mm across, and with petals soon turning backward (reflexed). Fruit a small berry, 6–10 mm in diam, red, and often dark-spotted when young. Fairly common; in cold bogs, swamps, and damp woods; Boreal forest. Syn.: *Vaccinium oxycoccus* L.

Often divided into two species (or two varieties of *V. oxycoccus*):

- Leaves 5–8 mm long, mostly elliptical; berry 8–10 mm long. *O. quadripetalus* Gilib.
- Leaves 3–5 mm long, mostly ovate; berry 6–8 mm long. *O. microcarpus* Turcz.

However, these characters and others are variable. *O. quadripetalus* (*V. oxycoccus* L. var. *oxycoccus*) is described as having peduncles 20–35 mm long, with small bracts at the base; in *O. microcarpus* (*V. oxycoccus* L. var. *microphyllum* (Lange) R. & R.) the peduncles are 10–20 mm long, and the bractlets are halfway up the peduncle. In many collections characters are mixed: long peduncles with small or very small leaves, short peduncles with larger leaves, and bractlets often missing.

Vaccinium blueberry

Usually shrubs or shrubby plants with alternate leaves, evergreen or deciduous. Fruit a blue or red many-seeded berry. Freely hybridizing between species.

1. Stems mostly creeping; leaves leathery, evergreen. *V. vitis-idaea*
 Stems mostly upright; leaves thin, deciduous. 2
2. Flowers in terminal racemes. *V. angustifolium*
 Flowers axillary, solitary or few together. 3
3. Flowers in clusters of 1–4, from axils of bud scales; leaves blunt, entire. *V. uliginosum*
 Flowers solitary in leaf axils; leaves toothed. 4
4. Branches round or nearly so, puberulent. *V. caespitosum*
 Branches more or less sharply angled. 5
5. Pedicels 3 mm long, about as long as the flowers; berries red. *V. scoparium*
 Pedicels longer than 3 mm, longer than the flowers; berries blue or black. 6
6. Shrubs to 40 cm high; leaves to 25 mm long. *V. myrtillus*
 Shrubs to 1 m or higher; leaves to 40 mm long. *V. membranaceum*

Vaccinium angustifolium Ait.

BLUEBERRY

Small shrubs to 30 or 40 cm high; young branches and twigs round or nearly so. Leaves 1–3 cm long, elliptic to lanceolate. Flowers white or pinkish, 3–5 mm long; pedicel recurved, about 6 mm long; berry blue, 4–7 mm in diam. Two varieties can be distinguished:

Leaves serrate, pubescent below on the veins, glabrous above; twigs and branches glabrous. var. *angustifolium*

Leaves entire or nearly so, pubescent on both sides; twigs and branches pubescent. var. *myrtilloides* (Michx.) House

Both varieties form large colonies in coniferous forest, particularly on light soils. The var. *angustifolium* is limited to southeastern Boreal forest; var. *myrtilloides* occurs throughout Boreal forest and Rocky Mountains. Syn.: *V. myrtilloides* Michx.

Vaccinium caespitosum Michx.

DWARF BILBERRY

A very low shrub 5–30 cm high, bearing thin obovate to oblanceolate leaves 10–25 mm long, green and shiny on both sides. Flowers pink or white, ovoid, about 5 mm long, and borne either singly or in groups of 3 or 4. Fruit an edible blue berry with a bloom, about 6 mm in diam. Found in pine woods; Boreal forest and Cypress Hills.

Vaccinium membranaceum Dougl.

BILBERRY

Shrubs to 1 m or higher, glabrous throughout; twigs and young branches slightly but clearly angled. Leaves 2–6 cm long, thin, ovate to oval; petioles 1–2 cm long. Flowers yellowish, 5–6 mm long; pedicel 10–15 mm long; berry black, 8–10 mm in diam. Coniferous forests; southern Rocky Mountains.

Vaccinium myrtilus L.

WHORTLEBERRY

Shrubs 15–30 cm high, with divergent branches; young branches and twigs sharply angular. Leaves 1–2 cm long, ovate to elliptic, rounded to subcordate at base. Flowers greenish white, 3–6 mm long, almost globular; pedicel the same length as or longer than flower; berry 6–10 mm in diam, at first red, later blue to purplish black, glaucous. Coniferous forest; southern Rocky Mountains.

Vaccinium scoparium Leib.

RED WHORTLEBERRY

Low shrubs 10–20 cm high, glabrous throughout; young branches and twigs conspicuously angled. Leaves 5–15 mm long, oval to broadly elliptic, serrulate. Flowers about 3 mm long; pedicels equaling or exceeding the flowers; fruit 3–5 mm in diam, red. Coniferous forests; southern Rocky Mountains.

Vaccinium uliginosum L.

BOG WHORTLEBERRY

Shrubs 10–50 cm high, much-branched; branches round and glabrous. Leaves 10–25 mm long, obovate to oval, firm, rounded or obtuse at the apex, conspicuously reticulate veined below. Flowers mostly in clusters of 2–4, pink, 5–7 mm long; berry 6–7 mm in diam, dark bluish black. Marshes and bogs; Boreal forest and Rocky Mountains.

A low shrub 10–20 cm high, with erect branches growing from a trailing stem. Leaves obovate with rolled edges, dark green and shiny above, paler with black dots beneath, 1–2 cm long. Bell-shaped flowers pink or white, 5–8 mm long, and borne in small terminal clusters. Fruit a dark red, acid berry about 6 mm in diam. Common; in swamps, muskegs, and sandy woodlands; throughout Boreal forest, Rocky Mountains. Syn.: *Vitis-idaea punctata* Moench.

DIAPENSIACEAE—diapensia family

Diapensia diapensia

Diapensia lapponica L.

NORTHERN DIAPENSIA

Dwarf shrubs with matted stems 5–10 cm long, ascending to 3–5 cm high. Leaves crowded, overlapping, 6–15 mm long, spatulate. Flowers solitary, white, on peduncles 1–4 cm long; capsule about 5 mm long. Dry tundra; northeastern Boreal forest.

PRIMULACEAE—primrose family

Herbs with simple leaves. Flowers perfect and regular, usually with 5 sepals, 5 petals, and 5 stamens that are borne opposite the petals.

- 1. Leaves all basal (but leafy bracts below umbels of flowers in *Androsace*). 2
 Leaves not all basal. 5
- 2. Flowers solitary or 1–3 on a peduncle, 1–3 cm high. *Douglasia*
 Flowers in umbels of few to many flowers. 3
- 3. Sepals and petals reflexed (turning backward). *Dodecatheon*
 Sepals and petals not reflexed. 4
- 4. Tube of corolla as long as or longer than calyx; flowers pink or lilac; perennials. *Primula*
 Tube of corolla shorter than calyx; flowers small and white; annuals. *Androsace*
- 5. Leaves mostly alternate, the lower ones sometimes opposite. *Centunculus*
 Leaves opposite or in whorls. 6
- 6. Petals absent, flower pink; low-growing plant of saline areas. *Glaux*
 Petals present. 7
- 7. Plants with scale-like lower stem leaves; upper leaves in a whorl below flowers; usually with sepals, petals, and stamens in 7's. *Trientalis*

- Plants with normal opposite stem leaves. 8
8. Flowers scarlet; plants annual,
procumbent. *Anagallis*
- Flowers yellow; plants perennial, erect. *Lysimachia*

Anagallis pimpernel

Anagallis arvensis L. SCARLET PIMPERNEL

Annual plants with decumbent 4-sided stems 10–30 cm long. Leaves sessile or clasping, oval to ovate, 5–20 mm long. Flowers 4–7 mm across, rotate; capsule 3–4 mm in diam. Introduced; a rare garden weed.

Androsace pygmyflower

1. Plants perennial; flowers large, crowded. *A. chamaejasme*
Plants annuals; flowers small; inflorescence diffuse. 2
2. Bracts below inflorescence elliptic, ovate,
and more or less oblong. *A. occidentalis*
Bracts below inflorescence lanceolate to
awl-shaped. *A. septentrionalis*

Androsace chamaejasme Host PYGMY SHIELDWORT

Plants with dense rosettes; leaves lanceolate, 5–10 mm long, ciliate. Scapes 2–10 cm high, villous. Flowers 8–10 mm across, white with yellow center. Rocky slopes, riverbanks, open forests, and disturbed areas; Rocky Mountains.

Androsace occidentalis Pursh WESTERN PYGMYFLOWER

A very small species, usually with few flowering stems. Uncommon; found on dry, sandy soil; throughout southern part of the Prairie Provinces.

Androsace septentrionalis L. (Fig. 171A) PYGMYFLOWER

Flowering stems few, usually only one well-developed. The var. *diffusa* (Small) Knuth. has many flowering stems, usually of almost equal lengths; calyx almost free of hairs. Occasionally found in eastern parts and in Riding Mountains. The var. *puberulenta* (Rydb.) Knuth. differs from var. *diffusa* by having a hairy calyx. A common species found on eroded and dry soils, and very plentiful on stubble fields and cultivated land in early spring. A common plant, but so small that it is often not noticed.

Centunculus chaffweed

Centunculus minimus L. CHAFFWEED

A small depressed annual 2–10 cm high, with obovate to spatulate leaves, almost stalkless, 3–8 mm long. Tiny pink flowers borne in axils of leaves, about 3 mm across. Rare; but has been found in wet places and lake edges; in Prairies.



Fig. 171. Primroses: *A*, pygmyflower, *Androsace septentrionalis* L.; *B*, mountain shootingstar, *Dodecatheon conjugens* Greene var. *beamishii* Boiv.; *C*, mealy primrose, *Primula incana* M. E. Jones; *D*, northern starflower, *Trientalis borealis* Raf.

Dodecatheon shootingstar

Perennial herbs with basal leaves. Nodding flowers very showy, with reflexed petal lobes in small umbel at head of leafless stem. Five stamens very prominent, united at base, and bearing long purple anthers. Fruit a cylindrical capsule containing many minute seeds.

- Leaves glandular pubescent. *D. conjugens*
- Leaves glabrous. *D. pulchellum*

Dodecatheon conjugens Greene var. *beamishii* Boiv. (Fig. 171B)

MOUNTAIN SHOOTINGSTAR

An early flowering, rather low-growing species with oblanceolate or spatulate leaves 4–10 cm long, pale green, and hairless. Flowers 1–5, borne in an umbel on a long stem from base, varying in color from purple to white, but usually pink; petals reflexed showing prominent purple stamens borne on orange filaments. Petals often having a yellowish inner base with a zigzag purple line, making flowers very handsome when closely examined. In the white phase, which is not unusual, anthers and petals are white with no contrasting colors. Common; in moist areas in grassland; Prairies, Cypress Hills, and Rocky Mountains. Syn.: *D. cylindrocarpum* Rydb.

Dodecatheon pulchellum (Raf.) Merr.

SALINE SHOOTINGSTAR

A midseason-flowering species, with long, spatulate, hairless basal leaves 3–10 cm long. Flowers pink or lilac with a yellowish throat, often with a purple wavy line, 1–12 in an umbel on a stem 10–30 cm high. Fairly common; in wet areas and around saline sloughs, often associated with *Primula incana* M. E. Jones; Prairies, Parklands, and Rocky Mountains.

Douglasia douglasia

Douglasia montana Gray

MOUNTAIN DWARF-PRIMULA

Small cushion plants with the leaves in rosettes. Leaves 5–15 mm long, thick, ciliate. Flowers on a peduncle 1–3 cm high, usually solitary or 1–3 in an umbel. Rare; alpine areas; southern Rocky Mountains.

Glaux sea-milkwort

Glaux maritima L.

SEA-MILKWORT

A low-growing rather succulent leafy perennial from a creeping rootstock. Leaves opposite, stalkless, oval to linear-oblong, 3–12 mm long. Small pinkish flowers, in the axils of the leaves, about 3 mm long. If closely examined, they will be seen to have no petals, merely one floral ring. Fairly common in moist saline locations; throughout the Prairie Provinces.

Lysimachia loosestrife

- 1. Plants not flowering but with bulblets in the upper leaf axils. *L. terrestris*
- Plants flowering with or without bulblets. 2

- 2. Flowers in short crowded spikes in axils of leaves, or terminal. 5
- Flowers borne on separate stalks in axils of leaves. 3
- 3. Leaves usually lanceolate, rounded or almost cordate at base; flowers 2–3 cm across. 4
- Leaves linear, tapering to base, sessile, in whorls; flowers 10–15 mm across. *L. quadrifolia*
- 4. Lower leaves with petioles 6–10 mm long, densely ciliate. *L. ciliata*
- Lower leaves with petioles 10–30 mm long, not ciliate. *L. hybrida*
- 5. Inflorescence terminal, loose; petals elliptic. *L. terrestris*
- Inflorescence axillary, dense; petals linear-lanceolate. *L. thyrsiflora*

Lysimachia ciliata L. (Fig. 172)

FRINGED LOOSESTRIFE

An erect plant 30–100 cm high, with opposite leaves 5–10 cm long, pale green, pointed at apex and rounded at base, borne on stalks 6–10 mm long with a row of hairs on one side of stalk. Flowers 15–25 mm across, with 5 yellow petals somewhat unevenly pointed at tips, and borne in twos or threes in upper leaf axils. There are 5 stamens and 5 infertile stamens (staminodia). Fruit a many-seeded ovoid capsule. Fairly common; in woodlands and in moist spots; throughout the Prairie Provinces. Syn.: *Steironema ciliatum* (L.) Raf.

Lysimachia hybrida Michx.

LANCE-LEAVED LOOSESTRIFE

Very similar in most respects to previous species, but leaves short-stalked and narrower, tapering to base. Found occasionally in moist meadows, but not nearly so common as *L. ciliata*. Syn.: *Steironema hybridum* (Michx.) Raf.

Lysimachia quadrifolia L.

WHORL-LEAVED LOOSESTRIFE

Stems erect, 30–70 cm high, glabrous to sparsely pubescent. Leaves usually in whorls of 4 (3–6). Flowers 10–15 mm across; petals yellow with dark lines. Rare; open woods; southeastern Boreal forest.

Lysimachia terrestris (L.) BSP.

SWAMP-CANDLES

Stems erect, 40–80 cm high, simple or branched. Leaves 5–10 cm long, narrowly lanceolate. Racemes erect, 10–30 cm long, many-flowered, loose, with pedicels to 20 mm long; flowers 10–15 mm across, yellow. Late in the season small elongate purplish bulblets appear in the upper leaf axils. Uncommon; swampy areas and lakeshores; southeastern Boreal forest.



Fig. 172. Fringed loosestrife, *Lysimachia ciliata* L.

An erect plant 20–50 cm high with lanceolate to linear-lanceolate stalkless leaves 3–10 cm long. Small flowers, about 6 mm or less in diam, borne in dense spike-like racemes in axils of leaves about halfway up stem. Common in swamps and moist places in Parklands and Boreal forest, but scarce in other areas. Syn.: *Naumburgia thyrsiflora* (L.) Duby.

Primula primrose

- 1. Leaves with sulfur yellow mealiness on undersides. *P. incana*
 Leaves green on both sides. 2
- 2. Leaves entire; flowers less than 1 cm across. *P. egaliksensis*
 Leaves dentate or crenate; flowers larger. 3
- 3. Pedicels 1–3 cm long, many times longer than the bracts. *P. mistassinica*
 Pedicels less than 1 cm long, at most twice as long as the bracts. *P. stricta*

Primula egaliksensis Wormsk.

Slender plants with stems to 20 cm high; leaves thin, ovate to oblong or spatulate, to 5 cm long. Flowers 5–9 mm across, greenish white to violet; calyx lobes glandular ciliate. Meadows and calcareous shores; Boreal forest.

Primula incana M. E. Jones (Fig. 171C)

MEALY PRIMROSE

A low plant with a basal rosette of spatulate to oval leaves 2–10 cm long, tapering at base to a stalk and blunt at apex. Upper sides green and underside covered with a sulfur yellow mealiness. Flowers pale lilac with a yellow center, 6–10 mm across, borne in an umbel-like cluster at head of a leafless stem 10–30 cm high. Found occasionally; in grass in saline meadows and moist spots; throughout the Prairies and Parklands.

Primula mistassinica Michx.

DWARF PRIMROSE

A small plant with a very slender stem 5–20 cm high. Leaves 1–5 cm long, oblanceolate to cuneate-obovate, dentate, green on both sides, or slightly mealy below. Flowers 10–20 mm across, pink, lilac, or white with yellow center. Marshy areas, shores, and bogs; Boreal forest, rare elsewhere.

Primula stricta Horn.

ERECT PRIMROSE

Plants with upright stems 10–30 cm high. Leaves 2–6 cm long, green on both sides, crenate, ovate to lanceolate. Flowers 8–12 mm across. Wet areas; northeastern Boreal forest.

Trientalis starflower

- Leaves almost entirely in a whorl at tip of stem, acute or acuminate; pedicels shorter than the leaves. *T. borealis*

Leaves not restricted to tip of stem, obtuse or rounded; pedicels as long as the leaves. *T. europaea*

Trientalis borealis Raf. (Fig. 171D)

NORTHERN STARFLOWER

A perennial from horizontal creeping roots, which send up simple stems 8–30 cm high, bearing a whorl of 5–10 lanceolate leaves 3–10 cm long and tapering at both ends. The few flowers are 10–12 mm across, white, with 7 pointed petals, and borne on slender stalks from the center of a leafy whorl. Found occasionally in damp woodlands; throughout Boreal forest. Syn.: *T. americana* Pursh.

Trientalis europaea L.

STARFLOWER

Similar to *T. borealis*, but with the leaves more or less spread out along the upper part of the stem; pedicels much longer, and also axillary from lower leaves. Muskeg and bogs; northwestern Boreal forest, Peace River.

OLEACEAE—olive family

Fraxinus ash

Hardwood trees with pinnate leaves. Inconspicuous flowers, usually unisexual, appearing about the same time as the leaves. Fruit, a samara, borne in pendulous clusters and consisting of a seed with a long membranous yellowish green wing.

Body or seed of samara flattened; wing extending to bottom of seed. *F. nigra*

Body or seed of samara almost circular in cross section; wing extending halfway down body of seed. *F. pennsylvanica* var. *austini*

Fraxinus nigra Marsh.

BLACK ASH

A tall tree with pinnate leaves of 7–11 leaflets; the difference between the wing and the body of the samara rather indistinct. Southeastern Parklands and Boreal forest.

Fraxinus pennsylvanica Marsh. var. *austini* Fern. (Fig. 173)

GREEN ASH

A tree to 10 m high with 5–7 leaflets. Branchlets densely short pubescent; the leaf rachis and underside of leaflets sparingly pubescent. Shores, thickets, and along rivers; southeastern Parklands and Boreal forest. Trees with glabrous branches and leaflets are var. *subintegerrima* (Vahl.) Fern., occurring along rivers and in ravines in the eastern part of the Prairies. Syn.: *F. campestris* Britt.

(Fig. 173 overleaf)



HERBARIUM OF THE PEPPER HARBOR STATION
 STATE COLLEGE, PA. N. Y.

SPECIES: *Fraxinus pennsylvanica* Marsh.
 var. *austinii* Fern.

LOCALITY: Turtle Mountain Community Pasture

COLLECTOR: J. Loeman

DATE:

LOCALITY: PA. N. Y.

DATE: 15 July 1970

ELEVATION: 400 ft.

HABITAT: woods




Fig. 173. Green ash, *Fraxinus pennsylvanica* Marsh. var. *austinii* Fern.

GENTIANACEAE—gentian family

Mostly low herbs with a bitter sap and, with one exception, opposite, simple, and stalkless leaves. Flowers regular and perfect, with a tubular corolla, usually with 4 or 5 lobes at mouth, with as many stamens as corolla lobes. Fruit a capsule opening by valves and containing many seeds.

- 1. Perennial bog plants with trifoliolate leaves (3 leaflets). *Menyanthes*
Plants with simple opposite leaves. 2
- 2. Flowers with 4 hollow spurs at base. *Halenia*
Flowers not spurred at base. 3
- 3. Corolla campanulate or bell-shaped. *Gentiana*
Corolla rotate. *Lomatogonium*

Gentiana gentian

- 1. Flowers with plaits or folds between lobes of corolla. 2
Flowers without plaits or folds between lobes of corolla. 8
- 2. Leaves with white margins; flowers solitary and terminal. *G. aquatica*
Leaves without white margins; flowers several, in leaf axils. 3
- 3. Mouth of corolla almost or quite closed; corolla lobes absent or very minute. *G. andrewsii*
Mouth of corolla open; corolla lobes distinct. 4
- 4. Leaves linear-oblong; flowers terminal and axillary. *G. linearis*
Leaves ovate to lanceolate. 5
- 5. Plants with basal rosettes present at flowering. *G. glauca*
Plants without basal rosettes at flowering. 6
- 6. Calyx lobes broadly ovate or oval. *G. calycosa*
Calyx lobes linear to linear-lanceolate. 7
- 7. Flowers 25–30 mm long; leaves glabrous. *G. affinis*
Flowers 35–45 mm long; leaves puberulent. *G. puberulenta*
- 8. Lobes of corolla fringed or toothed; flowers long-peduncled. 9
Lobes of corolla not fringed or toothed; flowers short-peduncled. 10
- 9. Calyx finely but distinctly glandular puberulent on the keels. *G. crinita*
Calyx glabrous. *G. detonsa*

10. Throat of corolla with a fringe of hairs. *G. amarella*
 Throat of corolla hairless. *G. propinqua*

Gentiana affinis Griseb.

OBLONG-LEAVED GENTIAN

A leafy-stemmed perennial 15–30 cm high, often prostrate, usually with several stems from a deep taproot. Leaves oblong to lanceolate, 10–35 mm long. Flowers dark blue or purple, 25–30 mm long, in raceme-like dense clusters on upper end of stems. Sometimes merely one or a few flowers to a stem. Fairly common; in sandy areas and moist, even saline, meadows; throughout Prairies and Parklands. Syn.: *Dasystephana affinis* (Griseb.) Rydb.; *G. interrupta* Greene.

Gentiana amarella L.

NORTHERN GENTIAN

An annual 15–50 cm high. Upper leaves lanceolate and rather sharply pointed; lower leaves usually spatulate or ovate and blunt at apex. Flowers varying from purplish and blue to greenish yellow or white, 10–20 mm, borne in clusters in upper leaf axils. Found fairly often; throughout the Prairie Provinces. Because of wide variations, authorities have made several separate species of various forms of this plant, such as *G. acuta*, *G. plebeia*, *G. scopulorum*, and *G. strictiflora*, but they are probably all local and variations of the same species.

Gentiana andrewsii Griseb.

CLOSED GENTIAN

An upright rather sturdy perennial herb 30–80 cm high. Leaves ovate to lanceolate, 5–10 cm long, with 3–7 veins. Flowers blue with whitish folds, or sometimes all white, 3–4 cm long, usually closed at the mouth and borne in clusters at the end of the stem or in axils of upper leaves. In wet meadows and among bushes; in eastern Parklands. Syn.: *Dasystephana andrewsii* (Griseb.) Small. Plants with white flowers are distinguished as f. *albiflora* Britt. Syn.: *G. flavida* Gray; *Dasystephana flavida* (Gray) Britt.

Gentiana aquatica L.

MOSS GENTIAN

A small annual or biennial, usually branching from base, 3–10 cm high. Leaves many, less than 6 mm long; basal leaves obovate; upper ones linear-lanceolate. Small purplish green flowers solitary at summit of stems, 5–8 mm long, and followed by capsule containing tiny seeds. Capsule opening into two valves and spreading outward at maturity. Found very rarely; around sloughs and marshy places; in Prairies and Parklands. Syn.: *Chondrophylla fremontii* (Torr.) A. Nels.; *G. fremontii* Torr.; *G. prostrata* Haenke.

Gentiana calycosa Griseb.

MOUNTAIN GENTIAN

Plants with simple or branched root crown, and several to many erect or ascending stems 15–40 cm high. Leaves 2–4 cm long, round-oval to ovate. Flowers solitary, or 1–3 at the tip of the stem, sometimes also 1 or 2 flowers in the upper axils. Corolla blue, 25–35 mm long; calyx tube 6–8 mm long, the oblong lobes equaling or exceeding the tube; capsule 12–16 mm long. Rare; alpine meadows; southern Rocky Mountains.

Gentiana crinita Froel.

FRINGED GENTIAN

An annual species 15–50 cm high, with somewhat clasping lanceolate leaves 2–5 cm long. Flowers large, 2–6 cm long, at end of stems, sky blue, with a conspicuous fringe around the lobes. Fairly common; in moist woods or low areas; in southeastern Parklands and Boreal forest. Syn.: *Anthopogon crinitus* (Froel.) Raf. Three varieties can be distinguished:

- 1. Leaves lanceolate, 5–20 mm wide. var. *crinita*
 Leaves linear, 3–5 mm wide. 2
- 2. Flowers large, mostly 4–6 cm long,
 corolla lobes strongly fringed. var. *browniana* (Hook.) Boiv.
 Syn.: *G. procera* Holm
- Flowers small, mostly 2–4 cm long,
 corolla lobes little fringed. var. *tonsa* (Lunell) Vict.
 Syn.: *G. macounii* Holm

Gentiana detonsa Rottb. var. *raupii* (Pors.) Boiv. NORTHERN FRINGED GENTIAN

Similar to *G. crinita*, but with the calyx glabrous. Stems 20–40 cm high, leafy below. Leaves lanceolate to linear-lanceolate, 2–5 mm wide. Corolla 3–5 cm long, with lobes short-fimbriate. Shores and marshes; Boreal forest.

Gentiana glauca Pall.

ALPINE GENTIAN

Low plants with rootstocks and basal rosettes; stems 2–10 cm high. Basal leaves 5–15 mm long, rather thick; stem leaves 2 or 3 pairs, about 1 cm long. Corolla 12–18 mm long, blue. Alpine meadows; southern Rocky Mountains.

Gentiana linearis Froel.

CLOSED MARSH GENTIAN

Rather stout plants with stems 30–70 cm high. Leaves 4–9 cm long, linear-oblong, to 1 cm wide. Flowers usually 2–4 in a terminal cluster, and solitary in the upper axils. Corolla 3–4 cm long, blue to white. Rare; marshy areas; southeastern Parklands and Boreal forest.

Gentiana propinqua Rich.

FELWORT

Annual plants, often branching from the base, 5–30 cm high. Basal leaves spatulate or oblanceolate; stem leaves lanceolate, 1–2 cm long. Flowers 1–3 in axils; corolla blue, 10–15 mm long; calyx lobes unequal: the outer two 5–7 mm long, ovate and the others 3–5 mm long, linear. Arctic and alpine grassland; northeastern Boreal forest, Rocky Mountains.

Gentiana puberulenta Pringle

DOWNY GENTIAN

Perennial, usually with a single stem 20–50 cm high, often covered with minute hairs. Leaves lanceolate, usually rough or minutely hairy on edges and midrib. Flowers blue, 3–4 cm long, in upper leaf axils. Very rare; grassland; southeastern Parklands. Syn.: *Dasystephana puberula* (Michx.) Small; *G. puberula* Michx.

Halenia spurred-gentian

Halenia deflexa (Smith) Griseb.

SPURRED-GENTIAN

An annual with slender upright stems 15–50 cm high. Basal leaves spatulate or obovate; stem leaves oblong to ovate, 2–5 cm long. Flowers in clusters at head of stems and in axils of upper leaves, about 6 mm long, purplish to greenish or yellowish white. Distinguished from gentians by 4 hollow spurs projecting downward from the base of flowers. Fairly common in moist woodlands; Parklands and Boreal forest.

Lomatogonium marsh felwort

Lomatogonium rotatum (L.) Fries

MARSH FELWORT

An erect slender annual 10–45 cm high, with spatulate basal leaves and linear to lanceolate stem leaves 1–5 cm long. White or bluish flowers 1–2 cm across, borne singly or in clusters in axils of leaves; corolla deeply cleft into 4 or 5 segments. Rare; occasionally reported from marshy land; throughout the Prairie Provinces. Syn.: *Pleurogyne rotata* (L.) Griseb.

Menyanthes buck-bean

Menyanthes trifoliata L.

BUCK-BEAN

A perennial bog plant arising from a thick scaly rootstock. Leaves trifoliate, of 3 elliptic leaflets 5–10 cm long, on basally sheathed stems 5–20 cm long. Flowers clustered in a raceme at the head of a separate stalk, whitish or pinkish purple, 10–15 mm long. Fruit an ovoid capsule containing a few shiny seeds. Common; in bogs and wet swampy places; in Boreal forest.

APOCYNACEAE—dogbane family

Perennial herbs with acrid milky sap, and entire opposite leaves. Flowers regular, perfect, and bell-shaped, with 5 sepals and 5 partly united petals. Fruits long, narrow follicles, in pairs, and containing many seeds; each seed bearing a tuft of hairs.

Apocynum dogbane

1. Petals fully twice as long as sepals; stem leaves drooping or spreading. 2
 Petals less than twice as long as sepals;
 stem leaves ascending. *A. cannabinum*
2. Petals three times as long as sepals; flower clusters at end of stems and in axils of leaves. *A. androsaemifolium*
 Petals twice as long as sepals; flower clusters at end of stems only. *A. medium*

Apocynum androsaemifolium L. var. *incanum* DC. (Fig. 174)

SPREADING DOGBANE

A somewhat bushy perennial from a horizontal rootstock, 30–150 cm high. Plant much branched, and stems when broken exuding a milky sap. Leaves opposite, ovate or oval, somewhat paler and often slightly hairy on lower side, 2–7 cm long. Flowers in clusters at ends of branches and in axils of leaves, pink, 6–8 mm long; lobes of petals spreading and often curved downward. Fruits in pairs of long narrow follicles or pods, to 10 cm long, tubular, containing many hairy-tipped seeds. Common; in woodland and on light sandy soil; throughout the Prairie Provinces.

Apocynum cannabinum L. var. *hypericifolium* Gray

INDIAN-HEMP

A deep-rooted perennial 30–150 cm high, with fairly erect branches. Leaves lanceolate-oblong or ovate-oblong, pale green above and often somewhat whitened beneath, 3–10 cm long, narrowed at either end. Flowers greenish white, about 3–5 mm long, in clusters at ends of branches and in leaf axils. Fruit similar to that of *A. androsaemifolium*. Found occasionally; in thickets; throughout the Prairie Provinces. Includes *A. sibiricum* Jacq.

Apocynum medium Greene

INTERMEDIATE DOGBANE

Similar to spreading dogbane, for which it is often mistaken, but having a shorter flower, about 6 mm or less in length, usually greenish or white with a pink tinge. Flower clusters only borne at ends of branches, not in leaf axils. A hybrid of the foregoing two species, sparingly found; throughout the Prairie Provinces where the parents are growing.


ASCLEPIADACEAE—milkweed family

Perennial herbs, usually with milky juice, and flowers borne in umbels. Flowers rather complicated, but having 5 corolla lobes with a 5-lobed crown joining the stamens and the corolla lobes. Corolla lobes usually reflexed or turned downward. Fruit a large and conspicuous follicle, or large pod, which opens down one side to release numerous seeds, each of which bears a tuft of silky hairs. Pollen grains in this family united into masses (pollinia), which are pear-shaped and attached in pairs. The peculiar structure of the flower causes these pairs of pollinia to adhere to visiting insects and to be transferred to other flowers, thereby ensuring cross-fertilization.

Asclepias milkweed

Coarse perennial herbs with deep taproots and white milky juice. Leaves opposite or whorled. Petals and sepals reflexed (turned downward). Fruit a large pod, or follicle, containing many seeds, each with a tuft of white silky hairs.

(Fig. 174 overleaf)



APOCYNACEAE
 Apocynum androsaemifolium L. var. incanum DC.

19918: Apocynum androsaemifolium L.

STATE: Turtle Mountain, Max Lake

COLL. J. Cooper

1970

COLL. NO. 11226

DATE: 15 July, 1970

PREPARED BY: J. Cooper

HOLOTYPE: Max Lake




Fig. 174. Spreading dogbane, *Apocynum androsaemifolium* L. var. *incanum* DC.

1. Hoods of the crown of flower without an incurved horn within. 2
Hoods of the crown of flower with an incurved horn within. 3
2. Umbels solitary, terminal. *A. lanuginosa*
Umbels several, lateral. *A. viridiflora*
3. Leaves linear and in whorls. *A. verticillata*
Leaves broader and opposite. 4
4. Flowers rose or red; plant almost hairless. *A. incarnata*
Flowers purplish or greenish; plants hairy or downy. 5
5. Leaves ovate or lanceolate, tapering to base; pods without tubercles. *A. ovalifolia*
Leaves blunt or almost cordate at the base, oblong or oval; pods with soft tubercles. 6
6. Hoods of corolla long and lanceolate, three times longer than stamens. *A. speciosa*
Hoods of corolla short and blunt, not much longer than stamens. *A. syriaca*

Asclepias incarnata L.

SWAMP MILKWEED

A tall slender-stemmed perennial, 50–150 cm high, almost devoid of hairs, with lanceolate opposite leaves 3–10 cm long. Small flowers usually rosy red but sometimes paler, in numerous many-flowered umbels. Pods of seeds borne upright, usually in pairs, 5–8 cm long, and almost smooth. Fairly common; in swamps, wet spots, and roadside ditches; throughout southeastern Parklands.

Asclepias lanuginosa Nutt.

HAIRY MILKWEED

Tall perennial, with stems 1–2 m high, and villous. Leaves linear-oblong or lanceolate, 4–7 cm long, pubescent on both sides. Umbel erect, short-peduncled, 4–5 cm across. Rare; open grasslands in sandhills; southeastern Parklands.

Asclepias ovalifolia Dcne.

DWARF MILKWEED

A low species 20–50 cm high, with ovate to lanceolate leaves narrowing or tapering to base, 3–7 cm long. Flowers greenish white, on long stalks in umbels. Found occasionally; on moist prairie; throughout Parklands and southern margin of Boreal forest.

Asclepias speciosa Torr. (Fig. 175)

SHOWY MILKWEED

A stout erect perennial 30–100 cm high, usually found in large colonies. Leaves broad and oval, 7–15 cm long, rounded or somewhat heart-shaped at base, and with a whitish downiness. Flowers flesh-colored or pinkish purple, 8–12 mm across, in very dense almost globular umbels 5–7 cm in diam. Inflorescence having a strong sweet smell, which may make a person sleepy. The nectar also appearing to have a stupefying effect on insects, which may often be found in a drowsy condition below the plants. Pods 6–10 cm long,




Fig. 175. Showy milkweed, *Asclepias speciosa* Torr.

densely white woolly, and covered with soft tubercles, borne on recurved stalks. The commonest milkweed; in moist places; throughout Prairies and Parklands.

Asclepias syriaca L.

SILKY MILKWEED

Very similar to *A. speciosa*, but with elliptic leaves, rounded and not heart-shaped at base. Flowers differ in having a short blunt hood on corolla. Fairly common; on moist sandy soil and riverbanks; southeastern Prairies and Parkland.

Asclepias verticillata L.

WHORLED MILKWEED

A slender-stemmed plant 20–50 cm high with very narrow linear leaves 3–7 cm long, borne in whorls of twos or fours up the stem. Small flowers greenish white, in small umbels. Found occasionally; on dry soil; extreme southeastern Parkland.

Asclepias viridiflora Raf.

GREEN MILKWEED

Perennial herb with stems sometimes reclining at base, 20–120 cm high. Leaves usually ovate-lanceolate, 2–8 cm long. Flowers borne in umbels at head of stem and also in leaf axils, greenish yellow to dull purple with purple hood. The var. *linearis* (A. Gray) Fern. has linear leaves. Found occasionally; in dry or sandy soil; eastern and south central parts of Prairies and Parkland. Syn.: *Acerates viridiflora* (Raf.) Eat.

CONVOLVULACEAE—convolvulus family

- Leafy twining plants with large funnel-like flowers. *Convolvulus*
 Parasitic twining plants with leaves reduced to scales. *Cuscuta*

Convolvulus morning-glory

Twining or climbing perennials from creeping roots. Leaves alternate, usually arrow- or spear-shaped. Flowers usually large, funnel-form or bell-shaped, with 5 stamens. Fruit a several-seeded capsule.

- Calyx not enclosed in two large bracts. *C. arvensis*
 Calyx enclosed in two large bracts. *C. sepium*

Convolvulus arvensis L. (Fig. 176A)

FIELD BINDWEED

A deep-rooted perennial with a very extensive system of white roots, and slender twining stems. Leaves 2–4 cm long, somewhat bluntly triangular with hastate bases. Flowers varying from pink to white, 20–25 mm across, borne either singly or 2–4 in a bunch. Fruit a round 2-celled capsule containing large, dark brown, angular seeds. An introduced weed; often found in gardens and roadsides, and very difficult to eradicate.




Fig. 176. A, Field bindweed, *Convolvulus arvensis* L.; B, wild morning-glory, *Convolvulus sepium* L. var. *americanus* Sims.

A perennial plant, naturally twining on bushes and shrubs, but in cultivated places often becoming a creeping field weed. Rhizomes white and creeping, sending up many shoots. Leaves varying in size and shape but roughly triangular with hastate or sagittate bases. Two large green bracts enclosing sepals and lower portion of flower. The large funnel-shaped flowers varying from pink to pure white. Fruit a capsule containing large angular brown or black seeds. The species has lobes of leaves directed downward, leaves 5–12 cm long. Flowers 4–5 cm long and up to 6 cm across. Probably introduced from Europe or Asia; found around bushes and waste places; especially in eastern Prairies and Parkland. The following key separates the two main varieties of this species.

- Leaves with basal lobes spreading sideways;
hairless or only slightly hairy. var. *americanus*
- Leaves with rounded or arrow-shaped, downward-pointing basal lobes; stems and leaves covered with dense fine hairs. var. *pubescens*

The var. *americanus* Sims (Fig. 176B), wild morning-glory, is a native twining perennial with broadly hastate leaves and either pink or white funnel-like flowers. When the area on which they grow is brought under cultivation, the roots are spread by tillage throughout the land, and the plant spreads rapidly by both root portions and seed, often taking full control of large areas and choking out crops. When growing in large masses, leaves and flowers often become much smaller, almost to suggest a different species. The seeds have a long dormant period and may remain viable in soil for many years, even under good germination conditions. Common; found climbing on bushes along water courses and in moist areas; throughout the Prairie Provinces. Syn.: *C. americanus* (Sims) Greene.

The var. *pubescens* (Gray) Fernald, inland bindweed, is a low-growing, twining or crawling perennial with leaves 25–35 mm long, rounded at angles, arrow-shaped or heart-shaped at base, and covered with fine downy hairs beneath. Flowers white, 3–4 cm long. Found occasionally; on sandy soils; throughout southern prairies. Syn.: *C. interior* House.

***Cuscuta* dodder**

Parasitic annual plants with leaves absent or reduced to scales, found twining on host plants. Dodders grow from seed, and after they become fastened to their host plants by aerial roots and suckers, the root and basal portion of the stem decays and sustenance is entirely derived from the host plant. Flowers pinkish or whitish, in clusters, with a bell-shaped or almost globular corolla of 4 or 5 overlapping lobes, and with small scales inside the throat, alternating with lobes. Fruits are somewhat globular capsules borne in clusters, each capsule containing 2–4 seeds. A rather difficult genus to identify, because the small flowers need to be closely examined for differing characteristics.

An orange- or yellow-stemmed species found twining on various coarse herbs and shrubs. Flowers 2–4 mm long, sessile or subsessile in dense clusters. Corolla lobes broadly ovate, shorter than the tube; calyx short, with lobes round-ovate to subrotund; capsules about 3 mm in diam. Rare; on various shrubs along shores and in moist areas; Prairies and Parklands. Includes *C. curta* (Engelm.) Rydb.; *C. pentagona* Engelm.; *C. planiflora* Tenor.

POLEMONIACEAE—phlox family

Usually low-growing annual or perennial herbs. Flowers perfect, generally regular, with 5 partly united sepals and 5 united petals. Flowers usually funnel-like or salverform (with a long tube, abruptly flattened at the end). Fruit a 3-celled capsule containing the seeds.

- 1. Leaves compound with many small leaflets. *Polemonium*
 Leaves simple. 2
- 2. Leaves not cleft or divided. 3
 Leaves divided or pinnately cleft. 4
- 3. Leaves opposite. *Phlox*
 Leaves alternate. *Collomia*
- 4. Leaves opposite. *Linanthus*
 Leaves alternate. 5
- 5. Calyx spine-tipped, and as long as corolla tube. *Navarretia*
 Calyx not as long as corolla tube. *Gilia*

Collomia collomia

Collomia linearis Nutt. (Fig. 177A) NARROW-LEAVED COLLOMIA

An erect annual herb 10–40 cm high, somewhat sticky hairy. Leaves alternate, lanceolate or linear-lanceolate, entire, 2–6 cm long; lower leaves usually shorter than upper ones. Flowers very small, pink or pale purple, in a dense head-like leafy cluster at top of stem. Common; on dry soils and sandy places; throughout the Prairie Provinces.

Gilia gilia

- Flowers usually red, about 2 cm long, in a panicle-like cluster. *G. aggregata*
- Flowers white, less than 1 cm long, in head-like clusters. *G. congesta*




Fig. 177. Phloxes: A, narrow-leaved collomia, *Collomia linearis* Nutt.; B, small navarretia, *Navarretia minima* Nutt.; C, moss phlox, *Phlox hoodii* Richardson.

Gilia aggregata (Pursh) Spreng.

SCARLET GILIA

An erect perennial 30–50 cm high, usually few-branched. Leaves pinnately divided into narrow segments, 2–7 cm long. Red or scarlet flowers very conspicuous, in a large, loose, narrow cluster, usually 2–4 cm, with petal lobes about 10 mm long. Very rare; southern Rocky Mountains.

Gilia congesta Hook.

CLUSTERED GILIA

A basally-branched perennial 10–15 cm high, usually covered with cobwebby hairs. Leaves pinnately cleft. White flowers in head-like clusters. Very rare, but may possibly be found; on dry, sandy soil; southern Rocky Mountains.

Linanthus linanthus

Linanthus septentrionalis H. L. Mason

LINANTHUS

A much-branched, very fine stemmed annual 5–30 cm high. Leaves divided to the base into very narrow thread-like segments, about 5–10 mm long, making them appear clustered. Flowers white, about 3 mm long, on long fine stalks. Very rare, but has been found; on sandy roadsides; Prairies. Syn.: *L. harknessii* (Curran) Greene var. *septentrionalis* (Mason) Jepson & Bailey.

Navarretia navarretia

Navarretia minima Nutt. (Fig. 177B)

SMALL NAVARRETIA

A low-growing depressed annual with much-branched stems 3–10 cm high. Leaves 20–35 mm long, deeply divided into needle-like segments. Flowers about 6 mm long, white, and almost hidden in round clusters of leaves and spiny sepals. Locally abundant; on bottom lands, sandy places, and slough margins; throughout southwestern Prairies.

Phlox phlox

- 1. Stem erect; leaves linear to lanceolate. 2
 Stem decumbent or tufted; leaves short
 and awl-shaped, usually not more than
 10–12 mm long. 3
- 2. Plants annual; stems 10–20 cm high;
 upper stem leaves alternate. *P. gracilis*
 Plants perennial; stems to 60 cm high;
 upper stem leaves opposite. *P. pilosa*
- 3. Leaves with cobwebby hairs; tube of
 corolla less than 12 mm long, slightly
 longer than calyx. *P. hoodii*
 Leaves without cobwebby hairs; tube of
 corolla more than 12 mm long, much
 longer than calyx. *P. alyssifolia*

Phlox alyssifolia Greene

BLUE PHLOX

A stout-stemmed prostrate plant with oblong or linear leaves 6–12 mm long, with sharp-pointed tips. Flowers few, purplish or bluish, with a tube

about 15 mm long. Rare; found occasionally on dry benchland; Prairies and southern Rocky Mountains.

Phlox gracilis (Hook.) Greene SLENDER PHLOX

Stems usually erect, simple below and sparsely to freely branching above; glandular pubescent above, sparsely tomentose below. Inflorescence branched, cymose, glandular; corolla 9–12 mm long, with lobes pinkish or purplish; calyx 5–6 mm long. Rare; moist, grassy slopes; southern Rocky Mountains.

Phlox hoodii Richardson (Fig. 177C) MOSS PHLOX

A low tufted mat-forming plant with coarse woody roots. Leaves awl-shaped, sharp-pointed, and somewhat imbricated or overlapping, 3–8 mm long. Flowers white or occasionally pale blue or purple, about 1 cm across, with 5 petal lobes, borne very freely in early spring. One of the most conspicuous of early spring prairie flowers, forming large masses of white on the plains and hillsides. After flowering, plants becoming rather inconspicuous but forming a large proportion of the ground cover, especially on eroded areas and in shallow soil. Common on open prairie throughout Prairies, less common in Parklands.

Phlox pilosa L. DOWNY PHLOX

An erect species 30–60 cm high, usually with soft downy hairs. Leaves linear or lanceolate, 2–10 cm long and stalkless. Flowers purplish pink or white, with a tube about 12 mm long, in a cymose cluster at summit of stem. Found occasionally; in sandy places; southeastern Parklands.

Polemonium Jacob's-ladder

- 1. Corolla lobes shorter than the tube;
flowers in dense, capitate cymes. *P. viscosum*
- Corolla lobes longer than the tube;
flowers in loose, open cymes. 2
- 2. Plants usually less than 30 cm high, sev-
eral to many stems. *P. pulcherrimum*
- Plants usually more than 30 cm high,
solitary. *P. occidentale*

Polemonium occidentale Greene WESTERN PHLOX

A leafy-stemmed erect perennial 30–80 cm high, somewhat glandular hairy above. Leaves of 15–27 ovate to lanceolate leaflets 15–35 mm long. Flowers bell-shaped, violet or blue, 8–12 mm long, in a narrow cluster. May be found occasionally, but not common; in open woodlands and valleys; Rocky Mountains.

Polemonium pulcherrimum Hook. (Fig. 178) SHOWY JACOB'S-LADDER

A glabrous many-stemmed perennial; leaves with 5–11 pairs of ovate to rotund leaflets 2–8 mm long. Inflorescence a cyme; flowers blue, 10–15 mm across; calyx glandular pubescent. A very showy species, found on slopes and in open areas; Rocky Mountains.



Fig. 178. Showy Jacob's-ladder, *Polemonium pulcherrimum* Hook.

Perennials with rootstock; stems 10–40 cm high. Leaves mostly basal; leaflets 3- to 5-lobed, with lobes 1–3 mm long, glandular pubescent. Inflorescence subcapitate; bracts pinnatifid, glandular pubescent; flowers blue, 20–25 mm across, with the calyx glandular pubescent. Rare; rocky slopes; southern Rocky Mountains.

HYDROPHYLLACEAE—waterleaf family

Mostly hairy annual herbs with a watery sap. Leaves lobed or pinnatifid and flowers perfect with 5 more or less united sepals and a 5-lobed bell-shaped corolla. Fruit a capsule.

- 1. Flowers solitary in leaf axils. 2
- Flowers several to many in cymes. 3
- 2. Leaves pinnately divided with 7–13 dentate segments. *Ellisia*
- Leaves with 3–5 obovate, entire-margined segments. *Nemophila*
- 3. Leaves palmately lobed; lobes deltoid. *Romanzoffia*
- Leaves pinnately lobed or parted; lobes elongate. 4
- 4. Flowers in dense capitate cymes. *Hydrophyllum*
- Flowers in simple or branched scorpioid cymes. *Phacelia*

Ellisia *ellisia*

Ellisia nyctelea L.

WATERPOD

A low-growing plant 10–30 cm high, with scattered hairs. Leaves opposite or alternate, pinnatifid with toothed segments, 2–10 cm long. Flowers bluish white, 6–12 mm across, on stalks opposite leaf axils. Calyx lobes (sepals) enlarging as fruiting capsule forms. Fruit a globular capsule about 6 mm in diam. Not common; found on river flats, shady spots, lake margins; throughout southern part of the Prairie Provinces.

Hydrophyllum waterleaf

Hydrophyllum capitatum Dougl.

WOOLLEN-BREECHES

Short-stemmed plants with a short rhizome, bearing finger-like tuberous roots. Leaves 5–12 cm long, 3–12 cm wide, with 5–7 primary lobes. Cymes one to several, globose; peduncles 1–5 cm long; flowers purplish blue to white, 6–10 mm across; calyx densely pubescent; capsule 4 mm long; seeds usually 2, 2–3 mm across. Rare; exposed areas; southern Rocky Mountains.

Nemophila baby-blue-eyes

Nemophila breviflora Gray

SMALL BABY-BLUE-EYES

Weak-stemmed plants 5–20 cm high, with stems sharply angled, bearing minute reflexed prickles. Leaves divided into 3–7 oblong-lanceolate lobes. Flowers short-pedicel; corolla 1.5–3 mm across, purplish blue to white; capsule 3–5 mm in diam, usually containing a single brick red regularly and deeply pitted seed. Rare; open disturbed soils; southern Rocky Mountains.

Phacelia scorpionweed

- 1. Stamens included in the corolla. *P. thermalis*
 Stamens exserted. 2
- 2. Leaves entire or shallowly dentate, suborbicular to lanceolate. 3
 Leaves deeply lobed to divided. 5
- 3. Leaves suborbicular, shallowly dentate or crenate. *P. campanularia*
 Leaves linear to lanceolate or elliptic. 4
- 4. Plants annual; leaves linear, the lower ones often with 3–5 linear divisions. *P. linearis*
 Plants perennial; leaves oblanceolate to elliptic. *P. hastata*
- 5. Leaves pinnately divided, the divisions entire to pinnatisect. 6
 Leaves more or less deeply lobed, but not divided. 7
- 6. Leaves divided into 7–15 acute lobes; the lobes shallowly incised to entire. *P. franklinii*
 Leaves divided into 5–9 oblong lobes; the lobes deeply incised to pinnatisect. *P. tanacetifolia*
- 7. Leaves divided halfway to the midrib; stamens twice as long as the corolla. *P. lyallii*
 Leaves divided almost to the midrib; stamens three times as long as the corolla. *P. sericea*

Phacelia campanularia Gray

DESERT BLUEBELLS

Annual plants 10–50 cm high, glandular hispid throughout. Basal leaves oblong-ovate to suborbicular, 15–75 mm long, 10–50 mm wide, truncate or cordate at base. Inflorescence a lax, open cyme, simple or few-branched; flowers 15–40 mm long, blue or rarely white; capsule 8–12 mm long. Introduced ornamental; occasionally reseeding itself, and locally established.

Phacelia franklinii (R. Br.) Gray

FRANKLIN'S SCORPIONWEED

A hairy erect annual 15–40 cm high. Leaves 3–7 cm long, pinnately divided into linear-oblong toothed segments. Flowers in a dense coiled or scorpioid raceme, blue or bluish white, about 8 mm long. Fairly common; on dry sandy soil; in the Boreal forest and in Riding Mountains.

Phacelia hastata Dougl.

SILVER-LEAVED SCORPIONWEED

Perennial plants with a branched caudex, ascending stems 15–30 cm long, canescent and appressed hirsute throughout. Leaves linear- or ovate-lanceolate, 3–10 cm long, mostly entire. Cymes 2–10 cm long, numerous, many-flowered; corolla white to light blue; capsule about 3 mm long. Sandy or rocky soil and eroded slopes; southern Rocky Mountains.

Phacelia linearis (Pursh) Holz.

LINEAR-LEAVED SCORPIONWEED

An annual 10–40 cm high with leaves 2–5 cm long, entire or divided into linear segments. Flowers bright blue, about 1 cm long, in a scorpioid panicle. Found occasionally; in valleys, on hillsides, and on light soil; in southern Rocky Mountains.

Phacelia lyallii (Gray) Rydb.

LYALL'S SCORPIONWEED

Perennial with sparingly hirsute stems 10–20 cm high; leaves oblanceolate in outline, 5–10 cm long, green, sparingly pubescent. Inflorescence short, dense; corolla dark blue, 7–10 mm long. Rare; alpine slopes; southern Rocky Mountains.

Phacelia sericea (Graham) Gray (Fig. 179)

SILKY SCORPIONWEED

Perennial with a woody caudex, densely silvery and silky pubescent, 10–40 cm high. Inflorescence an elongate panicle of short cymes, with numerous flowers; corolla dark bluish purple, rarely white; capsule 4–6 mm long. Mountain slopes and disturbed areas; Rocky Mountains.

Phacelia tanacetifolia Benth.

TANSY SCORPIONWEED

An annual plant 20–80 cm high, sparsely pubescent throughout. Basal leaves oblong-oval to ovate, 6–20 cm long, pinnately divided. Inflorescence densely short-pubescent and hispid, of several corymbosely branched cymes; flowers numerous; corolla 7–10 mm long, bluish purple; capsule 3–4 mm long. Introduced; occasional weed of gardens and roadsides.

Phacelia thermalis Greene

GLANDULAR SCORPIONWEED

An annual with stem branched from base, 5–30 cm high. Whole plant covered with whitish glandular hairs. Leaves pinnatifid with oblong lobes, 1–3 cm long. Flowers light blue with a paler tube; corolla about as long as calyx, approx 6 mm. A plant of the dry intermountain plains of Washington and Idaho that has been found in the southern Prairies.

Romanzoffia romanzoffia

Romanzoffia sitchensis Bong.

SITKA ROMANZOFFIA

A slender, few-branched perennial 5–25 cm high, slightly villous; leaf sheaths widened, ciliate. Leaves round-reniform, 10–25 mm across; petioles 1–6 cm long. Inflorescence of a few cymes; corolla 6–9 mm long, white, on pedicels 1–3 cm long; capsule 4–7 mm long. Rare; wet rocks and cliffs; southern Rocky Mountains.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, ALTA.

SPECIES: *Phacelia sericea* (Graham) Gray

LOCATION: Mile 12, Karasakle road

COLL: J. Looman

DATE:

COLL. NO: 17120

DATE: 26 July 1972

ELEVATION: 1800 m

HABITAT: Open woods on
river alluvium



Fig. 179. Silky scorpionweed, *Phacelia sericea* (Graham) Gray.

BORAGINACEAE—borage family

Rough-hairy entire-leaved herbs with perfect, usually regular flowers, with 5-lobed calyx and corolla. Stamens attached at their base to inside of corolla tube; style simple. Fruit usually of 4 nutlets.

1. Ovary of flower merely grooved, and style
at summit of the ovary. *Heliotropium*
Ovary deeply divided into 4 lobes, and
style arising from the center. 2
2. Nutlets with hooked prickles. 3
Nutlets with no prickles or, if prickly, the
prickles not hooked. 4
3. Nutlets prickly all over and spreading
horizontally. *Cynoglossum*
Nutlets usually prickly on edges and
borne vertically. *Lappula*
4. Flowers axillary or in leafy cymes. 5
Flowers in cymes, with small bracts or
bractless. 8
5. Upper leaves opposite or in threes. *Asperugo*
Upper leaves alternate. 6
6. Flowers white, small, 3–5 mm across. *Plagiobothrys*
Flowers light or dark yellow, larger, 8–15
mm across. 7
7. Leaves ovate-lanceolate; styles conspicu-
ously long exserted. *Onosmodium*
Leaves oblong-lanceolate; styles
included. *Lithospermum*
8. Inflorescence of cymes bracteolate to the
tip. 9
Inflorescence of bractless cymes, or only
the lower flowers bracted. 13
9. Flowers drooping; the pedicels recurved,
longer than the calyx. *Borago*
Flowers not drooping; the pedicels erect
or ascending. 10
10. Flowers white, usually less than 1 cm
across. *Cryptantha*
Flowers bluish or purplish, mostly more
than 1 cm across. 11
11. Stamens and style long exserted;
inflorescence racemose. *Echium*
Stamens and style included; inflorescence
not racemose. 12

12. Corolla tube curved in the middle, barely exceeding the calyx. *Lycopsis*
 Corolla tube straight, distinctly longer than the calyx. *Nonea*
13. Flowers yellow, small, about 7 mm long. *Amsinckia*
 Flowers white, blue, or reddish purple. 14
14. Plants to 1 m high; flowers 12–18 mm long, purplish. *Symphytum*
 Plants smaller; flowers smaller, white or blue. 15
15. Flowers blue. 16
 Flowers white. *Cryptantha*
16. Corolla tubular; inflorescence more or less congested. *Mertensia*
 Corolla not tubular; inflorescence elongate; flowers distant. *Myosotis*

Amsinckia fiddle-neck

Amsinckia menziesii Nels. & Macbr.

FIDDLE-NECK

Plants 20–50 cm high, simple or branched. Leaves ovate-lanceolate to linear-lanceolate, 2–7 cm long. Inflorescence elongate, 5–10 cm long, often strongly curled at the tip; flowers 7–12 mm long, pale yellow; nutlets wrinkled or papillate. Rare; disturbed areas, railways; Prairies and Parklands.

Asperugo madwort

Asperugo procumbens L.

MADWORT

A weak-stemmed annual, with stems 20–70 cm long, spreading or procumbent. Leaves oblanceolate, 3–6 cm long. Flowers solitary in the leaf axils, or sometimes 2–4 together; corolla blue, about 3 mm wide. Introduced; found occasionally as a weed.

Borago borage

Borago officinalis L.

OXTONGUE

An erect annual herb 20–60 cm high. Leaves elliptic to oblong or ovate, 3–10 cm long. Inflorescence in terminal many-flowered cymes; corolla bright blue to white, 15–25 mm across. Introduced; occasionally planted as ornamental, and escaped.

Cryptantha cryptanthe

1. Plants perennial; cymes in a raceme. *C. nubigena*
 Plants annual; cymes not in a raceme. 2
2. Cymes bracteolate to the tip. *C. minima*
 Cymes with only the lower flowers bracteolate. *C. fendleri*

Cryptantha fendleri (A. Gray) Greene

FENDLER'S CRYPTANTHE

A very hairy gray or whitish annual with linear leaves 2–6 cm long. Plant sometimes growing to 50 cm high, but usually staying crowded and short. Flowers small and white, borne at ends of branches in scorpioid clusters. Fruits 4 smooth shiny brown nutlets. Common on sand dunes; in western Prairies, sometimes forming ground cover of large areas.

Cryptantha minima Rydb.

SMALL CRYPTANTHE

A small several-stemmed annual, blooming when only 1 or 2 cm high. Leaves 5–15 mm long, spatulate; corolla 2.5–3 mm long. Rare or commonly overlooked; eroded soils; Prairies.

Cryptantha nubigena (Greene) Payson

CLUSTERED OREOCARYA

A low erect species 10–30 cm high, with lower leaves spatulate and upper ones linear. Basal leaves 2–5 cm long; upper ones often shorter. Whole plant covered with white bristly short hairs, often giving a grayish appearance. Flowers white, about 6 mm across, in compact spike-like clusters at ends of branches. Common on dry hillsides and prairies; throughout Prairies and southern edge of Parklands. Syn.: *C. bradburiana* Payson; *Oreocarya glomerata* (Pursh) Greene.

Cynoglossum hound's-tongue

Inflorescence leafy; flowers reddish to white. *C. officinale*

Inflorescence leafless; flowers blue. *C. boreale*

Cynoglossum boreale Fern.

WILD COMFREY

An erect unbranched perennial with stems 40–60 cm high. Basal leaves elliptic to lanceolate, 10–20 cm long, tapering to the petiole; stem leaves sessile, narrowed at the base or somewhat clasping. Racemes 1–3, terminating an elongate leafless peduncle; corolla blue, 6–8 mm across. Rare; in moist woods; southeastern Boreal forest.

Cynoglossum officinale L.

HOUND'S-TONGUE

A soft hairy biennial plant with erect leafy stems 40–80 cm high. Lower leaves oblong-lanceolate, 15–25 cm long, with slender stalks; upper leaves stalkless or clasping, and lanceolate. Flowers in scorpioid racemes at ends of branches, reddish purple, about 5–10 mm across. Fruit a pyramid of 4 nutlets 10–15 mm across. An introduced weed; found occasionally in pastures and waste places.

Echium viper's bugloss

Echium plantagineum L.

PURPLE BUGLOSS

Cymes elongate from the upper 1 or 2 leaf axils; flowers 20–25 mm, purple, with a large calyx. Reported as a very rare weed.

Echium vulgare L.

VIPER'S BUGLOSS

A coarse erect biennial, with stems 30–70 cm high and hairs somewhat prickly. Basal leaves to 15 cm long; stem leaves progressively smaller. Cymes

numerous, in the axils of upper stem leaves; corolla 12–20 mm long, blue. Introduced; weedy in several areas, but rare.

Heliotropium heliotrope

Heliotropium curassavicum L. var. *obovatum* DC.

SPATULATE-LEAVED HELIOTROPE

A perennial from thick fleshy white running roots, 5–30 cm high. Plants smooth with a slight bloom, giving them a somewhat waxy appearance. Leaves fleshy and spatulate, 2–5 cm long, with nerves very indistinct. Flowers white or faintly bluish-tinged, about 6 mm across, in several scorpioid spikes, 1–10 cm long, at ends of branches. Not common but very plentiful where found; in saline slough margins; Prairies and Parklands. Syn.: *H. spathulatum* Rydb.

Lappula bluebur

1. Plants grayish pubescent; leaves 2–6 cm long. *L. echinata*
Plants green, pubescent; leaves 5–15 cm long. 2
2. Flowers 8–12 mm across. *L. deflexa* var. *americana*
Flowers less than 8 mm across. 3
3. Branches of inflorescence strictly ascending, densely flowered. *L. floribunda*
Branches of inflorescence widely spreading, loosely flowered. *L. jessicae*

Lappula deflexa (Wahl.) Garcke var. *americana* (Gray) Greene

NODDING STICKSEED

A biennial, with downward-pointing hairs, 30–90 cm high. Leaves oblong-lanceolate; lower ones stalked and upper ones sessile (stalkless), 5–15 cm long. Inflorescence usually not leafy; pale blue flowers on reflexed stalks in a slender raceme. Found occasionally; in moist, shady woodlands; throughout Prairies and Parkland. Syn.: *L. americana* (Gray) Rydb.; *Hackelia americana* (Gray) Fern.

Lappula echinata Gilib.

BLUEBUR

A hairy annual or winter annual weed 15–60 cm high, much branching. Leaves 2–7 cm long, with only the lower ones stalked. Flowers pale blue, about 3 mm across, on erect leafy-bracted racemes at ends of branches. Fruit containing 4 nutlets with 2 rows of hooked prickles around margins. Whole plant has a strong smell resembling a mouse-infested building. Introduced from Europe, but now very common and widespread; on waste places in cultivated fields and in overgrazed pastures; throughout Prairies and Parkland.

The var. *occidentalis* (Wats.) Boiv. with nutlets having a single marginal row of hooked prickles. A fairly common native plant; on sandhills, light dry soils, and railway grades; throughout Prairies. Syn.: *L. occidentalis* (S. Wats.) Greene.

Lappula floribunda (Lehm.) Greene

LARGE-FLOWERED STICKSEED

A rough hairy biennial or perennial 40–100 cm high. Oblong-lanceolate or linear-lanceolate leaves 5–10 cm long; lower ones stalked and upper ones stalkless. Pale blue flowers in numerous erect racemes. Fairly common; in moist woodlands; throughout Western Parklands and Rocky Mountains. Syn.: *Hackelia floribunda* (Lehm.) Johnston.

Lappula jessicae McGregor

JESSICA'S STICKSEED

Stems erect or ascending, 30–60 cm high; basal leaves 8–15 cm long, 15–20 mm wide. Inflorescence an open panicle, with several racemes; flowers pale blue; corolla 3–5 mm across. Margins of woods and openings; southern Rocky Mountains.

Lithospermum puccoon

Herbs with narrow hairy alternate leaves. Flowers perfect, regular, usually bright yellow, funnellform or salverform with rounded spreading lobes. Fruit usually 4 very hard nutlets.

- 1. Annual plants; nutlets brown, wrinkled, and pitted. *L. arvense*
- Perennial plants; nutlets white, smooth. 2
- 2. Corolla tube not longer than calyx; flowers white or pale yellow. *L. officinale*
- Corolla tube longer than calyx. 3
- 3. Lobes of corolla either fringed or toothed. *L. incisum*
- Lobes of corolla entire, neither fringed nor toothed. 4
- 4. Leaves lanceolate; plant very leafy; flowers crowded, dull greenish yellow. *L. ruderale*
- Leaves linear to oblong; plant not very leafy; flowers bright yellow or orange. *L. canescens*

Lithospermum arvense L.

CORN GROMWELL

An erect usually branched plant, with stems 10–30 cm high. Leaves narrowly lanceolate to almost linear, 20–25 mm long. Flowers white; corolla 6–8 mm across. Introduced; found occasionally as a weed.

Lithospermum canescens (Michx.) Lehm.

HOARY PUCCOON

A softly hairy, somewhat hoary perennial growing erect to 15–50 cm high. Leaves linear-oblong, 1–4 cm long, without stalks. Flowers orange yellow, with tube 10–12 mm long, without stalks, in a rather compact cluster at summit of plant. Smooth white nutlets about 3 mm high. Fairly common; in bluffy country grasslands; eastern Parklands.

Lithospermum incisum Lehm. (Fig. 180)

NARROW-LEAVED PUCCOON

A deep taprooted perennial, often decumbent. Leaves linear, 1–5 cm long, covered with short stiff hairs. Early flowers stalked, bright yellow, with a tube about 2 cm long, and fringed spreading lobes. Later flowers smaller, self-fertilized while in the bud, and very fertile. Nutlets about 3 mm high, white, shiny,



Fig. 180. Narrow-leaved puccoon, *Lithospermum incisum* Lehm.

with minute pits, very hard. Common; on dry prairie land; throughout Prairies and Parklands. Syn.: *L. angustifolium* Michx.

Lithospermum officinale L.

GROMWELL

A finely haired perennial 15–60 cm high. Leaves ovate to ovate-lanceolate, 1–7 cm long. Flowers pale yellow or yellowish white, few, solitary in leaf axils, tube almost as long as calyx. Shiny nutlets about half as long as the sepals. An eastern weed, introduced from Europe, but has been found in southeastern Parklands and Boreal forest.

Lithospermum ruderale Lehm.

WOOLLY GROMWELL

A coarse hairy perennial from thick roots, with several erect stems; densely leafy. Leaves lanceolate with a prominent midrib, 5–10 cm long, stalkless, and crowded on stem. Numerous dull greenish yellow flowers crowded at top of stem in a leafy cluster. White, shining, ovoid nutlets about 3 mm high. Fairly common; in grasslands; Rocky Mountains and Cypress Hills.

Lycopsis bugloss

Lycopsis arvensis L.

SMALL BUGLOSS

An annual rough hairy plant 30–50 cm high. Lanceolate alternate leaves 2–5 cm long. Flowers crowded into somewhat scorpioid terminal racemes, bluish, up to 6 mm across. Tube of corolla slightly curved or bent, and throat closed with stiff fine hairs. The 4 nutlets shorter than calyx. An introduced weed; very uncommon; has been found in southeastern Parklands and in Prairies.

Mertensia lungwort

Perennial plants with alternate leaves and fairly large blue or purple funnelform flowers with lobes slightly spreading. Fruit erect wrinkled nutlets.

1. Plants entirely glabrous. *M. maritima*
Plants pubescent. 2
2. Plants 30–70 cm high; stem leaves 5–15
cm long, ovate to ovate-lanceolate. *M. paniculata*
Plants 10–30 cm high; stem leaves 2–10
cm long, oblong to lanceolate. 3
3. Plants with tuberous roots; flowers 15–25
mm long. *M. longiflora*
Plants with taproot; flowers 10–15 mm
long. *M. lanceolata*

Mertensia lanceolata (Pursh) DC.

LANCE-LEAVED LUNGWORT

A plant 15–30 cm high with linear to lanceolate leaves 5–10 cm long. Leaves usually hairless, but may occasionally be somewhat short hairy on upper side. Blue flowers 10–15 mm long, in few-flowered panicles at ends of branches. Form with no hairiness on upper surface sometimes called *M. linearis* Greene. Both forms fairly common; on open prairie and hillsides; Prairies.

Plants usually with a solitary stem, or rarely 2 or 3 stems, 10–30 cm high, from a shallow tuberous root. Basal leaves 2–5 cm long, mostly confined to sterile stems; stem leaves 2–7 cm long. Inflorescence often dense; flowers deep blue, usually about 2 cm long. Grasslands and slopes; southern Rocky Mountains.

Mertensia maritima (L.) S. F. Gray

Plants with spreading or decumbent stems to 60 cm long. Leaves 2–6 cm long, ovate to spatulate-ovate, fleshy. Cymes often numerous, lax, leafy-bracted; corolla 8–10 mm long, pinkish blue to white. Beaches; northeastern Boreal forest.

Mertensia paniculata (Ait.) G. Don (Fig. 181)

An erect species 30–70 cm high. Leaves lanceolate, 5–15 cm long, somewhat hairy on both sides. Flowers in few-flowered clusters at ends of stems, purplish blue, 10–15 mm long. Found in the woodlands and shady stream banks; throughout Parklands, Boreal forest, and Rocky Mountains.

Myosotis forget-me-not

- 1. Pubescence of the calyx of appressed straight hairs. 2
- Pubescence of the calyx of spreading hooked hairs. 3
- 2. Calyx lobes half as long as the tube; corolla lobes much larger than calyx lobes. *M. scorpioides*
- Calyx lobes equaling the tube; corolla lobes equaling calyx lobes. *M. laxa*
- 3. Flowers 5–8 mm across; plants perennial. *M. sylvatica* var. *alpestris*
- Flowers 2–3 mm across; plants annual or biennial. *M. arvensis*

Myosotis arvensis (L.) Hill

Plants with erect or ascending branching stems 15–40 cm high. Basal leaves 1–2 cm long; stem leaves to 3 cm long. Racemes loosely flowered; flowers blue or white, 2–3 mm across. Introduced; rare weed; in disturbed areas.

Myosotis laxa Lehm.

Perennial plants with decumbent or spreading stems, rooting at the nodes, 15–30 cm long. Leaves to 3 cm long, oblong or oblong-lanceolate. Racemes loosely flowered; corolla about 4 mm across. Introduced; a rare weed; in marshy areas.

Myosotis scorpioides L.

Perennial plants with slender rootstocks; stems decumbent or ascending, 15–40 cm long, rooting at the lower nodes. Leaves 2–8 cm long, oblong-lanceolate to oblanceolate. Racemes loosely flowered; corollas 6–8 mm across. Introduced; a rare weed; in cultivated and disturbed moist areas.



Fig. 181. Tall lungwort, *Mertensia paniculata* (Ait.) G. Don.

Myosotis sylvatica Hoffm. var. *alpestris* (F. W. Schmidt) Koch. FORGET-ME-NOT

Perennial plants with a short rootstock; stems usually upright, 10–20 cm high. Leaves 2–5 cm long, oblong-lanceolate to spatulate. Racemes mostly densely flowered; corolla 5–8 mm across. Alpine meadows and slopes; Rocky Mountains.

Nonea monk's-wort

Nonea vesicaria (L.) Reich. RED MONK'S-WORT

Perennial plants with stout roots; stems 10–40 cm high. Leaves 3–10 cm long, oblong-lanceolate. Racemes more or less densely flowered; corollas 10–15 mm across, dark red to brown red. Introduced; a very rare weed.

Onosmodium false gromwell

Onosmodium molle Michx. var. *hispidissimum* (Mack.) Cronq.
WESTERN FALSE GROMWELL

An erect coarse perennial 50–100 cm high, with rough hairy stems. Leaves lanceolate to ovate-lanceolate, 5–8 cm long, coarsely appressed hairy, with venation very prominent on undersurfaces. Flowers yellowish white or greenish, 10–20 mm long, in leafy terminal scorpioid spikes. Nutlets 3–4 mm long, distinctly constricted at the base. Margins of woods, openings, and grassland; southeastern Parklands and Boreal forest. Smaller plants, 30–60 cm high, with nutlets 2.5–3.5 mm long, not constricted at the base, are var. *occidentale* (Mack.) Cronq. Margins of woods and shrubbery; southeastern Parklands, southern Rocky Mountains.

Plagiobothrys allocarya

Plagiobothrys scouleri (H. & A.) Johnston var. *penicillatus* (Greene) Cronq.
SCOULER'S ALLOCARYA

A low, spreading much-branched annual herb, with linear leaves 1–5 cm long; the lower ones sometimes opposite but the upper ones alternate. Flowers white, very small, in a small scorpioid inflorescence, and also in axils of leaves. Fruits consisting of 4 nutlets, rough on back. Fairly common locally; in slough margins, sandy places, and moist bottomlands; throughout Prairies. Syn.: *Allocarya californica* (F. & M.) Greene.

Symphytum comfrey

Leaves all petiolate, not decurrent. *S. asperum*
Leaves decurrent, the upper ones sessile. *S. officinale*

Symphytum asperum Lepechin ROUGH COMFREY

A perennial herb with erect branched stems 50–100 cm high. Leaves 10–20 cm long, lanceolate to ovate, petioled; the petiole not decurrent on the stem. Flowers blue, 12–18 mm long, numerous in scorpioid cymes. Introduced; a rare weed in disturbed areas.

A large coarse hairy perennial 60–100 cm high. Leaves lanceolate to ovate-lanceolate, 8–25 cm long; lower ones with stalks decurrent along the stem; upper ones sessile or stalkless. Flowers purplish or yellowish, 10–15 mm long, in dense terminal scorpioid clusters. Fruit consists of shiny brown somewhat wrinkled nutlets. An introduced European plant; appears to be established in a few places.

VERBENACEAE—vervain family

Verbena vervain

Annual or perennial hairy herbs with 4-sided stems and opposite leaves. Flowers purplish or blue, tubular with 5 somewhat irregular lobes. There are 4 stamens, one pair longer than the other. Fruit of 4 linear nutlets.

- 1. Plants with decumbent stems; spikes conspicuously bracteose. *V. bracteata*
- Plants with erect stems; spikes not conspicuously bracteose. 2
- 2. Spikes densely flowered, thick; flowers and fruit overlapping. *V. hastata*
- Spikes loosely flowered, thin; flowers and fruit distant. *V. urticifolia*

Verbena bracteata Lag. & Rodr.

BRACTED VERVAIN

A prostrate or decumbent annual or short-lived perennial, much branched, with a 4-sided stem. Plants 30–80 cm across and forming mats on ground. Leaves 2–5 cm long, roughly spatulate, pinnately incised. Purplish blue flowers about 2 mm across, on dense spikes 5–10 cm long; conspicuous bracts about 6 mm long. Not common, but locally abundant; on prairie and waste places on lighter soils; throughout Prairies. Syn.: *V. bracteosa* Michx.

Verbena hastata L.

BLUE VERVAIN

A tall erect perennial with a 4-sided stem 60–150 cm high. Leaves opposite, lanceolate, sharply toothed, 5–10 cm long; lower leaves sometimes hastately lobed at base. Bluish white flowers on numerous terminal spikes 5–10 cm long; fruit densely overlapping on spikes. Fairly common; in woodlands and river valleys; eastern Parklands and Boreal forest.

Verbena urticifolia L.

NETTLE-LEAVED VERVAIN

Similar to *V. hastata*, but the leaves broadly lanceolate to oblong-ovate, rounded at base, and decurrent into the petiole; the inflorescence more open and more sparsely flowered; the corolla white. Thickets, moist fields, and marshes; southeastern Boreal forest.

LABIATAE—mint family

Usually scented plants with square stems. Leaves simple, opposite, or whorled, usually with small glandular pits. Flowers perfect, usually irregular in shape, with 2 lips to corolla. Stamens usually 4, one pair longer than the other, and sometimes only one pair bearing anthers. Fruit of 4 nutlets.

1. Inflorescences terminal. 2
 Inflorescences axillary. 13
2. Inflorescence a more or less globose head. *Monarda*
 Inflorescence a more or less elongated
 raceme. 3
3. Calyx distinctly 2-lipped; the upper lip 3-
 toothed, the lower one 2-lobed. 4
 Calyx indistinctly or not at all 2-lipped;
 one lobe may be larger than the others. 5
4. Plants with decumbent or creeping stems;
 spikes dense, erect. *Prunella*
 Plants with erect stems; spikes loose and
 lax. *Salvia*
5. Inflorescence a raceme of oppositely
 placed flowers. *Physostegia*
 Inflorescence a raceme of oppositely
 placed clusters of flowers. 6
6. Bracts much shorter than and differing
 from the stem leaves. 7
 Bracts grading into the stem leaves. 9
7. Plants annual; spikes strongly one-sided. *Elsholtzia*
 Plants perennial; spikes symmetrical. 8
8. Calyx longer on upper than lower side,
 somewhat oblique at throat. *Agastache*
 Calyx regular, clearly 5-lobed. *Mentha*
9. Leaves entire, linear to linear-lanceolate. *Hyssopus*
 Leaves crenate to dentate, broader. 10
10. Upper lip of the corolla inconspicuous,
 reduced to two small lobes. *Teucrium*
 Upper lip of corolla distinctly visible. 11
11. Upper lip of calyx much wider than the
 other 4. *Dracocephalum*
 Upper lip of calyx equal to at least the
 adjacent ones. 12
12. Flowers white; leaves deltoid or deltoid-
 ovate, cordate to subcordate at base;
 calyx oblique. *Nepeta*
 Flowers purplish; leaves oblong to ovate,
 obtuse to rounded at base; calyx
 regular. *Stachys*

13. Flowers solitary or in axillary racemes. *Scutellaria*
 Flowers in axillary clusters. 14
14. Leaves palmately 3-lobed. *Leonurus*
 Leaves not palmately lobed, crenate to
 serrate. 15
15. Calyx with 10 lobes, each hooked at the
 tip. *Marrubium*
 Calyx without hooks at tips of lobes. 16
16. Calyx distinctly 2-lipped, the upper part
 differing from the lower part. 17
 Calyx indistinctly 2-lipped or regular. 19
17. Upper calyx lobe 3–4 times wider than
 the other 4. *Dracocephalum*
 Upper 3 calyx lobes differing from the
 lower 2. 18
18. Upper 3 lobes reduced to teeth, lower
 lobes at least 3 times as long. *Melissa*
 Upper lobes short and broad, lower lobes
 long, awl-shaped. *Hedeoma*
19. Corolla 4- or 5-lobed, hardly, if at all,
 2-lipped. 20
 Corolla distinctly 2-lipped. 21
20. Stamens 2; flowers white, sessile. *Lycopus*
 Stamens 4; flowers blue or lavender,
 pedicellate. *Mentha*
21. Calyx lobes much shorter than the tube;
 stems decumbent or creeping. *Glechoma*
 Calyx lobes as long as, or longer than, the
 tube; stems erect or ascending. 22
22. Calyx lobes with sharp, spiny, glabrous
 tips. *Galeopsis*
 Calyx lobes herbaceous, pubescent to the
 tips. *Lamium*

Agastache giant-hyssop

Agastache foeniculum (Pursh) Ktze. (Fig. 182)

GIANT-HYSSOP

An erect, branched perennial, with smooth or minutely hairy stems 30–80 cm high. Leaves ovate or triangular-ovate, green above and pale below, 2–7 cm long, with short stalks. Flowers blue, 6–12 mm long, in a dense spike, sometimes interrupted, 2–10 cm long and 12–18 mm thick at ends of stems. Plant has a pleasant anise-like odor. Common; in open woodlands and semi-open prairies; throughout Parklands. Syn.: *A. anethiodora* (Nutt.) Britton.

Dracocephalum false dragonhead

- Inflorescence a dense terminal spike. *D. parviflorum*
 Inflorescence of loose interrupted racemes. *D. thymiflorum*




Fig. 182. Giant-hyssop, *Agastache foeniculum* (Pursh) Ktze.

An erect, usually branched, annual or biennial herb, with a finely hairy stem 30–50 cm high. Leaves oblong to lanceolate, 2–5 cm long, stalked, and with rather large pointed teeth. Flower clusters dense, at ends of branches, usually 2–5 cm long and 2–3 cm wide. Calyx membranous, stiff and spiny, giving heads a prickly stiffness. Light blue corollas scarcely longer than calyx. Locally abundant but not generally common; found in openings in woodlands and old pastures; throughout Parklands; rare in Prairies. Syn.: *Moldavica parviflora* (Nutt.) Britton.

Dracocephalum thymiflorum L.

Similar to *D. parviflorum*, but the leaves with more rounded teeth, and the inflorescence elongate. Introduced; a rare weed. Syn.: *Moldavica thymiflora* (L.) Rydb.

Elsholtzia

Elsholtzia ciliata (Thunb.) Hyl.

An annual herb with erect or ascending branched stems 30–50 cm high. Leaves 3–7 cm long, ovate or ovate-lanceolate, petioled. Inflorescence of terminal and axillary spikes, with the flowers crowded on one side of the axis. Introduced; a rare weed; in wet and shady places.

Galeopsis hemp-nettle

Corolla tube 2–3 times as long as the calyx,

2–3 cm long, yellow. *G. speciosa*

Corolla tube scarcely longer than the calyx,

1.5–2 cm long, purple. *G. tetrahit*

Galeopsis speciosa Mill.

An annual herb with stems 30–100 cm high. Leaves coarsely toothed, 3–8 cm long. Flowers large, yellow; the lower lip often with a purple spot. Introduced; a rare weed.

Galeopsis tetrahit L.

A coarse rough hairy weedy annual 30–100 cm high. Stems usually swollen below nodes where leaves and axillary flower clusters join. Leaves ovate, coarsely toothed, 5–10 cm long. Flowers in both terminal and axillary clusters, with sharp needle-pointed calyx teeth. Corolla 15–20 mm, purple or pink variegated with white. Seeds grayish brown, egg-shaped, about 3 mm long, somewhat similar to common hemp seed. Introduced from Europe; has become a field weed; in various parts of Parklands and Boreal forest.

Glechoma ground-ivy

Glechoma hederacea L.

A creeping perennial, rooting at nodes, with thin dull greenish purple stems 30–40 cm long. Leaves ovate-rounded, cordate at base, 2–4 cm across,

often with a purplish tinge. Flowers 15–20 mm long, light blue, in clusters of 2 or 3 in leaf axils. An introduced plant; has become common in waste places in southeastern Parklands and often found as a garden weed elsewhere.

Hedeoma mock pennyroyal

Hedeoma hispida Pursh ROUGH PENNYROYAL

A low-growing annual 10–20 cm high, with erect branched stems. Leaves narrowly linear, entire, 10–25 mm long. Numerous flower clusters in leaf axils. Flowers bluish purple, about 6 mm long. Found occasionally; on sandy soil, eroded slopes, and abandoned fields; Prairies.

Hyssopus hyssop

Hyssopus officinalis L. HYSSOP

A perennial with a stout woody rootstock, erect stems 30–60 cm high, and entire, lanceolate to oblanceolate leaves 1–3 cm long. Flowers 3–7 in a cluster, with the upper ones on each stem or branch forming a spike-like terminal inflorescence; corolla blue, about 1 cm long. Introduced; occasionally cultivated for medicinal purposes, and rarely weedy.

Lamium dead-nettle

Leaves all stalked, ovate; flowers white;
perennial. *L. album*

Upper leaves not stalked, roughly circular;
flowers red or purplish; annual or biennial. *L. amplexicaule*

Lamium album L. WHITE DEAD-NETTLE

A perennial from creeping rootstocks, 30–50 cm high. Leaves ovate, sometimes somewhat cordate at the base, slightly hairy, 2–7 cm long. Pure white flowers 20–30 mm long, in clusters of 7 or 8 in each leaf axil. An introduced plant, has been found in the Prairie Provinces, but is, as yet, rare.

Lamium amplexicaule L. HENBIT

A sparingly hairy annual or biennial with slender stems, branched from base, often somewhat decumbent, 15–50 cm high. Leaves almost round, with rounded teeth; the lower ones with slender stalks; the upper ones usually stalkless and somewhat clasping the stem, 1–5 cm across. Flowers red or purplish with spots on middle lower lobe, 10–20 mm long, in few-flowered axillary and terminal clusters. An introduced plant; uncommon; but found throughout the Prairie Provinces.

Leonurus motherwort

Calyx strongly 5-angled and 5-ribbed; bracts
small. *L. cardiaca*

Calyx not angled, 10-ribbed; bracts large. *L. sibiricus*

Leonurus cardiaca L. MOTHERWORT

An introduced perennial 50–150 cm high, with slightly hairy or downy stems. Leaves stalked, 2–10 cm long, decreasing in size toward upper part of

stem, deeply and irregularly cleft into 3–7 lobes. Pink flowers about 1 cm long, in dense whorls around axils of upper leaves. Upper lip of corolla densely white hairy. Occasionally persists as an intruder in shelterbelts.

Leonurus sibiricus L.

SIBERIAN MOTHERWORT

Biennial plants, with softly retrorse-pubescent stems to 1 m high. Leaves ovate or rotund in outline, deeply 3-parted, with the divisions toothed. Flowers purple, about 1 cm long; upper lip of the corolla pubescent. Introduced; a rare weed.

Lycopus water-horehound

Perennial plants of swamps and wet places, with flowers in dense axillary clusters.

1. Leaves more or less deeply incised or pinnatifid. *L. americanus*
Leaves toothed but not deeply incised. 2
2. Calyx lobes narrow and acute, longer than nutlets. *L. asper*
Calyx lobes ovate and blunt, not as long as nutlets. *L. virginicus*

Lycopus americanus Muhl.

WATER-HOREHOUND

An erect perennial 30–80 cm high. Leaves at definite intervals on stem, usually almost horizontally, giving plant an open and regular appearance. Leaves 2–8 cm long, lanceolate, short-stalked, and except for the upper ones, deeply cut or incised. Bluish white flowers about 3 mm across, in dense clusters around stem at axils of leaves. Fairly common; in moist places, stream banks, and swamps; throughout the Prairie Provinces.

Lycopus asper Greene

WESTERN WATER-HOREHOUND

A perennial 30–50 cm high; narrowly lanceolate leaves 2–7 cm long, with small-toothed margins. Leaves either stalkless or with a very short stalk. Tiny flowers in close clusters in leaf axils. Very common; in wet places and swamps in the western Prairies and Parklands, but comparatively scarce toward the eastern part.

Lycopus virginicus L. var. *pauciflorus* Benth.

NORTHERN WATER-HOREHOUND

An erect perennial 15–70 cm high, with somewhat toothed margined lanceolate leaves 2–7 cm long. Flowers about 3 mm long, in axils of leaves. Rare; found in wet areas; northwestern Boreal forest. Syn.: *L. uniflorus* Michx.

Marrubium horehound

Marrubium vulgare L.

COMMON HOREHOUND

A strongly aromatic perennial, with stout, erect, white pubescent stems 40–60 cm high. Leaves ovate, 3–5 cm long, with the lower ones petioled, white pubescent. Flowers about 6 mm long, densely crowded in upper axils; calyx

teeth awl-shaped, smooth and hooked at the apex, catching on fur and clothing. Introduced; occasionally cultivated and escaped.

Melissa balm

Melissa officinalis L.

HONEY BALM

Perennial plants with erect stems 40–80 cm high. Leaves 4–7 cm long, long-petioled, ovate to deltoid-ovate, coarsely crenate. Inflorescence of few-flowered clusters; corolla blue to white, 10–15 mm long. Introduced; occasionally cultivated for honey, flavoring, and medicinal purposes; rarely spreading.

Mentha mint

Flowers in axillary clusters. *M. arvensis*

Flowers in terminal spikes. *M. spicata*

Mentha arvensis L. var. *villosa* (Benth.) S. R. Stewart

FIELD MINT

Erect perennial herbs with a strong but pleasant mint odor, growing to 10–50 cm high. Square stems with a line of hairs running down each angle. Leaves almost hairless, but with minute glandular dots on both surfaces. Ovate to lanceolate in shape, 1–5 cm long, sometimes long-stalked and sometimes short-stalked. Flowers pink, about 3 mm long, in crowded whorls around stems at leaf axils. Common; in sloughs and wet places, often growing in water; found throughout the Prairie Provinces. Various authorities have split the mints into several species using hairiness of leaves or length of leaf stalks in relation to the size of flower clusters as distinguishing characteristics. It would appear, however, that the mints of the Prairie Provinces should at present be considered as a variety of one species.

Mentha spicata L.

SPEARMINT

Erect perennial herbs to 50 cm high, glabrous or nearly so. Leaves 2–6 cm long, sessile or subsessile, oblong-lanceolate, sharply serrate. Spikes several, terminating stem and short branches from the upper axils, 3–12 cm long, continuous or somewhat interrupted; flowers about 5 mm long. Introduced; cultivated for flavoring, and occasionally escaped.

Monarda wild bergamot

Monarda fistulosa L. (Fig. 183)

WILD BERGAMOT

Erect perennial plants with a strong but pleasant odor, 60–100 cm high. Leaves lanceolate or ovate-lanceolate, rounded or somewhat cordate based, 2–10 cm long. In the type species, leaves are only slightly hairy and have stalks 10–25 mm long. Inflorescence terminal and head-like, in clusters 3–6 cm across. Calyxes narrow green tubes with small purplish teeth. Corollas hairy, 20–25 mm long, protruding far above calyxes, pink or lilac, very conspicuous. Fairly common; on hillsides, thickets, and in shady places; eastern Parklands and Boreal forest, being replaced farther west by the variety.




Fig. 183. Wild bergamot, *Monarda fistulosa* L.

The var. *menthaefolia* (Graham) Fern., western wild bergamot, which is much more common than the species, differs by having very short leaf stalks, less than 6 mm long, and leaves often downy-hairy. Occasionally white-flowered specimens are found. Common; in edges of woods, along shady creek banks and coulees, and in shelter of shrubby patches on prairies; throughout almost all of the Prairie Provinces. Syn.: *M. menthaefolia* Graham.

Nepeta catnip

Nepeta cataria L.

CATNIP

A very hairy branching perennial 30–75 cm high, with ovate, stalked leaves, cordate at base, 2–7 cm long. Flowers white or pale purple with dark spots, hairy, about 12 mm long, in fairly dense terminal heads, somewhat similar to American dragonhead. An introduced plant; found rarely in disturbed areas.

Physostegia false dragonhead

Physostegia virginiana L. var. *formosior* (Lunell) Boiv. FALSE DRAGONHEAD

An erect, rarely branched perennial 30–100 cm high. Leaves usually rhomboid-lanceolate, 6–10 cm long, and stalkless, with short sharp teeth. Flowers in terminal spikes, purple, 20–25 mm long. Fairly common locally; along stream banks and moist places; southeastern Parklands and Boreal forest. Plants with narrower leaves, oblong-lanceolate to lanceolate, are var. *ledinghamii* Boiv.; Boreal forest. Plants with linear-lanceolate leaves, and flowers 10–15 mm long are var. *parviflora* (Nutt.) Boiv. Syn.: *P. parviflora* Nutt.; *Dracocephalum nuttallii* Britton.

Prunella selfheal

Prunella vulgaris L.

SELFHEAL

An introduced perennial plant growing from running rootstocks, 5–30 cm high. Leaves ovate-lanceolate, rather blunt at apex, 2–5 cm long. Flowers in short dense terminal spikes, violet purple, 8–12 mm long. Found occasionally; in woodlands and moist places in grassland; Parklands and Rocky Mountains.

Salvia sage

Flowers 2–4 at each node of the raceme. *S. reflexa*

Flowers numerous in whorls at the nodes. *S. nemorosa*

Salvia nemorosa L.

WOOD SAGE

Coarse densely downy perennial plants 30–80 cm high. Leaves oblong to ovate-lanceolate, 2–8 cm long; lower leaves with stalks; upper ones stalkless. Flowers in long, terminal, narrow spikes 5–15 cm long and 1–2 cm thick. Corollas deep violet blue, 8–12 mm long. Introduced in alfalfa seed; has been found on two or three occasions in widely scattered spots throughout the Prairie Provinces.

Salvia reflexa Hornem.

LANCE-LEAVED SAGE

Erect, much-branched annuals 30–60 cm high; stems finely pubescent with recurved hairs. Leaves 3–5 cm long, linear-lanceolate to lanceolate; petioles 1–2 cm long. Racemes 5–10 cm long, erect; flowers usually 2, rarely 3 or 4 at a node, about 1 cm long, blue; the corolla scarcely longer than the calyx. Found occasionally as a weed; in southeastern Parklands.

Scutellaria skullcap

Perennial herbs with few flowers, usually borne in leaf axils. Corolla has arched upper lip and is much longer than calyx, which has a crest-like protuberance on upper lip.

1. Flowers borne singly or in pairs in axils of leaves. 2
Flowers borne in several-flowered axillary or terminal racemes. *S. lateriflora*
2. Flowers 15–25 mm long. *S. galericulata*
Flowers 7–10 mm long. *S. parvula* var. *leonardii*

***Scutellaria galericulata* L.**

MARSH SKULLCAP

An erect-stemmed perennial from creeping roots, 30–60 cm high. Leaves oblong to oblong-lanceolate, wavy-margined, 2–6 cm long; lower ones short-stalked; upper ones stalkless. Flowers borne either singly or in pairs at axils of leaves, blue, 15–25 mm long. Fairly common; in wet places and along stream banks; throughout the Prairie Provinces. Syn.: *S. epilobiifolia* Hamilton.

***Scutellaria lateriflora* L.**

BLUE SKULLCAP

Very similar to *S. galericulata*, with ovate leaves. Flowers only about 6–10 mm long, in loose several-flowered racemes at leaf axils and occasionally at the end of the stem. Fairly common; along stream banks, in swamps, and wet places; southeastern Parklands.

***Scutellaria parvula* Michx. var. *leonardii* (Epling) Fern.**

SMALL SKULLCAP

Small perennials, with rootstocks deeply constricted between bead-like segments; stems 10–20 cm high. Main stem leaves sessile, 10–16 mm long, ovate-lanceolate to somewhat deltoid, glabrous or slightly scabrous. Flowers axillary, blue; calyx glabrous or somewhat scabrous. Very rare; in peaty soil over rock outcrops; southeastern Boreal forest.

Stachys hedge-nettle

***Stachys palustris* L. var. *pilosa* (Nutt.) Fern. (Fig. 184)**

MARSH HEDGE-NETTLE

A hairy, branched perennial, usually erect, but occasionally decumbent, 30–80 cm high. Leaves somewhat coarse, hairy, lanceolate to oblong-lanceolate, 2–10 cm long, generally rounded at the base, sometimes with very short stalks but usually without. Flowers pale purplish with darker spots, 10–15 mm long, in axillary clusters in the top portion of the plant, sometimes appearing as a leafy spike. Common; in moist places and along stream banks; throughout the Prairie Provinces.




Fig. 184. Marsh hedge-nettle, *Stachys palustris* L. var. *pilosa* (Nutt.) Fern.

Teucrium germander

Teucrium canadense L. var. *occidentale* (Gray) McCl. & Epl.

HAIRY GERMANDER

A very hairy, branching perennial 30–75 cm high. Leaves narrowly ovate to oblong-lanceolate, short-stalked, 2–7 cm long, white hairy beneath. Purplish flowers 6–10 mm long, in a spike-like terminal raceme. Very scarce, but has been reported from east central Saskatchewan.

SOLANACEAE—potato family

Herbs or vines with alternate leaves without stipules. Flowers perfect, and, with one exception, regular. Corolla varies from funnel-like and bell-like to rotate (wheel-shaped), with 5 stamens. Fruit either a berry or a capsule. Many native members of this family have either narcotic or poisonous properties, although some parts are edible.

- 1. Plants shrubby, often spiny. *Lycium*
Plants herbaceous. 2
- 2. Corolla rotate (wheel-shaped). *Solanum*
Corolla bell-shaped or funnel-shaped. 3
- 3. Fruit a berry, often enclosed in an
inflated bladder-like calyx. *Physalis*
Fruit a capsule. 4
- 4. Capsule prickly. *Datura*
Capsule not prickly, top falling off when
mature. *Hyoscyamus*

Datura stramonium

Datura stramonium L.

JIMSONWEED

An annual, **very poisonous**, weedy plant with stout smooth green to purplish stems, 50–150 cm high. Leaves simple, ovate, irregularly toothed, 7–20 cm long. Flowers funnel-shaped, white or purplish, 5–10 cm long and about 5 cm across, with slender-tipped lobes, borne in the axils of leaves. Fruit a large, prickly capsule 3–6 cm long, breaking into four segments to release the many seeds. A weed, probably escaped from gardens; has been found in two or three places; throughout the Prairie Provinces.

Hyoscyamus henbane

Hyoscyamus niger L. (Fig. 185)

BLACK HENBANE

Sticky, hairy, evil-smelling biennial plants 30–100 cm high from a spindle-shaped root. Leaves roughly ovate or oblong, irregularly toothed or lobed; upper ones clasping the stem, 7–20 cm long. Flowers funnelform, about 2 cm long and 25–35 mm across, greenish yellow with purplish veins, with purple anthers. Borne on one side of stem, crowded at head of stem. Fruit a capsule



Fig. 185. Black henbane, *Hyoscyamus niger* L.

almost 12 mm long, enclosed in a swollen calyx. An introduced **poisonous** plant, fortunately still rare; found in waste places and around gardens; in several locations throughout Prairie Provinces.

Lycium matrimony vine

Lycium halimifolium Mill.

MATRIMONY VINE

Shrubs with branches to 3 m long, glabrous or nearly so, arched or recurved, sometimes climbing; spines, if present, at the nodes. Leaves 2–5 cm long, lanceolate to spatulate, grayish green. Flowers pinkish violet, 10–12 mm wide, in clusters of 1–4; berries scarlet, ellipsoid, 1–2 cm long. Occasionally cultivated, and escaped.

Physalis ground-cherry

1. Corolla rotate, white; flowers 2–4 at a node. *P. grandiflora*
 Corolla funnelform, white or yellowish; flowers solitary. 2
2. Filaments slender, uniform; corolla usually less than 15 mm long; plants annual. 3
 Filaments flat, expanded; corolla usually more than 15 mm long; plants perennial. 4
3. Pubescence villous, with the hairs long and spreading. *P. pubescens*
 Pubescence, if present, of minute hairs. *P. ixocarpa*
4. Corolla white, distinctly 5-lobed; fruiting calyx bright red or orange. *P. alkekengi*
 Corolla yellow or greenish, not distinctly lobed; fruiting calyx brownish or green. 5
5. Pubescence of stem villous, of soft spreading hairs. *P. heterophylla*
 Pubescence of stem of short, stiff, deflexed or recurved hairs. *P. virginiana*

Physalis alkekengi L.

CHINESE LANTERN PLANT

Perennial with creeping rootstocks. Stems soft, short pubescent, usually several together, erect, unbranched, 40–60 cm high. Leaves petioled, ovate, to 15 cm long, to 8 cm broad, glabrous except on the veins. Fruiting calyx to 5 cm long, bright red or orange, and finely reticulately veined at maturity. Berry 10–15 mm in diam, reddish or orange. Introduced as an ornamental; occasionally escaped.

Physalis grandiflora Hook.

LARGE WHITE GROUND-CHERRY

A tall erect annual plant with hairy and somewhat sticky stems 40–60 cm high. Large ovate to lanceolate leaves entire and stalked, 6–12 cm long, somewhat hairy and sticky. Rotate or shallowly bell-shaped flowers, white with pale yellow centers, 25–35 mm across. Fruit a berry partly enclosed in the persistent calyx, about 12 mm long. In disturbed areas, open woodlands, especially on sandy soils; northeastern Boreal forest. Sometimes called *Chamaesaracha grandiflora* (Hook.) Fern.; *Leucophysalis grandiflora* (Hook.) Rydb.

Physalis heterophylla Nees

YELLOW GROUND-CHERRY

Plants with erect or spreading stems, usually with many branches; pubescence distinctly villous, the hairs spreading. Leaves mostly ovate to rhombic, 3–8 cm long, shallowly sinuate-dentate. Corolla 15–20 mm long; the pedicels about 1 cm in flower, elongating to 3 cm in fruit; fruiting calyx 3–4 cm long. Introduced; rarely found as a weed.

Physalis ixocarpa Brot.

TOMATILLO

Stems erect or spreading, branched; nearly glabrous, except on young shoots. Leaves ovate to rhombic, 2–6 cm long, glabrous below. Corolla 10–15 mm wide, yellowish, with a dark center; pedicels about 5 mm long, scarcely elongating in fruit; berry purplish. A rare garden weed.

Physalis pubescens L.

SMALL YELLOW GROUND-CHERRY

Stems branching near the base, spreading or prostrate. Upper part of stem and leaves densely pubescent. Leaves 4–8 cm long, sinuate-dentate, rounded or cordate at base. Corolla 6–10 mm long, yellow; pedicels 3–6 mm in flower, elongating to 1 cm in fruit; berry yellow. A rare garden weed.

Physalis virginiana Mill.

PRAIRIE GROUND-CHERRY

A perennial from creeping rootstocks, about 45 cm high, somewhat hairy. Leaves oblanceolate to spatulate, entire-margined and stalked, 25–50 mm long. Bell-shaped flowers, dull yellow with a brownish center, about 15 mm across. Fruit a yellow or greenish berry, entirely enclosed in the inflated bladder-like ovoid calyx. Rare; but found on sandy prairies and plains; in the eastern part of the Prairie Provinces.

Solanum nightshade

- 1. Climbing perennial vines with deep purple flowers. *S. dulcamara*
- Nonclimbing annual herbs with white or yellow flowers. 2
- 2. Plants prickly; flowers yellow. *S. rostratum*
- Plants not prickly; flowers white. 3
- 3. Leaves entire or merely wavy-margined; erect plants. *S. nigrum*
- Leaves pinnatifid or incised; much branched and decumbently spreading. *S. triflorum*

Solanum dulcamara L. (Fig. 186)

BITTERSWEET

A woody-based perennial climbing or twining vine, with branches several feet long. Leaves 5–10 cm long; upper leaves ovate, cordate at base, and pointed at apex; lower ones often 3-lobed, with a large ovate middle lobe and two small basal side lobes. Flowers in panicles or on compound cymes opposite a leaf stalk, somewhat rotate or wheel-shaped, with separate corolla lobes. Corolla purple with yellow anthers erect and pyramidal around female organ in center of flowers. Flowers 8–12 mm across, succeeded by oval or globose berries, which are red and about 1 cm long. This introduced plant is **very poisonous**, and as yet the only records of it in the area appear to be at Morden, Man., Eastend, Sask., and Edmonton, Alta.

Solanum nigrum L.

BLACK NIGHTSHADE

An erect annual weed 10–30 cm high, with entire-margined ovate leaves 2–5 cm long. White flowers rotate, about 1 cm across, in clusters of 3–10 flowers. Fruits green berries, about 5 mm across, turning black when ripe. An introduced species; now a fairly common weed; in gardens and waste places; across the Prairie Provinces. There appears to be considerable variation in the hairiness of this species, some specimens, especially from the west, being quite hairy, others almost hairless. Pubescent forms with black berries have been called *S. interius* Rydb.; densely pubescent plants with yellow or orange berries, *S. sarachoides* Sendt. Some cultivated forms of this species have edible fruit and are known as wonderberry or garden huckleberry. There is always a danger, however, that the fruit may be injurious to some people.

Solanum rostratum Dunal

BUFFALOBUR

An annual yellowish hairy prickly species 10–50 cm high. Leaves lobed and pinnatifid, 5–8 cm long, and yellowish hairy. Bell-shaped flowers yellow, about 2.5 cm across. Fruit a berry enclosed in a prickly calyx. A straggler from farther south, found occasionally; in Prairies. Syn.: *Androcera rostrata* (Dunal) Rydb.

Solanum triflorum Nutt. (Fig. 187)

WILD TOMATO

A low, spreading annual, forming mats 15–60 cm across. Leaves deeply lobed, oblong or ovate, 2–6 cm long, with scattered hairs. Flowers white, rotate, 6–10 mm across, usually in clusters of three. Fruit a smooth green berry about 12 mm in diam. A native plant found on disturbed areas such as gopher and badger mounds; in southern Prairies. With cultivation it persists as a very bad garden weed. If pulled and turned upside down, it will develop rootlets along the stems and continue to grow. Although it is enjoyed as preserves by some people, it causes violent sickness in others.

(Figs. 186 and 187 overleaf)



Fig. 186. Bittersweet, *Solanum dulcamara* L.




Fig. 187. Wild tomato, *Solanum triflorum* Nutt.

SCROPHULARIACEAE—figwort family

Herbs with opposite, alternate, or whorled leaves with no stipules. Flowers perfect, but generally irregular in shape, with petals partly united into a tube and mostly 2-lipped. Perfect stamens, often only 2, usually 4, rarely 5. Fruit a many-seeded capsule.

1. Five anther-bearing stamens; flowers rotate, yellow, almost regular, in dense spike-like racemes. *Verbascum*
 Only 2 or 4 stamens anther-bearing, others sterile or absent. 2
2. Corolla spurred at the base. 3
 Corolla not spurred at the base. 4
3. Flowers axillary, blue; low slender annuals. *Chaenorrhinum*
 Flowers in terminal racemes, yellow; perennials. *Linaria*
4. Four anther-bearing stamens and 1 long sterile stamen. *Penstemon*
 Two or four anther-bearing stamens and no sterile stamen. 5
5. Anther-bearing stamens 2, any others sterile. 6
 Anther-bearing stamens 4. 9
6. Sepals and petal lobes 5. *Gratiola*
 Sepals and petal lobes 4. 7
7. Petals absent; inflorescence very hairy. *Besseyia*
 Petals present. 8
8. Corolla rotate; stamens not protruding. *Veronica*
 Corolla short-tubular; stamens protruding; inflorescence a very long, narrow, spike-like raceme. *Veronicastrum*
9. Floral leaves or bracts brightly colored, usually red or yellow, or shades of these colors. *Castilleja*
 Floral leaves or bracts usually green, but may be brownish in *Pedicularis*. 10
10. Leaves pinnately lobed or cleft. *Pedicularis*
 Leaves simple, neither pinnately lobed nor cleft. 11
11. Leaves alternate. *Orthocarpus*
 Leaves opposite, whorled, or basal. 12
12. Leaves in a basal rosette; stem leafless; flower solitary; annuals of mud or water. *Limosella*
 Stems leafy; flowers not solitary. 13

13. Flowers large and showy, 10–35 mm long. 14
 Flowers less than 10 mm long. 17
14. Flowers in a terminal spike. *Chelone*
 Flowers in the axils of leaves or bracts. 15
15. Calyx 4-lobed; corolla dark purple; calyx
 and upper leaves violet-tinged. *Bartsia*
 Calyx 5-lobed; calyx and upper leaves or
 bracts green. 16
16. Leaves narrowly linear, 1–6 mm wide,
 entire. *Agalinis*
 Leaves lanceolate to ovate or rotund,
 dentate. *Mimulus*
17. Calyx membranous, veiny, and inflated in
 fruit. *Rhinanthus*
 Calyx not membranous and not much
 inflated in fruit. 18
18. Upper leaves stalkless. 19
 Leaves with stalks; flowers blue, about 6
 mm long. 21
19. Calyx and corolla 5-lobed; upper leaves
 and flowers whorled. *Collinsia*
 Calyx 4-lobed; all leaves and flowers
 opposite. 20
20. Leaves about as wide as long, palmately
 veined. *Euphrasia*
 Leaves lanceolate, pinnately veined. *Odontites*
21. Leaves sharply and coarsely toothed;
 flowers in terminal panicle-like cyme. *Scrophularia*
 Leaves barely toothed; flowers solitary in
 the upper leaf axils. *Melampyrum*

Agalinis *agalinis*

1. Pedicels all or nearly all shorter than the
 calyx. *A. purpurea*
 Pedicels all or nearly all longer than the
 calyx. 2
2. Pedicels very slender; calyx lobes 1–1.5
 mm long; corolla 10–15 mm long. *A. tenuifolia*
 Pedicels stout; calyx lobes 1.5–3 mm
 long; corolla 20–25 mm long. *A. aspera*

Agalinis aspera (Dougl.) Britt.

ROUGH AGALINIS

An annual herb 10–30 cm high, with leaves narrowly linear, very scabrous above, often with axillary clusters of reduced leaves. Flowers on pedicels to 3 cm long; corolla 20–25 mm long, pinkish, with the tube pubescent. Rare; moist grasslands; southeastern Parklands.

Agalinis purpurea (L.) Pennell var. *parviflora* (Benth.) Boiv. PURPLE AGALINIS

An annual herb 10–50 cm high, simple or branched; leaves to 4 mm wide. Flowers on short pedicels; corolla 15–25 mm long. Very rare; moist grassland; southeastern Parklands.

Agalinis tenuifolia (Vahl) Raf. var. *parviflora* (Nutt.) Pennell SLENDER AGALINIS

An annual herb 10–50 cm high, with stems very slender and usually much branched. Leaves linear, 1–6 mm wide. Pedicels filiform; corolla 10–15 mm long, pinkish. Rare; moist areas; southeastern Boreal forest.

Bartsia bartsia

Bartsia alpina L. VELVET BELLS

Plants with a somewhat woody rootstock; stems erect, 5–15 cm high, viscid-villous. Leaves sessile, ovate, 10–25 mm long, dentate, clasping the stem. Inflorescence a dense raceme; flowers to 2 cm long, glandular. Plants blackening on drying. Arctic meadows; northeastern Boreal forest.

Besseyia kittentails

Besseyia wyomingensis (A. Nels.) Rydb. KITTENTAILS

Softly hairy perennial plants 10–30 cm high, often with reddish-tinged leaves. Basal leaves stalked, ovate or oblong; stem leaves without stalks, alternate, and smaller. Inflorescence a dense terminal spike, 2–5 cm long in flower, and lengthening to 5–15 cm in fruit, very hairy, and usually with a purplish tinge. Flowers bearing 2 protruding stamens but no petals. Fruits many-seeded capsules. Found on open hillsides in the Foothills region and also on benchland in the Cypress Hills. Syn.: *B. cinerea* (Raf.) Pennell.

Castilleja Indian paintbrush

Annual, biennial, or perennial plants, usually partly parasitic on roots of other plants. Stem generally simple or with a few branches above. Leaves stalkless, alternate, lobed or entire. The bracts (leaves of the inflorescence) usually red or yellow. Flowers in terminal spikes with a 2-lipped corolla; the upper lip arched and called the galea, the lower one with 3 lobes.

1. Plants annual or biennial; inflorescence red or yellow. *C. coccinea*
Plants perennial, with more or less woody rootstock. 2
2. Flowers yellowish, 40–55 mm long, strongly curved. *C. sessiliflora*
Flowers smaller, usually straight and erect. 3
3. Inflorescence pinkish purple to deep purple. *C. raupii*
Inflorescence whitish to scarlet. 4

- 4. Upper stem leaves 3- to 5-lobed, with the lateral lobes linear. 5
- Upper stem leaves entire; floral leaves coarsely 3-lobed. 7
- 5. Inflorescence bright red to scarlet. *C. hispida*
- Inflorescence yellow or whitish. 6
- 6. Calyx 20–25 mm long, with the lobes rounded; corolla 18–22 mm long. *C. cusickii*
- Calyx 15–20 mm long, with the lobes acute; corolla 15–20 mm long. *C. lutescens*
- 7. Inflorescence reddish to scarlet. *C. miniata*
- Inflorescence whitish to yellow. 8
- 8. Flowers 20–30 mm long. *C. occidentalis*
- Flowers 15–20 mm long. 9
- 9. Stem stout, densely pubescent; leaves lanceolate. *C. lutescens*
- Stem slender, finely pubescent to glabrous; leaves narrowly linear-lanceolate. *C. pallida*

Castilleja coccinea (L.) Spreng.

SCARLET PAINTBRUSH

An annual or biennial species with hairy slender stems 30–50 cm high. Basal rosette leaves entire but stem leaves deeply divided into 3–5 linear divisions. Bracts crimson-tipped, 3- to 5-lobed, and usually about as long as flowers. Corolla greenish yellow, with the tube shorter than the calyx, and the upper lip much longer than the lower lip. Fairly common; in meadows and open woods; southeastern Prairies and Parklands.

Castilleja cusickii Greenm.

YELLOW PAINTBRUSH

A hairy simple-stemmed perennial species 20–30 cm high and very leafy. Leaves 2–4 cm long, hairy, and 3- to 5-ribbed; lower leaves lanceolate and entire; upper ones broader and 3- to 5-cleft at tip. The broad pale sulfur yellow bracts about as long as flowers. Calyx 15–20 mm long; corolla rarely protruding beyond calyx. Found occasionally; in extreme southwestern Alberta. Syn.: *C. lutea* Heller.

Castilleja hispida Benth.

HISPID PAINTBRUSH

Stems several, 20–40 cm high, hirsute, not glandular. Leaves lanceolate to ovate, with 2 or 3 pairs of ascending lobes. Inflorescence villose, scarlet to red; calyx cleft into oblong, rounded lobes. Montane grasslands and open woods; southern Rocky Mountains.

Castilleja lutescens (Greenm.) Rydb.

STIFF YELLOW PAINTBRUSH

Stems several, 30–50 cm high, scabrous to short-pubescent. Leaves linear-lanceolate to lanceolate, entire or the upper ones shallowly lobed. Inflorescence pale yellow, somewhat hirsute; calyx cleft into ovate acute-to-rounded lobes. Montane grasslands; southern Rocky Mountains.

A perennial growing to 40–60 cm high; stem usually without hairiness below inflorescence, sometimes branched above. Linear pointed leaves 2–5 cm long. Bracts broader than leaves, scarlet or bright red; the lower ones usually having 2 or 5 teeth near summit. Flowers longer than bracts, green with red margins, lip a little shorter than tube. Found in open pine woods; in the Foot-hills region and also in the Cypress Hills.

Castilleja occidentalis Torr.

LANCE-LEAVED PAINTBRUSH

A smooth-stemmed perennial 15–50 cm high. Leaves 5–10 cm long; lower leaves narrower than upper ones, and having 3–5 nerves. Yellowish, greenish white, or purple bracts oblong, oval or obtuse, sometimes with small teeth, and about as long as flower. Corolla 12–18 mm long; the upper lip 2–4 times as long as the lower. Fairly abundant; in open woodlands; southeastern Parklands. Syn.: *C. acuminata* (Pursh) Spreng.; *C. sulphurea* Rydb.

Castilleja pallida (L.) Spreng. var. *septentrionalis* (Lindl.) Gray

LABRADOR PAINTBRUSH

Plants finely pubescent to glabrous; stems several, 15–30 cm high, simple. Leaves linear-lanceolate, entire, with upper ones sometimes shallowly lobed. Inflorescence pale yellow; calyx cleft into lance-ovate, acute lobes; corolla puberulent above. Marshy areas; northeastern Boreal forest.

Castilleja raupii Pennell

PURPLE PAINTBRUSH

Stems usually several, 15–30 cm high, thinly pubescent. Leaves linear or linear-lanceolate, 2–5 mm wide. Inflorescence narrow to fairly dense, violet to deep violet purple; corollas shorter than bracts. Open areas, open forest, and forest margins; Boreal forest, Peace River district.

Castilleja sessiliflora Pursh

DOWNY PAINTBRUSH

A perennial pale ashy gray downy-haired species 10–30 cm high. Leaves 2–4 cm long; lower leaves generally linear and entire; upper ones cleft into narrow, spreading segments. Bracts green, shorter than flowers. Corolla 40–55 mm long, yellowish white; the upper lip about twice as long as the 3-lobed lower lip. Found occasionally; on dry hills and prairies; throughout Prairies, southeastern Parklands, and in Cypress Hills.

Chaenorrhinum small-snapdragon

Chaenorrhinum minus (L.) Lange

SMALL-SNAPDRAGON

A low glandular hairy annual 15–30 cm high, usually branched. Leaves alternate, linear to linear-spatulate, 10–25 mm long. Flowers blue, 6–8 mm long, with a short spur, borne on short stalks in axils of leaves. An introduced plant; not common, but found occasionally; on or in the vicinity of railroad grades; throughout the Prairies and Parklands.



Fig. 188. Red Indian paintbrush, *Castilleja miniata* Dougl.

Chelone turtlehead

Chelone glabra L. var. *linifolia* Colem. TURTLEHEAD

Perennial plants 50–80 cm high, with stems simple or branched above. Leaves linear or nearly so, serrate, short-petioled to sessile. Inflorescence 3–8 cm long; flowers greenish or yellowish white, 20–25 mm long. Rare; wet areas and shrubbery; southeastern Boreal forest.

Collinsia bluelips

Collinsia parviflora Dougl. BLUE-EYED MARY

A low much-branched spreading annual with slender purplish stems 10–30 cm long. Leaves linear to lanceolate, 10–25 mm long, opposite or with upper leaves in whorls of 3 or 5. Flowers blue, about 6 mm long, borne either singly or in clusters on stalks in axils of upper leaves. Not common, but plentiful locally; found in shady woods or openings in woodlands; throughout southern Parklands, Cypress Hills, and southern Rocky Mountains.

Euphrasia eyebright

Euphrasia arctica Lange NORTHERN EYEBRIGHT

Annual plants with slender stems 10–20 cm high. Leaves in 3–10 pairs, ovate, 5–15 mm long, with 3–5 teeth on each margin. Flowers white with purple lines, 4–7 mm long. In more or less disturbed soils; northern Boreal forest. In var. *dolosa* Boiv. the flowers only 3–4 mm long, and lacking the purple lines. Northern Boreal forest and Rocky Mountains.

Gratiola hedge-hyssop

Gratiola neglecta Torr. CLAMMY HEDGE-HYSSOP

Annual or perennial herbs 10–20 cm high, somewhat sticky hairy. Leaves opposite, linear to oblong-lanceolate, 1–5 cm long, without stalks. Flowers pale yellow to whitish, 6–10 mm long, borne singly on long stalks in axils of leaves. Found occasionally; in mud or shallow water; particularly throughout the Prairies and southern Parklands.

Limosella mudwort

Limosella aquatica L. MUDWORT

A low annual with stems that root at nodes, 7–10 cm high. Leaves linear or spatulate, 5–30 mm long, on long stalks from plant crown. White or purplish flowers solitary on short stalks from base of plants, about 3 mm long. Found occasionally; rooted in mud or floating in shallow water around lakes and in streams; throughout most of the Prairie Provinces.

Linaria toad-flax

1. Plants perennial, with extensive root-stocks; flowers yellow and orange. 2

- Plants annual; flowers white, pink, or purple, not yellow. 3
2. Stem leaves ovate to lanceolate, clasping. *L. dalmatica*
 Stem leaves linear to linear-lanceolate, not clasping. *L. vulgaris*
3. Lower lip of corolla violet; spur 6–10 mm long; pedicels 2–4 mm long. *L. canadensis*
 Lower lip of corolla with an orange spot; spur 8–15 mm long; pedicels 5–8 mm long. *L. maroccana*

Linaria canadensis (L.) Dum. var. *texana* (Scheele) Pennell FIELD TOAD-FLAX

Stems erect, slender, 20–60 cm high, glabrous, with several sterile shoots spreading from the base. Leaves 1–3 cm long. Racemes at first dense, later elongating; corolla 10–14 mm long, with the spur filiform; seeds distinctly rugose. Rare; an introduced weed.

Linaria dalmatica (L.) Miller BROAD-LEAVED TOAD-FLAX

An introduced perennial with a coarse branching growth 50–80 cm high. Leaves numerous, 1–5 cm long, ovate to lanceolate, often clasping the stem at the base. Flowers borne on a long spike-like raceme, 2–5 cm long, with long spurs, bright yellow, sometimes with orange-colored throat entrance. Grown as an ornamental in gardens, but has escaped in many places and is becoming a persistent weed.

Linaria maroccana Hook. f. MAROCCAN TOAD-FLAX

Plants to 50 cm high, with remote linear leaves. Raceme becoming lax; corolla violet; the lower lip with a large orange spot; the spur conical. Rare; an introduced weed.


Linaria vulgaris Miller BUTTER-AND-EGGS, YELLOW TOAD-FLAX

A perennial from creeping rootstocks, 20–60 cm high. Leaves linear, stalkless, alternate, 2–7 cm long. Flowers bright yellow with orange throat entrance and a long spur at base, mouth of flowers closed, 20–35 mm long, in dense terminal raceme. Introduced as a garden plant from Europe, and found as a persistent weed of fields and waste places; throughout the Prairies and Parklands.

Melampyrum cow-wheat

Melampyrum lineare Desr. (Fig. 189) COW-WHEAT

Annual plants 15–50 cm high with lanceolate to linear-lanceolate leaves 2–6 cm long. Lower leaves entire-margined; upper floral leaves more ovate and often bearing bristle-pointed teeth. Flowers 8–12 mm long, generally whitish with a yellowish lower lip, borne either solitary in upper leaf axils or in a terminal leafy spike. Fairly common; on dry, sandy soil in woodlands; Boreal forest.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Melampyrum lineare* Desr.

LOCATION: Waterhen River, Hwy 4

COLL: J. Loozen

DET:

COLL. NO: 13253

DATE: 13 August 1969

ELEVATION: 450 m

HABITAT: Open pine forest



Fig. 189. Cow-wheat, *Melampyrum lineare* Desr.

Mimulus monkeyflower

Perennial herbs of streams and very wet places, with large showy flowers. Calyx angled, and corolla open-throated with upper lip spreading or reflexed.

1. Flowers blue or crimson. 2
Flowers yellow. 3
2. Flowers blue to violet; plants glabrous. *M. ringens*
Flowers crimson to reddish; plants glandular pubescent. *M. lewisii*
3. Calyx regular or nearly so; corolla 5–12 mm long. *M. floribundus*
Calyx distinctly 2-lipped; corolla 12–25 mm long. 4
4. Stems mostly creeping; calyx 5–12 mm long; corolla to 25 mm long. *M. glabratus*
Stems mostly erect; calyx 12–17 mm long; corolla 25–45 mm long. *M. guttatus*

Mimulus floribundus Dougl.

MONKEYFLOWER

An annual herb, with stems weak, villosely glandular pubescent, much branched, 5–35 cm long. Leaves ovate, dentate, rounded or cordate at base, the larger ones 1–3 cm long. Flowers yellow; the corolla 5–12 mm long, on pedicels 5–25 mm long. Wet places; southern Rocky Mountains.

Mimulus glabratus HBK.

SMOOTH MONKEYFLOWER

Stems weak, glabrous or minutely pubescent. Leaves 1–4 cm long, ovate to rotund, rounded at base. Flowers on pedicels 1–2 cm long, bright yellow, pubescent in the throat. Rare; wet places, springs; southeastern Parklands.

Mimulus guttatus DC.

YELLOW MONKEYFLOWER

Perennial plants from basal branches that root at nodes. Stem 10–50 cm high, bearing opposite ovate or rounded leaves 1–5 cm long; lower leaves stalked; upper ones usually clasping the stem. Conspicuous flowers 25–45 mm long, bright yellow with an open mouth, somewhat hairy inside the lower lip; calyx inflating somewhat at maturity to contain the many-seeded capsule. Not common, but plentiful locally; found in running streams; in Prairies, southern Rocky Mountains, and in Cypress Hills. Syn.: *M. langsdorfii* Donn.

Mimulus lewisii Pursh

LEWIS MONKEYFLOWER

A somewhat sticky hairy perennial 30–60 cm high. Leaves lanceolate to oblong, 3–7 cm long. Crimson or reddish flowers 3–5 cm long and somewhat hairy inside open throat. Found along stream banks; in southern Rocky Mountains.

Mimulus ringens L.

BLUE MONKEYFLOWER

A hairless perennial from rootstocks 30–100 cm high. Leaves lanceolate to oblong, opposite, 5–10 cm long, without stalks. Flowers blue or violet, 20–35 mm long, with a narrow throat. Fairly common; in swamps, along streams and lakes; southeastern Parklands and Boreal forest.

Odontites eyebright

Odontites serotina (Lam.) Dum.

LATE-FLOWERING EYEBRIGHT

An annual herb with stems 10–40 cm high, finely retrorse-pubescent, usually branched. Leaves lanceolate, 1–3 cm long, roughly pubescent. Flowers nearly sessile, about 1 cm long, pubescent, light red. Introduced; a rare weed; along roads; in Boreal forest.

Orthocarpus owl's-clover

Orthocarpus luteus Nutt.

OWL'S-CLOVER

A short erect annual 10–30 cm high, sometimes with erect growing branches. Leaves 10–25 mm long, linear or narrowly lanceolate, crowded and ascending. Yellow flowers 10–15 mm long, on very leafy narrow terminal spikes, and followed by numerous many-seeded capsules. Common; on open dry prairie; throughout the Prairies and Parklands.

Pedicularis lousewort

Annual or perennial herbs with pinnately cleft or lobed leaves and flowers in dense terminal spikes or racemes.

- 1. Galea prolonged into a tubular recurved beak. 2
 Galea not prolonged, the apex broad. 5
- 2. Flowers purplish red or pink. *P. groenlandica*
 Flowers white or yellow. 3
- 3. Leaves serrulate to crenate. *P. racemosa*
 Leaves deeply lobed or divided. 4
- 4. Beak of galea bent into a half circle. *P. contorta*
 Beak of galea slightly bent, at most one-quarter of a circle. *P. lapponica*
- 5. Flowers in an elongating inflorescence, becoming axillary. *P. parviflora*
 Flowers in well-defined, more or less crowded racemes. 6
- 6. Stem leaves opposite or subopposite. *P. lanceolata*
 Stem leaves alternate. 7
- 7. Flowers 3–5 in a short raceme; corolla 30–35 mm long. *P. capitata*
 Flowers more numerous, usually less than 25 mm long. 8
- 8. Inflorescence glabrous. *P. flammea*
 Inflorescence pubescent to glandular pubescent. 9

9. Inflorescence pubescent in the rachis only. *P. lapponica*
 Inflorescence more pubescent. 10
10. Bracts of the inflorescence strongly ciliate. *P. bracteosa*
 Bracts not ciliate. 11
11. Inflorescence puberulent. *P. labradorica*
 Inflorescence more or less densely villous-lanate. 12
12. Flowers predominantly yellow. 13
 Flowers some shade of pink. 14
13. Calyx distinctly oblique and 2-lipped. *P. canadensis*
 Calyx not 2-lipped, the lobes subequal. *P. oederi* var. *albertae*
14. Stem leaves 1–3 below the middle of the stem. *P. sudetica*
 Stem leaves more or less numerous along entire stem. 15
15. Pubescence of calyx long, woolly, obscuring the veins. *P. lanata*
 Pubescence of calyx less dense; veins not obscured. *P. langsдорffii*

Pedicularis bracteosa Benth.

WESTERN LOUSEWORT

A perennial species with an erect stem 30–90 cm high. Leaves alternate, 10–30 cm long, divided so deeply into toothed leaflets that they appear pinnate. Flowers pale yellow, about 2 cm long, in a dense terminal spike 10–30 cm long. Bracts of inflorescence almost as long as flowers, but a shorter-bracted variety has been named *P. montanensis* Rydb. Fairly plentiful; among bushes and in moist places; Rocky Mountains and Cypress Hills.

Pedicularis canadensis L.

COMMON LOUSEWORT

A somewhat hairy, erect, perennial herb 15–50 cm high. Oblong leaves deeply incised or lobed, 7–15 cm long. Flowers 20–25 mm long, yellow or reddish, in a dense spike at head of stem. Fairly common; in woodlands; southeastern Parklands and Boreal forest.

Pedicularis capitata Adans.

LARGE-FLOWERED LOUSEWORT

Small plants, with the stems usually solitary, 10–20 cm high, leafless, glabrous to somewhat pubescent. Inflorescence a terminal raceme with 2–5 flowers, yellowish white to pinkish-tinged, 25–40 mm long. Alpine tundra; southern Rocky Mountains.

Pedicularis contorta Benth.

COILED-BEAK LOUSEWORT

Plants with stems 30–60 cm high, glabrous throughout. Leaves to 15 cm long, 2–3 cm wide. Inflorescence dense; the bracts the same length as or shorter than the flowers; flowers 15 mm long, white or pale yellow, with the hood of the galea recurved in a semicircle, finely purple-spotted. Dry montane and alpine slopes; southern Rocky Mountains.

Pedicularis flammea L.

FLAME-COLORED LOUSEWORT

Small plants with spindle-shaped roots; stems 5–20 cm high, with 1 or 2 stem leaves. Inflorescence few-flowered, 2–5 cm long; flowers 15–20 mm long, with the lower lip yellow, the galea purple to crimson. Arctic and alpine meadows; northeastern Boreal forest, northern Rocky Mountains.

Pedicularis groenlandica Retz. (Fig. 190)

ELEPHANT'S-HEAD

An erect hairless perennial 20–60 cm high, from rootstocks. Leaves lanceolate, very deeply incised; lower ones stalked, 5–15 cm long. Flowers borne on a terminal spike 5–15 cm long, usually purple or deep red, 10–15 mm long. Upper lip of corolla long, curved downward and then upward, making flowers resemble small red elephant heads. Found in swampy places and stream banks; in Boreal forest and Rocky Mountains.

Pedicularis labradorica Wirsing

LABRADOR LOUSEWORT

Low very branchy plants to 30 cm high; stems and foliage puberulent to retrorsely pilose. Inflorescence about 5 cm long; flowers about 15 mm long, at first yellow, and then fading purplish. Bogs and tundra; northeastern Boreal forest.

Pedicularis lanata C. & S.

WOOLLY LOUSEWORT

Small plants, with densely woolly stems; leaves 2–3 cm long, glabrous. Inflorescence 3–5 cm long, with flowers 20–25 mm long, reddish purple to rose. The lower lip the same length as the galea. Mountain meadows; southern Rocky Mountains.

Pedicularis lanceolata Michx.

SWAMP LOUSEWORT

A practically hairless perennial, with stout erect stems 30–90 cm high. Leaves both alternate and opposite, 5–12 cm long, with short marginal lobes. Yellow flowers 20–25 mm long with slight difference between lengths of upper and lower lips. Borne in a short dense terminal spike. Not common; but has been found in swamps; eastern Parklands and Boreal forest.

Pedicularis langsdorffii Fisch.

ARCTIC RATTLE

Similar to *P. lanata*, but not so woolly, the flowers deep pink, and the lower lip only half as long as the galea. Alpine slopes; southern Rocky Mountains.

Pedicularis lapponica L.

LAPLAND RATTLE

Plants 10–20 cm high, with stems and inflorescence densely retrorse-pubescent, otherwise glabrous. Inflorescence few-flowered, with the flowers 15–20 mm long, yellow. Tundra; northeastern Boreal forest.

Pedicularis oederi Vahl. var. *albertae* (Hulten) Boiv.

OEDER'S LOUSEWORT

Low plants 5–15 cm high, with glabrous stems and leaves; leaves 3–6 cm long. Inflorescence 3–10 cm long, woolly, and somewhat glandular; densely flowered; flowers 20–25 mm long, yellow, with the galea purple-tinged. Alpine tundra; southern Rocky Mountains.



Fig. 190. Elephant's-head, *Pedicularis groenlandica* Retz.

An annual or biennial species 15–50 cm high, much branched. Leaves 2–4 cm long, deeply divided. Flowers 10–15 mm long, purple, with galea rounded at apex. Uncommon; but has been found in marshlands and bogs; Boreal forest.

Pedicularis racemosa Dougl.

Plants with stems 30–50 cm high; leaves all on the stem, 4–7 cm long, 1–2 cm wide, glabrous. Inflorescence loose, leafy-bracted; the bracts as long as or longer than the flowers; flowers 10–15 mm long, pale violet to purple, with the galea strongly recurved. Moist open areas in montane woods; southern Rocky Mountains.

Pedicularis sudetica Willd.

Plants 10–20 cm high, with stems and leaves glabrous or somewhat short-pubescent. Leaves mostly basal, typically with a solitary stem leaf. Inflorescence 3–5 cm long, densely woolly; flowers 15–20 mm long, violet purple, with the lower lip pale pink, purple-dotted. Wet tundra; northeastern Boreal forest.

Penstemon beardtongue

Perennial herbs with opposite leaves and irregular entire flowers. There are 4 fertile stamens and 1 sterile stamen that is usually more or less bearded. Fruit an ovoid many-seeded capsule. Some authorities spell the generic name *Pentstemon* or *Pentastemon*.

- 1. Plants shrubby; stems woody, decumbent to ascending. 2
- Plants herbaceous; stems erect or nearly so. 3
- 2. Stems 20–50 cm high; leaves lanceolate. *P. fruticosus*
- Stems matted, to 20 cm high; leaves elliptic to orbicular. *P. ellipticus*
- 3. Flowers 8–12 mm long. 4
- Flowers 15–40 mm long. 5
- 4. Flowers dark bluish purple. *P. procerus*
- Flowers white to yellowish. *P. confertus*
- 5. Style yellowish, long-pilose, exserted. *P. eriantherus*
- Style not yellow-pilose, included. 6
- 6. Flowers 35–40 mm long. *P. lyallii*
- Flowers 15–25 mm long. 7
- 7. Flowers white, finely purplish lined or spotted. *P. albidus*
- Flowers blue or purplish. 8
- 8. Flowers blue; foliage glaucous; lower bracts suborbicular. *P. nitidus*

Flowers purple to violet; foliage not glaucous; bracts lanceolate. 9

9. Inflorescence glabrous; flowers bluish purple. *P. albertinus*

Inflorescence glandular pubescent; flowers pale lilac. *P. gracilis*

Penstemon albertinus Greene

BLUE BEARDTONGUE

Stems 10–30 cm high, mostly decumbent at the base, glabrous to the inflorescence. Leaves 2–4 cm long; lower ones petioled, glabrous; upper stem leaves denticulate. Inflorescence interrupted; flowers deep blue to purplish, 15–20 mm long; calyx 2.5–4.0 mm long. Slopes, openings in woods, and disturbed areas; southern Rocky Mountains.

Penstemon albidus Nutt.

WHITE BEARDTONGUE

A rather short, stout, clustered, and erect species 15–30 cm high, with fine hairs on stem. Lower leaves oblong to spatulate and stalked; upper ones stalkless and lanceolate, 2–7 cm long. Flowers about 2 cm long, white; whole flower somewhat sticky hairy. Inflorescence a narrow terminal raceme. Flowering in late May and early June, usually a little later than *P. nitidus*. Common; on dry prairies and hillsides; throughout the Prairies and southern parts of Parklands.

Penstemon confertus Dougl.

YELLOW BEARDTONGUE

A perennial species, usually with several slender stems 10–50 cm high, not hairy. Leaves 5–10 cm long, lanceolate to oblanceolate or linear, and entire-margined. Flowers yellow, 10–15 mm long, in dense clusters on an interrupted terminal spike. Common; on hillsides and dry areas; southern Rocky Mountains; introduced at Swift Current, Sask.

Penstemon ellipticus C. & F.

CREEPING BEARDTONGUE

A mat-forming species, woody at the base; flowering stems 10–20 cm high, puberulent. Leaves thick, elliptic to orbicular, 5–15 mm long. Inflorescence few-flowered, glandular pubescent; flowers 20–35 mm long, purple violet; calyx 7–10 mm long. Alpine slopes and ridges; southern Rocky Mountains.

Penstemon eriantherus Pursh

CRESTED BEARDTONGUE

A somewhat hairy-stemmed erect species 15–40 cm high. Upper portion of stem sticky hairy and glandular. Lower leaves oblong to spatulate and stalked; upper ones often somewhat clasping, acute, 2–5 cm long. Flowers red or purple, 20–25 mm long, with sterile stamens densely hairy; inflorescence a fairly dense, leafy, terminal raceme. Found occasionally; in grassland; southern Rocky Mountains.

Penstemon fruticosus (Pursh) Greene

SHRUBBY BEARDTONGUE

Plants forming dense clumps; stems 10–50 cm high, glabrous below the inflorescence. Leaves thick, leathery, narrowly lanceolate, 1–5 cm long, serrulate to denticulate. Inflorescence somewhat glandular pubescent, somewhat one-sided; flowers 25–40 mm long, purplish blue to violet. Slopes and ridges; southern Rocky Mountains.

Penstemon gracilis Nutt. (Fig. 191A)

LILAC-FLOWERED BEARDTONGUE

A slender erect species 15–40 cm high. Leaves linear-oblong to linear-lanceolate, slightly toothed at margins, 2–7 cm long. Usually flowering in mid-June; flowers borne in clusters of 2 or 3 in the axils of upper leaves, forming an open panicle, pale purple or lilac, 20–25 mm long. Fairly common; on moist prairie, slough margins; throughout Prairies and Parklands.

Penstemon lyallii Gray

LARGE-FLOWERED BEARDTONGUE

Plants with a woody caudex; stems 20–30 cm high, grayish pubescent. Leaves lanceolate to linear-lanceolate, 2–7 cm long, entire to finely denticulate. Inflorescence somewhat glandular pubescent; flowers 35–40 mm long, deep blue to purple; calyx 10–12 mm long. Mountain slopes; southern Rocky Mountains.

Penstemon nitidus Dougl. (Fig. 191B)

SMOOTH BLUE BEARDTONGUE

A stout-stemmed, often branching, hairless species 15–30 cm high. Lower leaves lanceolate to oblanceolate, 2–5 cm long; upper leaves smaller and ovate. Inflorescence a fairly dense, somewhat leafy raceme; flowers 15–20 mm long, usually deep blue, but also varying from purple through all shades of red and pink to pure white; earliest of penstemons, flowering in May. Very common; on dry hills, and eroded areas and banks; throughout Prairies, Parklands, and Rocky Mountains.

Penstemon procerus Dougl. (Fig. 191C)

SLENDER BEARDTONGUE

A slender-stemmed, low-growing perennial 15–30 cm high; stems often decumbent at base. Basal leaves stalked and usually oblanceolate; stem leaves not stalked, oblong to lanceolate, 2–7 cm long. Inflorescence a dense, terminal spike, often interrupted; usually flowering from end of May until early August; flowers crowded on the stem; corolla dark blue, 10–15 mm long. Common; usually found in large colonies around slough margins, in shelter of shrubs, and in openings in woodlands; Prairies, Parklands, and Rocky Mountains.

Rhinanthus yellowrattle

Rhinanthus crista-galli L.

YELLOWRATTLE

Somewhat branching annual plants 30–60 cm high. Leaves opposite, linear to lanceolate, 2–5 cm long, with finely toothed margins. Flowers pale yellow, 6–10 mm long, borne either in upper leaf axils or in a one-sided leafy terminal spike; calyx hairy, greenish yellow, compressed, membranous, and at maturity inflated, enclosing a capsule. Not common; has been found at widely scattered locations in moist places and open woodlands; throughout Parklands, Boreal forest, and Rocky Mountains.

Scrophularia figwort

Scrophularia lanceolata Pursh

HARE FIGWORT

An erect plant 50–200 cm high, with stems and inflorescence somewhat glandular. Leaves ovate to lanceolate, coarsely toothed, 2–15 cm long, on short stalks borne oppositely on stem. Flowers greenish, 8–12 mm long, on stalks in a tall terminal inflorescence. Rare; has been found on moist ground; Prairies.



Fig. 191. Beardtongues: *A*, lilac-flowered beardtongue, *Penstemon gracilis* Nutt.; *B*, smooth blue beardtongue, *Penstemon nitidus* Dougl.; *C*, slender beardtongue, *Penstemon procerus* Dougl.

Verbascum mullein

- 1. Lower stem leaves long-petioled, ovate-cordate. *V. nigrum*
 Lower stem leaves sessile, ovate-lanceolate. 2
- 2. Leaves long-decurrent; filaments of upper stamens 3-4 times as long as the anthers. *V. thapsus*
 Leaves short-decurrent, ovate; filaments of upper stamens 1.5-2 times as long as the anthers. *V. phlomoides*

Verbascum nigrum L.

BLACK TORCH

Biennial plants 50-150 cm high, pubescent. Leaves stellate-pubescent below; lower leaves long-petioled, mostly cordate at the base; upper stem leaves sessile. Inflorescence a raceme; flowers 2-5 together, 15-20 mm across, light yellow; filaments purplish pubescent. Introduced; a rare roadside weed.

Verbascum phlomoides L.

WOOLLY MULLEIN

Biennial plants with stems 50-200 cm high, densely yellow woolly pubescent with branched hairs. Inflorescence spike-like; flowers 30-35 mm across, yellow; stigma club-shaped. Introduced; a rare garden and roadside weed.

Verbascum thapsus L.

COMMON MULLEIN

Biennial plants, tall, erect, woolly, 30-200 cm high. Stems very stout and straight, bearing large, densely woolly spatulate to elliptic leaves 10-30 cm long. Flowers yellow, almost regular, 5-lobed, 20-25 mm across, on a dense woolly spike 10-50 cm high; lower flowers usually opening first. Rare; has been found on railway grades and waste places; at widely separated locations throughout Prairies and Parklands. Its scarcity is rather surprising, because in British Columbia and in Eastern Canada this species is a common roadside weed, appearing to thrive best in dry, dusty, and exposed locations.

Veronica speedwell

Annual or perennial herbs usually associated with wet places. Leaves opposite. Flowers perfect, 4-lobed, rotate, slightly irregular, with 2 stamens. Fruit a several-seeded capsule.

- 1. Calyx 5-lobed, with the upper 2 long and the lower 3 short. *V. teucrium*
 Calyx 4-lobed. 2
- 2. Flowers solitary in axils of leaves; annuals. 3
 Flowers in axillary or terminal racemes; perennials. 6
- 3. Flower stalks shorter than or equaling leaves. 4
 Flower stalks as long as or longer than leaves. 5

4. Leaves narrowly oblong to oblong-lanceolate; capsule glabrous, veinless. *V. peregrina* var. *xalapensis*
 Leaves oblong-ovate to almost rotund; capsule reticulate-veined. *V. agrestis*
5. Flowers 10–15 mm across; capsule reticulate-veined. *V. persica*
 Flowers 4–6 mm across; capsule glandular pubescent. *V. polita*
6. Flowers in terminal racemes. 7
 Flowers in axillary racemes. 9
7. Racemes well-defined, dense; leaves 5–15 cm long, often in whorls. *V. longifolia*
 Racemes not well-defined, loose; leaves mostly less than 5 cm long. 8
8. Leaves all sessile; capsule longer than wide, barely notched. *V. alpina* var. *unalaschensis*
 Lower leaves petioled; capsule as long as wide, deeply notched. *V. serpyllifolia* var. *humifusa*
9. Leaves ovate to oblong, all with short stalks. 10
 Leaves linear to lanceolate, those of flowering shoots without stalks. 11
10. Leaves and stems glabrous; semiaquatic plants. *V. americana*
 Leaves glabrous or pubescent, and stem with 2 rows of hairs; a plant of gardens and waste areas. *V. chamaedrys*
11. Leaves broadly lanceolate; fruit not much flattened, with its stem less than twice as long as fruit. *V. comosa* var. *glaberrima*
 Leaves linear to linear-lanceolate; fruit much flattened, with its stem several times as long as fruit. *V. scutellata*

Veronica agrestis L.

PROSTRATE SPEEDWELL

Annual plants with prostrate or ascending stems 10–30 cm long. Leaves ovate to rotund, crenately serrate, rounded or truncate at the base, 1–2 cm long. Flowers 6–8 mm across, blue or white, with blue veins; pedicels 6–10 mm long, elongating in fruit. Introduced; a rare garden weed.

Veronica alpina L. var. *unalaschensis* C. & S.

ALPINE SPEEDWELL

Stems erect or decumbent at the base, 10–20 cm high, more or less villous throughout. Leaves all sessile, elliptic to oblong, 15–35 mm long, pilose on both sides. Inflorescence densely villous, elongating during and after flowering; flowers about 5 mm across, blue; capsules 5–8 mm long, about 3–5 mm wide, barely notched. Alpine meadows and moist areas; Rocky Mountains.

Veronica americana (Raf.) Schwein.

AMERICAN SPEEDWELL

An aquatic or semiaquatic, rather weak-stemmed plant 10–50 cm long. Leaves all stalked, oblong-lanceolate to ovate, 2–6 cm long. Flowers blue or white, 3–5 mm across, in long loose racemes in axils of leaves. Not common; found in streams and around springs; Boreal forest, Rocky Mountains, and Cypress Hills.

Veronica chamaedrys L.

GERMANDER SPEEDWELL

A sparsely pubescent perennial; stems prostrate, 20–40 cm long. Leaves sessile or nearly so, ovate to deltoid-ovate, 2–3 cm long, serrate. Racemes few, erect from the upper axils, loosely 10- to 20-flowered, 8–15 cm long; flowers about 1 cm across, blue, with darker lines; capsules rarely produced, 4–5 mm wide, shallowly notched. Introduced; a rare garden weed.

Veronica comosa Richt. var. *glaberrima* (Pennell) Boiv.

WATER SPEEDWELL

A branched species 20–70 cm long, with stalkless, often clasping, lanceolate leaves 2–7 cm long. Flowers blue, 8–10 mm across, in loose axillary racemes. Uncommon; has been found in wet places and along rivers; Boreal forest and Cypress Hills. Syn.: *V. catenata* Pennell; *V. connata* Raf. ssp. *glaberrima* Pennell.

Veronica longifolia L.

SPIKED SPEEDWELL

Perennial plants with erect stems 40–100 cm high. Leaves opposite or in whorls of 3, lanceolate or ovate-lanceolate. Racemes erect, spike-like; axis pubescent; flowers 8–10 mm across, blue. Introduced ornamental; occasionally escaped and established.

Veronica peregrina L. var. *xalapensis* (HBK.) St. John & Warren

HAIRY SPEEDWELL

A glandular hairy erect species 10–30 cm high. Leaves spatulate to linear, 10–25 mm long. Flowers whitish, about 3 mm across, borne singly in axils of leaves. Not common but locally abundant; on moist sandy soils and in low areas; across the Prairie Provinces. Syn.: *V. xalapensis* HBK.

Veronica persica Poir.

BIRD'S-EYE

A low-spreading, very branching, finely hairy annual. Leaves oval, short-stemmed, 10–25 mm long, coarsely toothed. Flowers blue, about 8–15 mm across, borne singly on long stalks from axils of leaves. Introduced; becoming increasingly abundant as a garden weed.

Veronica polita Fr.

SMOOTH SPEEDWELL

A species similar to *V. persica*, but with flowers at most 6 mm across and usually smaller. Introduced; a rare garden weed.

Veronica scutellata L.

MARSH SPEEDWELL

A species with decumbent or ascending stems 20–50 cm high. Leaves linear to linear-lanceolate, sharp-pointed, 2–7 cm long. Flowers about 6 mm across, blue, in long loose racemes in axils of leaves. Fairly common; in moist meadows and swamps and around springs; throughout the Prairie Provinces.

Veronica serpyllifolia L. var. *humifusa* (Dicks.) Vahl. THYME-LEAVED SPEEDWELL

A slender-stemmed perennial 5–20 cm high, much-branched from base. Leaves ovate to oblong or almost orbicular, 5–15 mm long. Flowers white or pale blue, 5–8 mm across, in short narrow leafy terminal racemes. An eastern species, very rare; has been found in low swampy ground; Cypress Hills.

Veronica teucrium L.

BROAD-LEAVED SPEEDWELL

An erect finely pubescent perennial 30–60 cm high. Leaves sessile or nearly so, ovate to lanceolate, 2–4 cm long, serrate. Racemes 2–4, from the upper axils, 5–10 cm long, on pedicels to 10 cm long, rather densely flowered; flowers about 12 mm across, blue. Introduced ornamental; occasionally escaped and established.

Veronicastrum Culver's-root

Veronicastrum virginicum (L.) Farwell

CULVER'S-ROOT

A coarse erect leafy-stemmed perennial 30–150 cm high, with opposite or whorled leaves. Leaves oblong to lanceolate, 7–15 cm long, varying from 2 to 7 at a node. Flowers white, with prominent protruding stamens 4–6 mm long, crowded on long narrow terminal racemes 6–20 cm long. An eastern species; has been found on roadsides and meadows; southeastern Parklands and Boreal forest.

LENTIBULARIACEAE—bladderwort family

Small plants found growing in water or wet locations, carnivorous or insectivorous, catching small insects and aquatic life by means of sticky leaves or submersed bladders. Leaves basal. Flowers irregular, perfect, with 2 stamens.

Calyx with 5 sepals; land plants; leaves entire,
basal; flowers solitary. *Pinguicula*

Calyx with 2 sepals; plants floating on or sub-
mersed in water; leaves dissected, bearing
small bladders; flowers several in a raceme
above water. *Utricularia*

Pinguicula butterwort

1. Stem villous below; flower 6–8 mm long. *P. villosa*

Stem glandular pubescent; flower more
than 17 mm long. 2

2. Lower corolla lip over 12 mm long; flower
dark purple with a straight spur. *P. macroceras*

Lower corolla lip less than 12 mm long;
flower pale purple with slightly
recurved spur. *P. vulgaris*

Pinguicula macroceras Willd.

WESTERN BUTTERWORT

A low bog plant with a basal rosette of sticky leaves. Flower single, 2–3 cm long, dark purple, on a stalk 5–10 cm high. Not common; found occasionally in bogs and swamps; southern Rocky Mountains.

Pinguicula villosa L.

SMALL BUTTERWORT

A low bog plant, similar to *P. macroceras*, but even smaller, 2–5 cm high; stem densely villous below, glandular puberulent above. Tundra and subarctic bogs; northern parts of Boreal forest.

Pinguicula vulgaris L.

BUTTERWORT

A low plant with leaves in a basal rosette. Leaves oval or elliptic, 2–3 cm long, usually somewhat rolled at edge, and having a sticky secretion. Flower single, 12–18 mm long, pale purple, on a stalk 3–10 cm high. Very rare; has been found in bogs and cold swampy lands; Boreal forest and Rocky Mountains.

***Utricularia* bladderwort**

- 1. Bladders minute; leaves thread-like, simple, not readily seen. *U. cornuta*
Bladders readily seen; leaves finely dissected. 2
- 2. Stems either submersed or floating; flowers 15–25 mm long. *U. vulgaris* var. *americana*
Stems creeping on bottom in shallow water; flowers 3–10 mm long. 3
- 3. Flower stalks ascending; flowers 10–15 mm long. *U. intermedia*
Flower stalks recurved; flowers 3–8 mm long. *U. minor*

Utricularia cornuta Michx.

HORNED BLADDERWORT

A delicate terrestrial plant; stems creeping underground. Leaves linear-filiform, with bladders in the margins. Scapes erect, 3–15 cm high, bearing 1–3 yellow flowers 15–25 mm long; spur 7–12 mm long. Rare; peaty or muddy shores and bogs; Boreal forest.

Utricularia intermedia Hayne

FLAT-LEAVED BLADDERWORT

An aquatic plant creeping on mud in shallow water, with somewhat floating branches 6–15 cm long. Leaves scattered and very finely dissected into thread-like segments; some branches leafless, bearing several bladders 3–10 mm long. Flowers yellow, with a large lower lip, about 10 mm long, in a 1- to 4-flowered raceme on a leafless upright stem. Found occasionally; in bogs and shallow water; Parkland, Boreal forest, and Cypress Hills.

Utricularia minor L.

LESSER BLADDERWORT

A small submersed plant with alternate leaves. Leaves having a few thread-like divisions, very minute, less than 6 mm long, bearing a few bladders

slightly longer than 1.5 mm. Flowers pale yellow, less than 8 mm long, on curved stalks on an upright flowering stem. Very rare; has been found in bogs and shallow water; Boreal forest.

Utricularia vulgaris L. var. *americana* Gray (Fig. 192) GREATER BLADDERWORT

A species often floating on the surface of the water, with leaves submerged. Stems 30–80 cm long, branched. Many much-divided leaves 2–5 cm long, bearing numerous bladders 3–5 mm long. Flowers yellow, 15–25 mm long, in a raceme of 6–15 flowers on a long, flowering stem 10–30 cm high. Fairly common; in lakes and sloughs; throughout the Prairie Provinces. Syn.: *U. macrorhiza* Le Conte.

OROBANCHACEAE—broom-rape family

Low pinkish herbs without green foliage, parasitic on roots of other plants (particularly *Artemisia*). Leaves scale-like. Flowers perfect and irregular; calyx 5-lobed; corolla 2-lipped; stamens 4. Fruit a many-seeded capsule.

Calyx deeply cleft on the lower side only;
glabrous throughout. *Conopholis*

Calyx nearly regular or cleft on both upper
and lower sides; glandular puberulent. *Orobanche*

Conopholis squawroot

Conopholis americana (L.) Wallr. CANCERROOT

Stems stout, erect, 5–20 cm high, pale brown or yellowish throughout, covered by ovate fleshy leaf scales to 2 cm long. Spike about half the height of the stem, 15–20 cm thick; corolla 10–15 mm long. Very rare; parasitic on shrubs and trees; southeastern Parklands and Boreal forest.

Orobanche broom-rape

1. Plants with a solitary flower; scale leaves
blunt. *O. uniflora*

Plants with more than one flower; scale
leaves acuminate. 2

2. Flowers borne singly on a naked stalk,
with no bracts beneath each flower. *O. fasciculata*

Flowers borne in a racemose cluster, with
1 or 2 long bracts beneath each flower. *O. ludoviciana*

Orobanche fasciculata Nutt. CLUSTERED BROOM-RAPE

A low pinkish-stemmed plant 3–10 cm high, parasitic on roots of *Artemisia* and other Compositae. Leaves scale-like. Flowers 1–10, purplish to yellow, about 25 mm long, each on a naked unbracted stalk. Common; wherever *Artemisia* is found; throughout Prairies. Syn.: *Anoplanthus fasciculatus* (Nutt.) Walp.




Fig. 192. Greater bladderwort, *Utricularia vulgaris* L. var. *americana* Gray.

A stout, somewhat sticky, glandular hairy, pink-stemmed plant 5–25 cm high, parasitic on roots of *Artemisia* and other Compositae. Leaves scale-like. Flowers purplish, 1–2 cm long, in a dense terminal spike, with 1 or 2 long bracts beneath each flower. Not so common as *O. fasciculata*; found on sandy soil; throughout Prairies. Syn.: *Myzorrhiza ludoviciana* (Nutt.) Rydb.

A small thin-stemmed plant; stem almost entirely underground, only 1–3 cm showing, bearing a few overlapping, glabrous, oblong-obovate scale leaves. Pedicels 1–4, usually 2, erect, 6–20 cm long, finely glandular pubescent, each bearing a single flower, about 2 cm long, white to violet. Rare; margins of woods and wooded slopes; southern Rocky Mountains and Cypress Hills.

PHRYMACEAE—lopseed family

Phryma lopseed

Perennial plants with stems 50–100 cm high, erect, simple or with a few branches. Petioles to 5 cm long in lower leaves, shorter in higher leaves, and absent or nearly absent in uppermost leaves; leaves ovate, 6–15 cm long, coarsely toothed. Inflorescence a spike-like raceme; flowers 6–8 mm long, pale purple to white, opposite, subtended by 3 small bracts; fruiting calyx to 1 cm long, reflexed-appressed. Rare; moist woods; southeastern Boreal forest.

PLANTAGINACEAE—plantain family

Plantago plantain

Plants with leaves usually having several prominent longitudinal ribs. Flowers borne in long narrow spikes, having a 4-lobed calyx, a 4-lobed corolla, and 2 or 4 stamens. Fruit a pyxis (capsule), with a conical top falling off at maturity and releasing the seeds.

- 1. Plants with stem leaves opposite or whorled. *P. psyllium*
- Plants with all leaves basal. 2
- 2. Leaves coarsely toothed to pinnately lobed. *P. coronopus*
- Leaves all entire, or distantly and shallowly toothed. 3
- 3. Leaves lanceolate to ovate. 4
- Leaves linear to filiform (thread-like). 8

- 4. Leaves lanceolate, gradually tapering to stalks. 5
 Leaves ovate, abruptly joining stalk. 7
- 5. Spikes of inflorescence long and narrow; plants of saline areas. *P. eriopoda*
 Spikes of inflorescence short, dense, and oblong; stamens protruding and very conspicuous. 6
- 6. Stem and leaves densely pubescent; flower bracts smaller than buds, acute. *P. canescens* var. *cylindrica*
 Stems and leaves not densely pubescent; flower bracts larger than buds, blunt. *P. lanceolata*
- 7. Leaves glabrous to sparsely pubescent, long-petioled; peduncle slightly longer than leaves; filaments white, inconspicuous. *P. major*
 Leaves short-pubescent, short-petioled; peduncle many times longer than leaves; filaments lilac, conspicuous. *P. media*
- 8. Plants densely white silky-woolly. *P. patagonica*
 Plants not densely woolly. 9
- 9. Flower spike densely long-pubescent. *P. aristata*
 Flower spike not long-pubescent. 10
- 10. Leaves thick, fleshy; flowers pubescent. *P. maritima*
 Leaves thread-like, not fleshy; flowers glabrous. *P. elongata*

Plantago aristata Michx.

BUCKHORN

An annual plant with scapes 15–25 cm high, more or less villous. Leaves linear or very narrowly lanceolate, 10–15 cm long, more or less pubescent. Spikes cylindric, 3–6 cm long; bracts linear, with the lowest ones to 2 cm long and the upper ones shorter, hirsute to long-villous. Introduced; a rare weed in disturbed soils; Prairies.

Plantago canescens Adams var. *cylindrica* (J. M. Macoun) Boiv.

WESTERN RIBGRASS

A perennial species with a woolly-pubescent scape about 15 cm high. Leaves narrowly lanceolate, 10–15 cm long, woolly-pubescent, especially below. Spike about 5 cm long, rather loose in fruit. Dry grasslands; southern Rocky Mountains.

Plantago coronopus L.

BUCKHORN PLANTAIN

A biennial or short-lived perennial; scapes 5–20 cm high. Leaves 3–15 cm long, with several pairs of linear lobes 5–20 mm long, pubescent. Spike 3–8 cm long, dense; flowers puberulent. Introduced; a rare weed; southeastern Parklands.

Plantago elongata Pursh

LINEAR-LEAVED PLANTAIN

An annual plant 6–15 cm high. Leaves very narrow, thread-like, 3–12 cm long, about 2 mm wide, single-ribbed. Inflorescence a loosely flowered very narrow spike 5–10 cm long. Not common; found occasionally in wet places and low flats; throughout Prairies.

Plantago eriopoda Torr.

SALINE PLANTAIN

A somewhat fleshy perennial species from a long coarse rootstock. Leaves oblanceolate, 5–20 cm long, with several longitudinal ribs borne on long stalks usually arising from a mass of long brown hairs at crown of root. Inflorescence a somewhat dense narrow spike 2–10 cm long, on a long stem. Common; on saline or alkaline soils, river flats, and slough margins; throughout Prairies and Parklands.

Plantago lanceolata L.

RIBGRASS

An introduced biennial or perennial species with a short rootstock. Leaves numerous, narrowly lanceolate, 3- to 5-ribbed, 5–30 cm long, tapering to very short stalks, usually some tufts of brownish hair at crown of root. Flowering stems much longer than leaves, 30–50 cm; inflorescence a short thick dense spike 2–8 cm long; stamens much protruding, often forming a conspicuous yellow or white ring around flower head. Found occasionally where imported lawn grass has been sown and as a weed in newly seeded lawns.

Plantago major L. (Fig. 193)

COMMON PLANTAIN

A perennial species from a short thick rootstock. Leaves numerous, very dark green, oval or ovate, 3–25 cm long, with many conspicuous longitudinal ribs contracting abruptly at the base into a long stalk. Flowering stems fairly long; inflorescence a dense narrow spike 7–30 cm long. Common; a weed in lawns, waste places, and yards; throughout the Prairie Provinces. Possibly introduced, but, if so, having spread across the country.

The var. *asiatica* (L.) Dcne., with tapering leaf bases and more upright leaves, has been found in Boreal forest.

Plantago maritima L.

SEASIDE PLANTAIN

A short-lived perennial with scapes 5–20 cm high. Leaves thick and fleshy, almost 3-angled in cross section, entire or somewhat distantly toothed. Spikes 2–10 cm long, often somewhat interrupted below, with the axis pubescent; sepals and corolla tube pubescent. Shores and saline areas; Boreal forest, rare in southern edge of Boreal forest and Parklands.

Plantago media L.

HOARY PLANTAIN

Perennial plants. Leaves elliptic to obovate, 10–20 cm long, narrowed at the base to a petiole. Scapes 20–40 cm high, somewhat pubescent; spikes at first narrowly conic, elongating to short-cylindric, 3–10 cm long. Introduced; a rare weed of disturbed soils and waste areas.



HERBARIUM OF THE MICHIGAN STATE UNIVERSITY
 WEST LANSING, MICH.

DATE: 1927

LOCALITY: Michigan

COLLECTOR: S. P. ...

NO.:

DATE: 1927

ELEVATION:

HABITAT:




Fig. 193. Common plantain, *Plantago major* L.

Plantago patagonica Jacq.

PURSH'S PLANTAIN

An annual plant, very pale green and whitish silky-woolly all over. Leaves 2–6 cm long, linear, with 1–3 nerves. Inflorescence a dense spike 2–12 cm long, on a stem 7–10 cm long, and very woolly. Found in sandy soil, on river flats, and dry soils; throughout southwestern part of the Prairie Provinces. An indicator of an overgrazed condition. Syn.: *P. purshii* R. & S.; *P. spinulosa* Dcne.

Plantago psyllium L.

SAND PLANTAIN

Annual plants with erect stems, 10–50 cm high, pubescent throughout. Leaves linear, 3–8 cm long. Heads 10–15 mm long, on peduncles 3–5 cm long arising from leaf axils; bracts broadly scarious-margined. Introduced; a rare weed of disturbed soils.

RUBIACEAE—madder family

Plants with opposite or whorled leaves, stipules, and perfect regular flowers. As many stamens as corolla lobes. Fruit a capsule, berry, or drupe.

- 1. Leaves opposite, in pairs; low plants. *Houstonia*
 Leaves apparently whorled; the stipules
 resembling the leaves; stems square. 2
- 2. Inflorescence of open cymes. *Galium*
 Inflorescence a head, subtended by
 bracts. *Asperula*

Asperula woodruff

Asperula arvensis L.

FIELD WOODRUFF

Annual plants, with stems 20–30 cm high, erect; the angles retrorsely scabrous. Leaves in whorls of 4–8, lanceolate or the lower ones obovate. Flowers subsessile in a terminal head, blue to purplish; bracts pubescent. Introduced; a very rare weed of disturbed areas.

Galium bedstraw

- 1. Ovary and fruit bristly or hairy. 2
 Ovary and fruit glabrous. 4
- 2. Principal stem leaves in whorls of 4. *G. boreale*
 Principal stem leaves in whorls of 6 or 8. 3
- 3. Leaves mostly in whorls of 8; plants
 annual. *G. aparine*
 Leaves mostly in whorls of 6; plants
 perennial. *G. triflorum*
- 4. Flowers yellow; leaves strongly revolute,
 in whorls of 6–12. *G. verum*
 Flowers white or greenish; leaves in
 whorls of 2–6. 5

5. Flowers in many-flowered cymes, short-pedicellate. *G. palustre*
 Flowers solitary or in few-flowered cymes, long-pedicellate. *G. trifidum*

Galium aparine L.

CLEAVERS

A trailing or decumbent annual with a square stem 30–100 cm long and covered with retrorse (backward-pointing) bristly hairs. Leaves in whorls of 6 or 8, oblong-linear to oblanceolate, sharp-tipped, rough bristly on margins and midrib, and 2–5 cm long. Long-stalked cream-colored flowers in axillary clusters of 4–9. Fruits in pairs, bearing hooked hairs. Found often; in moist woodlands and along riverbanks; a troublesome weed in Parklands. Syn.: *G. aparine* L. var. *vaillantii* (DC.) Koch; *G. vaillantii* DC.

Galium boreale L. (Fig. 194)

NORTHERN BEDSTRAW

An erect perennial 20–50 cm high, from thin brown rootstocks, with a slender square stem. Leaves in whorls of 4 around stem, linear to lanceolate, 3-ribbed, 2–6 cm long. Inflorescence a terminal leafy panicle, fairly dense, many-flowered; flowers white, about 3 mm across, with 4 corolla lobes, and faintly fragrant. Fruits in pairs, about 1.5 mm long, covered with short whitish hairs. Common; in openings in woodlands, along roadsides and moister places on prairies; almost throughout the Prairie Provinces. In some years, almost the dominant roadside flower. Most of the plants of this species found in the prairies are var. *intermedia* DC., with short curved hairs on fruit. Syn.: *G. septentrionale* R. & S.

Galium palustre L.

MARSH BEDSTRAW

A slender perennial with simple or branched stems 20–60 cm long, sparsely retrorse-scabrous on the angles. Leaves 5–15 mm long, linear to narrowly oblanceolate, commonly in whorls of 4 or 6. Inflorescences many-flowered, often forked; pedicels widely spreading; corolla white, about 4 mm wide. Rare; wet places, bogs, and marshes; Boreal forest.

Galium trifidum L.

SMALL BEDSTRAW

A slender-stemmed perennial with decumbent or erect weak stems 20–40 cm long. Leaves borne in whorls of 4, linear to spatulate, 6–12 mm long. Flowers terminal or axillary, on long stalks, greenish white, in 2's or 3's, very small, usually with 3 blunt corolla lobes. Fruits in pairs, smooth, without any hairiness. Found occasionally; in moist places; Parkland, Boreal forest, and Cypress Hills.

Galium triflorum Michx.

SWEET-SCENTED BEDSTRAW

A spreading, branched perennial 30–100 cm long, often trailing or decumbent, with an almost hairless stem. Leaves usually in whorls of 6, narrowly oval or oblanceolate, with a sharp, pointed tip, 2–6 cm long. Flowers long-stalked, greenish white, in clusters of 3, with 4 corolla lobes. Fruits in pairs, covered with long hooked hairs. Fairly common; in moist places and damp woodlands; throughout the Prairie Provinces.



Fig. 194. Northern bedstraw, *Galium boreale* L.

A perennial with a woody rootstock; stems erect, 30–80 cm high, finely pubescent. Leaves linear, 1–3 cm long, mostly in whorls of 8, often deflexed at maturity. Inflorescences several from the upper axils, many-flowered. Introduced; a very rare weed; mostly along roadsides; southeastern Parkland.

Houstonia bluets

Houstonia longifolia Gaertn.

LONG-LEAVED BLUETS

Low, tufted perennials 6–25 cm high, usually with a purplish square stem. Leaves opposite, linear to linear-oblong, 10–25 mm long, with small whitish or purplish stipules at base. Flowers pinkish or purple, 3–5 mm long, in very leafy terminal clusters. Fruit an ovoid capsule. Found occasionally; on grasslands and sandy soils; Parklands and Boreal forest.

CAPRIFOLIACEAE—honeysuckle family

Trees, shrubs, vines, or perennial herbs with opposite leaves without stipules. Flowers perfect; corolla varying from rotate or campanulate to urn-shaped; usually having 5 stamens, except for one genus, *Linnaea*, with only 4. Fruit a berry, drupe, or capsule.

- 1. Leaves pinnately compound. *Sambucus*
 Leaves simple, at most deeply lobed. 2
- 2. Low, trailing evergreen shrub; flowers
 with 4 stamens. *Linnaea*
 Shrubs or small trees; flowers with 5
 stamens. 3
- 3. Corolla rotate; inflorescence all terminal
 in compound cymes; fruit a drupe con-
 taining a flattened stone. *Viburnum*
 Corolla tubular or bell-shaped;
 inflorescence axillary as well as termi-
 nal; fruit a capsule or berry with 2 or
 more seeds. 4
- 4. Leaves with finely toothed margins; calyx
 with 5 linear lobes; fruit a capsule. *Diervilla*
 Leaves with entire or wavy margins; calyx
 with short lobes; fruit a berry. 5
- 5. Flowers regular, bell-shaped, in clusters;
 fruit a white berry. *Symphoricarpos*
 Flowers often slightly irregular, tubular,
 usually in pairs; fruit a red or black
 berry. *Lonicera*

Diervilla bush-honeysuckle

Diervilla lonicera Mill.

BUSH-HONEYSUCKLE

A shrub 50–100 cm high, with opposite simple leaves. Leaves short-stalked, ovate to oval, finely toothed on margins, 5–12 cm long. Flowers narrow, funnel-shaped, about 2 cm long, yellow, in axillary and terminal clusters of 1–5. Fruit a slender capsule with linear stamens remaining on the end. Not common; found in rocky woodlands; along the eastern Boreal forest.

Linnaea twinflower

Linnaea borealis L. var. *americana* (Forbes) Rehder (Fig. 195A) TWINFLOWER

A very low, creeping or trailing evergreen plant 15–75 cm long. Leaves opposite, short-stalked, oval or orbicular, 10–15 mm across, with a somewhat wavy margin. Flowers pink, fragrant, funnellform, about 1 cm long, with only 4 stamens, in pairs, pendent or hanging downward from the top of a stem 3–10 cm high. Common; in cool woodlands and often found on decayed, fallen tree trunks; throughout the entire wooded parts of the Prairie Provinces. Syn.: *L. americana* Forbes.

Lonicera honeysuckle

1. Somewhat twining shrub; inflorescence a terminal cluster. 2
Erect shrubs; leaves not connate-perfoliate; inflorescence axillary and flowers in pairs. 3
2. Uppermost connate leaves glaucous, obtuse or acute. *L. dioica* var. *glaucescens*
Uppermost connate leaves green, pointed at tip. *L. hirsuta* var. *schindleri*
3. Bracts below flowers large, broad, and leaf-like; berries of the 2 flowers separate. *L. involucrata*
Bracts below flowers thin or minute; berries of the 2 flowers more or less united. 4
4. Ovaries completely fused, ripening into a single bluish black berry. *L. caerulea* var. *villosa*
Ovaries only partly united, with berries distinguishable. 5
5. Pith of branches brown, not filling the entire core. *L. tatarica*
Pith of branches white, filling the entire core. 6
6. Leaves densely puberulent, at least below. *L. oblongifolia*
Leaves subglabrous to lightly pilose below. *L. utahensis*




Fig. 195. A, Twinflower, *Linnaea borealis* L. var. *americana* (Forbes) Rehder; B, twining honeysuckle, *Lonicera dioica* L. var. *glaucescens* (Rydb.) Butt.

Lonicera caerulea L. var. *villosa* (Michx.) T. & G. BLUE FLY HONEYSUCKLE

Low erect shrubs 30–100 cm high; twigs sometimes finely hairy. Leaves oval to obovate, slightly downy when young, very short-stalked, 2–4 cm long. Flowers in pairs in axils of leaves, yellow, about 12 mm long. Fruit an oblong or globose bluish black berry, covered with a bloom, edible. Fairly common in moist or swampy places in woodlands; Boreal forest and Rocky Mountains. Syn.: *L. villosa* (Michx.) R. & S. var. *solonis* (Eat.) Fern.

Lonicera dioica L. var. *glaucescens* (Rydb.) Butt. (Fig. 195B) TWINING HONEYSUCKLE

A more or less twining shrub with light-colored shreddy bark. Leaves obovate or oval, all opposite, often connate (joined at bases) and perfoliate, especially the upper ones, 5–8 cm long, pale, and often hairy below, particularly on veins. Flowers in a terminal cluster, yellow, later turning reddish, 20–25 mm long. Fruits in clusters of red berries. Fairly common; in woodlands; throughout the Prairie Provinces. Syn.: *L. glaucescens* Rydb.

Lonicera hirsuta Eat. var. *schindleri* Boiv. HAIRY HONEYSUCKLE

Similar to *L. dioica*, but with leaves dull green, and, except for the upper ones, short-petioled; young stems glandular pubescent. Rare; moist woods; southeastern Boreal forest.

Lonicera involucrata (Richards.) Banks INVOLUCRATE HONEYSUCKLE

A shrub 1–3 m high, sometimes with downy stems. Leaves oblong to oval, 5–10 cm long, somewhat glandular-dotted and downy below, with a prominent midrib. Flowers in pairs, yellow, with large leaf-like bracts at bases. Fruits 2 large berries, very dark purple to black, about 8 mm across, surrounded by persistent bracts. Fairly common; in woodlands; Boreal forest and Rocky Mountains. Syn.: *Distegia involucrata* (Richards.) Cockerell.

Lonicera oblongifolia (Goldie) Hook. SWAMP FLY HONEYSUCKLE

A shrub 50–150 cm high, with ascending branches. Leaves oblong, downy below when young, 2–5 cm long. Flowers yellow with a purplish tinge inside, 10–15 mm long, 2-lipped, in pairs on a long stem. Fruits in pairs of more or less united purplish red berries. Found occasionally; in swamps and marshy lands; eastern Boreal forest. Syn.: *Xylosteon oblongifolium* Goldie.

Lonicera tatarica L. TARTARIAN HONEYSUCKLE

A shrub 1–3 m high, with thin, ovate, somewhat cordate-based leaves 2–8 cm long. Flowers pink or white, about 15 mm long, in pairs, very numerous. Fruit an orange or yellow berry about 6 mm in diam. An introduced species, commonly used for ornamental planting and hedges. Has become established in woodlands and moist spots; in several locations in the Prairie Provinces. Syn.: *Xylosteon tataricum* (L.) Medic.

Lonicera utahensis Wats. UTAH HONEYSUCKLE

A low shrub 60–150 cm high; leaves oblong to oblong-ovate, 2–5 cm long, entire, thin glabrous or sparsely pubescent below, obtuse to blunt at the base. Flowers in pairs; peduncle slender, 10–15 mm long; bracts 2, very narrow, 1–3

mm long; corolla 15–20 mm long. Berries red. Moist, coniferous woods; southern Rocky Mountains.

Sambucus elder

Sambucus racemosa L.

RED ELDERBERRY

Shrubs 1–6 m high, with 3–7 leaflets and yellowish brown pith. Inflorescence ovate in outline, 5–8 cm long, 3–5 cm wide; flowers somewhat greenish white to yellowish.

In var. *pubens* (Michx.) Wats., twigs and leaflets pubescent, inflorescence rather oblong-ovate, berries small, and stems rarely more than 3 m high. Moist open woods; eastern Boreal forest, rarely in eastern Parklands.

In var. *arborescens* (T. & G.) Gray, leaves and twigs subglabrous, inflorescence more compact and broadly ovate, berries large and deep red, and stems to 6 m high. Moist woods; southern Rocky Mountains.

Symphoricarpos snowberry

Low shrubs with strongly creeping roots. Leaves opposite, simple, short-stalked. Flowers perfect, small, pink and white, campanulate (bell-shaped). Fruit a small, round, 2-seeded, white berry.

Stamens and styles not protruding from corolla. *S. albus*
Stamens and styles protruding from corolla. *S. occidentalis*

Symphoricarpos albus (L.) Blake

SNOWBERRY

An erect shrub 50–150 cm high. Leaves thin, oval 1–5 cm long, sometimes slightly toothed. Flowers in several-flowered clusters. Berry waxy, white, 6–10 mm across. Found occasionally; eastern Parklands and Boreal forest. The var. *pauciflorus* (Robbins) Blake, few-flowered snowberry, a low-growing spreading shrub 10–50 cm high. Leaves oval or round, 1–3 cm long, softly hairy beneath. Flowers borne singly in upper leaf axils or in terminal clusters of 2 or 3, about 6 mm long. Berries white, waxy, about 6 mm in diam. This variety is more common than the species; in rocky and sandy woodlands; throughout bush areas of the Prairie Provinces and in Cypress Hills. Syn.: *S. pauciflorus* (Robbins) Britt.

Symphoricarpos occidentalis Hook.

WESTERN SNOWBERRY, BUCKBRUSH

A shrub 50–100 cm high, growing from creeping roots. Leaves oval, ovate, or almost round, somewhat softly hairy beneath, 2–6 cm long. Flowers pink and white, with styles and stamens conspicuously projecting from corolla, in rather dense terminal and axillary spikes. Fruits snow white waxy berries, often in large numbers. One of the most widespread and commonest shrubs; in dense stands on open prairie, in ravines, coulees, and woodlands; throughout the Prairie Provinces.

Viburnum bush-cranberry

Shrubs or small trees with opposite leaves and flowers all in terminal clusters. Corolla rotate or wheel-shaped to short bell-shaped, regular, 5-lobed; flowers perfect, with 5 stamens and a short 3-cleft style. Fruits single-seeded edible drupes; stone somewhat compressed or flattened.

- 1. Leaves usually 3-lobed, with veins radiating from base. 2
- Leaves not lobed, with veins pinnate from midrib. 3
- 2. Outer flowers of cluster large and neutral (sexless), with large lobes. *V. opulus* var. *americanum*
- None of flowers large and neutral. *V. edule*
- 3. Leaves densely hairy beneath, with stalks not more than 6 mm long. *V. rafinesquianum*
- Leaves not hairy beneath, with slender stalks more than 6 mm long. *V. lentago*

Viburnum edule (Michx.) Raf. (Fig. 196) LOW BUSH-CRANBERRY

A rather straggly shrub 50–200 cm high. Leaves usually shallowly 3-lobed, 4–10 cm across; leaf bases usually flat or cordate. Inflorescence few-flowered, 1–3 cm across; flowers all perfect. Fruit a red drupe about 3 mm long. Fairly common; in rich moist woodlands; throughout the heavily wooded areas of the Prairie Provinces. Syn.: *V. eradiatum* (Oakes) House.

Viburnum lentago L. NANNYBERRY

A tall shrub or small tree up to 6 m high. Leaves 4–10 cm long, ovate, with sharp, small, marginal teeth tapering to a point at apex. Flowers in large clusters 5–12 cm across. Fruit an edible bluish black berry 8–12 mm long, covered with a bloom. Common; in woodlands; southeastern Boreal forest.

Viburnum opulus L. var. *americanum* (Mill.) Ait. HIGH BUSH-CRANBERRY

A shrub 1–4 m high, with smooth branches. Leaves palmately veined, broad, 3-lobed, 5–10 cm across. Outer flowers in clusters 10–15 mm across, with 5 large petals, but neutral (without fully formed styles or stamens); inner flowers smaller, creamy white, and perfect. Fruit a red berry, about 1 cm in diam, very acid. Fairly common; in woodlands throughout heavily wooded areas. Syn.: *V. trilobum* Marsh.

Viburnum rafinesquianum Schultes DOWNY ARROWWOOD

A shrub 50–150 cm high, with many slender gray branches. Leaves coarsely toothed, oval or ovate, slightly cordate at base, 2–7 cm long; under-surface densely velvet-hairy. Inflorescence rounded, 2–7 cm across; flowers all perfect, white. Fruits in large clusters, almost black, about 6 mm long; stone grooved slightly on both sides. Fairly common; in woodlands; eastern Boreal forest. Syn.: *V. affine* Bush var. *hypomalacum* Blake.



HERBARIUM OF THE UNIVERSITY OF MANITOBA
 SWIFT CURRENT, MAN.

SPECIES: *Viburnum edule* (Michx.) Raf.

LOCATION: Sattler Lake

COLL: J. Logan

DET:

COLL NO: 15085

DATE: 30 July 1970

ELEVATION: 100 m

HABITAT: woods along lake




Fig. 196. Low bush-cranberry, *Viburnum edule* (Michx.) Raf.

ADOXACEAE—moschatel family

Adoxa moschatel

Adoxa moschatellina L.

MOSCHATEL

A dwarf perennial plant from a scaly rootstock, 10–20 cm high, with several long-stalked basal leaves divided into 3 long-stalked, somewhat 3-cleft leaflets. Flowers very small, in a cluster about 6 mm across, greenish or yellowish, at head of slender stem bearing 2 leaves. Found occasionally; in cool mossy woodlands; along margins of Boreal forest.

CUCURBITACEAE—gourd family

Echinocystis balsam-apple

Echinocystis lobata (Michx.) T. & G.

WILD CUCUMBER

An annual, succulent, trailing vine 3–6 m long, climbing by means of tendrils over bushes and shrubbery. Stem somewhat angled, bearing long, spirally twisted tendrils. Leaves thin, pale green, rough on both sides, palmately veined, with 3–7 large lobes 5–12 cm across. Flowers greenish white, unisexual; male flowers in panicles or racemes; female flowers 1 or 2, in axils of leaves. Fruit a pepo (a large, ovoid, fleshy berry with a thick skin), pale green, covered with weak spines. Fruits 4–10 cm long, containing several large, flat, roughened, almost black seeds. Fairly common; in moist locations among bushes; southeastern Parklands and Boreal forest. Syn.: *Micrampelis lobata* (Michx.) Greene.

CAMPANULACEAE—bluebell family

Campanula bellflower

Perennial plants with rootstocks and alternate simple leaves. Flowers perfect, usually blue, campanulate or funnelform, in panicles; stamens 5; stigmas 3–5. Fruit an ovoid capsule containing many seeds.

1. Stem simple, bearing a solitary terminal flower. 2
- Stem usually bearing many flowers. 3
2. Leaves sharply dentate; corolla 15–30 mm long. *C. lasiocarpa*
- Leaves entire or glandular denticulate; corolla 10 mm long. *C. uniflora*
3. Flowers in part crowded into a head. *C. glomerata*
- Flowers not crowded. 4
4. Stem leaves broad, ovate to ovate-lanceolate. *C. rapunculoides*
- Stem leaves narrow, linear to filiform. 5

5. Basal leaves ovate or cordate, soon withering; flowers more than 12 mm long. *C. rotundifolia*
 Basal leaves linear or narrowly lanceolate; flowers not over 12 mm long. *C. aparinoides*

Campanula aparinoides Pursh

MARSH BELLFLOWER

A very thin slender-stemmed plant 20–50 cm high. Stem 3-angled, somewhat roughened, with downward-pointing stiff short hairs. Leaves lanceolate to linear-lanceolate, 2–5 cm long. Flowers at ends of branches, usually white or with a faint bluish tinge, 10–15 mm long. Found often; in wet meadows and marshes; eastern Parklands and Boreal forest. Includes *C. uliginosa* Rydb.

Campanula glomerata L.

CLUSTERED BLUEBELL

An erect perennial with a short rootstock and glabrous to softly pubescent stems 30–70 cm high. Leaves 2–5 cm long, oblong-lanceolate; lower leaves petiolate; upper ones sessile to more or less clasping. Flowers 2–3 cm long, violet blue, sessile in a terminal head subtended by leafy bracts. Introduced ornamental; locally escaped and established.

Campanula lasiocarpa Cham.

ALASKA BLUEBELL

Small perennial plants 4–15 cm high, from a creeping rootstock. Leaves 2–7 cm long, spatulate to narrowly oblanceolate; lower leaves petioled; upper ones sessile, with margin laciniate-denticulate. Flowers solitary and terminal; calyx long villous; lobes lancinate. Mountain meadows; southern Rocky Mountains.

Campanula rapunculoides L.

CREeping BLUEBELL

Erect perennial plants with a creeping rootstock; stems 40–100 cm high, usually unbranched. Leaves coarse, serrate, usually sparsely pubescent below. Lower leaves petioled; upper ones sessile or subsessile, 3–7 cm long. Inflorescence one-sided; flowers 2–3 cm long, blue. Introduced ornamental; locally escaped and established.

Campanula rotundifolia L. (Fig. 197)


HAREBELL

A perennial growing from rootstocks, to 10–50 cm high, often with many stems. Basal leaves soon disappearing, ovate or deeply cordate-based, 10–25 mm long, on long stalks; stem leaves linear or linear-oblong, 1–5 cm long. Flowers blue, campanulate, 15–25 mm long, occasionally solitary, but more often in a raceme of 3 or 4 flowers. Common; throughout the Prairie Provinces.

Campanula uniflora L.

ALPINE BLUEBELL

A dwarf perennial 10–20 cm high, with a single flower terminating the stem. Leaves and calyx lobes entire or finely glandular denticulate. Arctic and alpine meadows; northeastern Boreal forest and Rocky Mountains.



A.C. Budd

Fig. 197. Harebell, *Campanula rotundifolia* L.

LOBELIACEAE—lobelia family

Annual or perennial herbs often with a bitter, milky sap. Leaves alternate or basal. Flowers perfect, irregular, with a 2-lipped corolla split to the base on one side in one genus; stamens 5. Fruit a many-seeded capsule.

Annual plants; corolla not split; fruit 1-celled. *Downingia*
Perennial plants; corolla split; fruit 2-celled. *Lobelia*

Downingia downingia

Downingia laeta Greene

DOWNINGIA

Low-growing annual herbs 5–15 cm high. Leaves alternate, oblong, to 15 mm long. Flowers in axils of upper leaves or bracts, about 1 cm long, light blue with darker blue veins and with a whitish throat spotted with yellow. Very scarce; has been found in moist and saline places; southwestern Prairies. Syn.: *Bolelia laeta* Greene.

Lobelia lobelia

1. Plants aquatic, with leaves all submerged. *L. dortmanna*
Plants terrestrial. 2
2. Stem leaves linear; plants of moist marshy places; racemes few-flowered. *L. kalmii*
Stem leaves lanceolate to ovate; plants of drier soils; racemes many-flowered. *L. spicata* var. *hirtella*

Lobelia dortmanna L.

WATER LOBELIA

A perennial plant with a submerged rosette of hollow basal leaves 2–9 cm long and fleshy, linear. Stem 40–70 cm high, hollow, with minute filiform stem leaves. Raceme emersed, few-flowered; flowers usually about 15 mm long, pale blue to white. Shallow lakes; Boreal forest.

Lobelia kalmii L.

KALM'S LOBELIA

A leafy-stemmed biennial or perennial branching plant 10–30 cm high. Lower leaves spatulate, 10–25 mm long; upper leaves linear. Flowers light blue, about 1 cm long, in loose racemes; each flower stalk usually bearing 2 small bracts or 2 glands. Fairly common; in bogs and wet meadows; throughout Boreal forest, rare in Parklands. Syn.: *L. strictiflora* (Rydb.) Lunell.

Lobelia spicata Lam. var. *hirtella* A. Gray

SPIKED LOBELIA

A perennial with a simple erect stem 30–100 cm high. Lower leaves spatulate, 2–10 cm long; stem leaves lanceolate to spatulate. Flowers on a long, spike-like terminal inflorescence, pale blue, about 1 cm long. In the type, sepals and bracts slightly or not at all hairy. Found occasionally; in dry sandy soil; throughout eastern Parklands and Boreal forest.

VALERIANACEAE—valerian family

Valeriana valerian

Perennial herbs with scented roots. Leaves opposite, without stipules, varying; lower leaves often entire and spatulate; upper ones pinnately lobed with a large terminal lobe. Flowers usually rose or white, 5-lobed, small, in many-flowered clusters, lengthening and becoming paniculate as the plant matures. Seed an achene.

Stem leaves with 9–15 segments; basal leaves
oblong to spatulate, petioled. *V. dioica* var. *sylvatica*

Stem leaves with 3–7 segments; basal leaves, if
present, pinnate. *V. sitchensis*

Valeriana dioica L. var. *sylvatica* (Sol.) Gray (Fig. 198) NORTHERN VALERIAN

An erect rather weak-stemmed plant 20–60 cm high. Basal leaves entire, usually spatulate; stem leaves pinnate, with a large terminal lobe pinnately veined. Flowers white, about 3 mm across, in dense terminal clusters, later lengthening into a short cymose panicle. Fairly common; in wet places; throughout Boreal forest. Syn.: *V. septentrionalis* Rydb.

Valeriana sitchensis Bong. MOUNTAIN HELIOTROPE

A robust species with a stout rootstock; stems 40–100 cm high, glabrous or nearly so. Stem leaves 10–15 cm long, 3- to 5-pinnate; leaflets 3–5 cm long, with those of the lower leaves ovate and those of the upper ones lanceolate and coarsely sinuate-dentate; basal leaves similar to stem leaves or simple, usually absent at flowering time, sparingly pilose to glabrous. Inflorescence in dense terminal clusters; corolla 6–8 mm long, white or pinkish. In var. *scouleri* (Rydb.) M. E. Jones, basal leaves larger than stem leaves, leaflets less deeply toothed, and corolla 5–6 mm long. Both varieties in mountain meadows and moist woods; southern Rocky Mountains.

DIPSACACEAE—teasel family

Knautia bluebuttons

Knautia arvensis (L.) Duby FIELD SCABIOUS

An erect hairy perennial 30–100 cm high. Lower leaves stalked, oblong-lanceolate, 9–25 cm long; upper stem leaves opposite, pinnatifid or lobed, without stalks. Flowers pale lilac or blue, in heads 2–4 cm broad, with involucre bracts resembling heads of Compositae. Introduced from Europe; found occasionally; western Alberta and near Winnipeg. Syn.: *Scabiosa arvensis* L.

(Fig. 198 overleaf)



Fig. 198. Northern valerian, *Valeriana dioica* L. var. *sylvatica* (Sol.) Gray.

COMPOSITAE—composite family

Compositae is a very large and diverse family (Fig. 199). The flower head is composed of many florets or small flowers borne on a common receptacle, which is surrounded by an involucre composed of one or more rows of bracts. There are 2 main types of floret: tubular (or disk) and ray (or ligulate). Tubular florets are regular and tube-shaped; a flower head entirely composed of this type of floret is called discoid. Ray florets are irregular in shape and have a single strap-like petal; a head entirely composed of this type of floret is called ligulate. When both forms of floret are present, the flower head is called radiate. In such cases, the tubular florets occupy the center (or disk) and the ray florets form the margins. In ligulate and discoid flowers the florets are all perfect (having both pistil and stamens). In radiate flowers the tubular florets are usually perfect and the ray florets are either female (pistil only) or neutral (without either pistil or stamens). Often there are bracts or scales, which are called the chaff, among the florets on the top of the receptacle. If there are no bracts, the receptacle is said to be naked. There are 5 stamens, usually with their anthers united to form a ring around the pistil. In some genera the flowers are unisexual, the male and female florets being borne in separate flowers. The fruit is an achene, sometimes enclosed in a bur-like closed involucre, sometimes provided with a pappus (or tuft of hairs) to aid in dissemination, and sometimes having awns or scales. In one group all the species contain latex, a sticky, often milky sap. In this publication, for convenience in making identifications, this large family has been divided into three groups.

1. Stamens not united to form a tube around the pistil. Group 1, p. 683
Stamens united to form a tube around the pistil. 2
2. Some or all of the florets tubular. Group 3, p. 703
None of the florets tubular, heads ligulate; plants with a milky or sticky sap. Group 2, p. 689

GROUP 1

The plants in this group are annual or perennial herbs, usually with alternate leaves. There are no perfect florets, and the flowers are inconspicuous. In one genus, both male and female florets are borne on the same head, but in the others they are borne on separate heads in different parts of the plant. Heads composed entirely of male florets and those with both male and female florets hang downward; this trait is helpful in recognizing this group. When the male and female florets are borne in separate heads, the fruit is often enclosed in a bur-like or nut-like involucre. Pollen from flowers of this group is often responsible for causing hay fever.

1. Both male and female florets on one head; achenes not contained in bur-like involucre. *Iva*
Male and female florets on separate heads; involucre of female flowers bur-like or nut-like. 2




Fig. 199. Characteristics of composite flowers and fruit.

- 2. Leaves not lobed or dissected; female heads with 2 florets; involucre forming an oblong bur with hooked prickles. *Xanthium*
 Leaves lobed or divided; female heads usually with one floret; prickles on involucre not hooked. 3
- 3. Involucre of female heads with a single series of prickles or tubercles, and with spines short. *Ambrosia*
 Involucre of female heads with several rows of spines, and with spines long and prominent. *Franseria*

***Ambrosia* ragweed**

- 1. Leaves all opposite, some entire, some with 3–5 lobes. *A. trifida*
 Leaves opposite or alternate, all once- or twice-divided. 2
- 2. Annuals; leaves with stalks; fruit with sharp spines or tubercles. *A. artemisiifolia* var. *elatior*
 Perennials with running rootstocks; leaves without stalks; fruit without sharp spines. *A. psilostachya* var. *coronopifolia*

Ambrosia artemisiifolia L. var. *elatior* (L.) Descourtils COMMON RAGWEED

An erect annual 30–90 cm high. Leaves much-divided, stalked, somewhat grayish hairy below. Many male heads borne in long narrow racemes; fewer female heads borne in small clusters. Common; a weed of roadsides and waste places; in eastern Parklands, but scarcer farther west. Syn.: *A. elatior* L.

Ambrosia psilostachya DC. var. *coronopifolia* (T. & G.) Farwell
PERENNIAL RAGWEED

A grayish hairy erect perennial 30–90 cm high, from running rootstocks. Leaves once- or twice-divided, without stalks. Inflorescence similar to *A. artemisiifolia*. Fruit almost devoid of spines. Fairly common; along roadsides and waste places; eastern Parklands, but scarcer farther west. Syn.: *A. coronopifolia* T. & G.

Ambrosia trifida L. GREAT RAGWEED

A rather stout-stemmed erect annual 60–150 cm high, with a rough hairy stem. Leaves all opposite, 5–25 cm across, stalked, mostly 3- to 5-lobed, with 3 main palmate veins. Male flowers in long terminal racemes; female flowers in clusters in axils of upper bract-like leaves. Common; along roadsides and in waste places and fields; eastern Parklands, but scarce elsewhere.

***Franseria* bur-ragweed**

Franseria acanthicarpa (Hook.) Coville BUR-RAGWEED

An annual, usually rather decumbent and spreading plant 15–60 cm high, much-branched. Leaves doubly divided, 5–10 cm long. Male flowers in long

terminal racemes; female flowers clustered in leaf axils; involucre with long straight spines making the plant very prickly. May be plentiful locally; only found in sand dunes; Prairies.

Iva marsh-elder

Low-growing perennials with creeping woody roots; flowers solitary on stalks in leaf axils. *I. axillaris*

Tall-growing coarse annuals with fibrous roots; flowers in terminal and axillary spike-like panicles. *I. xanthifolia*

Iva axillaris Pursh

POVERTYWEED

An erect herbaceous perennial 10–50 cm high, with woody running roots. Leaves stalkless, entire, 3-nerved, obovate to linear-oblong, 1–4 cm long; lower leaves opposite; upper ones alternate. Flowers small, yellow, reflexed in leaf axils. A serious weed in many farming areas where soil conditions favor its growth. Very plentiful locally; in heavy somewhat saline soils; Prairies.

Iva xanthifolia Nutt. (Fig. 200)

FALSE RAGWEED

A tall, erect, branching annual with a rough downy stem 0.5–2 m high. Leaves broadly ovate, rough above and downy below, long-stalked, 5–15 cm long. Heads small, inconspicuous, crowded on terminal and axillary panicles. Plant looks like a sunflower except for its inflorescence. Very common; in vacant lots and waste places in towns and settlements; throughout the Prairie Provinces, a common weed along roadsides and fields in Prairies and Parklands. Syn.: *Cyclachaena xanthifolia* (Nutt.) Fresn.

Xanthium cocklebur

Annual herbs with alternate leaves. Male flowers in terminal spikes or racemes; female flowers in axils of leaves. Fruit a bur with 2 stout beaks at one end, covered with hooked prickles, and containing 2 long flat seeds; one seed germinating rapidly, but the other having a long-delayed germinating period.

Xanthium strumarium L. (Fig. 201)

COCKLEBUR

An annual herb, rather decumbent, coarse-stemmed, much-branched, 15–60 cm high. Leaves roughly ovate, with wavy or slightly lobed margins, 2–8 cm long. Male flowers clustered at ends of branches; female flowers in clusters below; involucre of female flower closed, ending in 2 beaks; involucre forming a bur over fruit. Fruit with 2 terminal beaks, and covered with stout hairy hooked prickles. Common; around slough margins and low places, especially somewhat saline areas; throughout entire southern portion of the Prairie Provinces. Cocklebur seedlings considered **poisonous** to swine. Authorities disagree on the identification of species of *Xanthium*: *X. echinatum* Murr., *X. commune* Britt., *X. italicum* Moretti, *X. macounii* Britt., and *X. glanduliferum* Greene have been reported for Western Canada, but the identification is doubtful.




Fig. 200. False ragweed, *Iva xanthifolia* Nutt.




Fig. 201. Cocklebur, *Xanthium strumarium* L.

GROUP 2

The plants in this group have a ligulate (strap-like) irregular corolla in each of the florets. All the florets are perfect (with both pistils and stamens), and the anthers of the stamens are united in a ring around the pistil. The leaves are either basal or alternate, and all the plants contain a milky sap. The primary leaves, those appearing after the cotyledons or seed leaves of seedlings, are a single leaf instead of a pair of equal leaves.

1. Plants with leaves all basal or with a few reduced leaves on the stem. 2
 Plants with normal stem leaves. 7
2. Leaves tomentose ciliate; achene about 8 mm long. *Microseris*
 Leaves glabrous, or if pubescent, not tomentose. 3
3. Flower heads solitary on a long stalk. 4
 Heads several on a flowering stalk. 5
4. Bracts of the involucre imbricated (overlapping like shingles). *Agoseris*
 Main bracts of the involucre equal, with shorter spreading bracts at their base. *Taraxacum*
5. Pappus hairs plumose (with fine hairs on either side like a feather); bracts of involucre few. *Hypochoeris*
 Pappus hairs simple, not plumose. 6
6. Bracts in a single series, all equal with their midribs thickened. *Crepis*
 Bracts in series of 1-3, with their midribs not thickened. *Hieracium*
7. Pappus not hairy, merely covered with scales. 8
 Pappus hairy. 9
8. Flowers blue. *Cichorium*
 Flowers yellow. *Lapsana*
9. Pappus bristles plumose (branched). 10
 Pappus bristles simple. 13
10. All leaves long, narrow, grass-like; heads solitary on the stem. *Tragopogon*
 Not all leaves grass-like; more than one head on the stem. 11
11. Ligules pink. *Stephanomeria*
 Ligules yellow to red. 12
12. Stem leaves few, small, mostly basal. *Hypochoeris*
 Stem leaves many, reaching to the inflorescence. *Picris*

13. Pappus double, the outer ring consisting of 5 small scale-like appendages, the inner ring bristles. *Krigia*
 Pappus simple, consisting of bristles only. 14
14. Leaves narrowly linear or scale-like; flowers pink. *Lygodesmia*
 Leaves broader and leaf-like; flowers yellow, white, or purplish. 15
15. Seeds flattened; pappus usually white. 16
 Seeds not flattened; pappus tawny or white. 17
16. Seeds with a long beak at the apex. *Lactuca*
 Seeds without a long beak at the apex. *Sonchus*
17. Stems leafy at the base only; leaves mostly basal. *Microseris*
 Stems leafy throughout or at least to midway up the stem. 18
18. Pappus white; involucre of one series of equal bracts. *Crepis*
 Pappus brownish or tawny; involucre usually of unequal bracts. 19
19. Heads borne erect; leaf margins entire or merely toothed; bracts of involucre narrow and green. *Hieracium*
 Heads usually nodding; leaves divided or lobed; bracts of involucre broad and colored. *Prenanthes*

Agoseris false dandelion

Perennial herbs from a taproot, with leaves all basal and usually long and narrow. Large yellow flower heads, about 2 cm across, borne singly on a long stalk or scape. Pappus composed of white hairs.

1. Achene beaked, with the beak 2–4 times as long as the body; outer bracts much broader than the inner ones. *A. grandiflora*
 Achenes with the beak less than twice as long as the body. 2
2. Ligules yellow; beak half as long as the body or less. *A. glauca*
 Ligules orange; beak longer than the body. *A. aurantiaca*

***Agoseris aurantiaca* (Hook.) Greene**

ORANGE FALSE DANDELION

A species with narrowly linear leaves 5–35 cm long, 1–30 mm wide, and entire or with a few teeth or a short lobe. Involucral bracts narrow, often purplish-dotted; ligules deep orange, usually drying to purple or pinkish. Achenes 5–9 mm long; beak thin, from shorter to longer than the body. Alpine meadows and slopes; southern Rocky Mountains.

Agoseris glauca (Pursh) Raf.

FALSE DANDELION

A hairless species with linear-oblongate sometimes toothed leaves up to 25 cm long. Flowers 2–5 cm across, light yellow when young but turning pinkish at maturity, on a stem 15–45 cm high. Common; on moist prairie; throughout most of the Prairie Provinces, but occasionally replaced by var. *dasycephala* (T. & G.) Jepson, with leaves oblongate to linear-oblongate, 10–30 cm long, usually entire-margined, and pubescent. Involucre of large flower head 2–3 cm across; bracts hairy, at least along the margins. Common; on hillsides and prairies; Cypress Hills, Rocky Mountains, Peace River.

Agoseris grandiflora (Nutt.) Greene

LARGE-FLOWERED FALSE DANDELION

Stems 25–60 cm high, from a branched caudex. Leaves linear to spatulate, entire to irregularly lobed, glabrous to canescent, usually pubescent along the midribs. Scapes tomentose below; flower heads 3–6 cm across; involucre bracts in 4 or 5 series; outer bracts conspicuously ciliate; ligules yellow, shorter than the bracts. Beak of achene filiform. Moist meadows; western Parklands, Peace River.

Cichorium chicory

- Upper stem leaves large, triangular or rounded, entire. *C. endivia*
- Upper stem leaves small, lanceolate, dentate. *C. intybus*

Cichorium endivia L.

ENDIVE

Annual or biennial plants, with stems 15–60 cm high. Leaves mostly basal, 10–20 cm long, deeply toothed to pinnatifid; upper stem leaves 3–5 cm long, clasping. Flowers 3–5 cm across, bright blue, in clusters of 3 or 4 along upper part of stem. Introduced, cultivated for salad; occasionally escaped.

Cichorium intybus L.

CHICORY

A perennial plant from a long thick deep root, with branching hairy stems 60–100 cm high. Leaves mostly near base of stem, somewhat spatulate, but deeply toothed or pinnatifid with backward-pointing lobes, 7–15 cm long; upper stem leaves small, lanceolate, entire or lobed, stalkless. Flowers bright blue, almost 5 cm across, in clusters of 3 or 4 without stalks, at intervals along upper leafless part of stems. Seed without hairy pappus, but having a series of short blunt scales. An introduced weedy plant; has become fairly common in some localities in Parklands and at scattered locations in Prairies.

Crepis hawk's-beard

- 1. Annuals; stem with many stalkless leaves. 2
- Perennials; few and smaller stem leaves. 3
- 2. Basal rosettes persistent; stems several to many. *C. capillaris*
- Basal rosettes absent when flowering; stems solitary. *C. tectorum*
- 3. Plants dwarfed, with stems 5–10 cm high. *C. nana*
- Plants with well-developed stems. 4

4. Stems and foliage glabrous and glaucous,
with stems much-branched. *C. elegans*
Stems or foliage or both pubescent. 5
5. Plants glabrous except for the rosette
leaves. *C. runcinata*
Plants lightly to densely pubescent
throughout. *C. occidentalis*

Crepis capillaris (L.) Wallr.

GREEN HAWK'S-BEARD

An annual or biennial plant with several stems from a taproot. Stems 2–60 cm high, slender, somewhat pubescent at the base. Basal leaves 3–20 cm long, lanceolate, runcinate-pinnatifid, glabrous or somewhat pubescent with yellow hairs. Inflorescence corymbiform, with several to many heads 10–15 mm across, light yellow. Achenes at first gray, later black. Introduced; a rare weed of open areas; southern Rocky Mountains.

Crepis elegans Hook.

YOUNGIA

Many-stemmed plants from a deep taproot; stems 10–25 cm high. Basal leaves petiolate, 1–3 cm long, lanceolate to spatulate, entire to crenate; stem leaves numerous, linear-lanceolate to linear. Flower heads numerous, 6–8 mm high, about 1 cm across. Gravel flats along creeks and rivers; Rocky Mountains.

Crepis nana Rich.

DWARF HAWK'S-BEARD

Stems 5–10 cm high, leafless, as long as or shorter than the leaves. Basal leaves to 7 cm long, long-petioled, ovate to suborbicular, glaucous, often purplish, clustered on the long thick taproot. Heads 10–13 mm high; ligules short, yellow tinged with purple. Rare; loose gravel and slides; southern Rocky Mountains.

Crepis occidentalis Nutt.

SMALL-FLOWERED HAWK'S-BEARD

A gray hairy or scurfy plant 20–60 cm high. Basal leaves deeply lobed or divided, 10–15 cm long; stem leaves smaller and less divided. Several yellow heads, 12–20 mm across, borne near top of flowering stem. Uncommon; may be found on hillsides; Prairies and southern Rocky Mountains. Syn.: *C. intermedia* A. Gray.

Crepis runcinata (James) T. & G.

SCAPOSE HAWK'S-BEARD

A species with a stem 20–80 cm high. Basal leaves oblanceolate to spatulate, 5–15 cm long, usually entire, but occasionally somewhat toothed. Flower heads 20–25 mm across, on stalks near top of flowering stem. Found in meadows, wooded areas, and low places; throughout the Prairie Provinces.

The var. *glauca* (Nutt.) Boiv. (Fig. 202), smooth hawk's-beard, with involucre bracts neither hairy nor glandular. Found in low often alkaline areas; Prairies. The var. *hispidulosa* Howell, larger than the typical form, with leaves usually glandular along the midrib. Grasslands; southern Rocky Mountains and Cypress Hills.




Fig. 202. Smooth hawk's-beard, *Crepis runcinata* (James) T. & G. var. *glauca* (Nutt.) Boiv.

An introduced annual with a slender, branched, leafy stem 10–50 cm high. Stem leaves linear and stalkless; basal leaves 10–15 cm long, usually with backward-pointing teeth. Many small yellow flowers, 10–15 mm across. Plentiful as a weed; on light soils and roadsides; in Boreal forest and Parklands, and spreading farther southward.

Hieracium hawkweed

Perennial herbs with alternate or basal leaves. Bracts of involucre in series of 1–3 with a few smaller bracts at their base. Pappus composed of brownish or tawny hairs.

- 1. Flowers white or cream. *H. albiflorum*
Flowers yellow or orange. 2
- 2. Rootstock long, slender, with offshoots;
flowers orange. *H. aurantiacum*
Rootstock short, stout, without offshoots;
flowers yellow. 3
- 3. Bracts of involucre imbricate (overlap-
ping like shingles). *H. umbellatum*
Bracts of involucre almost in a single row. 4
- 4. Involucres and flower stalks with dense
long yellow hairs. *H. cynoglossoides*
Involucres blackish or glandular, with
very few yellow hairs. *H. triste* var. *gracile*

Hieracium albiflorum Hook.

WHITE HAWKWEED

A species growing to 30–60 cm high. Basal leaves stalked, hairy, spatulate, 5–12 cm long; stem leaves smaller and without stalks. Flowers white or pale cream, about 12 mm across, in an open-branched cluster at the head of the stem. Found fairly often; in woodlands; southern Rocky Mountains and Cypress Hills.

Hieracium aurantiacum L.

ORANGE HAWKWEED

A perennial with slender running roots. Basal leaves hairy, spatulate, 5–15 cm long. Stem glandular hairy, 15–50 cm high, and bearing a cluster of orange red flowers 20–25 mm across and sometimes 1 or 2 small leaves. Rare; has been found in scattered locations in the Prairie Provinces. A common weed in Eastern Canada.

Hieracium cynoglossoides Arv.-Touv.

WOOLLY HAWKWEED

A species 30–50 cm high, with long white or yellowish hairs. Leaves 5–15 cm long; lower leaves having winged stalks; upper ones oblong to lanceolate, stalkless, hairy. Not common; found occasionally on hillsides and in woodlands; southern Rocky Mountains. Syn.: *H. albertinum* Farr.

Hieracium triste Willd. var. *gracile* (Hook.) Gray

ALPINE HAWKWEED

A plant very similar to *H. cynoglossoides*, but smaller, to 30 cm high, with the foliage glabrous or nearly so, and with the involucre densely black pubescent. Mountain meadows; Rocky Mountains.

Hieracium umbellatum L.

CANADA HAWKWEED

An erect plant 30–100 cm high, with a leafy stem. Leaves ovate to lanceolate, 3–8 cm long. Numerous yellow heads 20–25 mm across. Fairly common; in dry woodlands; Boreal forest and Rocky Mountains. Syn.: *H. canadense* Michx.

Hypochoeris cat's-ear

Hypochoeris radicata L.

CAT'S-EAR

A perennial with stems 20–50 cm high and with a basal rosette of lobed, somewhat hairy leaves. Leaves 5–15 cm long, oblanceolate, with backward-pointing lobes or teeth; the few stem leaves scale-like. Flowers yellow, stalked, 3 or 4 at summit of stems; flower heads 20–25 mm across. Achene (seed) with a long beak and a pappus of plumose white hairs. Introduced; a rare weed; Prairie Provinces.

Krigia dwarf dandelion

Krigia biflora (Walt.) Blake

DWARF DANDELION

Perennial plants with fibrous roots; stems 20–80 cm high. Leaves mostly basal, 3–25 cm long, oblanceolate to broadly elliptic, entire or toothed or occasionally lobed; stem leaves few, sessile, clasping, usually small. Heads several; involucre 7–14 mm high; ligules orange yellow, about 15 mm long. Rare; woods, openings, and fields; southeastern Parklands and Boreal forest.

Lactuca lettuce

Tall leafy plants, annual, biennial, or perennial, with yellow, blue, or white flowers and alternate leaves. Involucres in 1 or 2 rows of almost equal inner main bracts, and with smaller bracts at base. Seeds with white or brown pappus hairs.

- 1. Pappus brown; achenes without a beak. *L. biennis*
- Pappus white; achenes with a distinct beak. 2
- 2. Flowers blue; achenes with a short beak; plant smooth, with a bluish bloom, or blue. *L. pulchella*
- Flowers yellow or blue; achenes with a long slender beak; annual or biennial. 3
- 3. Heads with 6–12 florets. *L. serriola*
- Heads with 12–20 florets. 4
- 4. Leaves with hairs or bristles on underside of midrib. *L. ludoviciana*
- Leaves smooth, without hairs or bristles. 5

5. Pappus on a long beak; flowers yellow; inflorescence crowded. *L. canadensis*
 Pappus sessile; flowers blue; inflorescence open. *L. floridana*

Lactuca biennis (Moench) Fern.

TALL BLUE LETTUCE

An annual or biennial plant 1–2.5 m high, with a leafy stem. Leaves large, 10–25 cm long, deeply lobed or pinnatifid, with sharp-toothed margins. Flowers very numerous, about 5 mm across, blue or creamy white, in a large dense panicle. Found occasionally; in swampy or moist places; throughout Boreal forest. Syn.: *L. spicata* (Lam.) Hitchc.

Lactuca canadensis L.

TALL YELLOW LETTUCE

An annual or biennial plant, smooth, hairless, 1–3 m high, very leafy. Leaves with wavy or lobed margins, lanceolate, 5–20 cm long. Numerous flowers pale yellow, about 6 mm across, in an open, long, terminal panicle. Rare; has been reported from eastern Boreal forest. Native to Eastern Canada.

Lactuca floridana (L.) Gaertn.

ANNUAL BLUE LETTUCE

An annual or biennial plant with stems 50–150 cm high. Leaves mostly petiolate, often pubescent on the underside of the main veins, toothed and often also pinnatifid, 8–30 cm long. Heads numerous in a large panicle; involucre 10–15 mm high; ligules bluish. Margins of woods; southeastern Boreal forest.

Lactuca ludoviciana (Nutt.) Riddell

WESTERN LETTUCE

A biennial species, hairless, 50–150 cm high, with a leafy stem. Leaves ovate-oblong, 5–10 cm long, wavy-lobed; lobes often spiny-tipped. Flowers numerous, yellow or pale lilac, 5–10 mm across, in an open panicle. Not common; found occasionally along riverbanks; southeastern Parklands.

Lactuca pulchella (Pursh) DC.

BLUE LETTUCE

A pale bluish green, smooth, glaucous perennial 30–100 cm high, with white running rootstocks. Leaves usually linear-lanceolate, often with backward-pointing lobes. Flowers bright blue, nearly 25 mm across, in a few-flowered panicle. Seed with a short thick beak and a pappus of white hairs. Common; a weed in cultivated lands and along roadsides; throughout the Prairie Provinces.

Lactuca serriola L.

LOBED PRICKLY LETTUCE

An annual, winter annual, or biennial plant, erect, 30–150 cm high. Leaves 5–20 cm long, deeply lobed; upper leaves clasping stem and often eared at base; underside of midrib bearing a row of short stiff prickles. Flowers 6–8 mm across, yellow, with 6–12 florets; many florets on a large open panicle. Seeds long-beaked, bearing a pappus of white hairs. Introduced, but very common; a weed along roadsides, slough margins, waste places, and on cultivated land; throughout the Prairie Provinces.

The var. *integrata* Gren. & Godr., dentate prickly lettuce, is similar to the species in every way except the leaves are neither lobed nor pinnatifid, except for those at the base. As plentiful in most locations as the type; both often found growing together. Syn.: *L. virosa* L.

Lapsana nipplewort

Lapsana communis L. COMMON NIPPLEWORT

An annual plant with hirsute to subglabrous stems 15–150 cm high. Leaves 3–10 cm long, thin, ovate to subrotund, toothed or occasionally lyrate. Inflorescence corymbiform or paniculate, with several to many heads; involucre 5–8 mm high; ligules yellow. Very rare; a weed of shaded places; southeastern Boreal forest.

Lygodesmia skeletonweed

Stiff-stemmed branching plants with leaves very narrow or reduced almost to scales. Flower heads pink to rose, with 3–10 florets. Pappus with white or very light brown hairs.

Perennials from deep tough rootstocks; leaves small or scale-like. *L. juncea*

Annuals from a tough taproot; leaves long and very narrow; flowers racemose among branches. *L. rostrata*

Lygodesmia juncea (Pursh) D. Don (Fig. 203) SKELETONWEED

A much-branched skeleton-like perennial herb 10–40 cm high, from deep, tough, sticky rootstocks. Lower leaves linear-lanceolate, sometimes 36 mm long; upper ones smaller or reduced to scales. Flower heads with 3–5 florets, pink, 12–15 mm across, borne singly at ends of branches. Seeds short, without a beak, and bearing a pale brownish pappus. Common; on light sandy soil and in sandhills; throughout the southern and central portions of the Prairie Provinces.

Lygodesmia rostrata A. Gray ANNUAL SKELETONWEED

A branching thin-stemmed annual 10–60 cm high, from a tough thin root. Leaves linear, 3-nerved, sharp-tipped, 7–16 cm long; upper leaves much smaller than lower ones. Flowers numerous, with 6–10 florets, pink, about 12 mm across, borne in racemes toward ends of branches. Seeds almost 12 mm long, with a tapering beak and a pappus of white hairs. Somewhat rare; has been found in sandhills and similar locations; in southwestern part of the Prairie Provinces.

Microseris scorzonella

A plant very similar to *Agoseris*, but with the achenes beakless and the pappus sessile.

Leaves all basal; scape with a solitary flower. *M. cuspidata*

Leaves not all basal, usually at least one stem leaf. *M. nutans*




Fig. 203. Skeletonweed, *Lygodesmia juncea* (Pursh) D. Don.

Microseris cuspidata (Pursh) Schultz-Bip.

PRAIRIE FALSE DANDELION

Leaves linear, 10–20 cm long, somewhat hairy, very narrow, tapering to a long narrow point; margins somewhat crinkled. Flowering stem usually no longer than leaves, and bearing a single head with involucre about 20 mm high. Rare, has been found occasionally; eastern Parklands. Syn.: *Agoseris cuspidata* (Pursh) Raf.

Microseris nutans (Hook.) Schultz-Bip.

NODDING SCORZONELLA

A perennial with fleshy taproot; stems few, erect or curved at base, 10–50 cm high, usually branched and leafy above. Leaves 10–30 cm long, linear or linear-lanceolate, entire to toothed or pinnatifid. Inflorescence with 1 to several heads, nodding before flowering; involucre 8–22 mm high; bracts glabrous to scurvy puberulent; ligules yellow, 3–8 mm long. Mountain slopes; southern Rocky Mountains.

Picris bitterweed

Picris echioides L.

BRISTLY OXTONGUE

Coarse spiny-hispid annual plants, with stems 30–80 cm high. Leaves 10–20 cm long and 7 cm wide, toothed or entire; lower leaves oblanceolate, petioled; upper ones oblong to lanceolate, sessile, often clasping. Involucre 1–2 cm high; bracts in two series; inner bracts narrow, with spiny tip; ligules about 1 cm long, yellow. Introduced; a rare weed of waste places; southern fringe of Boreal forest, Peace River.

Prenanthes rattlesnakeroot

Perennial plants with alternate, usually stalked leaves. Heads clustered, usually nodding or pendent. Bracts of involucre usually colored, not green. Pappus hairs brownish or tawny.

- 1. Bracts of involucre hairy; basal leaves tapering to winged stalks. *P. racemosa*
- Bracts of involucre not hairy; basal leaves cordate- or hastate-based, abruptly joining stalk. 2
- 2. Flower heads nodding; leaves lobed or cleft. *P. alba*
- Flower heads erect; leaves dentate. *P. sagittata*

Prenanthes alba L.

WHITE LETTUCE

A smooth hairless plant with an erect stem 50–150 cm high, sometimes purplish. Stalkless lower leaves hastate, cordate, or triangular, sometimes lobed, 2–15 cm long; upper leaves lanceolate and without stalks. Heads in a long terminal panicle, drooping, greenish or yellowish white with purplish bracts, slightly scented. Pappus cinnamon brown, very conspicuous. Found often; in open woodlands; throughout the Prairie Provinces. Syn.: *Nabalus albus* (L.) Hook.

Prenanthes racemosa Michx.

GLAUCOUS WHITE LETTUCE

A very erect stout perennial 50–150 cm high; stem usually covered with a white bloom. Lower leaves oblanceolate to spatulate, 10–20 cm long, tapering to a winged stalk; upper leaves smaller and somewhat clasping. Heads in a long spike of crowded clusters, purplish, about 12 mm long and 6 mm broad, usually not as pendent as in *P. alba*. Pappus straw-colored. Fairly abundant; in wooded areas; across the Prairie Provinces. Syn.: *Nabalus racemosus* (Michx.) DC.

Prenanthes sagittata (Gray) Nels.

PURPLE RATTLESNAKEROOT

An erect glabrous perennial with stems 30–70 cm high. Lower leaves petioled, sagittate or hastate, dentate, glabrous, with blades 10–15 cm long; upper leaves sessile, lanceolate. Heads in a narrow panicle, at first erect, later spreading; involucre about 12 mm high; ligules purplish-tinged. Mountain woods; northwestern Boreal forest, Rocky Mountains.

Sonchus sow-thistle

- 1. Flowering heads 3–5 cm across; perennials from creeping rootstocks. *S. arvensis*
 Flowering heads not over 2.5 cm across; annuals growing from deep taproots. 2
- 2. Ears at base of clasping leaves rounded. *S. asper*
 Ears at base of clasping leaves acutely pointed. *S. oleraceus*

Sonchus arvensis L. (Fig. 204)

PERENNIAL SOW-THISTLE

A weedy plant 50–150 cm high, with vigorous creeping rootstocks. Stems usually hollow and slightly branched. Lower leaves runcinate-pinnatifid or with backward-pointing lobes, 10–25 cm long, and narrowed to a short stalk; upper leaves less lobed and without stalks; teeth of leaves spiny-pointed. Flowers numerous, showy, bright yellow, on bristly stalks in a corymbose panicle; involucre glandular hairy. Very common; in moister districts and wet places; throughout the Prairie Provinces.

The var. *glabrescens* Guenth., Grab. & Wimm., smooth perennial sow-thistle, differing from the species in having a smooth, hairless, not glandular involucre. Common; in many locations; throughout the Prairie Provinces. Syn.: *S. uliginosus* Bieb.

Sonchus asper (L.) Hill

PRICKLY SOW-THISTLE

An annual weedy plant 50–150 cm high. Leaves clasping stem at base, slightly lobed or divided; leaf lobes usually having spine-tipped teeth; basal lobes rounded. Flowers 12–25 mm across, pale yellow. Fairly common; a weed of gardens and roadsides; throughout the Prairie Provinces.

Sonchus oleraceus L.

ANNUAL SOW-THISTLE

An annual plant 50–200 cm high, slightly branched. Leaves deeply lobed, with rather soft prickles; lower leaves stalked; upper ones clasping the stem.



Fig. 204. Perennial sow-thistle, *Sonchus arvensis* L.

Heads pale yellow, 12–25 mm across. Fairly common; in gardens and waste places. Leaves undivided in f. *integrifolius* (Wallr.) G. Beck.

Stephanomeria skeletonweed

Stephanomeria runcinata Nutt.

RUSH-PINK

Plants similar to *Lygodesmia*, with creeping rootstocks. Leaves toothed or pinnatifid, well-developed. Heads terminal; ligules 10–15 mm long, pink. Pappus plumose. Rare; badlands, loose shale beds; Prairies.

Taraxacum dandelion

1. Involucral bracts all appressed or ascending. *T. ceratophorum*
Involucral bracts reflexed. 2
2. Leaves deeply divided, almost to midrib, into narrow segments, with terminal lobe small; seeds reddish. *T. laevigatum*
Leaves shallowly divided, with terminal lobe large; seeds greenish or brownish yellow. *T. officinale*

Taraxacum ceratophorum (Ledeb.) DC.

DANDELION

Plants similar to *T. officinale*, but usually less robust; leaves less lobed, commonly glabrous or nearly so. Flower heads 15–20 mm across; involucre about 15 mm high; outer bracts appressed to spreading, often with horned tip. Open ground; throughout the Prairie Provinces.

Taraxacum laevigatum (Willd.) DC.

RED-SEEDED DANDELION

A stemless perennial plant, with all leaf and flower stalks arising from a fleshy root crown, and growing from a deep taproot. Leaves 10–20 cm long, deeply divided into narrow segments. Flowers 25–35 mm across, with many florets, bright yellow, borne singly on stems or scapes 10–30 cm high. Seeds bright red, with a white hairy pappus on a long thin beak. Fairly common; in waste places and roadsides; throughout the Prairie Provinces. Introduced from Europe. Syn.: *T. erythrospermum* Andrz.

Taraxacum officinale Weber

DANDELION

An introduced stemless perennial with deep fleshy taproots. Leaves coarsely incised, with triangular lobes and a large terminal lobe; leaf stalks and flowering stems arising from a root crown. Flower heads yellow, 35–50 mm across, with very many florets. Seed greenish buff, with a long beak and a pappus of white hairs. Often the earliest plant blooming in spring. Very common; on lawns, along roadsides, and in waste places; throughout the Prairie Provinces.

Tragopogon goat's-beard

A fairly tall biennial or perennial plant, with deep fleshy taproots and grass-like leaves. Flowering heads large. Seeds and plumose seed heads very

large; seeds narrow and long, with 5–10 ribs, ending in a long beak terminating in a pappus of plumose hairs.

1. Flowers purple; involucre bracts much longer than the florets. *T. porrifolius*
- Flowers yellow; involucre bracts varying in length. 2
2. Involucre bracts longer than florets; stem usually thickened below flower head. *T. dubius*
- Involucre bracts not longer than florets; stem slightly thickened below flower head. *T. pratensis*

Tragopogon dubius Scop.

YELLOW GOAT'S-BEARD

A coarse biennial plant from a deep fleshy taproot 30–60 cm high. Leaves narrow, erect, grass-like, 10–30 cm long, stalkless, clasping at base. Flowers sulfur yellow, 3–5 cm across; involucre bracts 10–14, longer than yellow florets; heads borne singly on summit of an erect scape or stem that is decidedly thickened just below head. Seed heads plumose, very conspicuous, usually 7–10 cm in diam; seeds long, tapering to a long beak, having ribs of minute tubercles and a pappus of plumose white hairs; body of seed about 12 mm long, with beak slightly longer. An introduced plant, apparently coming from southwestern United States through Colorado. Common; along roadsides and in waste places; almost throughout the Prairie Provinces.

Tragopogon porrifolius L.

SALSIFY

A perennial or biennial with an edible fleshy taproot 30–80 cm high. Leaves grass-like. Flower heads 5–10 cm across; florets purple; bracts much exceeding the florets. A garden plant sometimes found as a weed; in southeastern Parklands and southern Rocky Mountains.

Tragopogon pratensis L.

GOAT'S-BEARD

A biennial very similar to *T. dubius*, but somewhat smaller. Florets chrome yellow; bracts of involucre usually 8 or 9, not longer than florets. Seed somewhat shorter than that of *T. dubius*, but similar in appearance. Not as aggressive and rapid-spreading as *T. dubius*, which came into the Prairies from the west. A fairly common weed; in eastern Parklands.

GROUP 3

This very large group consists of plants in which some or all florets are tubular. The tubular (central) florets are usually perfect or bisexual, whereas the marginal or ligulate (strap-shaped) florets are either female or neutral. Plants are without milky sap. The primary leaves (those following the cotyledons or seed leaves) are usually in pairs.

1. Flower heads with all florets tubular (discoid heads). 2
- Flower heads with florets both tubular and ray (radiate heads). 33

2. Bracts of involucre dry, parchmenty or membranous, not green.	3
Bracts of involucre not dry or parch- menty, usually green.	7
3. Some florets bisexual, with both male and female florets on same plant.	4
No fertile bisexual florets, with male and female florets on separate plants.	6
4. Leaves alternate; flowers all perfect.	<i>Gnaphalium</i>
Leaves opposite.	5
5. Pappus consisting of terminal awns; inner bracts about twice as long as outer ones.	<i>Thelesperma</i>
Pappus absent; heads subtended by foli- age leaves.	<i>Psilocarphus</i>
6. Tall plants; stems leafy.	<i>Anaphalis</i>
Low plants; leaves mostly basal, with stem leaves reduced.	<i>Antennaria</i>
7. Pappus none or inconspicuous.	8
Pappus present, distinct.	11
8. Leaves large, deltoid-ovate.	<i>Adenocaulon</i>
Leaves elliptic, lobed or divided.	9
9. Inflorescence glandular sticky; leaves linear-elliptic.	<i>Madia</i>
Inflorescence not glandular; leaves lobed or divided.	10
10. Heads small, numerous, in open or spike- like panicles.	<i>Artemisia</i>
Heads larger, 4–7 mm across.	11
11. Leaves 2–3 times pinnatifid, or pinnately divided into toothed segments, inflorescence has no ray florets.	<i>Tanacetum</i>
Leaves entire or toothed, not pinnately divided. Inflorescence usually with white ray florets.	<i>Chrysanthemum</i>
12. Involucral bracts spiny-tipped or with hooked bristles.	13
Involucral bracts not spiny-tipped and without hooked bristles.	16
13. Involucral bracts with hooked bristles; leaves large, cordate, and not prickly.	<i>Arctium</i>
Some bracts spiny-tipped.	14
14. Pappus hairs not longer than achene.	<i>Centaurea</i>
Pappus hairs much longer than achene.	15

15. Pappus hairs plumose or feathery.	<i>Cirsium</i>
Pappus hairs roughened but not plumose.	<i>Carduus</i>
16. Florets purplish white, purple, or rose.	17
Florets yellow or greenish yellow.	25
17. Pappus consisting of scales; flower heads	
globose.	<i>Echinops</i>
Pappus consisting of bristles or hairs.	18
18. Pappus hairs not longer than achene.	<i>Centaurea</i>
Pappus hairs much longer than achene.	19
19. Flowers in terminal, more or less flat-	
topped clusters or solitary.	20
Flowers in elongated racemes or spikes.	22
20. Leaves alternate; flowers purple.	21
Leaves opposite or whorled; flowers pur-	
plish white or purple.	<i>Eupatorium</i>
21. Pappus plumose, white.	<i>Saussurea</i>
Pappus barbed, purple.	<i>Vernonia</i>
22. Florets purple.	<i>Liatris</i>
Florets whitish.	23
23. Involucre 3–4 mm high.	<i>Erigeron</i>
Involucre 8–15 mm high.	24
24. Leaves linear.	<i>Aster</i>
Leaves deltoid or cordate.	<i>Brickellia</i>
25. Low grayish shrubs, with linear leaves.	<i>Chrysothamnus</i>
Herbs.	26
26. Leaves with spiny-tipped teeth.	<i>Haplopappus</i>
Leaves without spiny-tipped teeth.	27
27. Most of the leaves opposite.	28
Leaves alternate.	29
28. Pappus a series of rough bristles.	<i>Arnica</i>
Pappus of barbed teeth.	<i>Bidens</i>
29. Flowers appearing before the large felty-	
pubescent leaves.	<i>Petasites</i>
Flowers appearing after the leaves.	30
30. Heads not over 5 mm across, in open or	
spike-like panicles.	<i>Artemisia</i>
Heads over 5 mm across, in more or less	
flat-topped clusters.	31
31. Stems and leaves usually white woolly.	<i>Hymenopappus</i>
Stems and leaves green and scarcely	
hairy.	32
32. Foliage and flowers not scented.	<i>Senecio</i>
Foliage and flowers scented, especially	
when bruised.	<i>Matricaria</i>

33. Ray florets yellow.	34
Ray florets not yellow, usually blue, purple, or white.	53
34. Pappus composed of hairs.	35
Pappus, when present, composed of scales or bristles.	39
35. Involucral bracts usually one series, not or scarcely imbricated (overlapping like shingles).	36
Involucral bracts in 2 or more series, well-imbricated.	37
36. Leaves opposite.	<i>Arnica</i>
Leaves alternate.	<i>Senecio</i>
37. Heads small and numerous, in terminal or axillary clusters; leaves simple, not pinnatifid or divided.	<i>Solidago</i>
Heads usually solitary at ends of branches.	38
38. Stems leafy almost to inflorescence; pappus double, with the outer series composed of scales or small bristles.	<i>Chrysopsis</i>
Leaves mostly basal, with stem leaves reduced; pappus not double.	<i>Haplopappus</i>
39. Receptacle (disk to which florets are attached) bare and naked, without chaffy scales but sometimes bristly.	40
Receptacle with chaffy scales between florets.	45
40. Flower heads not more than 6 mm across, in terminal clusters.	<i>Gutierrezia</i>
Flower heads more than 6 mm across.	41
41. Involucre of flower heads sticky and gummy.	<i>Grindelia</i>
Involucre not sticky and gummy.	42
42. Heads large, usually more than 35 mm across; disk florets purple; ray florets yellow with usually a purplish base.	<i>Gaillardia</i>
Heads usually less than 35 mm across; all florets yellow.	43
43. Involucral bracts decidedly reflexed or turned backward when mature; leaves lanceolate.	<i>Helenium</i>
Involucral bracts not reflexed; leaves usually pinnatifid with narrow lobes.	44
44. Lower leaves opposite; seeds long, the length four times the width.	<i>Bahia</i>
All leaves alternate; length of seeds less than four times the width.	<i>Hymenoxys</i>

45. Pappus absent, or present merely as a short crown.	46
Pappus composed of scales, bristles, or barbs.	50
46. Ray florets neutral, sterile.	47
Ray florets female.	48
47. Receptacle flat; flowers not over 25 mm across; ray florets orange; leaves pinnately divided into narrow segments.	<i>Coreopsis</i>
Receptacle convex; flowers 5 cm or wider across; ray florets yellow; leaves not divided into narrow segments.	<i>Rudbeckia</i>
48. Annual plants with glandular sticky inflorescence; plants strongly scented; leaves linear.	<i>Madia</i>
Perennial plants; leaves broad.	49
49. Leaves basal, large, and white woolly beneath; roots thick, exuding a sticky balsam.	<i>Balsamorhiza</i>
Leaves opposite, very rough.	<i>Heliopsis</i>
50. Achenes (seeds) with barbed awns.	51
Achenes without barbed awns.	52
51. Aquatic floating plants, with the submerged leaves very finely divided; perennial.	<i>Megalodonta</i>
Plants not aquatic, but often found standing in water or mud; mostly annual.	<i>Bidens</i>
52. Plants with entire, undivided leaves; receptacle flat.	<i>Helianthus</i>
Plants with pinnately divided leaves; receptacle cylindrical and tall.	<i>Ratibida</i>
53. Pappus composed of numerous hairs.	54
Pappus absent or composed of a few hairs.	59
54. Ray florets not conspicuous.	55
Ray florets conspicuous.	56
55. Bracts nearly equal; flowers purplish.	<i>Aster</i>
Bracts unequal; flowers white.	<i>Erigeron</i>
56. Leaves all basal, appearing after the flowers.	<i>Petasites</i>
Leaves appearing before the flowers.	57
57. Plants low, stemless, tufted; leaves usually extending beyond the flowers.	<i>Townsendia</i>
Plants usually with leafy stems; leaves not extending beyond flowers.	58

58. Ray florets 50 or more; bracts in 1 or 2 series. *Erigeron*
 Ray florets 10–50; bracts in several series. *Aster*
59. Receptacle with chaffy scales. 60
 Receptacle naked, without chaffy scales. 63
60. Leaves opposite. *Galinsoga*
 Leaves alternate. 61
61. Flowers large and purple; leaves undivided. *Echinacea*
 Ray florets white; leaves divided. 62
62. Heads small, numerous, in dense almost flat-topped clusters; plant not strongly scented. *Achillea*
 Heads fewer, about 25 mm across; plant unpleasantly scented. *Anthemis*
63. Pappus composed of bristly scales or with 2–4 slender bristles. *Boltonia*
 Pappus absent, or composed of merely a border. 64
64. Leaves entire, not divided; heads solitary on stem, 25–50 mm across; plants not scented. *Chrysanthemum*
 Leaves much divided into narrow segments; heads several to a stem, less than 25 mm across; plants usually somewhat scented. *Matricaria*

Achillea yarrow

1. Leaves pinnately dissected into linear-lanceolate segments. *A. millefolium*
 Leaves subentire to incised, not pinnately dissected. 2
2. Leaves subentire to serrate; ligules 3–5 mm long. *A. ptarmica*
 Leaves incised; ligules 1–2 mm long. *A. sibirica*

Achillea millefolium L.

YARROW, MILFOIL

A perennial from rootstocks, with stems 20–100 cm high, and somewhat aromatic. Leaves pinnately dissected, with the ultimate segments 1–2 mm wide; blades 3–15 cm long and to 25 mm wide. Heads numerous, in a flat-topped inflorescence, white to pinkish. Moist meadows, woods, and openings; Parklands, Boreal forest, and Rocky Mountains. In f. *purpurea* (Gouan) Schinz & Thell., ligules purple.

The var. *occidentalis* DC., woolly yarrow, a perennial from shallow rootstocks, 10–30 cm high, usually covered with silky hairs. Leaves 3–10 cm long, finely divided into segments. Flowers in a compact round-topped cluster at head of stem, white, rarely pink, about 6 mm across; involucre bracts with

straw-colored margins. Common; on prairies and along roadsides; throughout the Prairie Provinces. Syn.: *A. lanulosa* Nutt.

In var. *nigrescens* E. May., tips and margins of bracts dark brown to blackish. This form more common than the others in the northern part of Boreal forest. Syn.: *A. borealis* Bong.

Achillea ptarmica L. (Fig. 205)

SNEEZEWEED

A perennial with creeping rootstocks; stems 30–60 cm high, villous-tomentose above, subglabrous below. Leaves glabrous or nearly so, 8–10 cm long, 2–6 mm wide, closely serrate to subentire. Heads several to numerous in a corymbiform inflorescence; involucre 4–5 mm high; ligules white, 3–5 mm long. Introduced; commonly the “double-flowered” form, occasionally escaped and established; Boreal forest.

Achillea sibirica Ledeb.

MANY-FLOWERED YARROW

A perennial 30–60 cm high. Leaves linear, deeply toothed, 3–7 cm long. Flowers very similar to those of *A. ptarmica*, but bracts having dark brown margins. Found occasionally; in wooded areas and disturbed places; Boreal forest. Syn.: *A. multiflora* Hook.

Adenocaulon

Adenocaulon bicolor Hook.

TRAILPLANT

A slender fibrous-rooted perennial 30–70 cm high. Leaves mostly basal, long-petioled, 3–15 cm wide, deltoid-ovate to cordate, subglabrous above, white lanate below. Inflorescence an open panicle; heads discoid; involucre bracts about 2 mm long, reflexed in fruit. Moist shady woods; southern Rocky Mountains.

Anaphalis pearly everlasting

Anaphalis margaritacea (L.) C. B. Clarke

PEARLY EVERLASTING

A woolly-stemmed perennial growing in a cluster from numerous running rootstocks; stem usually erect, 30–80 cm high, often branched near top. Leaves stalkless, alternate, linear-lanceolate, white woolly beneath, grayish above, 5–12 cm long. Flowers numerous, in a fairly dense terminal cluster up to 15 cm across; each flower head discoid, 6–8 mm across, white, with pearly white papery bracts. Found in open woodlands; in eastern Parklands.

The var. *subalpina* Gray, with very congested inflorescence, found in southern Rocky Mountains and in Cypress Hills.

Antennaria everlasting

Perennial herbs, usually mat-forming, with leaves mostly basal and usually woolly or hairy. Flowers unisexual, sometimes with both sexes on the same plant, but usually on separate plants; female heads sometimes producing fertile seed without pollination; involucre bracts papery. Achenes bearing a white, hairy pappus.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Achillea ptarmica* L.

LOCATION: The Narrows

COLL: J. Loomer

DET:

COLL.NO: 15190

DATE: 3 August 1970

ELEVATION: 240 m

HABITAT: Waste area




Fig. 205. Sneezeweed, *Achillea ptarmica* L.

1. Plants with well-developed stem leaves; basal leaves, if present, upright around the stem; leaves mostly linear-lanceolate. 2
 - Plants with greatly reduced stem leaves; basal leaves in well-developed appressed rosettes; leaves usually small, ovate to spatulate. 6
2. Plants low, densely cespitose; flowering stems 2–5 cm high; heads solitary. *A. dimorpha*
 - Plants not low and densely cespitose; flowering stems to 50 cm high; heads several on a stem. 3
3. Involucres 4–5 mm high; bracts scarious throughout, glabrous. *A. luzuloides*
 - Involucres 6–10 mm high; bracts scarious only at the tip, woolly at the base. 4
4. Bracts of involucre conspicuously dark green or brown below; plants usually 10–20 cm high. *A. lanata*
 - Bracts of involucre with or without a brown spot; plants 20–50 cm high. 5
5. Involucral bracts in 6 or 7 series; base with a large dark spot; tip brownish or yellowish. *A. pulcherrima*
 - Involucral bracts in 3 or 4 series; base with or without a small dark spot; tip white. *A. anaphaloides*
6. Heads on long peduncles; inflorescence open, spread out along the stem. *A. racemosa*
 - Heads on short peduncles or subsessile; inflorescence more or less crowded at the top of the stem. 7
7. Rosette leaves densely pubescent on both sides. 8
 - Rosette leaves glabrous to subglabrous above, densely pubescent below. 19
8. Terminal part of involucral bracts greenish or light to dark brown. 9
 - Terminal part of involucral bracts colorless, white, or pink. 15
9. Terminal part of involucral bracts pale to dark brown. 10
 - Terminal part of involucral bracts greenish. 12
10. Involucre 4–5 mm high; tips of bracts pale brown. *A. umbrinella*
 - Involucre 6–7 mm high; tips of bracts pale or dark brown. 11

11. Rosette leaves 5–10 mm wide; tips of bracts golden brown. *A. russellii*
 Rosette leaves 1–5 mm wide; tips of bracts dark brown. *A. alpina*
12. Involucre 8–10 mm high; leaves narrowly oblanceolate. 13
 Involucre 4–7 mm high; leaves oblanceolate. 14
13. Leaves green, glabrous or nearly so above. *A. glabrata*
 Leaves grayish tomentose above, especially those of rosettes. *A. angustata*
14. Flower heads typically solitary; bracts squarrose. *A. monocephala*
 Flower heads typically 3–5 on a stem; bracts appressed. *A. alpina*
15. Terminal part of involucre bracts roseate to deep pink, even at maturity. *A. rosea*
 Terminal part of involucre bracts whitish or light pink when young. 16
16. Involucre bracts with a large dark spot at the base. *A. corymbosa*
 Involucre bracts green or whitish at the base. 17
17. Rosette leaves to 50 mm long, 20 mm wide. *A. neodioica*
 Rosette leaves to 25 mm long, usually less than 10 mm wide. 18
18. Involucre 5–7 mm high; stems with 8–12 leaves. *A. parvifolia*
 Involucre 8–13 mm high; stems with 5–7 leaves. *A. aprica*
19. Rosette leaves clearly 3- to 5-veined, often to 50 mm long by 30–40 mm wide. 20
 Rosette leaves with only the midrib clearly visible, and usually smaller than 50 mm long by 40 mm wide. 21
20. Rosette leaves glabrous; involucre 5–9 mm high. *A. plantaginifolia*
 Rosette leaves at first somewhat pubescent above; involucre 7–10 mm high. *A. howellii*
21. Rosette leaves glabrous or glabrate, 20–40 mm long; involucre 7–10 mm high. *A. parlinii*
 Rosette leaves usually thinly pubescent above to glabrate, 10–50 mm long; involucre 6–9 mm high. *A. neodioica*

Antennaria alpina (L.) Gaertn.

ALPINE EVERLASTING

A mat-forming perennial with more or less leafy stolons; stems 3–15 cm high. Basal leaves 5–20 mm long, 1–5 mm wide, oblanceolate, loosely pubescent on both sides. Heads 3–5 in a small cyme; involucre mostly 4–7 mm high; bracts pointed with terminal part dark brown or blackish green. Rare; alpine meadows, openings, and slopes; southern Rocky Mountains.

The common form is var. *canescens* Lange, with leaves persistently pubescent on both sides.

Antennaria anaphaloides Rydb.

TALL EVERLASTING

A tall species, not mat-forming, 20–50 cm high, with oblanceolate basal leaves 5–10 cm long and long narrow stem leaves. Many-flowering heads in an open corymb; bracts with white tips, and with or without very small dark spots. Found often; in southern Rocky Mountains and locally in Cypress Hills.

Antennaria angustata Greene

PUSSY-TOES

A mat-forming perennial without stolons. Rosette leaves linear-oblanceolate, tomentose, often glabrescent above. Heads usually solitary; involucre 8–10 mm high; tips of bracts greenish black. Alpine slopes; southern Rocky Mountains.

Antennaria aprica Greene

LOW EVERLASTING

A low mat-forming perennial usually less than 15 cm high. Spatulate- or wedge-shaped rosette leaves 1–2 cm long, densely whitish woolly on both sides; stem leaves 5–7, linear, about 1 cm long. White flower heads occasionally with a faint pinkish tinge, in compact, short-stalked clusters on a stem 5–15 cm high. Common; on dry prairies; throughout the Prairie Provinces.

Antennaria corymbosa E. Nels.

CORYMBOSE EVERLASTING

A medium-tall species with stems 20–30 cm high. Rosette leaves oblanceolate, 1–3 cm long, grayish, finely woolly; stem leaves linear and sharp-tipped. Flower heads stalked, in a terminal corymb; bracts having white upper portions. Found in mountain meadows; in southern Rocky Mountains and in open woodlands in Cypress Hills.

Antennaria dimorpha Nutt.

CUSHION EVERLASTING

A densely caespitose grayish tomentose perennial with a many-branched caudex forming small mats or cushions. Stems 1–4 cm high, leafy, terminated by a solitary flower head; involucre 10–15 mm high. Rare, but probably often overlooked; more or less open grassland; Prairies and southern Rocky Mountains.

Antennaria glabrata (Vahl) Greene

PUSSY-TOES

Possibly only a much less pubescent, often almost glabrous, form or variety of *A. angustata*. Rare; moist slopes; southern Rocky Mountains.

A tall species 20–35 cm high. Basal leaves ovate, wedge-shaped, 2–5 cm long, closely silky-woolly beneath, but bright green above; stem leaves small and narrow. Heads in a corymb. Three varieties are distinguished in this species:

- 1. Rosette leaves 1–2 cm wide. var. *howellii*
 Rosette leaves 5–10 mm wide. 2
- 2. Stems to 20 cm high when in flower, elongating to 35 cm at maturity. var. *athabascensis* (Greene) Boiv.
 Stems 15–20 cm high at maturity. var. *campestris* (Rydb.) Boiv.

The var. *howellii* is commonly found in pine woods and open coniferous forests in Boreal forest, Riding Mountain, and Cypress Hills; var. *athabascensis*, in Parklands and dry open areas in Boreal forest; var. *campestris*, in moist grasslands in Prairies and on slopes in Parkland.

Antennaria lanata (Hook.) Greene

A perennial with a rather thick caudex or rootstock. Stems densely grayish tomentose, stout, erect, leafy, 10–20 cm high. Basal leaves in tufts, linear-oblong to oblanceolate, 2–6 cm long, 3–15 mm wide, obscurely to distinctly 3-nerved. Heads 6–12 in a dense cyme; involucre densely tomentose, 5–8 mm high; outer bracts with a dark brown tip. Alpine meadows; southern Rocky Mountains.

Antennaria luzuloides T. & G.

A perennial with a somewhat woody caudex and erect or ascending leafy stems 10–50 cm high. Basal leaves erect in tufts, 3–10 cm long, 1–5 mm wide, thinly tomentose. Heads often numerous, in a dense cyme or cymose panicle; involucre 4–5 mm high, glabrous or nearly so; bracts pale green or brownish with white or somewhat pinkish tips. Alpine meadows and slopes; southern Rocky Mountains.

Antennaria monocephala DC.

Plants with dense basal rosettes and stems 5–10 cm high. Rosette leaves 8–15 mm long, silky tomentose below; stem leaves linear, 5–10 mm long. Heads usually solitary; involucre about 5 mm high; bracts with greenish brown tips, squarrose. Alpine meadows and slopes; southern Rocky Mountains.

Antennaria neodioica Greene

A fairly tall species with rosette leaves indistinctly 3-ribbed, oblanceolate to obovate, woolly, 2–3 cm long, yellowish green, more or less tomentose above. Flower heads with rather long stalks, on a leafy stem 15–25 cm high, strongly stoloniferous. Syn.: *A. obovata* E. Nels. Plants with the leaves glabrous above distinguished as var. *randii* (Fern.) Boiv. Syn.: *A. canadensis* Greene. Both forms found in dry open woods, shrubbery, and moist grassland; throughout Parklands and southern fringes of Boreal forest.



HERBARIUM OF THE RESEARCH STATION
 WINDY CREEK, ALA.
 SPECIES: *Antennaria howellii* Greene
 var. *howellii*
 LOCATION: 11.89, 1000 ft. above
 COLLECTOR: L. S. Brown
 HT: 47
 COLL. NO: 1700
 DATE: 13 June 1949
 ELEVATION: 1000 ft.
 HABITAT: 1000 ft. above

Fig. 206. Howell's everlasting, *Antennaria howellii* Greene.

Antennaria parlinii Fern.

EASTERN EVERLASTING

A mat-forming stoloniferous plant with stems 20–40 cm high. Rosette leaves 2–6 cm long, 1–4 cm wide, bright green and glabrous above, obovate-spatulate. Heads 4–8 in a corymb; involucre 7–10 mm high. Open woods; southeastern Boreal forest.

Antennaria parvifolia Nutt.

SMALL-LEAVED EVERLASTING

A mat-forming perennial 15–25 cm high. Rosette leaves very small, barely exceeding 15 mm long, white, densely woolly on both sides, somewhat angular but roughly spatulate. Stem leaves 7–12, about 10 mm long, linear. Flower heads nodding or pendent when young. Very plentiful; on dry prairie and in saline meadows; throughout Prairies.

Antennaria plantaginifolia (L.) Hook.

PLANTAIN-LEAVED EVERLASTING

A tall species; basal leaves oval to spatulate, 3–7 cm long, woolly above and below. Flower heads on an erect leafy stem 20–50 cm high. Found occasionally; southeastern Boreal forest.

Antennaria pulcherrima (Hook.) Greene

SHOWY EVERLASTING

Perennials, not mat-forming plants, 30–50 cm high. Basal leaves oblanceolate, 5–10 cm long, 3-ribbed, woolly; stem leaves narrow and smaller. Heads in a cluster at the top of the stem; bracts usually brownish with gray white tips. Found occasionally; in moist soils; Boreal forest and Rocky Mountains.

Antennaria racemosa Hook.

RACEMOSE EVERLASTING

Plants with elongate stolons and tufts of basal leaves; stems 10–60 cm high, erect, leafy. Basal and stolon leaves elliptic to elliptic-obovate, 2–8 cm long, 1–5 cm wide, 1- to 3-nerved, glabrous or nearly so above, more or less tomentose below. Heads in open racemes; lower ones on peduncles 2–5 cm long; involucre 6–8 mm high. Coniferous woods; southern Rocky Mountains.

Antennaria rosea Greene

ROSY EVERLASTING

A somewhat mat-forming white woolly perennial 15–50 cm high. Basal rosette leaves 10–25 mm long, oblanceolate to spatulate, stalked, white woolly, pointed at apex. Flower heads in a close terminal cluster, having conspicuous pinkish-tipped involucral bracts. Found in meadows and moist hillsides in favored localities; throughout the Prairie Provinces, but more particularly in Cypress Hills and Rocky Mountains.

Antennaria russellii Boiv.

RUSSELL'S EVERLASTING

A species very similar to *A. neodioica*, but with the tips of the involucral bracts golden brown or straw-colored, and with the involucre smaller (only 6–7 mm high). Open woods; Cypress Hills.

Antennaria umbrinella Rydb.

BROWN-BRACTED MOUNTAIN EVERLASTING

A low mat-forming species 2–6 cm high, with small, whitish, silky rosette leaves and linear-lanceolate stem leaves. Bracts of the few small flower heads dark brown. Very local, but found on dry hillsides; southern Rocky Mountains and Cypress Hills. Syn.: *A. aizoides* Greene.

Anthemis mayweed

Anthemis cotula L.

STINKING MAYWEED

A hairless annual weed, much-branched, 30–70 cm high, having a fetid and unpleasant odor. Leaves deeply dissected into very narrow lobes, 2–5 cm long. Flowers about 2 cm across, with numerous yellow disks; ray florets 10–18, white, borne on heads of stems, forming a large flat cluster. An introduced weed, plentiful in Eastern Canada, rare in southeastern and northwestern Parklands. Most plants recently reported in the Prairie Provinces as stinking mayweed now have been identified as *Matricaria maritima* var. *maritima* (*M. inodora*). Syn.: *Maruta cotula* (L.) DC. Another species, *A. tinctoria* L., with yellow flowers and somewhat broader leaf divisions, also reported, but very rare.

Arctium burdock

Large, coarse, biennial plants with broad, oval, or cordate leaves, which are long-stalked and paler beneath. Flowers discoid; involucre globose and much-imbricated; bracts stiff, hook-tipped.

1. Bracts of involucre densely cottony. *A. tomentosum*
Bracts of involucre not cottony, sometimes slightly woolly. 2
2. Involucre 2 cm or more across; inner bracts as long as, or longer than, flower head. *A. lappa*
Involucre 10–25 mm across; inner bracts not longer than flower head. *A. minus*

Arctium lappa L.

COMMON BURDOCK

A coarse, branching plant 1–3 m high. Leaves broadly ovate, stalked, pale beneath, often cordate-based, up to 45 cm long. Flowers purple, discoid, with a globose involucre 25–40 mm across; bracts tipped with hooked bristles. An introduced weed becoming increasingly plentiful; in waste places; eastern Parklands and southeastern Boreal forest.

Arctium minus (Hill) Bernh.

LESSER BURDOCK

A coarse, tall, branching biennial 1–2 m high, from deep thick taproots. Leaves large, cordate, pale and downy beneath, up to 30 cm long. Flower heads numerous, usually in a leafy one-sided raceme, discoid with purple florets, 15–25 mm across. Involucre somewhat globose, green to purplish, bracts with hooked bristles. Burs (seed-bearing heads) very prickly, and can be carried on clothing and the coats of animals. An introduced and widely distributed weed, very plentiful locally; as soon as it becomes established, becoming very common.

Arctium tomentosum Mill.

COTTON BURDOCK

A species similar to *A. minus*, but with involucre slightly larger and covered with a cottony web. Involucral bracts having hooked tips. Introduced; found in a few locations, widely separated, but still quite rare.

Arnica arnica

Perennial plants from rootstocks, with opposite leaves and large radiate yellow or orange heads. Pappus a single series of rough bristles.

1. Leaves cordate to oblong. 2
Leaves oblong-lanceolate to linear. 4
2. Stem leaves long-petioled; the large ones
cordate or ovate. *A. cordifolia*
Stem leaves with the petiole shorter than
the blade, oblong. 3
3. Pappus pale brown; lower stem leaves
deltoid, dentate. *A. diversifolia*
Pappus white; lower stem leaves
elliptic-lanceolate. *A. latifolia*
4. Stem leaves numerous, in 4–8 pairs. 5
Stem leaves less numerous, in 1–3 pairs. 8
5. Lower stem leaves ovate or deltoid,
sharply dentate. *A. diversifolia*
Lower stem leaves lanceolate, entire to
remotely dentate. 6
6. Heads solitary or few. *A. mollis*
Heads several to many. 7
7. Bracts with a distinct tuft of hairs at the
tip. *A. chamissonis*
Bracts not with a tuft of hairs, and with
the tip acute. *A. longifolia*
8. Heads discoid; pappus tawny to
brownish. *A. parryi*
Heads radiate; pappus white to brownish. 9
9. Pappus straw-colored or brownish;
flowering stems without tufts of basal
leaves. 10
Pappus white; basal leaves mostly
present. 11
10. Heads usually several, narrow,
top-shaped. *A. diversifolia*
Heads usually solitary or few, nearly
half-round. *A. mollis*
11. Basal leaves usually petiolate, ovate to
cordate. *A. latifolia*
Basal leaves linear to lanceolate or
oblanceolate. 12
12. Lower leaves mostly more or less
long-petioled. 13
Lower leaves sessile to short-petioled. 15
13. Leaves dentate or denticulate; heads 1–5,
top-shaped. *A. lonchophylla*

- Leaves entire; heads mostly solitary,
half-round. 14
14. Base of stem with a tuft of brown hairs;
leaves lanceolate. *A. fulgens*
- Base of stem without brown hairs; leaves
linear-lanceolate. *A. sororia*
15. Involucres 9–11 mm high; stem leaves
remotely dentate. *A. rydbergii*
- Involucres 10–15 mm high; stem leaves
entire. 16
16. Achenes densely pubescent; involucral
bracts villose-glandular. *A. alpina* var. *ungavensis*
- Achenes glabrous or nearly so; involucral
bracts glabrous, at least above the
middle. *A. louiseana*

Arnica alpina (L.) Olin var. *ungavensis* Boiv.

ALPINE ARNICA

Plants with stems 10–30 cm high. Stem leaves usually 2 or 3 pairs, linear-lanceolate, acuminate, entire. Heads usually solitary; involucre villous and glandular; ligules distinctly toothed at apex. Pappus white. Alpine and arctic tundra; northeastern Boreal forest, southern Rocky Mountains.

The var. *vestita* Hulten. densely soft woolly pubescent, especially on the involucral bracts. Alpine tundra; southern Rocky Mountains.

Arnica chamissonis Less.

LEAFY ARNICA

A species 30–60 cm high, with a very leafy softly hairy stem. Leaves oblong-lanceolate; lower leaves tapering to a winged stalk clasping stem at base, 6–15 cm long; upper leaves stalkless and opposite, usually in several pairs. Flowers lemon yellow, 2–5 cm across, in cluster at the head of the stem. Widespread, but nowhere very common; in moist places; throughout Boreal forest and Peace River district.

Arnica cordifolia Hook. (Fig. 207A)

HEART-LEAVED ARNICA

A species 20–50 cm high, with cordate (heart-shaped) leaves. Basal leaves long-stalked, 2–8 cm long; stem leaves smaller, usually without stalks. Flowers lemon yellow, 3–7 cm across, on the top of the stem. Plentiful; in wooded areas; in Rocky Mountains and Cypress Hills, rarer in Boreal forest.

Arnica diversifolia Greene

LAWLESS ARNICA

A perennial with freely branching, creeping rootstocks; stems solitary or in small tufts, 15–40 cm high, glandular pubescent to subglabrous. Stem leaves 4–8 cm long, ovate or deltoid to elliptic; upper leaves sessile; lower ones wing-petioled. Heads several; ligules 15–20 mm long; bracts glandular. Slopes and creek banks; southern Rocky Mountains.

Arnica fulgens Pursh (Fig. 207B)

SHINING ARNICA

A plant 20–40 cm high. Basal leaves stalked, oblanceolate, 5–8 cm long, usually 3-ribbed, and entire-margined. Stem leaves linear-lanceolate, smaller,



Fig. 207. A, Heart-leaved arnica, *Arnica cordifolia* Hook.; B, shining arnica, *Arnica fulgens* Pursh.

opposite, in 2 or 3 pairs. Flower heads orange yellow, 2–5 cm across, usually solitary on stem, but occasionally 2 or 3. Fairly abundant; in meadows and slightly moister spots on prairie; throughout Prairies and Parklands. Plentiful some years and scarce in others.

Arnica latifolia Bong.

MOUNTAIN ARNICA

Plants with long creeping rhizomes; stems solitary or few together, 10–60 cm high. Basal leaves long-petioled, often on separate sterile shoots; stem leaves 3–10 cm long, ovate or elliptic, more or less dentate, the upper ones sessile, the lower ones petiolate. Heads usually 1–3, top-shaped; ligules 10–15 mm long; bracts 10–15 mm high, more or less glandular pubescent. Moist to wet montane forests; southern Rocky Mountains.

Arnica lonchophylla Greene

SPEAR-LEAVED ARNICA

Plants with dark, scaly, branching rootstocks; stems usually solitary, 15–40 cm high, finely stipitate-glandular, and more or less pubescent. Leaves 5–15 cm long, lance-elliptic to lanceolate, petiolate, mostly finely stipitate-glandular, denticulate to dentate. Heads 1 to several; ligules 1–2 cm long; bracts 8–12 mm high, glandular pubescent. Lakeshores and riverbanks, particularly on calcareous soils; Boreal forest.

Arnica longifolia Eat.

LONG-LEAVED ARNICA

Plants with a short rootstock or branching caudex; stems clustered, usually 20–40 cm high, with many short sterile shoots at the base. Leaves 5–12 cm long, narrowly lanceolate or lance-elliptic, the lower ones connate or nearly so, the others shortly petiolate to connate. Heads several to numerous; ligules 1–2 cm long; bracts 7–10 mm high, glandular puberulent. Along montane and alpine creeks; southern Rocky Mountains.

Arnica louiseana Farr

ROCK ARNICA

Perennial plants with long, creeping, densely scaly rootstocks; stems solitary or several together, 10–25 cm high, softly villous, and usually more or less glandular. Leaves 2–6 cm long, oblanceolate to elliptic, entire or distantly denticulate, mostly subglabrous to sparsely villous, and more or less glandular. Heads solitary; ligules 1–2 cm long; bracts about 1 cm high, glabrous above the middle, somewhat pubescent toward the base. On rock cliffs and shale slides at high altitudes; southern Rocky Mountains.

Arnica mollis Hook.

CORDILLERAN ARNICA

Plants with branching rootstocks; stems 20–40 cm, puberulent to long pubescent, and glandular. Stem leaves 3–6 cm long, ovate to elliptic, irregularly dentate to entire, sessile or the lower ones short petiolate. Heads solitary or few; ligules 15–25 mm long; bracts 10–15 mm high, pubescent below, glandular toward the tips. Wet, boggy areas; southern Rocky Mountains. Plants with very leafy stems and 5–12 pairs of stem leaves, as compared with 3–5 pairs in the typical variety, are distinguished as var. *aspera* (Greene) Boiv.

Arnica parryi Gray

NODDING ARNICA

Plants with creeping rhizomes; stems usually solitary, 20–40 cm high, often woolly-pubescent toward the base, somewhat glandular above. Stem

leaves 5–15 cm long, lanceolate to lance-ovate, the lowermost petiolate. Heads usually several, nodding in bud; discoid bracts 10–15 mm high, usually glandular to glandular pubescent. Mountain meadows; southern Rocky Mountains.

Arnica rydbergii Greene

NARROW-LEAVED ARNICA

Plants with short, branched, scaly rootstocks; stems 10–30 cm high, glandular pubescent to subglabrous. Basal leaves petiolate, oblanceolate or spatulate, to 7 cm long; stem leaves 3–10 cm long, lanceolate to spatulate, entire or nearly so. Heads solitary or few; rays 1–2 cm long; bracts 10–15 mm high, glandular, and sparsely pubescent to subglabrous. High slopes and shale slides; southern Rocky Mountains.

Arnica sororia Greene

TWIN ARNICA

Plants with slender, freely rooting rootstocks; stems 10–40 cm high, glandular pubescent. Basal leaves lanceolate to narrowly lanceolate, glandular pubescent. Heads solitary, rarely 2 or 3; ligules 15–25 mm long; bracts 10–15 mm high, sparingly pubescent and glandular. Montane grasslands and forest openings; southern Rocky Mountains.

Artemisia wormwood

Biennial or perennial shrubs or herbs, usually with a conspicuous odor. Leaves alternate, varying from entire to much dissected, green and hairless to dense white woolly. Flowers small, usually in spike-like panicles, discoid with only tubular florets. Seeds bearing no pappus. The wind-borne pollen of some species often causing hay fever.

- 1. Shrubs with woody bases and branches. 2
 Herbs or subshrubs with bases not woody
 and branches usually herbaceous. 3
- 2. Leaves entire, lanceolate to oblanceolate. *A. cana*
 Leaves 3-lobed at the tip, cuneate. *A. tridentata*
- 3. Plants glabrous or nearly so. 4
 Plants hairy, silky, or woolly. 6
- 4. Leaves mostly entire and undivided,
 rarely 3-cleft. *A. dracunculus*
 Leaves pinnatifid and divided. 5
- 5. Leaves cleft into very narrow linear
 divisions. *A. campestris*
 Leaves cleft into lanceolate toothed
 divisions. *A. biennis*
- 6. Leaves hairy or silky hairy, but not
 woolly. 7
 Leaves white woolly, at least on
 underside. 8
- 7. Leaves 2 or 3 times divided. *A. frigida*
 Leaves only once or twice divided. *A. absinthium*

8. Leaves entire or at most coarsely lobed. 9
 Leaves much divided; lobes narrow,
 mostly linear. 11
9. Leaves white woolly on both sides. *A. ludoviciana*
 Leaves white woolly only on lower side. 10
10. Leaves entire or sharply toothed. *A. tilesii* ssp. *unalaschensis*
 From coarse, woody roots; leaves 5–10
 cm long and usually with rolled
 margins. *A. longifolia*
11. Ultimate leaf segments usually less than 1
 mm wide, entire or subentire. 12
 Ultimate leaf segments usually more than
 1 mm wide, usually toothed or lobed. 13
12. Plants usually sterile; herbaceous; ultimate
 leaf segments often 1- or
 2-toothed. *A. pontica*
 Plants usually fertile; subshrubs; ultimate
 leaf segments entire. *A. abrotanum*
13. Pubescence of leaves loosely woolly or
 absent; most or all of the leaves basal. *A. norvegica* var. *saxatilis*
 Pubescence densely woolly; most of the
 leaves cauline. 14
14. Leaves deeply dissected to near midrib
 into linear divisions with rolled
 margins. *A. michauxiana*
 Leaf segments doubly divided with broad
 ultimate divisions. *A. vulgaris*

Artemisia abrotanum L.

LEMONTREE

A perennial plant, somewhat shrubby at the base, 50–200 cm high. Leaves 3–6 cm long, lemon-scented when bruised, thinly tomentose below, green and subglabrous above. Inflorescence ample; involucre 2–3.5 mm high. Introduced; occasionally cultivated and locally escaped.

Artemisia absinthium L. (Fig. 208)

ABSINTHE

A somewhat shrubby plant 50–100 cm high. Stems with many branches, finely hairy. Leaves 5–10 cm long, several times divided into ovate to oblong segments, finely grayish hairy. Inflorescence a large, many flowered, somewhat spike-like panicle; heads stalked, drooping, about 5 mm across. Found in waste places where it has escaped from gardens.

Artemisia biennis Willd.

BIENNIAL WORMWOOD

An annual or biennial plant 30–100 cm high, with hairless coarse stems, usually reddish for about half their length. Early leaves twice or three times dissected into toothed segments, usually forming a rosette on the ground. Stem leaves and upper leaves 2–7 cm long, once or twice divided into narrow segments, and hairless. Flowers in short compact spikes in axils of upper leaves, forming a dense leafy spike-like panicle. Common; in moist places, slough margins, roadsides, and cultivated fields; throughout the Prairie Provinces.




Fig. 208. Absinthe, *Artemisia absinthium* L.

Artemisia campestris L.

PLAINS WORMWOOD

A very variable glabrous to pubescent perennial or biennial, usually 10–80 cm high. Most of the leaves in a basal rosette, 3–10 cm long, 2 or 3 times divided into narrow segments. Stem leaves, when present, sessile and smaller than basal leaves. Inflorescence a leafy panicle with numerous heads; involucre 2–4.5 mm high. Three varieties are distinguished:

- 1. Stems usually 10–30 cm high; foliage glabrous or nearly so; most leaves basal. var. *wormskjoldii* (Bess.) Cronq.
 Stems higher; foliage more pubescent; stems leafier. 2
- 2. Stems 30–80 cm high; leaves green, sparsely pubescent, mostly cauline; basal leaves withering. var. *scouleriana* (Bess.) Cronq.
 Stems 20–40 cm high; leaves grayish pubescent. var. *douglasiana* (Bess.) Boiv.

The var. *wormskjoldii* occurs on subarctic shores and slopes in northern fringe of Boreal forest. Syn.: *A. borealis* Pall. The var. *scouleriana*, usually biennial, found in sandy woods, shores, and openings in Boreal forest and Parklands. Syn.: *A. canadensis* Michx.; *A. caudata* Michx. The var. *douglasiana* is the common, usually short-lived perennial of grasslands and disturbed areas in Prairies and Parklands. Syn.: *A. camporum* Rydb.; *A. bourgeauiana* Rydb.

Artemisia cana Pursh

HOARY SAGEBUSH

A shrub with somewhat gnarled and twisted shreddy-barked woody stems 30–150 cm high. Leaves 1–3 cm long, silvery hairy on both sides, linear to linear-lanceolate, usually entire, or rarely with toothed points. Yellow flowers crowded into a leafy panicle. Very common; on lighter soils; Prairies, rare in Parklands.

Artemisia dracunculus L.

LINEAR-LEAVED WORMWOOD

An entirely hairless perennial 50–100 cm high. Leaves 1–6 cm long, narrowly linear, usually entire, but basal leaves occasionally cleft or divided. Inflorescence in a leafy compound panicle. Common; on dry prairie; throughout Prairies.

Artemisia frigida Willd. (Fig. 209)

PASTURE SAGE

A densely silky hairy silvery gray perennial with a somewhat woody base, 15–50 cm high. Leaves 1–3 cm long, 2 or 3 times divided into linear segments. Plants having a distinct odor when handled. Numerous yellowish heads borne in terminal somewhat leafy racemes. Common; on unforested land and in overgrazed pastures; throughout the Prairies and Parklands. Abundant on prairies, its unpalatability enabling it to increase with heavy grazing at the expense of more palatable plants; therefore a useful indicator of overgrazing.

Artemisia longifolia Nutt.

LONG-LEAVED SAGE

A perennial from a coarse woody much-branched root crown, with densely white woolly stems 40–80 cm high. Leaves white on both sides, 5–10



Fig. 209. Pasture sage, *Artemisia frigida* Willd.

cm long, linear or linear-lanceolate, and often with rolled margins. Upper sides of leaves sometimes losing the tomentum or woolliness with age. Inflorescence a narrow leafy spike-like panicle. Found occasionally; in saline areas, badlands, and shaly outcrops; throughout Prairies.

Artemisia ludoviciana Nutt. var. *ludoviciana*

PRAIRIE SAGE

A white woolly often much-branched perennial 15–60 cm high, very variable in form and size. Leaves 1–7 cm long, white woolly on both sides, but usually slightly less so on upper side. Lower leaves oblanceolate, usually entire, but occasionally with a few lobes; upper leaves lance-linear and entire. Inflorescence in dense axillary spikes, making a leafy panicle; flower heads brownish. Very common; on prairie, especially where conditions are a little moist; throughout the Prairie Provinces. The var. *gnaphalodes* (Nutt.) T. & G., slender sage, is a low perennial from creeping rootstocks, 15–40 cm high, but often growing prostrate on the ground. Stems white woolly and slender. Leaves white woolly or more often pale yellowish woolly on both sides, 2–5 cm long, narrowly linear-lanceolate, usually entire, but the lower ones sometimes toothed and usually conduplicate (folded lengthwise). Flower heads small, in narrow terminal spike-like inflorescences. Common; along slough margins and moister prairie; throughout Prairies and Parklands.

Artemisia michauxiana Besser

MICHAUX'S SAGE

A perennial from a much-branched root crown, 20–50 cm high. Leaves 1–5 cm long, green above, white woolly beneath, and cleft into narrow linear sometimes toothed segments, with margins often rolled. Rather large flower heads in a sparingly leaved terminal spike-like panicle. Found on hillsides and along rivers; southern Rocky Mountains.

Artemisia norvegica Fries var. *saxatilis* (Besser) Jepson

MOUNTAIN SAGE

Plants with a branching caudex; stems 15–50 cm high, loosely villous to subglabrous. Leaves 2–10 cm long, pinnately dissected, green and glabrous above, green or more or less tomentose below. Inflorescence spike-like; heads 4–6 mm high. Alpine slopes; southern Rocky Mountains.

Artemisia pontica L.

ROMAN WORMWOOD

Perennial with a creeping rootstock, often somewhat shrubby, 40–100 cm high. Leaves 1–3 cm long, fragrant, white tomentose on both sides, or glabrate above, 2- or 3-pinnate, with the ultimate segments usually less than 1 mm wide. Inflorescence rarely present; if so, rather narrow, elongate, with involucre 2–3 mm high, more or less tomentose. Introduced and cultivated; occasionally escaped.

Artemisia tilesii Ledeb. ssp. *unalaschensis* (Besser) Hulten

HERRIOT'S SAGE

An erect perennial from a coarse woody base, 30–90 cm high, with a simple fine-hairy stem. Leaves usually linear, 5–15 cm long, sometimes with a few sharp linear teeth, densely white woolly beneath, and smooth and green above. Numerous heads in a dense spike-like panicle. Quite unusual; but found in badlands and river breaks; southern Rocky Mountains. Syn.: *A. herriotii* Rydb.

An evergreen shrub 30–150 cm high, strong-scented, with a single short trunk. Leaves 1–4 cm long, wedge-shaped; the apex usually with 3 blunt teeth, but sometimes 4- to 9-toothed. Inflorescence equaling or exceeding the sterile branches, panicle-like, narrow to fairly broad; heads 3–4 mm high. Grasslands; southern Rocky Mountains; very rare, possibly introduced in Prairie.

Artemisia vulgaris L.

A coarse weedy perennial 30–100 cm high, with a much-branched stem. Leaves 2–8 cm long, dark green above, and densely white woolly below, several times divided; segments somewhat broad and oblanceolate. Numerous heads in erect leafy panicles. An introduced plant; escaped from gardens.

Aster aster

Perennial or biennial herbs with alternate leaves and usually showy blue or white radiate heads. Ray florets and disk florets perfect (bisexual). Ray florets usually 10–50 with bracts in several series. Pappus of capillary bristles or hairs. Asters usually flower in late summer and early fall.

- 1. Plants annual, with fibrous roots. 2
 Plants perennial, with rootstocks or root
 crowns. 3
- 2. Ray florets lacking or inconspicuous. *A. brachyactis*
 Ray florets purple; involucre bracts glandular sticky. *A. canescens*
- 3. Involucres and usually branches of the
 inflorescence glandular. 4
 Involucres and branches not glandular. 9
- 4. Leaves to 20 cm long and 10 cm wide,
 coarsely toothed or closely serrate. 5
 Leaves smaller, entire or distantly serrate. 6
- 5. Lower leaves petioled, cordate or subcordate at the base, serrate. *A. macrophyllus*
 Lower leaves sessile, somewhat clasping,
 coarsely toothed to subentire. *A. conspicuus*
- 6. Leaves 3–8 cm long, 3–8 mm wide; ray
 florets 5–8 mm long. 7
 Leaves 5–15 cm long, 8–20 mm wide; ray
 florets 10–20 mm long. 8
- 7. Plants glabrous except for the glandular
 pubescence; leaves rather thick,
 grass-like. *A. pauciflorus*
 Plants glandular pubescent more or less
 throughout; leaves not thick or
 grass-like. *A. campestris*
- 8. Ray florets numerous, usually 45–100,
 10–20 mm long; leaves entire. *A. novae-angliae*

- Ray florets fewer, usually 20–40, 10–15 mm long; leaves serrate. *A. modestus*
9. Heads solitary or 1–3 on a stem. 10
 Heads few to many on a stem. 11
10. Stem leaves greatly reduced; stem and leaves densely pubescent; plants with a root crown. *A. alpinus*
 Stem leaves not greatly reduced; stem and leaves sparsely pubescent; plants strongly rhizomatous. *A. sibiricus*
11. Basal and lower stem leaves petioled or narrowed to a petiolar base. 12
 Basal and lower stem leaves sessile or subsessile. 16
12. Lower leaves cordate or subcordate to rounded at the base, usually ovate. 13
 Lower leaves not cordate or rounded at the base, usually lanceolate to linear-lanceolate. 14
13. Lower leaves narrowly ovate; stem pubescent in lines above middle. *A. ciliolatus*
 Lower leaves lanceolate; stem glabrous except in the inflorescence. *A. maccallae*
14. Upper stem leaves clearly serrate, 10–25 mm wide; heads 25–30 mm across; some foliaceous involucre bracts often present. *A. subspicatus*
 Upper stem leaves entire or obscurely toothed, 5–10 mm wide; heads 15–25 mm across; foliaceous involucre bracts absent. 15
15. Stem leaves linear, the upper ones not reduced in size; branches of the inflorescence strongly ascending. *A. adscendens*
 Stem leaves linear-lanceolate, the upper ones reduced in size; branches of the inflorescence spreading. *A. occidentalis*
16. Plants glabrous, except sometimes in the inflorescence; leaves clasping, glaucous. *A. laevis*
 Plants more or less pubescent. 17
17. Leaves silvery silky on both sides, entire. *A. sericeus*
 Leaves not silvery silky. 18
18. Pappus double, with the inner series of firm long bristles, the outer series of bristles about 1 mm long. *A. umbellatus*
 Pappus single. 19

19. Involucre usually with some foliaceous
outer bracts. *A. eatoni*
Involucre without foliaceous bracts. 20
20. Stem leaves auriculate clasping; plant
rough pubescent. *A. puniceus*
Stem leaves sessile or somewhat clasping;
plant not rough pubescent. 21
21. Leaves narrowly linear, seldom more than
6–7 mm wide. 22
Leaves broadly linear to linear-lanceolate
or lanceolate-ovate, usually more than
10 mm wide. 24
22. Plants very slender; stems arising from
slender rhizomes; leaves glabrous;
stem pubescent in lines below the leaf
bases. *A. junciformis*
Plants coarse; leaves and stems more or
less densely pubescent. 23
23. Heads solitary or few at the end of
branches, 20–25 mm across. *A. falcatus*
Heads numerous, more or less secund
along the branches, 10–15 mm across. *A. ericoides*
24. Ray florets 10–15 in a head. 25
Ray florets 20–40 in a head. 26
25. Heads 20–40 mm across; leaves in the
inflorescence ovate-lanceolate. *A. engelmannii*
Heads 12–18 mm across; leaves in the
inflorescence narrowly linear. *A. lateriflorus*
26. Leaves serrate, 10–35 mm wide; involucre
4–5.5 mm high; ray florets white. *A. simplex*
Leaves entire to subentire, seldom more
than 18 mm wide; involucre 5–8 mm
high; ray florets blue. *A. hesperius*

Aster adscendens Lindl.

WESTERN ASTER

A slender erect species 15–50 cm high, with the stem sometimes decumbent at base. Leaves 2–7 cm long, with the lower ones short-stalked and spatulate, the upper ones stalkless and more or less clasping and linear. Heads about 25 mm across, not very numerous, in a panicle. Bracts oblanceolate, blunt-pointed, in series of 3–5. Found occasionally; in valleys and favored localities; Prairies and southern Rocky Mountains.

Aster alpinus L.

ALPINE ASTER

Stems 5–20 cm high, pilose, and somewhat glandular throughout. Basal leaves numerous, 2–5 cm long, oblong to lanceolate, rounded at the apex. Heads solitary, 3–5 cm across; ligules violet to whitish; involucre pilose. Mountain slopes and forests; southern Rocky Mountains.

Aster brachyactis Blake

RAYLESS ASTER

A slender purplish-stemmed somewhat branching annual 15–60 cm high. Leaves linear, 2–7 cm long, without hairs except for some on margin. Flower heads numerous, often almost hiding the foliage and appearing to be mainly composed of white hairy pappus, 8–12 mm broad in a raceme; the few florets purple and occasionally with a few rudimentary ligulate florets. Involucral bracts very narrow, often purplish. Common; in saline soil and moist places; throughout most of the Prairies and Parklands. Syn.: *Brachyactis angusta* (Lindl.) Britton; *A. angustus* (Lindl.) T. & G.

Aster campestris Nutt.

MEADOW ASTER

Plants with slender creeping rootstocks; stems 10–50 cm high, simple or branched, leafy, densely stipitate glandular. Stem leaves linear to linear spatulate, 2–5 cm long, almost uniform in size, sessile and often slightly clasping, entire, glabrous to somewhat hispid. Heads 1 to many, 15–20 mm wide, in a narrow cyme or panicle; the ligules 5–8 mm long, violet to purple; involucral bracts densely glandular. Montane grasslands; southern Rocky Mountains.

Aster canescens Pursh

CANESCENT ASTER

A low, branching biennial, from a taproot, 10–30 cm high; stems covered with short fine hairs. Leaves 10–35 mm long, with the lower ones spatulate or oblanceolate, the upper ones often linear, varying from entire to slightly toothed. Teeth and apex of leaves bearing a short bristle-like tooth. Flowers 15–25 mm across, bluish purple, usually very numerous. Common; especially along roadsides; Prairies. Syn.: *Machaeranthera canescens* (Pursh) A. Gray; *M. pulverulenta* (Pursh) A. Gray.

Aster ciliolatus Lindl. (Fig. 210)

LINDLEY'S ASTER

A stout erect-stemmed species 30–75 cm high. Leaves thick, sometimes slightly hairy on veins. Basal leaves ovate or cordate, 5–10 cm long, with long stalks; the lower stem leaves with wing-margined stalks, the upper ones often stalkless. Flower heads usually few, blue to violet, 15–30 mm across. Common; in woodlands; throughout Parklands and Boreal forest. Syn.: *A. lindleyanus* T. & G.

Aster conspicuus Lindl. (Fig. 211)

SHOWY ASTER

An erect coarse species 40–100 cm high, with rough and hairy stems. Leaves ovate to obovate, coarsely toothed, rough on upper surface and finely hairy below, 10–15 cm long. Inflorescence a large corymb of violet to blue flowers, each about 4 cm across; bracts and flower stems glandular. Very palatable to livestock. Common; in woodlands; Parklands and Boreal forest, except in the eastern part, and also found abundantly in Cypress Hills.

Aster eatoni (A. Gray) Howell

EATON'S ASTER

A branching species 30–80 cm high. Leaves linear to lance-linear, 5–10 cm long, stalkless, and entire, but not clasping. Outer bracts large and leaf-like. Heads 20–25 mm across, with pinkish or white ray florets. Not common; has been found in river valleys; southern Rocky Mountains. Syn.: *A. mearnsii* Rydb.



HERBARIUM OF THE RESEARCH INSTITUTE
 SIVTSEVSKAYA, SAKH.
 PLANTS OF SIBERIA, and the Institute of
 Botany, Siberian Academy of Sciences.
 LOCATION: Siberian Academy of Sciences
 TOMSK, Siberia, 1950
 FIELD NO. 10000, DATE OF COLLECTION 1950
 VARIETY: *Aster ciliolatus* Lindl.

Fig. 210. Lindley's aster, *Aster ciliolatus* Lindl.




Fig. 211. Showy aster, *Aster conspicuus* Lindl.

Aster engelmannii (Eat.) Gray

ELEGANT ASTER

Plants with woody root crowns; stems tufted, 30–100 cm high, branched in the inflorescence. Leaves 4–10 cm long, elliptic to oblong, rounded or narrowed at the base, sessile, entire or nearly so, glabrous above except along the midrib, sparsely pilose on the veins below. Heads few to many, mostly terminal on short branches, 2–4 cm across; involucre 7–10 mm high, glabrous or somewhat pubescent. Mountain meadows; southern Rocky Mountains.

Aster ericoides L.

MANY-FLOWERED ASTER

A branching perennial from a thick tufted rootstock, 20–60 cm high. Stems finely hairy. Leaves linear to narrowly linear-lanceolate, 1–5 cm long, entire-margined. Heads numerous, 8–12 mm across, white, usually on one side of the recurved branches. Common; on open prairie, roadsides; throughout Prairies and Parklands. Syn.: *A. pansus* (Blake) Cronquist; *A. multiflorus* Ait.; *A. adsurgens* Greene.

Aster falcatus Lindl.

WHITE PRAIRIE ASTER

A much-branched perennial, from running rootstocks, 30–80 cm high. Stem rough hairy. Leaves linear to linear-oblong, 1–4 cm long, stalkless, and entire-margined. Inflorescence of a few or single heads at ends of branches; flowers white, 20–25 mm across. Bristle-tipped bracts almost equal in length, with the outer ones at least as high as the involucre. Fairly common; throughout Prairies and Parklands. Syn.: *A. commutatus* (T. & G.) A. Gray.

Aster hesperius A. Gray

WILLOW ASTER

A species 40–80 cm high, with a much-branching stem. Leaves narrow, 5–15 cm long, sometimes entire-margined and sometimes somewhat toothed. Flower heads numerous, in a branching panicle, varying from white to violet or pink. A very variable species, known under many names. Common; along stream banks, ditches, and moist spots; throughout the Prairie Provinces. Syn.: *A. osterhoutii* Rydb.

Aster junciformis Rydb.

RUSH ASTER

A slender-stemmed, erect, little-branched species 20–60 cm high. Leaves 2–8 cm long, narrow, and linear, usually with entire margins, stalkless, and having clasping stems. Inflorescence an open panicle; flower heads white, 15–20 cm across. Fairly common; in swamps and bogs; throughout Parklands, Boreal forest, and in the Cypress Hills.

Aster laevis L. (Fig. 212)

SMOOTH ASTER

A stout-stemmed hairless species 30–100 cm high. Leaves thick, 2–10 cm long, often toothed, ovate or lanceolate, and hairless, with the basal leaves on wing-margined stalks but the upper ones stalkless and often clasping. Numerous flower heads 2–3 cm across, with blue ray florets; pappus somewhat tawny-colored. Common; on moist prairie, around bluffs or clumps of shrubby, and in open woodlands; throughout the Prairie Provinces.

Aster lateriflorus (L.) Britt.

WOOD ASTER

A perennial with short stout rhizomes or a branching caudex; stems several, 30–90 cm high, curly villous to subglabrous. Leaves scabrous to subglab-



HERBARIUM OF THE UNIVERSITY OF CALIFORNIA,
 BERKELEY, CALIF.

NO. 1010 Aster laevis L.
 COLLECTED BY J. LOGAN
 CALIFORNIA 1891
 HERBARIUM OF THE UNIVERSITY OF CALIFORNIA
 BERKELEY, CALIF.

Number	1010
Collector	J. Logan
Date	1891
Locality	California
Plant	Aster laevis L.
Other	

Fig. 212. Smooth aster, *Aster laevis* L.

rous above, glabrous below, except on the midrib; the principal leaves 5–12 cm long, to 3 cm wide, broadly linear or linear-lanceolate. Inflorescence mostly branched, or rarely simple; heads 10–15 mm across, with rays white or purple-tinged; involucre 4–5.5 mm high. Open woodlands, openings, or beaches; southeastern Parklands and Boreal forest.

Aster maccallae Rydb.

MACCALLA'S ASTER

Plants with creeping rootstocks; stems 30–60 cm high, often purplish; glabrous below the inflorescence, pubescent in lines above. Lower leaves 8–15 cm long; petioles winged; blades lanceolate, usually serrate, glabrate or nearly so. Inflorescence with few heads; ligules about 15 mm long, blue or purplish; involucre 8–10 mm high, glabrous. Open areas, along mountain streams, and in grassland; southern Rocky Mountains.

Aster macrophyllus L.

WHITE WOOD ASTER

A tufted perennial with a zigzag twisted brittle stem 40–100 cm high. Leaves thin, slender-stalked, coarsely toothed, with the lower ones cordate-based, 2–10 cm long. Flower heads usually white, 2–3 cm across, usually in a flattish wide corymb. Found occasionally; in extreme southeastern Boreal forest.

Aster modestus Lindl.

LARGE NORTHERN ASTER

Plants with a long creeping rootstock; stems usually solitary, 30–80 cm high, branching in the inflorescence, densely glandular above or throughout. Leaves rather uniformly 6–15 cm long, lanceolate, entire to sharply serrate, with the base more or less clasping. Heads 1 to many, 2–3 cm wide; ligules about 1 cm long, purple or violet; involucre about 7 mm high, densely stipitate-glandular. Wet areas and bogs; western Parklands and Boreal forest.

Aster novae-angliae L.

NEW ENGLAND ASTER

A tall erect stout-stemmed species 60–200 cm high. Leaves lanceolate to oblong, entire, 3–12 cm long, stalkless, and clasping stem by a cordate or an eared base. Flower heads numerous, 2–5 cm across, with reddish purple to violet purple ray florets in compact clusters at ends of branches. Fairly common; in moist woodlands and low ground; in eastern Parklands and Boreal forest.

Aster occidentalis (Nutt.) T. & G.

WESTERN MOUNTAIN ASTER

Plants with creeping rootstocks; stems mostly less than 50 cm high, glabrous below, pubescent in lines above. Lower leaves 4–12 cm long, oblanceolate, tapering to a petiolate ciliate base. Heads 1 to few, about 25 mm across; ligules 6–10 mm long; involucre about 6 mm high. Mountain meadows; Rocky Mountains.

Aster pauciflorus Nutt.

FEW-FLOWERED ASTER

A hairless species, growing from creeping rootstocks, much-branched, 15–50 cm high. Leaves entire-margined, somewhat fleshy, 2–8 cm long, with the upper ones linear and stalkless, the lower ones linear-lanceolate and stalked. Flower heads few, 15–25 mm across, with blue or white ray florets. Rare; found in saline soil; Prairies and Parklands.

Aster puniceus L.

PURPLE-STEMMED ASTER

A stout purplish-stemmed branching species from a thick rootstock, 60–200 cm high. Leaves 7–15 cm long, lanceolate to oblong-lanceolate, often hairy on midrib below, usually sharp-toothed. Flower heads numerous, 25–35 mm across, with light violet to pale purple ray florets. Fairly common; in swamps and marshlands; throughout Boreal forest.

Aster sericeus Vent.

WESTERN SILVERY ASTER

A slender-stemmed branching species 30–60 cm high. Leaves 10–35 mm long, covered with dense silvery white silky hairs on both sides; the lower leaves with short winged stalks and the upper ones stalkless, oblanceolate to oblong. Numerous heads about 35 mm across, with reddish violet to violet blue ray florets and tawny pappus. Rare; southeastern Parklands.

Aster sibiricus L.

ARCTIC ASTER

Plants with slender creeping rootstocks; stems 3–30 cm high, simple or with few branches, usually purple, sparingly pubescent. Leaves 1–6 cm long, obovate or elliptic, entire to serrate above the middle, with the lower ones short petiolate, the upper ones sessile and narrowly clasping. Heads solitary to few in a leafy cymose panicle, with the lateral heads on axillary peduncles 2–3 cm across; ligules 7–12 mm long, purple to violet or sometimes whitish; involucre 5–10 mm high. Open slopes, creeks, and riverbanks; Rocky Mountains, Peace River.

Aster simplex Willd.

SMALL BLUE ASTER

Plants with stout, long creeping rootstocks; stems stout, 60–150 cm high, glabrous below, pubescent in lines above. Leaves 8–15 cm long, lanceolate to linear, glabrous on both sides, serrate, rarely entire, sessile, narrowly clasping. Heads more or less numerous in a leafy inflorescence; ligules 5–10 mm long; involucre 3–6 mm high, glabrous, with ciliolate margins. Moist areas and shores; eastern Parklands and Boreal forest.

Aster subspicatus Nees

LEAFY-BRACTED ASTER

An erect species 30–100 cm high. Lower leaves 10–15 cm long, oblanceolate, entire-margined, and with wing-margined stalks; upper leaves shorter with clasping bases. Flowers violet, 15–25 mm across. A mountain species, which may be found in Cypress Hills. Syn.: *A. frondeus* (A. Gray) Rydb. Smaller plants, usually less than 25 cm high, with leaves narrowly oblanceolate and bracts purple-tipped are var. *apricus* (Gray) Boiv. Syn.: *A. foliaceus* Lindl. var. *apricus* Gray.

Aster umbellatus Mill.

FLAT-TOPPED WHITE ASTER

A tall erect species, from a woody rootstock, 60–200 cm high. Leaves 6–15 cm long, narrowly elliptic to lanceolate, tapering to apex and at base to a short stalk. Very numerous flower heads 12–20 mm across, with white ray florets in a large flat-topped terminal cluster. Pappus has an outer row of short stiff bristles, the inner bristles being long and hair-like. Common; in moist woodlands; throughout Manitoba. Syn.: *Doellingeria umbellatus* (Mill.) Nees. The var. *pubens* Gray differs from the species by the hairy undersides of leaves. Found

occasionally in woodlands; eastern Parklands and Boreal forest. Syn.: *Doellingeria pubens* (A. Gray) Rydb.

Bahia picradeniopsis

Bahia oppositifolia (Nutt.) DC.

PICRADENIOPSIS

A much-branched perennial, from creeping rootstocks, 10–25 cm high, with a somewhat woody base. Stem very finely hairy and very leafy. Leaves 1–3 cm long, gray green, very finely hairy, and often several times divided into narrow linear segments. Yellow flower heads, about 12 mm across, at ends of branches. Many tubular florets, but only a few short ray florets. Quite unusual, but has been found on a few occasions in Prairies as a rather persistent weed in cultivated land. It occurs on saline flats and dry plains in the USA. Syn.: *Picradeniopsis oppositifolia* (Nutt.) Rydb.

Balsamorhiza balsamroot

Balsamorhiza sagittata (Pursh) Nutt.

BALSAMROOT

A low perennial from a thick edible spindle-shaped root (often more than 5 cm thick) that exudes a balsam or sticky substance with a turpentine-like odor. Long-stalked leaves mostly basal, 10–25 cm long, varying in shape from cordate to hastate or sagittate (arrow-shaped). White woolly on both sides, densely below, sparsely above. Flowers bright yellow, usually borne singly on stems about 30 cm long arising from root crown. Bracts lanceolate and densely white woolly; seeds without pappus. Fairly plentiful; on hillsides and prairie; southern Rocky Mountains.

Bidens beggarticks

Annual herbs, with leaves usually opposite. Heads discoid or radiate, yellow or orange. Achenes (seeds) bearing 2 or 4 retrorsely (downward-pointing) barbed awns, by which the seed attaches to coats of animals or clothing and is disseminated.

1. Plants aquatic; the submerged leaves filiform-dissected. *B. beckii*
Plants mud or land plants, not aquatic. 2
2. Ray florets conspicuous; leaves stalkless and clasping. *B. cernua*
Ray florets absent or inconspicuous; leaves usually with short stalks, not clasping. 3
3. Leaves simple, deeply incised or 3-lobed. *B. tripartita*
Leaves 1- to 3-pinnatifid, with the terminal leaflet usually petioled. *B. frondosa*

Bidens beckii Torr.

WATER-MARIGOLD

A perennial, with the submerged leaves finely filiform-dissected and the emergent leaves simple, lanceolate, sessile, and serrate. Heads terminal, soli-

tary, about 1 cm wide; ligules 10–15 mm long; outer involucre bracts several; pappus of 3–6 awns, longer than the achene. In shallow waters; eastern Parklands and Boreal forest. Syn.: *Megalodonta beckii* (Torr.) Greene.

Bidens cernua L.

SMOOTH BEGGARTICKS

An erect hairless annual 30–80 cm high. Leaves opposite, 5–15 cm long, toothed, linear-lanceolate, stalkless, clasping the stem, somewhat paler on underside. Flower heads usually nodding, 20–35 mm across; ray florets conspicuous. Long outer bracts usually as long as or longer than ray florets and reflexed. Common; in water and very wet soil; throughout almost the entire Prairie Provinces. Syn.: *B. glaucescens* Greene.

Bidens frondosa L.

COMMON BEGGARTICKS

An erect often branching species 30–100 cm high, usually with a purplish stem. Leaves slender-stalked, 5–10 cm long, usually divided into 3 or 5 lanceolate segments, slightly hairy below, and toothed. Flowers orange, 12–20 mm across, with 4–8 large outer involucre bracts. Fairly common; in wet places and along stream banks; southeastern Parklands and Boreal forest. The var. *puberula* Wieg. is coarser and has 10–16 outer bracts.

Bidens tripartita L.

TALL BEGGARTICKS

Annual plants with stems 30–200 cm high, glabrous or nearly so. Leaves simple, usually deeply 3-lobed, serrate, 3–15 cm long. Heads erect; disk about 10–25 mm wide; pappus of 2–4 awns. Introduced; very rare; southwestern Prairies.

Boltonia boltonia

Boltonia asteroides (L.) L'Hér.

BOLTONIA

A rather stout-stemmed perennial 60–200 cm high. Leaves linear to lanceolate, 5–8 mm long, without stalks. Flower heads numerous, radiate, 20–35 mm across; ray florets varying from white to pink. Achenes bearing a pappus of short scales, often with 2–4 slender bristles, differing from asters, which have a hairy pappus. Rare; has been found in moist soil; eastern Parklands and Boreal forest.

Brickellia brickellia

Brickellia grandiflora (Hook.) Nutt.

LARGE-FLOWERED BRICKELLIA

Perennial plants with long spindle-shaped rootstocks; stems 30–70 cm high, minutely hirsute throughout. Leaves 3–12 cm long, deltoid to cordate, toothed. Heads several to many, discoid; flowers creamy or greenish white; disks 8–12 mm across and high. Wet places and rocky areas; southern Rocky Mountains.

Carduus plumeless thistle

Carduus nutans L.

NODDING THISTLE

A branching biennial 60–100 cm high. Leaves deeply divided, lanceolate, 7–15 cm long, very prickly. Flower heads borne singly on long stems, nodding

or drooping, 35–65 mm across, purple, rarely white. Involucral bracts in many series, each with a prominent midrib prolonged into a spine. Pappus hairs about 25 mm long, white, roughed but not feathery, thus distinguishing this genus from *Cirsium*, which has plumose or feathery pappus hairs. An introduced weed found, though rarely, at various widely separated locations.

Centaurea cornflower

Plants somewhat resembling thistles, with a globular involucre, but pappus hairs either very short or absent.

1. Bracts of involucre definitely spine-tipped; flowers cream or yellow. 2
 Bracts of involucre not definitely spine-tipped; flowers pink or purple. 3
2. Stems winged by extensions of leaf bases (decurent); spines of involucre about 12 mm long. *C. solstitialis*
 Stems not winged; spines of involucre not 12 mm long. *C. diffusa*
3. Most stem leaves deeply cleft into long narrow segments; bracts of involucre fringed at end. *C. maculosa*
 Most stem leaves entire or dentate; bracts of involucre not fringed. *C. repens*

Centaurea diffusa Lam.

DIFFUSE KNAPWEED

A much-branched introduced annual weed 15–60 cm high, sometimes with a fine web-like covering on stems. Leaves once or twice pinnately divided into very narrow segments, some of the uppermost ones entire. Numerous flowering heads in terminal panicles, usually pale yellowish or cream, about 1 cm high. Bracts of involucre spiny-margined, with a terminal spine about 3 mm long. This weed, though not reported in the area, should be looked for because it is a noxious weed in south central British Columbia.

Centaurea maculosa Lam.

SPOTTED KNAPWEED

An introduced biennial species 30–100 cm high; not as much branched as the preceding species. Leaves pinnately divided into narrow lobes, except those of the inflorescence, which are smaller and entire. Flower heads on long stems; flowers purplish, occasionally white. Bracts of involucre tipped with a short dark fringe, not stiff and spiny. A common weed of roadsides in southern British Columbia; it is expected to invade prairie regions.

Centaurea repens L.

RUSSIAN KNAPWEED

An erect perennial 30–100 cm high, growing from coarse, woody, running roots. Stems grooved and ridged; young stems covered with whitish woolly hairs. Leaves 12–75 mm long, pale green, sometimes woolly when young, linear to lanceolate. Lower leaves deeply lobed, upper ones entire-margined. Flower heads numerous at ends of branches, with a hard globular involucre 8–12 mm high and wide, with broad pale green to almost white bracts, with

membranous tips. Florets all tubular, pale pink or sometimes purplish. Seeds white. An extremely persistent introduced weed; has been found in shelter-belts and fields; various localities throughout Prairies. Syn.: *C. picris* Pall.

Centaurea solstitialis L.

YELLOW STAR-THISTLE

An annual plant, with branching stems bearing cottony hairs and growing 30–60 cm high. Basal leaves deeply lobed, up to 12 cm long; upper leaves entire, lanceolate to linear, 10–25 mm long. Involucre has yellowish bracts, many of them tipped by a yellow spine 12–20 mm long. Corollas yellow and all florets tubular. A rare, introduced weed; has been found in gardens; throughout Prairie Provinces.

Chrysanthemum ox-eye daisy

1. Heads with tubular florets only. *C. balsamita*
Heads with ray florets. 2
2. Leaves long, wedge-shaped, tapering into
the petiole. *C. arcticum*
Leaves petiolate below, with the middle
and upper stem leaves sessile. *C. leucanthemum*

Chrysanthemum arcticum L. var. *polaris* (Hulten) Boiv.

ARCTIC DAISY

Plants with creeping rootstocks; stems 10–20 cm high, glabrous or nearly so. Leaves fleshy and mostly basal, 4–8 cm long. Heads 2–3 cm across; bracts with conspicuous black tips. Arctic coasts and gravel banks; northeastern Boreal forest.

Chrysanthemum balsamita L.

COSTMARY

A fragrant perennial with stems 50–100 cm high, strigose above, glabrous below. Leaves 10–25 cm long, silvery strigose when young, becoming glabrate. Heads in a corymbiform inflorescence; disk 4–7 mm wide; rays mostly lacking, but if present 5–7 mm long, white; bracts with a conspicuous hyaline tip. The rayless form is *tanacetoides* (Boiss.) Boiv. Occasionally spreading from cultivation.

Chrysanthemum leucanthemum L.

OX-EYE DAISY

An erect perennial with few branches, 30–60 cm high. Lower leaves somewhat stalked, obovate to spatulate, toothed or incised, 2–8 cm long; upper leaves not stalked, clasping, oblong, and toothed near base. Heads usually borne singly at summit of stem, 2–5 cm across, radiate, with yellow disk florets and white ray florets. Seeds lacking pappus. Plants found in the Prairie Provinces are var. *pinnatifidum* Lecoq & Lamotte. An introduced plant, escaped from gardens and found in meadows and moist roadsides adjacent to forested areas; Parklands and Boreal forest. Syn.: *Leucanthemum vulgare* Lam.

Chrysopsis golden-aster

Perennial much-branched plants with a tufted root crown, usually low-growing and decumbent. Stems very leafy, with stalkless, alternate, and usually entire-margined leaves. Flowers radiate, medium in size, with yellow ray

florets. Pappus of the hairy achenes double, the inner ones consisting of rough hairs and the outer ones of small scales or minute bristles.

Chrysopsis villosa (Pursh) Nutt.

HAIRY GOLDEN-ASTER

A much-branched species 15–60 cm high, from a woody, branching taproot. Stems covered with coarse stiff hairs. Leaves numerous, grayish green, oblong or oblanceolate, alternate, 2–5 cm long, covered with short stiff appressed hairs; the lower leaves occasionally with a short stalk; the upper ones usually stalkless. Flower heads not numerous, radiate, with bright yellow ray florets, 25 mm or more across. This species has been divided into several species or varieties by various authorities, a linear-leaved variety being sometimes found in southern Alberta. Common; on dry sandy prairies and hill-sides; throughout Prairies and less common in Parklands. Including *C. hispida* (Hook.) DC.

Chrysothamnus rabbitbrush

Chrysothamnus nauseosus (Pall.) Britt.

RABBITBRUSH

A low shrubby much-branched plant with a very coarse thick woody root often protruding some distance above the surface of the soil, and appearing very large for the size of the plant. Growing 20–60 cm high with white woolly upright branches bearing very narrowly linear pale grayish green leaves 1–5 cm long, usually erect. Inflorescence dense, in terminal panicles; flower heads discoid, with no ray florets; flowers pale yellow, about 10–15 mm high, with bracts in 2 or 3 series. Copious pappi consisting of dull white hairs. A very local plant; abundant on badlands, eroded hillsides, and occasionally on saline clay flats; Prairies. Syn.: *C. frigidus* Greene.

Cirsium thistle

Stout erect biennial or perennial herbs with alternate lobed or dentate very prickly spiny leaves. Involucres ovoid or globose with imbricated bracts, which are usually spine-tipped. Flower heads discoid, with all florets tubular; achenes bearing a pappus of plumose (feathery) hairs.

1. Bracts of involucre covered with cobwebby hairs, and all bracts spine-tipped; upper leaf surface with short stiff bristles. *C. vulgare*
Bracts of involucre only slightly, if at all, cobwebby, and if so the inner bracts twisted and not spine-tipped. 2
2. Perennial plants from deep creeping rootstocks; involucres rarely over 12 mm wide; male and female flowers on separate plants. *C. arvense*
Plants without creeping rootstocks; involucres usually more than 12 mm wide; male and female florets on the same plant or flower head. 3

3. Inner bracts of involucre with loose, twisted, and spineless tips. *C. drummondii*
 Inner bracts of involucre without loose, twisted, and spineless tips. 4
4. Plants stemless or with stems; flowering heads exceeded by the foliage. 5
 Plants always with stems; flowering heads overtopping the foliage. 6
5. Bracts straight, with appressed tips; foliage long villous. *C. foliosum*
 Bracts squarrose, with spreading spiny tips; foliage thinly tomentose. *C. hookerianum*
6. Leaves green on both sides when mature; bracts usually without spiny tips. *C. muticum*
 Leaves white woolly beneath; bracts with spiny tips. 7
7. Leaves, upper ones at least, entire or only slightly lobed. *C. flodmanii*
 Leaves pinnately divided or deeply cleft. 8
8. Leaves deep green and bristly on upper sides. *C. altissimum* var. *discolor*
 Leaves pale green or gray, somewhat woolly above. 9
9. Perennial from deep roots; involucre rarely over 25 mm high; occasionally some lower leaves entire-margined. *C. flodmanii*
 Biennial; involucre usually over 25 mm high; no entire-margined leaves. *C. undulatum*

Cirsium altissimum (L.) Spreng. var. *discolor* (Muhl.) Fern. FIELD THISTLE

A tall perennial 90–150 cm high, with a grooved stem. Leaves deeply cleft into linear-lanceolate lobes, stalkless, and large, the lower ones sometimes 30 cm long, prickly, with rolled margins, deep green above, and white woolly beneath. Flower heads 35–50 mm across, with florets pale purple or pink, occasionally white. Rare; has been found in rich soil; southeastern Boreal forest.

Cirsium arvense (L.) Scop. (Fig. 213) CANADA THISTLE

A persistent perennial from deep running rootstocks, 30–100 cm high, usually in large patches. Leaves stalkless, often somewhat clasping, curled and wavy-surfaced, 5–15 cm long, roughly lanceolate, but deeply incised with toothed prickly segments; basal leaves sometimes stalked. Numerous flower heads in large loose corymbs at tops of stems and bearing purple or occasionally white florets. Plants bearing florets of only one sex; some all-male florets with heads often 25 mm across and others about 12 mm across; female florets bearing large quantities of seed. Achene bearing a pappus of white plumose hairs. A common introduced weed, found in great quantities; in waste places,




Fig. 213. Canada thistle, *Cirsium arvense* (L.) Scop.

fields, and roadsides; across the Prairie Provinces. Caterpillars of the painted lady (or thistle butterfly) and other closely related butterflies feed on thistles and will occasionally almost eradicate it by continual defoliation. The var. *integrifolium* Wimm. & Grab., entire-leaved Canada thistle, differs from the species by having flat almost entire leaves, with spiny margins, the lower leaves being slightly lobed. Found occasionally; throughout the Prairie Provinces.

Cirsium drummondii T. & G.

SHORT-STEMMED THISTLE

A low-growing species 10–30 cm high, with a hairy, slightly cobwebby stem. Leaves oblanceolate, green on both sides, with triangular lobes and weak spines, occasionally somewhat cobwebby when young, but not white woolly. Heads purple or rose purple, 35–50 mm high, with twisted inner bracts and spiny outer ones. Found occasionally; on somewhat open prairie in wooded areas; Rocky Mountains, Parklands, and Boreal forest.

Cirsium flodmanii (Rydb.) Arthur

FLODMAN'S THISTLE

A slender-stemmed perennial 40–100 cm high, from a deep root. Stem usually branched and with loose cottony hairs. Leaves deeply cleft into lanceolate spiny lobes, white cottony or woolly beneath, and somewhat cottony above, 5–15 cm long. Flower heads rose to rose purple, 3–4 cm across. Not common; found on moist prairie and valleys; Parklands and Prairies. An albino form with cream-colored flowers can be found occasionally. This species can be distinguished from *C. undulatum*, which it resembles, by its perennial habit and by the lower and newer basal leaves often being entire-margined, and its numerous new shoots around the base of the older plant.

Cirsium foliosum (Hook.) DC.

DWARF THISTLE

Plants biennial or short-lived perennials with a thick taproot, stemless, or stems to 1 m high, very leafy; the uppermost leaves much overtopping the inflorescence. Rosette leaves to 50 cm long, 5 cm wide, entire or shallowly lobed, with margins spinulose-ciliate. Heads solitary or in clusters at the top of the stem, 3–5 cm high; florets whitish to pink or rose. Mountain meadows; Rocky Mountains.

Cirsium hookerianum Nutt.

HOOKER'S THISTLE

A short-lived perennial with deep taproot; almost stemless or to 1.5 m high; the uppermost leaves overtopping the inflorescence. Foliage thinly tomentose; basal leaves to 30 cm long, to 15 cm wide, pinnately lobed or divided; lobes with stiff marginal spines and spinulose-ciliate. Inflorescence racemose or subspicate, or densely clustered; heads 3–4 cm high, with flowers whitish or pale yellowish. Grasslands and slopes; southern Rocky Mountains.

Cirsium muticum Michx.

SWAMP THISTLE

A biennial 60–150 cm high, with a branched leafy stem. Leaves 10–30 cm long, deeply cleft into oblong or lanceolate segments with slender spines, white woolly beneath when young, but becoming hairless when more mature. Basal leaves stalked, the upper ones smaller and stalkless. Flower heads few, about 35 mm across, purple; bracts of involucre not spiny, but somewhat sticky and hairy or cobwebby. Found in wet marshy lands; eastern Parklands and Boreal forest.

A white woolly biennial species 30–100 cm high, with a stout, branched, leafy stem. Leaves oblong or lanceolate, with triangular lobes, very prickly, with the lower leaves stalked and the upper ones stalkless, often continuing for some distance down the stem, usually densely white woolly on both sides, although with age the upper surface often becoming bare. Flower heads solitary at ends of the branches, purple or pink, and 3–8 cm across. Very common; on dry prairies and roadsides; throughout Prairies and Parklands. A variety with much larger flowers and the involucre over 4 cm high has been named var. *megacephalum* (Gray) Fernald. Found occasionally; Parklands.

Cirsium vulgare (Savi) Tenore

A stout-stemmed biennial, more or less woolly, 50–150 cm high, branched, and leafy up to the heads. Leaves 7–15 cm long, dark green, hairy on both sides, deeply cleft, and very prickly. Leaves usually continuing down the stem, forming prickly lobed wings. Numerous flower heads at ends of branches, purple, 3–5 cm broad and high. Involucral bracts cobwebby, all spine-tipped. An introduced species, found occasionally; on waste land and field borders; throughout Prairies and Parklands. Syn.: *C. lanceolatum* Scop.

Coreopsis tickseed

Coreopsis tinctoria Nutt.

A much-branched annual plant with a slender hairless stem 30–70 cm high. Leaves once or twice divided into linear segments, with the very uppermost ones entire, the lower ones sometimes stalked. Numerous flower heads on slender stalks, about 25 mm across, with brownish disk florets and 6–10 broad yellow ray florets having brownish bases. Seeds without a pappus and somewhat resembling small insects, from which the common name of the plant was derived. Very plentiful locally; in moist places, slough margins, low clay flats, irrigation ditches; throughout Prairies. The plant may be very abundant for one season in a particular location and then apparently disappear for several seasons. Often used as a garden flower. Most of the western tickseeds are *C. atkinsoniana* Dougl., distinguishable by a narrow wing around the seed.

Echinacea purple coneflower

Echinacea angustifolia DC.

An erect perennial plant 30–60 cm high, with a stiff hairy stem. Lower leaves lanceolate, 2–20 cm long, pointed at apex, and narrowed at base to a slender stalk; upper leaves stalkless or short-stalked, and all leaves either stiffly short hairy or at least very rough to touch. Flowers borne singly at the head of the stems; the conic disk bearing awned, stiff, purplish chaff that almost hides disk florets; ray florets purple, 20–25 mm long, often somewhat reflexed. Found fairly often; on dry benchland; southeastern Prairies and Parklands.

Echinops globe-thistle

- Stems white tomentose, densely so above. *E. exaltatus*
- Stem glandular tomentose, with the hairs long
and colored. *E. sphaerocephalus*



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Cirsium undulatum* (Nutt.) Spreng.

LOCATION: West of Lac Pelletier

COLL: J. Looman

DET:

COLL. NO: 13449

DATE: 2 September 1969

ELEVATION: 730 m

HABITAT: Moist prairie



Fig. 214. Wavy-leaved thistle, *Cirsium undulatum* (Nutt.) Spreng.

A perennial herb with stem 60–150 cm high, simple or with few branches above. Leaves to 40 cm long, 20 cm wide, tomentose and pale below. Inflorescence 3–4 cm in diam; heads 15–20 mm long, lavender blue. Cultivated, and rarely escaped.

Echinops sphaerocephalus L.

A coarse, branching perennial to 2.5 m high, with the stem spreading hairy. Leaves to 35 cm long, 20 cm wide, white tomentose below. Inflorescence 3–6 cm in diam; heads 15–20 mm long. Cultivated, and occasionally escaped.

Erigeron fleabane

Herbs with basal or alternate leaves. Flower heads usually radiate with many ray florets, usually more than 50. Involucral bracts usually in 1 or 2 series and imbricated (overlapping). Disk florets yellow and ray florets white to pink or purple, although in one rare species they are yellowish. A very large and widespread genus.

- 1. Heads discoid or with short inconspicuous ray florets. 2
 Heads with well-developed ray florets. 8
- 2. Leaves deeply dissected into at least 2 or 3 lobes. *E. compositus*
 Leaves entire. 3
- 3. Heads solitary; plants usually less than 10 cm high. 4
 Heads 2 or more; plants usually taller. 5
- 4. Involucre 6–8 mm high, densely blue lanate. *E. uniflorus* var. *unalaschkensis*
 Involucre 4–6 mm high, pubescence of hyaline hairs. *E. scotteri*
- 5. Involucre glabrous, 2.5–5 mm high. *E. canadensis*
 Involucre pubescent to glandular pubescent, higher. 6
- 6. Involucre finely glandular, not or sparsely pubescent. *E. acris* var. *asteroides*
 Involucre pubescent, not glandular. 7
- 7. Leaves linear-oblong to oblong, shorter than peduncles of lower flower heads. *E. elatus*
 Leaves narrowly long linear, exceeding the lower heads in the inflorescence. *E. lonchophyllus*
- 8. Plants usually less than 20 cm high; heads solitary. 9
 Plants usually more than 20 cm high; heads 2 to many, or both. 19
- 9. Stem leaves 0–3, mainly basal. 10
 Stem leaves numerous. 19

10. Leaves deeply dissected into at least 2 or 3 lobes. *E. compositus*
 Leaves entire or merely 3-toothed at apex. 11
11. Ray florets bright to deep yellow. *E. aureus*
 Ray florets white to pink or bluish. 12
12. Involucres usually more than 10 mm high, densely lanate. 13
 Involucres usually less than 8 mm high, not lanate, pubescence short or pilose. 14
13. Ray florets white; some or all leaves 3-toothed at apex. *E. lanatus*
 Ray florets reddish purple to whitish; leaves entire. *E. grandiflorus*
14. Leaves long linear; ray florets broad. 15
 Leaves narrowly obovate to oblanceolate or spatulate. 16
15. Leaves usually 1–3 cm long; stems usually leafless. *E. radicans*
 Leaves usually 3–7 cm long; stems with 1–3 leaves. *E. ochroleucus* var. *scribneri*
16. Leaves in part with 3- to 5-toothed apex. *E. pallens*
 Leaves all entire. 17
17. Plants long stoloniferous; leaves strigose. *E. flagellaris*
 Plants not stoloniferous; leaves pilose with spreading hairs. 18
18. Ray florets inconspicuous, about 3 mm long, very narrow; plants usually less than 5 cm high. *E. scotteri*
 Ray florets about 10 mm long; plants usually about 10 cm high. *E. arthurii*
19. Ray florets white. 20
 Ray florets colored. 25
20. Stem leaves at middle of stem longer than those below or above. *E. hyssopifolius*
 Stem leaves gradually reduced upward. 21
21. Ray florets with a minute pappus. *E. annuus*
 Ray florets with a normal pappus. 22
22. Leaves 1–3 mm wide, pilose; stem leaves all about 2 mm wide. *E. pumilus*
 Leaves more than 3 mm wide; stem leaves reduced in width and length. 23
23. Stem leaves usually more than 10; plants with thick taproot. *E. caespitosus*
 Stem leaves usually 5–7; plants with caudex, no taproot. 24

24. Leaves scabrous or pilose on both sides. *E. asper*
 Leaves glabrous below, lightly pubescent
 above. *E. glabellus*
25. Stem leaves at middle of stem longer than
 those above and below. *E. hyssopifolius*
 Stem leaves gradually reduced upward. 26
26. Leaves all or mostly clasping and
 auriculate. *E. philadelphicus*
 Leaves not clasping or auriculate. 27
27. Ligules filiform, less than 1 mm wide. 28
 Ligules wider than 1 mm. 29
28. Stem leaves usually 5–7; involucre
 pubescent. *E. glabellus*
 Stem leaves numerous; involucre
 glandular. *E. speciosus*
29. Stem leaves usually 3–5, sparsely
 pubescent. *E. grandiflorus*
 Stem leaves numerous, with the margins
 ciliate, only the midrib pubescent. *E. peregrinus*
 ssp. *callianthemus* var. *scaposus*

Erigeron acris L. var. *asteroides* (Andrz.) DC. NORTHERN DAISY FLEABANE

Biennial or short-lived perennial plants, with a simple or branched caudex; stems 30–80 cm high, hirsute to subglabrous, glandular in the inflorescence. Basal leaves 5–15 cm long, lanceolate; stem leaves narrower, sessile. Inflorescence more or less corymbiform, with several to numerous heads; ligules inconspicuous; involucre glandular to glandular hirsute. Open areas and moist woods; Boreal forest.

Erigeron annuus (L.) Pers. WHITETOP

Annual or sometimes biennial plants; stems 30–80 cm high, sparsely to copiously long pubescent below the inflorescence. Leaves 4–10 cm long, elliptic to broadly ovate, coarsely toothed. Inflorescence leafy, with more or less numerous heads; ligules 5–10 mm long, white or bluish; involucre finely glandular. Open woods and waste places; Boreal forest.

Erigeron arthurii Boiv. CLIFF DAISY

A caespitose perennial plant, with stems about 10 cm high. Rosette leaves 2–10 cm long, spatulate to oblanceolate; stem leaves 1–4, smaller than basal leaves, more or less densely glandular pubescent. Heads solitary; ligules about 1 cm long, white or pinkish; involucre 6–7 mm high, with tips purple, finely glandular. Alpine gravel slopes; southern Rocky Mountains.

Erigeron asper Nutt. ROUGH FLEABANE

A woody-rooted perennial 20–30 cm high; stems erect, covered by short stiff hairs. Basal leaves linear-oblongate, 2–10 cm long, with short stiff hairs. Upper leaves linear to linear-lanceolate. Heads 1–4, 20–25 mm across, with

100–150 very narrow white ray florets. Common on dry prairies and hillsides in Prairies and Parklands. By some authorities considered merely a form of *E. glabellus*.

Erigeron aureus Greene

YELLOW DAISY

Perennial plants with fibrous roots and branched caudex; stems 2–15 cm high; leaves 2–5 cm long, elliptic to obovate, finely pubescent, with hairs appressed or somewhat loose; stem leaves few and reduced. Heads solitary; ligules 6–10 mm long, yellow; involucre 5–8 mm high, sometimes purplish, sparsely to densely villous; hairs often with purple cross walls. Alpine slopes and meadows; southern Rocky Mountains.

Erigeron caespitosus Nutt. (Fig. 215)

TUFTED FLEABANE

A deep-rooted tufted perennial 10–20 cm high. Leaves finely hairy, with the lower ones spatulate, stalked, 2–8 cm long, the upper ones smaller, oblong, and stalkless. Flower heads borne singly or 3 or 4 to a stem, 2–3 cm across, with many narrow white ray florets. Involucre has 3 or 4 series of unequal bracts, thickened on back. Plentiful; on dry hillsides and prairie; throughout Prairies and Parklands.

Erigeron canadensis L.

CANADA FLEABANE

A slender bristly hairy-stemmed annual 10–100 cm high, usually with many branches toward the top. All leaves usually somewhat hairy, the lower ones spatulate, short-stalked, 2–10 cm long, and slightly toothed, the upper ones linear, entire, stalkless, and smaller. Flower heads very numerous, in a large open panicle, small, not over 5 mm across; florets, though numerous and white, usually hidden in the pappus. A common native weed; on dry soils, slough margins, and fields; throughout the Prairie Provinces. Syn.: *Leptilon canadense* (L.) Britt.

Erigeron compositus Pursh

COMPOUND FLEABANE

A low, tufted perennial from a woody root crown, 2–15 cm high, but usually very low. Leaves mostly basal, crowded, and usually twice divided into 3 linear or spatulate divisions. Flower heads 10–15 mm across, with white or rarely violet ray florets, borne singly on short stems. Found occasionally; on eroded hillsides, badlands, and dry or gravelly ridges; Prairies and southern Rocky Mountains.

Erigeron elatus (Hook.) Greene

TALL FLEABANE

Plants perennial, with slender stems 5–20 cm high, sparingly hirsute. Rosette leaves 2–7 cm long, oblanceolate; stem leaves 3–8, linear-oblong or oblanceolate. Heads 1–8, on flexuous or arching hirsute peduncles; ligules about 3 mm long; involucre 7–8 mm high, hirsute. On light soils; Prairies and Parklands.

Erigeron flagellaris Gray

CREEPING FLEABANE

Perennial plants, with stems 10–20 cm high, stoloniferous, strigose. Basal leaves 2–4 cm long, spatulate to oblanceolate, strigose; stem leaves 1–3, small, linear. Heads on peduncles 4–10 cm long; ligules about 5 mm long, white to purplish; involucre about 4 mm high, hirsute. Mountain meadows; southern Rocky Mountains.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Erigeron caespitosus* Nutt.

LOCATION: Frenchman Creek, north of Glinax

COLL: J. Looman

DET:

COLL. NO: 12607

DATE: 24 June 1969

ELEVATION: 960 m

HABITAT: Active prairie,
Astragalus cectinatus



Fig. 215. Tufted fleabane, *Erigeron caespitosus* Nutt.

Erigeron glabellus Nutt.

SMOOTH FLEABANE

A perennial, from a somewhat tufted rootstock, 15–40 cm high. Stems either hairless or sparingly hairy, and usually somewhat decumbent or horizontal at base. Basal leaves 5–10 cm long, oblanceolate and hairless; the upper leaves much smaller and linear-lanceolate. Leaves have only one prominent nerve. Flower heads 1–3, 1–2 cm across, with numerous narrow purple ray florets. Fairly common; in moist woods; Parklands and Boreal forest.

Erigeron grandiflorus Hook.

LARGE-FLOWERED DAISY

Perennials with stems to 10 cm high, white villous. Basal leaves 2–4 cm long; stem leaves 1–2 cm long, oblanceolate, long pilose. Heads solitary, 3–4 cm across; ligules 8–10 mm long, reddish purple to white; involucre 1 cm high, densely white lanate. Alpine prairies; southern Rocky Mountains.

Erigeron hyssopifolius Michx.

WILD DAISY

Perennial with extensive fibrous roots; stems solitary to several, 15–35 cm high, subglabrous to occasionally densely spreading villous. Leaves thin, lax; the lowermost leaves linear, reduced, and scale-like, the higher ones to 3 cm long, often with axillary leafy shoots. Heads solitary, rarely 2–5, borne on long peduncles; ligules 4–8 mm long, white to rose purple; involucre 4–6 mm high. Open woods and grassy openings; Boreal forest, rare in Parklands.

Erigeron lanatus Hook.

WOOLLY DAISY

Low perennials with black scaly rootstocks and branching caudex; stems 3–10 cm high, densely villous. Leaves 1–2 cm long, with the apex rounded; some leaves 3-toothed, densely woolly. Heads solitary; ligules about 1 cm long, white or purplish; involucre 10–12 mm high, densely woolly. Alpine meadows and slopes; southern Rocky Mountains.

Erigeron lonchophyllus Hook.

HIRSUTE FLEABANE

A somewhat hairy-stemmed biennial with one to several bunched stems 15–60 cm high. Lower stem leaves stalked and narrow oblanceolate, 5–12 cm long; upper stem leaves linear and shorter. All leaves usually smooth and hairless, but often with lower margins hairy. Few heads, about 15 mm across, with short white ray florets about 3 mm long borne erect on head. Not common; found occasionally in wet places; throughout the Prairie Provinces.

Erigeron ochroleucus Nutt. var. *scribneri* (Canby) Cronq. YELLOW ALPINE DAISY

Perennial plants with a caespitose caudex; stems about 10 cm high, grayish strigose. Leaves 3–7 cm long, narrowly linear, strigose, and ciliate. Heads usually solitary; ligules about 1 cm long, light yellow to white or lavender; involucre about 5 mm high, with tips purple, squarrose, somewhat lanate. Alpine meadows; southern Rocky Mountains.

Erigeron pallens Cronq.

PALE DAISY

Low perennial plants, with a caespitose caudex; stems about 10 cm high, long villous, and somewhat glandular throughout. Leaves spatulate, 2–4 cm long, mostly 3-toothed at apex. Heads solitary; ligules about 1 cm long, white; involucre 4–8 mm high, yellowish lanate. Alpine shale slides; southern Rocky Mountains.

Erigeron peregrinus (Pursh) Greene ssp. *callianthemus* (Greene) Cronq.

var. *scaposus* (T. & G.) Cronq.

WANDERING DAISY

Perennial plants with short rhizomes or caudex; stems 20–40 cm high, glabrous. Leaves linear-oblong to spatulate, tapering to the petiole; stem leaves much reduced; stem often subscapose. Heads solitary; ligules 10–15 mm long; involucre 7–10 mm high, densely glandular. Openings and open woods; southern Rocky Mountains.

Erigeron philadelphicus L.

PHILADELPHIA FLEABANE

A slender-stemmed upright perennial 30–60 cm high, with a stem either smooth or downy. Lower and basal leaves spatulate, blunt-rounded, toothed, 2–8 cm long, tapering to a short stalk, with upper leaves lanceolate, stalkless, partly clasping, and shorter. Flower heads in a terminal corymb 12–25 mm across, with many narrow pinkish or white ray florets. Fairly common; in moist places, open woodlands; throughout the Prairie Provinces.

Erigeron pumilus Nutt.

HAIRY DAISY

Usually a short-lived perennial with a taproot; stems 10–20 cm high, simple; spreading-pubescent leaves 3–7 cm long, linear-oblong, copiously villous; cauline leaves somewhat smaller. Heads mostly solitary or few on axillary peduncles; ligules 6–15 mm long, white or occasionally rose purple; involucre 4–7 mm high, densely villous. Dry grassland, especially on light soils; Prairies and Parklands.

Erigeron radicans Hook.

DWARF FLEABANE

A perennial with erect stem 3–10 cm high, somewhat stiff hairy. Basal leaves 10–35 mm long, linear to oblong, with stem leaves few and linear. Flower heads few or solitary, 1–2 cm across, with white ray florets. Rare; slopes and crests of eroded hills; Prairies and southern Rocky Mountains.

Erigeron scotteri Boiv.

DWARF DAISY

Perennial with a short taproot; stems 2–10 cm high, solitary or few together. Rosette leaves petiolate, entire, ciliate, more or less pubescent. Heads solitary; ligules about 3 mm long, pinkish; involucre 5–6 mm high, densely glandular. Alpine slopes; Rocky Mountains.

Erigeron speciosus (Lindl.) DC.

SHOWY FLEABANE

An erect-growing species, with smooth or slightly hairy leafy stems 30–40 cm high. Basal leaves 5–10 cm long, linear-oblong, tapering at base to a winged stalk, each with 3 ribs and a fringe of marginal hairs; stem leaves linear-lanceolate, clasping at broad base, with a marginal fringe of hairs. Flower heads large, 25–35 mm across, with numerous, narrow, blue or violet ray florets. Southern Rocky Mountains.

Erigeron uniflorus L. var. *unalaschkensis* (DC.) Boiv.

PURPLE DAISY

Low perennial plants with short rootstocks; stems 3–10 cm high, villous with blackish hairs. Leaves mostly basal, 1–4 cm long, spatulate, hirsute. Heads solitary; ligules 2–3 mm long, purplish or white, involucre black purple, densely pubescent with long hairs with blackish purple cross walls. Arctic and alpine tundra; northeastern Boreal forest and southern Rocky Mountains.

Eupatorium thoroughwort

Tall erect perennials with opposite or whorled leaves. Inflorescence a large terminal corymb-like cluster of discoid flower heads (without ray florets), either white or purplish, with several series of overlapping involucre bracts, unequal in length, outer ones quite small.

Leaves stalked and usually in whorls of 3–6 around the stem; flowers purple or purplish-tinged. *E. maculatum*

Leaves connate-perfoliate (leaves opposite with their bases united around the stem); flowers white. *E. perfoliatum*

Eupatorium maculatum L.

SPOTTED JOE-PYE WEED

An erect-stemmed somewhat branching plant 50–200 cm high; stem somewhat purplish or purple-spotted, with soft short hairs near the top. Leaves ovate to ovate-lanceolate, sharp-pointed at the apex, coarsely toothed, somewhat hairy beneath, 10–15 cm long, short-stalked, in whorls of 3–6 at internodes of stems. Inflorescence consisting of many flower heads forming rounded to flat-topped clusters up to 15–20 cm across, very conspicuous. Flower heads pinkish or purplish and consisting of pinkish lilac tubular florets; each head about 1 cm high and about 6 mm across. Pappus of white hairs. Pinkish involucre bracts unequal in size. Frequent in moist ground and in low moist woodland openings; southeastern Parklands and Boreal forest. The var. *bruneri* (Gray) Breitung, bruner's trumpetweed, very similar to the species, but with the underside of somewhat narrower leaves densely covered with a fine, soft, velvety hairiness. Stem hairy in its entire length. Found in similar locations, and of same distribution as the species, possibly advancing slightly farther west. Syn.: *E. bruneri* Gray.

Eupatorium perfoliatum L.

BONESET

An erect stout hairy-stemmed perennial 50–150 cm high. Lanceolate leaves 10–20 cm long, opposite, each pair joined at base and encircling stem. Leaves wrinkled, hairy on undersides. Inflorescence a large terminal cluster of white flowers. Flower heads about 6 mm high, with all tubular florets and bracts of unequal length. Fairly common; in wet places; southeastern Parklands and Boreal forest.

Gaillardia gaillardia

Gaillardia aristata Pursh (Fig. 216)

GREAT-FLOWERED GAILLARDIA

An erect-stemmed perennial 20–60 cm high, with a somewhat hairy stem. Lower leaves oblong to spatulate, sometimes lobed or pinnatifid, grayish hairy, 5–12 cm long, tapering to a stalk; upper leaves stalkless, smaller, and usually entire or slightly lobed. Flower heads terminal, 3–7 cm across, radiate, with a rounded purple disk. Ray florets 10–18, wedge-shaped, with 3 short triangular lobes at apex, yellow, but often with a purplish tinge at base. After flowers fall, receptacle somewhat globose and bearing achenes or seeds, which have a pappus of lanceolate papery bristled scales. Common; on dry prairie; a roadside plant throughout entire unforested portion of the Prairies and Parklands.



Fig. 216. Great-flowered gaillardia, *Gaillardia aristata* Pursh.

Galinsoga galinsoga

Galinsoga ciliata (Raf.) Blake

GALINSOGA

An erect or spreading annual 30–60 cm high, with many branches; stems often rooting at nodes when decumbent. Leaves opposite, stalked, ovate with rounded teeth, 15–50 mm long. Numerous terminal or axillary flower heads, 3–6 mm across, with 4 or 5 very small ray florets. Introduced from South and Central America, a weed that is invading the southeastern part of the Prairies and Parklands as a garden weed. Another species, *G. parviflora* Cav., yellowweed, subglabrous to strigose, has been found once in the southeastern Parklands.

Gnaphalium cudweed

Annual or biennial woolly and sometimes glandular herbs, with narrow entire alternate leaves. Flower heads discoid, with no ray florets, borne in panicles of crowded clusters on the stem. Bracts of involucre usually dry and membranous, and in several overlapping series. Achenes bearing a pappus of white hairs.

- 1. Tall plants without leafy bracts below flower heads; bracts of involucre well overlapping, white, dry, and membranous. 2
- Low plants with leafy bracts under flower heads; bracts of involucre slightly overlapping, yellowish or white. 3
- 2. Stem and leaves glandular; leaves continuing down the stem and forming a wing (decurent). *G. viscosum*
- Stem and leaves not glandular; leaves not decurent. *G. microcephalum*
- 3. Leaves broad, oblanceolate to spatulate; plant loosely woolly tufted. *G. palustre*
- Most of leaves narrow, linear, or oblanceolate; plant covered with short appressed hairs. *G. uliginosum*

Gnaphalium microcephalum Nutt.

COMMON CUDWEED

A perennial white tomentose plant, with an erect stem 30–60 cm high. Leaves 3–8 cm long, stalkless, linear-lanceolate, densely white woolly beneath. Flower heads about 6 mm high, in numerous clusters at head of stem. Involucral bracts membranous and dry, white tinged with brown. In open places in moist wood; southern Rocky Mountains.

Gnaphalium palustre Nutt.

WESTERN MARSH CUDWEED

A low-growing much-branched loosely woolly annual plant 5–20 cm high. Leaves 1–2 cm long, spatulate to oblong, stalkless, and loosely white woolly. Heads small, about 3 mm high, with woolly white or yellowish involucre, in leafy-bracted small clusters. Found occasionally; in sloughs and wet places; Prairies and Parklands.

Gnaphalium uliginosum L.

LOW CUDWEED

Somewhat similar to *G. palustre*, but with narrow linear leaves; the woolliness or hairiness appressed and not loose and spreading. Rare in the Prairies and Parklands, but found in wet places in Boreal forest.

Gnaphalium viscosum HBK.

CLAMMY CUDWEED

A biennial, from a taproot, 30–70 cm high, with a glandular hairy stem. Leaves linear to lanceolate, 5–10 cm long, bright green but glandular hairy above and white woolly beneath, stalkless and decurrent, continuing as a narrow wing for a distance down the stem. Flower heads discoid, about 6 mm high, white, with dry membranous white or straw-colored involucre bracts. Heads borne in several globose clusters. Rare; has been found in southern Rocky Mountains. Syn.: *G. macounii* Greene.

Grindelia gumweed

Grindelia squarrosa (Pursh) Dunal

GUMWEED

Biennial or perennial branching smooth-stemmed plants 20–60 cm high. Leaves oblanceolate, hairless, stalkless, finely and closely toothed, 1–4 cm long, alternate. Plants may be found occasionally with long narrowly spatulate basal leaves up to 7 cm long. Flower heads 2–3 cm across, with an involucre of many series of very sticky gummy bracts, in large numbers at heads of stems. Ray florets bright yellow. Achenes or seeds bearing 2 or 3 awns. Common; on dry prairie, roadsides, and especially on somewhat saline flats and slough margins; throughout entire unwooded part of the Prairie Provinces. Syn.: *G. perennis* A. Nels.

Gutierrezia broomweed

Erect many-stemmed perennials, with many branches from a deep woody taproot, to 50 cm long. Leaves narrow; flowers small, but very numerous. Native plants, unpalatable to livestock, increasing with overgrazing.

Gutierrezia sarothrae (Pursh) Britt. & Rusby

COMMON BROOMWEED

A low erect many-stemmed perennial, from a branching crowned woody taproot, 10–30 cm high. The numerous leaves stalkless and entire, narrowly linear, 1–4 cm long. The very numerous small flowers about 3 mm high, in close clusters at the ends of the branches. Unpalatable to livestock and therefore tending to increase in abundance when native pastures are heavily grazed. Extremely drought-tolerant, with deep roots and narrow leaves. Common; on dry prairie lands; throughout Prairies and Parklands. Syn.: *G. diversifolia* Greene.

Haplopappus ironplant

- 1. Flowers having only tubular florets. *H. nuttallii*
- Flowers with both tubular and ray florets. 2
- 2. Leaves pinnately dissected. *H. spinulosus*
- Leaves not pinnately dissected. 3

3. Leaves entire-margined; plants low, tufted with woody base; pappus white. 4
 Leaves usually somewhat toothed; plants with stem dying down to base annually; pappus reddish brown or yellow. *H. lanceolatus*
4. Stem leaves few, reduced; plants not glandular. *H. armerioides*
 Stem leaves several, not reduced; plants glandular throughout. *H. lyallii*

Haplopappus armerioides (Nutt.) A. Gray NARROW-LEAVED STENOTUS

A tufted hairless perennial from a large branching woody root crown, 10–15 cm high. Leaves mostly basal, linear or very narrowly spatulate, 2–7 cm long, pale greenish. Flowering stems almost leafless or with 2 or 3 narrow reduced leaves, 4–8 cm high; flower head about 25 mm across, yellow, with 8–10 ray florets. Not common, but found occasionally on dry eroded hills and badlands; Prairies. Syn.: *Stenotus armerioides* Nutt.

Haplopappus lanceolatus (Hook.) T. & G. LANCE-LEAVED PYRROCOMA

A herbaceous leafy-stemmed plant 10–40 cm high, with stems practically hairless. Basal leaves lanceolate, 2–7 cm long, long-stalked, with a few spine-pointed teeth; stem leaves shorter and stalkless. Flowers in a raceme or panicle of 3–15 heads, 20–25 mm across, yellow. Involucral bracts in 2 or 3 series of unequal length, dry, white at base, but with a greenish tip, sharp-pointed. Not uncommon; in meadows and moist saline areas; throughout Prairies. Syn.: *Pyrrcoma lanceolata* (Hook.) Greene.

Haplopappus lyallii Gray STICKY STENOTUS

A perennial plant with a large underground branching root crown, with few to several erect leafy stems 5–15 cm high. Basal leaves tufted, oblanceolate to obovate, 1–5 cm long, entire; cauline leaves several. Heads solitary; ligules short; involucre 8–12 mm high, like the stem and leaves densely stipitate glandular. Rocky ridges and slopes; southern Rocky Mountains.

Haplopappus nuttallii T. & G. TOOTHED IRONPLANT

A low tufted perennial from a deep woody root, 10–30 cm high. Leaves either oblong or lanceolate to spatulate, 1–3 cm long, grayish green, with short spiny teeth. Flowers borne either singly or in groups of 2 or 3 at ends of branches, about 12 mm across, and without ray florets. Bracts of involucre yellowish with a faint green tip. Fairly common; on dry eroded hillsides and plains; Prairies. Syn.: *Sideranthus grindelioides* (Nutt.) Britt.

Haplopappus spinulosus (Pursh) DC. (Fig. 217) SPINY IRONPLANT

A much-branched perennial from a thick woody root, 15–40 cm high. Leaves bluish green, often finely hairy, 1–4 cm long, and very deeply dissected into narrow segments, which have bristle-pointed teeth. Heads yellow, 6–15 mm across, often very numerous, and with narrow ray florets. Pappus soft, faintly tawny white. Plentiful on dry plains and hillsides; throughout Prairies and Parklands. Syn.: *Sideranthus spinulosus* (Pursh) Sweet.




Fig. 217. Spiny ironplant, *Haplopappus spinulosus* (Pursh) DC.

Helenium sneezeweed

Helenium autumnale L. var. *montanum* (Nutt.) Fern. MOUNTAIN SNEEZEWEED

A perennial 20–70 cm high, with a stout erect stem. Leaves 4–10 cm long, lanceolate, sometimes slightly toothed, stalkless, and continuing down the sides of the stem. Numerous flower heads at ends of branches, 2–3 cm across, with a high rounded yellow disk, and yellow ray florets up to 12 mm long. Bracts of involucre short, reflexed; ray florets usually somewhat reflexed. Achenes, which remain on almost globose receptacle after maturity, bearing a pappus of several sharp-pointed long scales. Common in low meadows, beside water courses, and in low places; Prairies, less common in Parklands. Thought to be somewhat **poisonous** to livestock and producing a taint in milk. Syn.: *H. montanum* Nutt.

Helianthus sunflower

Tall, coarse, erect annual or perennial herbs with opposite or alternate undivided leaves. Flowers large, with yellow ray florets. Receptacle or disk flat or rounded, broad, and chaffy. Disk florets yellow, brown, or purple. Achenes with a pappus of small scales or awns soon falling off. A difficult genus to classify.

- 1. Annual plants. 2
- Perennial plants. 3
- 2. Lower leaves usually cordate-based and toothed; bracts of involucre conspicuously hairy on margins; scales between disk florets not very hairy. *H. annuus*
- Leaves not cordate-based, entire; bracts of involucre not conspicuously hairy on margins; scales between disk florets each with tuft of hairs. *H. couplandii*
- 3. Disk of flower heads dark brown or purple. *H. laetiflorus* var. *subrhomboideus*
- Disk of flower heads yellow or light brown. 4
- 4. Leaf blades ovate; rootstocks bearing tubers. *H. tuberosus*
- Leaf blades lanceolate or linear-lanceolate; rootstocks not tuber-bearing. 5
- 5. Leaves rough on both sides and usually somewhat folded lengthwise. *H. maximilianii*
- Leaves rough above but somewhat hairy beneath, not folded lengthwise. *H. nuttallii*

Helianthus annuus L. var. *giganteus* Hort. SUNFLOWER

The sunflower with large flower head 15–30 cm across, cultivated for its seed, and occasionally escaped. More common is *H. annuus* f. *lenticularis* (Dougl.) Boiv., showy sunflower. A tall stout-stemmed erect annual 50–200 cm high. Stem rough, sometimes with short bristly hairs. Leaves coarsely toothed,

ovate, and often cordate-based, usually alternate, 10–20 cm long. Heads 7–15 cm across, with a dark brown or purple disk, usually about 5 cm across. Common; on clay and heavier soils and plentiful along roadsides; Prairies. Syn.: *H. annuus* L. ssp. *lenticularis* (Dougl.) Cockerell.

Helianthus couplandii Boiv.

PRAIRIE SUNFLOWER

An annual species 30–100 cm high, with a somewhat hairy stem. Leaves 2–7 cm long, with fairly long stalks, entire-margined, ovate to ovate-lanceolate, rough on both sides, usually with somewhat wedge-shaped bases, never cordate-based. Flower heads 3–8 cm across, usually with a rather small brown raised disk. Scales between disk florets each bearing a small tuft of hairs, which aid in distinguishing this annual species from *H. annuus*. Some authorities refer to this species as *H. aridus* Rydb. This species is found only on sandy soils. Plentiful; along roadsides in sandy areas and on sand dunes. Often where sandy soils merge into heavier ones, the demarcation line can be seen by the change from this species to the showy sunflower. Syn.: *H. petiolaris* AA.

Helianthus laetiflorus Pers. var. *subrhomboideus* (Rydb.) Fern.

BEAUTIFUL SUNFLOWER

An erect perennial from a rootstock, 30–100 cm high, with a sparingly hairy stem often tinged with red. Leaves opposite, 3-veined, 5–10 cm long, somewhat rhombic (obliquely 4-sided); the upper leaves rhombic-ovate or rhombic-lanceolate, the lower ones somewhat spatulate. Flower heads 2–7 cm across, with ray florets 15–35 mm long, disk purplish. Found occasionally; on moist grassland; Prairies and Parklands.

Helianthus maximilianii Schrad. (Fig. 218)

NARROW-LEAVED SUNFLOWER

A plant spreading by underground stems. A short-stemmed perennial 50–200 cm high. Stem rough, usually with coarse stiff hairs on upper part. Leaves stalkless or very short-stalked, 7–15 cm long, narrowly lanceolate, usually folded lengthwise along midrib, rough on both sides; the lower leaves opposite, the upper ones alternate. Yellow-disked flower heads 5–8 cm broad, on short stalks. Common; along roadsides, prairies, and valleys; throughout eastern Parklands and Boreal forest.

Helianthus nuttallii T. & G.

TUBEROUS-ROOTED SUNFLOWER

A perennial with creeping rootstocks, often with spindle-shaped fleshy roots, 1–3 m high. Leaves lanceolate, stalkless or very short-stalked, 5–15 cm long, somewhat narrow, and very rough on both sides. Flower heads on long stalks, 35–65 mm across, with a yellowish disk. Common; in moist and saline soils; throughout almost entire Prairie Provinces, especially in the southern portion. The var. *subtuberosus* (Britt.) Boiv. is distinguishable by having leaves lanceolate to linear-lanceolate, 6–15 cm long; the lower leaves opposite but the upper ones sometimes alternate. Flowers with yellowish brown disk, 3–9 cm across. A plant of moist meadows, slough margins, and wet roadsides; sometimes found in Prairies.



Fig. 218. Narrow-leaved sunflower, *Helianthus maximilianii* Schrad.

A perennial with fleshy edible tubers on roots, 1–3 m high. Stems hairy and with many branches. Ovate or oblong leaves 10–20 cm long, rough above and finely hairy below, tapering at the base to a stalk. Numerous heads 5–8 cm across, with a yellow disk and 12–20 yellow ray florets. Often cultivated for edible tubers. Fairly common; in moist soil, river flats, and other alluvial soil; southeastern Parklands and Boreal forest.

Heliopsis ox-eye

Heliopsis helianthoides (L.) Sweet var. *scabra* (Dunal) Fern.

ROUGH FALSE SUNFLOWER

A perennial from a cluster of fibrous roots, 40–100 cm high. Leaves ovate to ovate-lanceolate, opposite, 3-ribbed, 5–12 cm long, rough on both sides, abruptly narrowed at base to a short stalk, with large sharp teeth on margins. The few flower heads usually borne singly on long stalks, 5–6 cm across, with yellow ray florets 15–25 mm long. Found occasionally; on dry soil, banks, and sandhills; southeastern Parklands and Boreal forest.

Hymenopappus hymenopappus

Hymenopappus filifolius Hook.

TUFTED HYMENOPAPPUS

A perennial from a deep woody root, 15–30 cm high. Numerous stems from root crown, usually tufted, white woolly when young, becoming smooth and purple with age. Leaves 2–7 cm long, white woolly when young, turning smooth and pale green when older. Lower leaves stalked, each divided several times into very narrow almost thread-like segments. Flower heads usually few, borne terminally, 12–20 mm across, discoid, yellow. Densely woolly bracts in 1 or 2 series. A very drought-tolerant plant found on badlands, eroded slopes, and gravelly hills; Prairies and southern Rocky Mountains. Mostly considered to be var. *polycephalus* (Osterh.) B. L. Turn.

Hymenoxys rubberweed

Heads solitary on a long scape or stem; leaves

all basal and entire. *H. acaulis*

Heads several to many; leaves alternate and

divided into linear segments. *H. richardsonii*

Hymenoxys acaulis (Pursh) Parker

STEMLESS RUBBERWEED

A low grayish perennial from a coarse taproot, 10–15 cm high. Linear-ob-lanceolate leaves, all basal and clustered, 1–5 cm long, silky hairy, entire-mar-gined. Yellow flowers 2–3 cm across, with 10–15 rather broad ray florets, borne singly on leafless stems 10–15 cm long. Not common; found on eroded hill-sides; western Prairies. Syn.: *Tetraneris acaulis* (Pursh) Greene.

Hymenoxys richardsonii (Hook.) Cockerell (Fig. 219) COLORADO RUBBERWEED

A perennial, 10–20 cm high, from a coarse woody taproot, often with a divided woolly crown. Stems almost hairless. Leaves alternate, mostly basal, 5–10 cm long, divided into very narrow linear lobes, their divisions 2–3 cm




Fig. 219. Colorado rubberweed, *Hymenoxys richardsonii* (Hook.) Cockerell.

long. Flowers usually in a flat-topped cluster at ends of branches, yellow, about 2 cm across. Unpalatable to livestock, therefore tending to increase with heavy grazing. Common; on open prairie; throughout Prairies. Syn.: *Actinea richardsonii* (Hook.) Kuntze.

Liatrix blazingstar

Perennial plants from a globular corm or tuber-like root, with narrow undivided leaves. Flowers purple, in racemes or spikes, discoid, very showy; all florets tubular; bracts in several overlapping series; pappus of white, purplish, or tawny hairs.

Flower heads with 4–6 florets; pappus plumose or feathery. *L. punctata*

Flower heads with more than 15 florets; pappus barbed but not plumose. *L. ligulistylis*

Liatrix ligulistylis (A. Nels.) K. Schum. MEADOW BLAZINGSTAR

An erect plant 30–60 cm high, from a corm or tuber. Leaves 3–10 cm long, linear-oblongate, bright green with a rather conspicuous whitish midrib. Heads 2–3 cm across, reddish purple, on short stalks in a long raceme. Bracts of involucre in many series, green with purple tips; the rounded tops irregularly jagged, as if torn. Fairly common; in moist places, slough margins, forest openings, and sandhills; throughout Parklands, rare in Prairies.

Liatrix punctata Hook. (Fig. 220) DOTTED BLAZINGSTAR

A perennial from a stout, often corm-like rootstock, 10–30 cm high, often decumbent. Stiff linear leaves 5–15 cm long, densely covered with minute dots or depressions, smooth, but with a marginal fringe of short white hairs. Heads in a dense crowded spike, usually about 15 mm wide, with 4–6 pinkish purple tubular florets, usually showing many white plumose pappus hairs. A common species; on dry hillsides and grasslands; Prairies and Parklands.

Madia tarweed

Madia glomerata Hook. TARWEED

A very sticky, strong, and peculiarly scented annual 20–50 cm high, with glandular hairy stem and leaves. Leaves alternate, linear, entire, 2–7 cm long. Heads about 6 mm high and 3 mm wide, in close congested terminal and axillary clusters, with up to 5 small yellow ray florets, often none, usually hidden by the sticky glandular hairy involucre bracts. Found occasionally; in moist open spots in woodlands or slough margins; Prairies, especially in areas adjacent to Rocky Mountains and Cypress Hills.

Matricaria chamomile

1. Ray florets absent; leaf segments linear; plants with a pleasant pineapple-like odor. *M. matricarioides*

Ray florets present; leaf segments thread-like. 2



Fig. 220. Dotted blazingstar, *Liatris punctata* Hook.

2. Plant with distinct odor; disk conical *M. chamomilla*
 Plant odorless; disk merely rounded. *M. maritima*

Matricaria chamomilla L.

WILD CHAMOMILE

A branching annual plant 15–60 cm high, with finely divided leaves. Flower heads about 2 cm across, with white ray florets and a conical disk of yellow disk florets. Whole plant has a distinctive odor, faintly resembling pineapple, but well known to older people, who were dosed with chamomile tea in their childhood. An introduced weed, which has been reported from several locations in the Prairie Provinces.

Matricaria maritima L. (Fig. 221)

SCENTLESS CHAMOMILE

An annual with a much-branched hairless stem 20–70 cm high. Numerous leaves stalkless and several times divided into narrow thread-like segments. Flower heads numerous, at the ends of the branches, 10–25 mm across, radiate, with yellow disk and white ray florets. An introduced plant found occasionally; in waste places and roadsides; across the Prairie Provinces, and becoming more common in Parklands. Syn.: *M. inodora* L.

Matricaria matricarioides (Less.) Porter

PINEAPPLEWEED

A hairless annual plant 5–40 cm high, with leaves several times divided into linear segments. Leaves usually compact and copious, much more so than in *M. maritima*. Numerous flower heads, discoid, with tubular florets only, at the ends of the branches, about 6 mm across, conical, with yellow florets and greenish yellow involucre bracts with membranous margins. Plants, when squeezed, have a strong pleasant odor of pineapple. An introduced plant, which has become very persistent and plentiful locally; in waste places, and especially around farm or ranch yards and driveways; rapidly spreading throughout the Prairie Provinces. Syn.: *Chamomilla suaveolens* (Pursh) Rydb.

Petasites colt's-foot

Perennial, usually woodland plants, with thick creeping rootstocks and scaly-bracted flowering stems, which usually appear before the leaves. Leaves long-stalked and all basal, usually white woolly on underside. Flower heads in terminal clusters on stem, radiate, often unisexual, with male and female florets on separate plants. Achenes bearing a pappus of soft white hairs.

1. Leaves cordate to triangular, toothed but not deeply cleft. *P. sagittatus*
 Leaves round, kidney-shaped, or triangular and deeply cleft. 2
2. Leaves almost round in outline, deeply cleft almost to base. *P. palmatus*
 Leaves somewhat triangular, cleft only half or one-third of the way to midrib. *P. vitifolius*




Fig. 221. Scentless chamomile, *Matricaria maritima* L.

Petasites palmatus (Ait.) A. Gray (Fig. 222A) PALMATE-LEAVED COLT'S-FOOT

A species with leaves almost circular, 7–30 cm across, deeply cleft almost to the base into several divisions, green and smooth above, somewhat white woolly below when young, but losing the woolliness with age. Flowering stem appearing before the leaves, stout and scaly-bracted, 15–50 cm high, with heads in a corymbose cluster on long stalks. Flower heads white, 8–12 mm across, with male and female flowers on separate plants. Common; in moist woodlands; Parklands and Boreal forest.

Petasites sagittatus (Pursh) A. Gray (Fig. 222B) ARROW-LEAVED COLT'S-FOOT

This species has triangular-ovate to cordate leaves, with rounded marginal teeth, but not cleft or divided. Leaves 10–30 cm long, dull green above, densely white woolly below, on long stalks from root crown. Flowers in a dense terminal cluster at head of stem 20–50 cm high, which is scaly and somewhat bracted, appearing before the leaves. Fairly plentiful; in wet places, slough margins; Parklands and Boreal forest.

Petasites vitifolius Greene (Fig. 222C) VINE-LEAVED COLT'S-FOOT

A species with leaves somewhat triangular, usually with hastate basal lobes, and cleft as much as halfway to midrib, 7–15 cm long, green on both sides or sometimes white woolly below. Flowers nearly white, in a corymbose cluster at head of a scaly-bracted stem 20–30 cm high. Found occasionally; in wet places in woods; Boreal forest.

Psilocarphus woolly-heads

Psilocarphus elatior Gray TALL WOOLLY-HEADS

An annual species with gray woolly stems 5–15 cm high, erect, sometimes branched; depauperate plants sometimes no more than 1 cm high. Leaves 1–3 cm long, 2–5 mm wide, linear or linear-oblong. Heads solitary or clustered at the tip of stems and axillary branches 4–8 mm thick; involucre densely woolly or tomentose. Rare; damp areas, dried-up sloughs; Prairies.

Ratibida coneflower

Ratibida columnifera (Nutt.) Woot. & Standl. (Fig. 223)

LONG-HEADED CONEFLOWER

A perennial from a taproot, 30–70 cm high, usually branched from near base. Stems somewhat stiff hairy, with longitudinal grooves and angles. Leaves 5–10 cm long, very deeply pinnately divided into narrow segments. Flower heads borne at ends of long stalks, conspicuous by the cylindrical disk or receptacle, gray to purple, about 6 mm wide, 10–35 mm high. Ray florets yellow, 15–25 mm long, usually reflexed. Common; on dry prairie and roadsides; throughout Prairies, less common in Parklands. Syn.: *Lepachys columnifera* (Nutt.) Rydb. The f. *pulcherrima* (DC.) Fern., brown coneflower, similar to the species, but with ray florets partly or entirely brownish purple. This form has been found occasionally, but is very rare.





Fig. 222. A, Palmate-leaved colt's-foot, *Petasites palmatus* (Ait.) A. Gray; B, arrow-leaved colt's-foot, *Petasites sagittatus* (Pursh) A. Gray; C, vine-leaved colt's-foot, *Petasites vitifolius* Greene.



A.C. Budd

Fig. 223. Long-headed coneflower, *Ratibida columnifera* (Nutt.) Woot. & Standl.

Rudbeckia rudbeckia

Leaves neither lobed nor divided; disk or
flower heads dark brown. *R. hirta*

Leaves, at least the lower ones, deeply lobed
and divided; disk of flower heads greenish
yellow. *R. laciniata*

Rudbeckia hirta L.

BLACK-EYED SUSAN

A rather coarse hairy biennial 30–60 cm high, with erect sometimes tufted stems. Leaves lanceolate to oblanceolate, 5–15 cm long, 3-ribbed and hairy, with the lower ones stalked, but the upper ones stalkless and smaller. Flowers borne singly on long stalks, 5–8 cm across, with a dark brown hemispheric disk and 10–20 orange yellow ray florets. Fairly common; on prairies and woodlands; Parklands and Boreal forest. Some botanists consider the western form with oblanceolate to linear-lanceolate leaves as a separate species, *Rudbeckia serotina* Nutt., and reserve the name *R. hirta* L. for the eastern plants with ovate leaves.

Rudbeckia laciniata L.

TALL CONEFLOWER

A branching perennial 1–2 m high, with a smooth stem. Leaves, except the uppermost, deeply divided into 3–7 segments and stalked. Leaves varying in size up to about 20–25 cm long, with the upper ones progressively smaller. Stems usually bearing several long-stalked heads 5–10 cm across, with a greenish yellow disk, and 6–10 bright yellow ray florets. Frequent in open woodlands and forest edges; southeastern Parklands and Boreal forest.

Saussurea sawwort

Plants usually less than 20 cm high; involucre
bracts equal, acuminate at tip. *S. nuda* var. *densa*

Plants to 40 cm high; involucre bracts un-
equal, with the tip dilated. *S. glomerata*

Saussurea glomerata Poir.

TALL SAWWORT

Plants with rootstocks; stems 20–40 cm high, pubescent. Leaves 5–10 cm long, lanceolate to linear-lanceolate, densely brown glandular punctate below. Heads 10–15 mm across, with the outer bracts much shorter than the inner ones, which have a pink petaloid segment 1–2 mm wide. Rare; introduced; Peace River.

Saussurea nuda Led. var. *densa* (Hook.) Hulten

DWARF SAWWORT

Plants with short thick leafy stems 5–20 cm high, more or less densely woolly or tomentose. Leaves 5–10 cm long; the lower ones with winged petioles; the upper ones sessile, lanceolate, loosely pubescent when young, somewhat dentate. Heads crowded; inflorescence 4–7 cm across; involucre about 1 cm high and wide; inner and outer bracts equal. Alpine meadows and slopes; Rocky Mountains.

Senecio groundsel

Annual or perennial plants with alternate leaves. Flowers yellow, usually radiate, although one introduced species is discoid. Bracts is one series or a few basal bracts forming an outside row. Pappus consisting of many soft white hairs.

- 1. Flower heads solitary. 2
Flower heads few to many. 3
- 2. Heads 15–20 mm across; leaves entire to denticulate. *S. megacephalus*
Heads 10–15 mm across; leaves toothed to pinnatifid. *S. resedifolius*
- 3. Annual plants with taproots. 4
Perennial plants, usually with rootstocks. 6
- 4. Stems and leaves densely glandular pubescent. *S. viscosus*
Stems and leaves not glandular. 5
- 5. Plants branching; flower heads discoid. *S. vulgaris*
Plants with few branches, tall; flower heads radiate. *S. congestus*
- 6. Leaves all or mostly subentire to dentate. 7
Leaves all or mostly coarsely lobed to pinnately divided. 13
- 7. Flower heads solitary or with a small head at base. *S. megacephalus*
Flower heads 2 to many. 8
- 8. Stem leaves and basal leaves equal in size and shape. *S. triangularis*
Stem leaves different from basal leaves. 9
- 9. Middle and upper stem leaves much reduced in size. 10
Middle and upper stem leaves as large as or slightly smaller than lower leaves. 11
- 10. Inflorescence mostly terminal; axillary pedicels short. *S. integerrimus*
Inflorescence mostly axillary; pedicels long. *S. foetidus*
- 11. Leaves all cauline; basal leaves entirely lacking. *S. fremontii*
Basal leaves present. 12
- 12. Leaves and stems somewhat tomentose, at least till flowering. *S. tridenticulatus*
Leaves and stems glabrous; leaves rather succulent. *S. streptanthifolius*
- 13. Basal and stem leaves all deeply pinnately lobed or divided; plants often 1 m high. *S. eremophilus*

Basal leaves, if present, dentate or shallowly lobed.	14
14. Basal leaves entire, white woolly.	<i>S. canus</i>
Basal leaves mostly toothed or pinnatifid.	15
15. Leaf blades thin, smooth, and hairless.	<i>S. pauperculus</i>
Leaf blades thick to fleshy and hairy to woolly.	16
16. Basal leaves wedge-shaped at base; 3- to 5-toothed toward apex.	<i>S. streptanthifolius</i>
Basal leaves cordate to rounded or truncate at base.	17
17. Heads discoid, or with very short inconspicuous ray florets.	18
Heads with well-developed ray florets.	19
18. Heads few, mostly 2-6; basal leaves 1-3 cm long.	<i>S. pauciflorus</i>
Heads mostly 6-40; basal leaves 2-6 cm long.	<i>S. indecorus</i>
19. Plants low, less than 30 cm high; leaves thick and somewhat fleshy.	<i>S. tridenticulatus</i>
Plants taller, usually over 30 cm; leaves thin and not fleshy.	<i>S. aureus</i>

Senecio aureus L.

GOLDEN RAGWORT

A plant 30-80 cm high, with long-stalked cordate-based almost circular leaves 3-10 cm long. Stem leaves pinnatifid, and reduced in size. Flowers deep yellow, 6-8 mm high, in a terminal cluster. Found in meadows and moist places; southeastern Parklands and Boreal forest, Cypress Hills, and southern Rocky Mountains.

Senecio canus Hook. (Fig. 224)

SILVERY GROUNDSEL

White woolly perennial from horizontal rootstocks, 10-30 cm high. Basal leaves entire-margined, oval to spatulate, 3-7 cm long, stalked, and somewhat white woolly; stem leaves much smaller, stalkless, oblong, and often somewhat pinnatifid or lobed. Heads in a terminal cluster 10-25 mm across. Not common; on dry hills; across the Prairies and Parklands.

Senecio congestus (R. Br.) DC.

MARSH RAGWORT

A coarse hollow-stemmed annual 15-60 cm high, with fleshy stems somewhat cobwebby when young but hairless when mature. Lower leaves lanceolate to spatulate, 5-15 cm long, with wavy margins, and winged stalks; upper leaves smaller, stalkless, somewhat lobed or dentate, linear-lanceolate, and clasping the stem. Flower heads in a very crowded, dense, terminal cluster, pale yellow, 1-2 cm across. Fairly common; around sloughs, stream banks, and lakes, often forming a solid belt around a small lake; throughout the Prairie Provinces. Syn.: *S. palustris* (L.) Hook.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

Senecio canus Hook.

STATION: 2410, Swift Current, Sask.

DATE: 1910

COLLECTOR: ...

... ..

... ..

... ..

... ..



Fig. 224. Silvery groundsel, *Senecio canus* Hook.

Senecio eremophilus Richards. (Fig. 225)

CUT-LEAVED RAGWORT

A leafy-stemmed species with a stout often purplish stem 30–100 cm high. Leaves lobed or pinnatifid, 5–15 cm long; the lower leaves stalked; the upper ones without stalks, slightly smaller than the lower ones. Stems leafy up to inflorescence. Numerous yellow flower heads 10–25 mm across, with black-tipped bracts, in compact terminal clusters. Found occasionally; on moist soil; Parklands, more common in Boreal forest.

Senecio foetidus Howell

MARSH BUTTERWEED

A fibrous-rooted perennial with solitary or sometimes clustered stems 30–100 cm high, glabrous throughout. Basal leaves 7–20 cm long, to 7 cm wide, rather sharply dentate; stem leaves progressively reduced, sessile. Inflorescence dense, with the heads several to many, more or less corymbosely arranged; ligules about 8 mm long; involucre 6–9 mm high, with the bracts black-tipped. Moist woods; southern Rocky Mountains.

Senecio fremontii T. & G.

MOUNTAIN BUTTERWEED

A glabrous perennial with taproot and branching caudex, freely branching; stems decumbent, 10–20 cm high. Leaves thick, 1–4 cm long, obovate to spatulate, rather sharply dentate. Heads terminating short branches; ligules 6–10 mm long; involucre 7–12 mm high. Slopes and open areas in mountain woods; southern Rocky Mountains.

Senecio indecorus Greene

RAYLESS RAGWORT

A fibrous-rooted perennial with a simple or occasionally branched caudex and glabrous stems 30–80 cm high. Leaves rather thin, lightly floccose when young; basal leaves elliptic or ovate, tapering to truncate at base, serrate or somewhat incised, petiolate; stem leaves incised to pinnatifid, with toothed lobes. Heads several to many; involucre 7–10 mm high; bracts often purple-tipped. Moist meadows; Boreal forest and occasionally in Parklands.

Senecio integerrimus Nutt.

ENTIRE-LEAVED GROUNDSEL

A stout-stemmed perennial from coarse, fleshy, fibrous roots, 20–60 cm high. When young, the stem is somewhat hairy, but later the plant is entirely smooth and hairless. Leaves entire, thick and fleshy, 5–20 cm long; basal leaves oblong to lanceolate with long stalks; upper leaves stalkless, lanceolate to linear, and reduced in size. Flower heads often 12–20 mm across, in dense terminal clusters.

The species is represented by three varieties:

- 1. Involucral bracts green to tip. var. *integerrimus*
 Involucral bracts with a black tip. 2
- 2. Black tip about 1 mm long. var. *exaltatus* (Nutt.) Cronq.
 Black tip about 3 mm long, triangular. var. *lugens* (Richards.) Boiv.

The var. *integerrimus* is found throughout Prairies and Parklands, var. *exaltatus* in Cypress Hills and southern Rocky Mountains, and var. *lugens* in Rocky Mountains, northwestern Parklands, and Peace River.



HERBARIUM OF THE RESEARCH STATION
SWIFT CURRENT, SASK.

SPECIES: *Senecio eremophilus* Richards.

LOCATION: Rusty Lake

COLL: J. Looman

DET:

COLL. NO: 13354

DATE: 14 August 1969

ELEVATION: 500 m

HABITAT: Lakeshore in
reed vegetation




Fig. 225. Cut-leaved ragwort, *Senecio eremophilus* Richards.

Senecio megacephalus Nutt.

LARGE-FLOWERED RAGWORT

Plants with a thick rootstock, and simple stout stems 10–20 cm high, more or less floccose pubescent. Leaves linear-oblong to lanceolate, entire or denticulate. Heads solitary or seldom 2 or 3, 20–25 mm high; ligules orange, 15–20 mm long; involucre with the outer bracts often surpassing the inner ones. Alpine slopes or ridges; southern Rocky Mountains.

Senecio pauciflorus Pursh

FEW-FLOWERED RAGWORT

A fibrous-rooted perennial with a simple caudex; stems 10–30 cm high, glabrous, or somewhat floccose-tomentose when young. Basal leaves 2–4 cm long, elliptic-ovate to subrotund, abruptly contracted or truncate at base, long-petioled; stem leaves reduced, with the upper ones sessile, toothed to pinnatifid. Heads usually 2–6, discoid or rarely with a few short rays; involucre usually 6–8 mm high; bracts more or less purplish. Arctic and alpine meadows; Boreal forest and Rocky Mountains.

Senecio pauperculus Michx.

BALSAM GROUNDSEL

A slender-stemmed perennial with a woolly-based stem 20–40 cm high. Basal leaves stalked, oblong or oval, with wavy margins, 2–8 cm long; stem leaves smaller, not stalked, and lobed or toothed. Leaf blades hairless and thin. Few flower heads, in a loose cluster on thin stalks at the head of the stem, 10–15 mm across.

Three varieties of this species can be found:

- 1. Involucre 4–5 mm high; basal leaves about 1 cm wide. var. *pauperculus*
 Involucre 5–9 mm high; basal leaves 1–2 cm wide. 2
- 2. Involucre 5–7 mm high; basal leaves 10–15 mm wide. var. *firmifolius* Greenman
 Involucre 6–9 mm high; basal leaves 15–20 mm wide. var. *thomsoniensis* (Greenman) Boiv.

The var. *pauperculus* is found in wet, often somewhat saline meadows throughout Prairies and Parklands; the other varieties are usually found in wet meadows, especially on sandy soils, in Parklands and Boreal forest.

Senecio resedifolius Less.

ARCTIC BUTTERWEED

Plants with a rather thick rhizome and branched caudex; stems 5–30 cm high, essentially glabrous. Leaves thick, 1–2.5 cm long, ovate or reniform to suborbicular, toothed or entire; stem leaves reduced, with the upper ones sessile. Heads solitary or rarely 2; ligules to 15 mm long or sometimes missing; involucre 6–8 mm high, purplish. Boggy mountain meadows and slopes; southern Rocky Mountains.

Senecio streptanthifolius Greene

NORTHERN RAGWORT

A fibrous-rooted perennial with a short rootstock; stems 10–50 cm high, glabrous or lightly pubescent. Lower leaves 2–7 cm long, oval, long-stalked,

and usually toothed or lobed; upper leaves stalkless, reduced, linear-lanceolate, and deeply toothed; most of the leaves somewhat fleshy. Orange yellow flower heads 1–2 cm across, in a rather loose terminal cluster on long stalks. Found occasionally; on moist soils; in valleys in Parklands and lightly wooded parts of northwestern Boreal forest.

Senecio triangularis Hook.

BROOK RAGWORT

A leafy-stemmed often tufted perennial, from fleshy fibrous roots, 30–60 cm high. Stem hairless and leafy to the inflorescence. Leaves almost all stalked, 2–8 cm long, with conspicuous teeth, triangular to lanceolate. Flower heads 1–2 cm across, in a terminal cluster. Along streams and in wet places; southern Rocky Mountains.

Senecio tridenticulatus Rydb.

COMPACT GROUNDSEL

A low slender plant from a short rootstock, 15–30 cm high, slightly white hairy at the base of the stem and at the axils of the leaves. Lower leaves 2–7 cm long, short-stalked, oblanceolate, and often somewhat toothed; upper leaves linear, stalkless, much smaller, and lobed. Flower heads few in number, 1–2 cm across, with rather pale yellow ray florets, in a fairly compact terminal cluster. Found occasionally; on moist prairie; southeastern Parklands.

Senecio viscosus L.

STINKING GROUNDSEL

An annual weed with taproot; stems 10–60 cm high, usually branched. Herbage densely glandular pubescent, strong-scented. Leaves pinnatifid, 3–12 cm long, 1–5 cm wide. Heads several to numerous, long pedicellate; ligules inconspicuous; involucre 6–8 mm. Introduced, rare; southeastern Parklands.

Senecio vulgaris L.

COMMON GROUNDSEL

A low hollow-stemmed much-branched annual, 15–40 cm high. Leaves lobed, 5–15 cm long; lower leaves stalked and upper ones clasping at base. Flower heads discoid (without any ray florets), about 6 mm across, often with some black-tipped bracts, in clusters at ends of branches. An introduced weed, found often in gardens and around settlements; across the Prairie Provinces, but nowhere very common.

Solidago goldenrod

Perennial herbs from rootstocks, with alternate entire leaves. Inflorescence of terminal panicles, in racemes or corymbose clusters, with many radiate yellow flower heads. Two species are white or cream. Pappus of rough bristly hairs in one or two series. Most goldenrods bloom in mid- or late summer and are good honey-producing plants.

1. Inflorescence corymbose, with the branches slanting upward, forming a rounded or flat-topped inflorescence. 2
- Inflorescence a panicle or spike. 6
2. Ray florets white or pale yellow, large, 5–8 mm long. *S. ptarmicoides*

Ray florets yellow, usually less than 5 mm long.	3
3. Leaves firm, very rough, gray pubescent; stems rigid and gray puberulent.	<i>S. rigida</i> var. <i>humilis</i>
Leaves not rough or gray pubescent.	4
4. Stem leaves, at least the lower ones, petioled, the upper ones much reduced.	<i>S. multiradiata</i>
Stem leaves sessile or subsessile, all about the same size.	5
5. Leaves finely glandular pubescent, flat; inflorescence with 10–50 heads.	<i>S. graminifolia</i>
Leaves not glandular, commonly folded lengthwise; inflorescence mostly with more than 50 heads.	<i>S. riddellii</i>
6. Inflorescence racemose, with branches ascending along the stem; inflorescence more or less cylindrical or narrowly oval.	7
Inflorescence a panicle with more or less spreading or arching branches; heads often secund on the branches.	11
7. Flowers white or cream.	<i>S. bicolor</i>
Flowers yellow.	8
8. Leaves and stems densely hairy; lower leaves ovate, with the petiole as long as the blade.	<i>S. bicolor</i> var. <i>concolor</i>
Leaves and stems not densely hairy; lower leaves not ovate, or if so, not long-petioled.	9
9. Leaves obovate, subsessile, 2.5–3 times as long as wide, grayish green.	<i>S. mollis</i>
Leaves lanceolate, at least the lower ones petioled, more than 4 times as long as wide.	10
10. Branches of the inflorescence shorter than or as long as the subtending stem leaves; inflorescence cylindrical, more or less interrupted.	<i>S. spathulata</i> var. <i>spathulata</i>
Branches of the inflorescence much longer than the subtending stem leaves; inflorescence more or less oval, not interrupted.	<i>S. uliginosa</i>
11. Basal leaves much longer than stem leaves, often forming a rosette.	12
Basal leaves not much longer than stem leaves, seldom forming a rosette.	14
12. Stem and leaves minutely gray pubescent.	<i>S. nemoralis</i>
Stem and leaves not gray pubescent.	13

13. Lower and middle stem leaves distinctly 3-nerved, to 10 cm long, 15 mm wide, entire or obscurely dentate. *S. missouriensis*
 Lower and middle stem leaves not distinctly nerved, usually 20 cm or more long, 20–75 mm wide, clearly serrate. *S. juncea*
14. Leaves obovate, sessile, 2.5–3 times as long as wide, grayish green. *S. mollis*
 Leaves lanceolate to oval, more than 4 times as long as wide, green. 13
15. Stem below the inflorescence glabrous or very sparsely pubescent. *S. gigantea* var. *serotina*
 Stem below the inflorescence densely pubescent. *S. canadensis*

Solidago bicolor L.

PALE GOLDENROD

A stout plant with white hairy stems 20–60 cm high. Lower leaves long-stalked, obovate to oblong, 5–10 cm long; upper leaves stalkless, smaller and narrow; all leaves hairy on both sides. Inflorescence narrow, 5–15 cm high, with short branches bearing flower heads, which have white or pale cream ray florets. Found occasionally; on dry soil; southeastern Boreal forest and Parklands. The var. *concolor* T. & G. differs only in having yellow ray florets. Found in dry to moist woods; throughout Parklands and Boreal forest. Syn.: *S. hispida* Muhl. var. *lanata* (Hook.) Fern.

Solidago canadensis L. var. *canadensis* (Fig. 226)

GRACEFUL GOLDENROD

A slender-stemmed leafy plant from a horizontal rootstock, 30–80 cm high. Leaves narrowly lanceolate, 3-veined, 5–10 cm long, with fine teeth, slightly hairy along the veins on the underside, green. Inflorescence a broad, pyramid-like panicle; flower heads small, borne on one side of spreading branches. Found commonly; in woodlands and forests; Parklands and Boreal forest, also in Cypress Hills. The var. *gilvocanescens* Rydb., canescent goldenrod, is a slender-stemmed yellowish gray species from a horizontal rootstock, 25–60 cm high, with a finely downy-haired stem. Leaves 2–5 cm long, 3-veined, narrowly lanceolate, entire or somewhat toothed, with the lower ones falling off early. Inflorescence pyramid-shaped with somewhat recurved branches. Fairly common; on disturbed moist areas; Prairies. Syn.: *S. pruinosa* Greene.

Solidago gigantea Ait. var. *serotina* (Ait.) Cronq.

LATE GOLDENROD

A stout erect species from stout rootstocks, 60–150 cm high, with a smooth leafy stem. Leaves thin, lanceolate, 2–12 cm long, without stalks, 3 nerves, usually sharply toothed, smooth on both sides. Inflorescence large, pyramidal; heads crowded on spreading recurved branches. Common; around bluffs, woodlands, and also in coulees and low areas; throughout Prairie Provinces. Syn.: *S. serotina* Ait.

Solidago graminifolia (L.) Salisb. var. *graminifolia*

FLAT-TOPPED GOLDENROD

A much-branched plant from a long rootstock, 30–60 cm high. Leaves numerous and somewhat crowded, 2–10 cm long, linear-lanceolate, 15–20



Fig. 226. Graceful goldenrod, *Solidago canadensis* L. var. *canadensis*.

times as long as wide, and pointed at both ends. Small yellow flowers, in very short-stemmed terminal clusters, forming a somewhat flat-topped inflorescence. Fairly common; along stream banks and moist places; eastern Parklands and Boreal forest. The var. *major* (Michx.) Fern. has wider leaves and the length-to-width ratio is usually 8–10:1; plants smaller; inflorescence with fewer heads. The variety is more common in the western part of the distribution area.

Solidago juncea Ait.

SHARP-TOOTHED GOLDENROD

A stout smooth-stemmed species from a horizontal rootstock, 40–80 cm high. Lower leaves single-ribbed, broadly oblanceolate, stalked, hairless, 15–30 cm long; upper leaves stalkless, smaller, and narrower. Inflorescence a somewhat flat-topped panicle with recurved spreading branches; flower heads borne on one side of branches. Rarely found; a species of woodlands and banks; southeastern Boreal forest.

Solidago missouriensis Nutt.

LOW GOLDENROD

A rather low smooth-stemmed species from horizontal rootstocks, 15–50 cm high. Stems hairless, often tufted, usually somewhat reddish. Leaves 3-ribbed, linear-lanceolate, 2–10 cm long, often reddish and hairless except for sparse short marginal hairs. Inflorescence a compact terminal panicle with erect branches. Earlier flowering than other goldenrods. Common on dry prairies and hillsides; throughout Prairies and southern fringe of Parkland. The var. *fasciculata* Holzinger, low goldenrod, differs from the species by having branches of inflorescence recurved and spreading instead of erect. Probably the eastern form of the species. Common; on dry prairie; in south central parts of the distribution area. Syn.: *S. glaberrima* Martens.

Solidago mollis Bartl.

VELVETY GOLDENROD

A stout, low, erect plant from a horizontal rootstock, 20–50 cm high. Whole plant covered with very fine, short, velvety hairs. Leaves pale green, almost entire-margined, obovate to oval (or the upper leaves elliptic), 3-nerved, 2–7 cm long; lower leaves short-stalked; upper leaves stalkless and crowded on stem. Inflorescence a pyramid-shaped dense panicle. Common; on dry prairie land and roadsides; probably the most common species in Prairies.

Solidago multiradiata Ait.

ALPINE GOLDENROD

A rather small species with a short rootstock or branching caudex; stems 5–40 cm high, pilose above and in the inflorescence. Heads rather large; involucre 4–8 mm high, few, seldom more than 10–20. Boreal and montane or alpine meadows; Boreal forest and Rocky Mountains.

Solidago nemoralis Ait.

SHOWY GOLDENROD

A species growing in clumps of several stems from a thick rootstock, 30–50 cm high. Stems often somewhat decumbent, reddish, covered with fine downy hairs. Basal leaves narrowly oblanceolate, entire, 5–10 cm long; upper leaves oblong or linear; all leaves somewhat ashy gray, with very minute hairs. Inflorescence narrow, usually bent over or somewhat nodding at top. Plentiful; on sandy soil and in sandhills; throughout Prairies and Parklands.

Solidago ptarmicoides (Nees) Boiv.

UPLAND WHITE GOLDENROD

A somewhat tufted plant from creeping rootstocks, 20–60 cm high. Leaves linear-lanceolate, 2–15 cm long, sometimes slightly toothed, firm, and shiny. Flower heads numerous, 10–25 mm across, white, in a somewhat flat-topped terminal cluster. Rare, but found occasionally; in dry, saline, or gravelly soil; eastern Parklands and Boreal forest. Syn.: *Unamia alba* (Nutt.) Rydb.; *Aster ptarmicoides* (Nees) T. & G. Plants with somewhat smaller heads and pale yellow ligules are considered to be a hybrid between *S. ptarmicoides* and *S. rigida* var. *humilis*, and have been named \times *S. lutescens* (Lindl.) Boiv. Syn.: *Aster ptarmicoides* (Nees) T. & G. var. *lutescens* (Lindl.) A. Gray.

Solidago riddellii Frank

RIDDELL'S GOLDENROD

A tall species with a more or less developed caudex and sometimes short rootstocks; stems 40–100 cm high, glabrous below the inflorescence. Leaves 5–15 cm long, glabrous, with margins scabrous, entire, tending to be 3-nerved; stem leaves numerous, somewhat clasping or sheathing at the base. Inflorescence corymbiform; branches puberulent; heads numerous, rarely less than 50, often several hundred; involucre 5–6 mm high. Rare; swamps and wet meadows; southeastern Parklands and Boreal forest.

Solidago rigida L. var. *humilis* Porter (Fig. 227)

STIFF GOLDENROD

An erect stout-stemmed species, with a densely fine-hairy rough stem, from a thick woody rootstock, 10–40 cm high. Basal leaves long-stalked, oval, 4–10 cm long, thick, and densely fine-hairy on both sides; stem leaves oval, stalkless, and smaller. Inflorescence a dense corymbose cluster, somewhat flat-topped, with an involucre 6–8 mm high. Common; on prairies and openings in woodlands; throughout the Prairie Provinces.

Solidago spathulata DC. var. *spathulata* (Fig. 228)

MOUNTAIN GOLDENROD

Rather low plants 20–50 cm high. Stems decumbent at base, often reddish-tinged. Lower leaves spatulate, 2–10 cm long, usually blunt-tipped, often with rounded teeth; stem leaves smaller and entire; basal leaves usually somewhat crowded at root crown. Inflorescence a narrow erect panicle; heads fairly large, usually about 8 mm high. Common; in grasslands; throughout Prairies, Parklands, and southern Boreal forest; often mistaken for *S. missouriensis*, with smaller flowers and triple-veined leaves. Syn.: *S. decumbens* Greene; *S. oreophila* Rydb.

Solidago uliginosa Nutt.

MARSH GOLDENROD

A large species with a long much-branched rootstock or caudex; stems 60–150 cm high, glabrous below the inflorescence. Basal and lower stem leaves 6–35 cm long, lanceolate to narrowly elliptic, subentire to dentate, tapering to a long winged petiole; upper stem leaves gradually reduced. Inflorescence puberulent, elongate; branches straight or recurved-secund; involucre 3–5 mm high. Bogs and marshes; southeastern Boreal forest.



Fig. 227. Stiff goldenrod, *Solidago rigida* L. var. *humilis* Porter.



Fig. 228. Mountain goldenrod, *Solidago spathulata* DC. var. *spathulata*.

Tanacetum tansy

Leaf segments very fine, about 1 mm wide. *T. huronense*

Leaf segments much coarser, 3–10 mm wide. *T. vulgare*

Tanacetum huronense Nutt.

INDIAN TANSY

A perennial with long rootstocks; stems 10–60 cm high, more or less vil-
lous throughout. Leaves 5–20 cm long, 2–8 cm wide, 2- or 3-pinnatifid. Heads
1–15, with the disk 10–18 mm across; ligules inconspicuous, or rarely to 4 mm
long. Rare; sandy lakeshores; Boreal forest.

Tanacetum vulgare L. (Fig. 229)

TANSY

A stout erect-stemmed perennial 30–100 cm high. Leaves 5–25 cm long,
pinnately divided into narrow toothed segments, very aromatic when bruised.
Flower heads discoid (without ray florets), 6–8 mm across, in a somewhat flat-
topped cluster. An introduced garden plant, which has become established
and often forms extensive colonies in roadsides, waste areas, and ditches; in
many locations throughout the Prairie Provinces.

Thelesperma tickseed

Thelesperma marginatum Rydb.

TICKSEED

A perennial with rootstock or woody caudex; stems 10–20 cm high, leafy
at the base. Leaves once or twice pinnately divided, with the divisions linear,
1–2 mm wide. Peduncles mostly 7–10 cm long; heads solitary or few, about 1
cm wide; involucre bracts fused; lobes with a distinct white margin. Rare;
eroded slopes; Prairies.

Townsendia townsendia

Tufted plants with clustered basal alternate entire leaves and large aster-
like radiate heads. The pappus consisting of bristly hairs.

Heads on a stalk from the root crown. *T. parryi*

Heads stalkless, growing directly on the root
crown. *T. exscapa*

Townsendia exscapa (Rich.) Porter

LOW TOWNSENDIA

An almost stemless plant from a deep woody branching root. Leaves nar-
rowly spatulate to linear, 2–5 cm long. Flower heads stalkless, borne amongst
rosettes of leaves, 20–35 mm across, and with bluish or white ray florets.
Found occasionally; on eroded prairies and dry stony hillsides; throughout
Prairies and southern fringe of Parklands. Syn.: *T. sericea* Hook. A similar
species, *T. hookeri* Beaman, is smaller, with narrower leaves, and involucre
bracts very densely ciliate throughout and tipped with a tuft of hairs. How-
ever, the pubescence and size of plants in any location can vary greatly, and
there seems little reason to assume the presence of two species of low townsen-
dia.



Fig. 229. Tansy, *Tanacetum vulgare* L.

A biennial plant, usually cushion-like, with crowded basal leaves. Leaves spatulate and thick, 2–5 cm long; stem leaves few, very small. Flower head on a short stem 2–15 cm high, about 5–7 cm across, with many narrow violet or purplish blue ray florets and a wide disk about 25–30 mm across. Found occasionally; on open benchland; southern Rocky Mountains.

Vernonia ironweed

Vernonia fasciculata Michx. var. *corymbosa* (Schwein.) Schub.

A coarse erect perennial, usually with a red smooth stem 40–100 cm high. Leaves stalkless, lanceolate to ovate-lanceolate, 7–15 cm long, smooth, and sharply toothed. Inflorescence a loose terminal cluster. Flower heads discoid (with no ray florets), about 6 mm across, dark purple. Rare; has been found beside sloughs and in river valleys; along southeastern Parklands and Boreal forest.

Tabular classification

The heading “varieties” includes subspecies, forms, and small or included species in addition to varieties. Plants represented only by a subspecies or variety are included under species.

	Families	Genera	Species	Varieties
Pteridophyta	7	21	58	2
Spermatophyta				
Gymnospermae	3	8	22	3
Angiospermae				
Monocotyledoneae	16	125	429	66
Dicotyledoneae	92	436	1465	149
Totals	118	590	1974	220



Fig. 230. Parry's townsendia, *Townsendia parryi* D. C. Eaton.

Glossary

- acaulescent** Stemless, or having a very short stem.
- achene** A 1-celled, 1-seeded, dry hard fruit that does not open when ripe (Fig. 9, p. 19).
- acrid** Pungent, bitter.
- acuminate** Gradually tapering to a point.
- acute** Somewhat abruptly tapering to a point.
- adnate** Of a plant part, united with or attached to another part, usually of a different kind.
- alternate** Distributed, as leaves, at different positions on the stem, not opposite each other (Fig. 5, p. 16).
- ament** A scaly spike of flowers of one sex only; also called catkin.
- androgynous** Of spikes in *Carex*, having both staminate and pistillate flowers, the latter at the base.
- angiosperm** A plant bearing seed in a closed ovary.
- annual** A plant germinating, flowering, and ripening seed in 1 year.
- annular** Arranged in rings.
- anther** The pollen container of a stamen or male organ (Fig. 7, p. 17).
- anthesis** In full flower, usually referring to the flowering period.
- apetalous** Without petals.
- apex** The summit or point (Fig. 5, p. 16).
- appressed** Lying flat and close to some part of a plant, usually referring to hairs.
- approximate** Close together, but not overlapping.
- aquatic** Living in water.
- articulation** Natural separation, joint.
- ascending** Growing upward or turned up.
- astringent** Binding, contracting.
- attenuate** Becoming very narrow.
- auricle** An ear-shaped appendage, or the ear at the base of a leaf (Fig. 5, p. 16).
- auriculate** Having auricles.
- awl-shaped** Broad at base, tapering to a sharp point.
- awn** A bristle, often found on grass flowers.
- axil** The upper angle formed where a leaf stalk or a branch joins a stem (Fig. 5, p. 16).
- axillary** In an axil.
- axis** The central line of an organ.
- barb** A short, stiff point, or short bristle, often bent backward.
- basifixed** Attached at or near the base, at one end only.
- beak** A tip or point, somewhat resembling the beak of a bird.
- bearded** Hairy, often used of a stamen or of the throat of a flower.
- berry** A pulpy fruit with several seeds, as that of currant and grape (Fig. 5, p. 16).
- biennial** Of 2 years' duration.
- bifid** Two-cleft.
- bilabiate** Two-lipped.

- bisexual** Having both stamens and pistils.
- blade** The expanded part of a leaf (Fig. 5, p. 16).
- bloom** A whitish, powdery covering.
- bract** A small leaf or scale, often borne below a flower or flower cluster.
- bracteose** Having numerous or conspicuous bracts.
- bractlet** A secondary bract, as one borne on the pedicel of a flower.
- bud** The rudimentary state of a stem or branch; an unexpanded flower.
- bulb** A thick underground organ composed of successive fleshy layers.
- bulblet** A small bulb, borne on the stem or inflorescence.
-
- callus** A small, hard protuberance.
- calyx** The outer floral ring, or sepals, usually green, but sometimes brightly colored (Fig. 7, p. 17).
- campanulate** Bell-shaped.
- canescent** Densely fine pubescent, giving a gray appearance.
- capitate** Gathered into a head, as a cluster at the end of a stem.
- capsule** A dry fruit, as that of the gentian, consisting of more than one chamber, and opening at maturity.
- carpel** An ovule-bearing chamber at the base of the pistil or female organ of a flower.
- caryopsis** A grain, as in the grasses.
- catkin** As used in this book, a scaly spike of flowers of one sex.
- caudex** The woody base of a plant from which the stems arise.
- caulescent** Having a well-defined stem above the ground.
- cauline** Belonging or attached to the stem.
- cell** A chamber of the ovary or anther; an individual unit of plant structure.
- cespitose** Growing in tufts; matted or turf-forming; also spelled caespitose.
- chaff** A small, thin, dry, and membranous scale or bract.
- chartaceous** Having the texture of paper, papery.
- chlorophyll** The green coloring matter within the cells of a plant.
- choripetalous** Having petals separated from each other.
- ciliate** Having marginal hairs (Fig. 5, p. 16).
- circumscissile** Splitting all the way around, as the lid of a fruit, such as that of purslane.
- clasping** Partly or entirely surrounding the stem; used of leaf bases (Fig. 5, p. 16).
- cleft** Deeply lobed.
- cleistogamous** Of flowers, small, inconspicuous, permanently closed, and hence self-fertilizing.
- compound** Of a leaf, composed of 2 or more leaflets; of a branch, composed of 2 or more parts, forming a common whole.
- compressed** Flattened, especially laterally.
- cone** A dense, usually elongated collection of flowers or fruits borne beneath scales; or a collection of spore-bearing leaves on an axis, the whole mass forming a fruit-like body.
- coniferous** Cone-bearing.
- connate** Of leaves, united at the base (Fig. 5, p. 16); joined together.
- convex** Curved outward, as the surface of a sphere.
- convolute** Rolled together, coiled.

- cordate** Of a leaf, heart-shaped, with the point away from the base (Fig. 3, p. 15).
- corm** A thick enlarged base of a stem, as found in crocus and gladiolus.
- corolla** The petals or inner floral ring (Fig. 7, p. 17).
- corymb** A cluster of flowers in which the flower stalks arise from different points on the stem; the cluster has a flat or rounded top (Fig. 6, p. 17).
- cosmopolite** Occurring in all or most parts of the world.
- cotyledon** The first leaf from the seed, sometimes called the seed leaf.
- crenate** Having the margin cut into rounded scallops.
- crown** The place where stem and root meet.
- culm** The stem of a grass or sedge.
- cuneate** Of a leaf, broadly rounded at the apex and tapering rather abruptly toward the point of attachment (Fig. 3, p. 15).
- cylindric** In the form of a cylinder.
- cyme** A cluster of flowers in which the central flowers open first (Fig. 6, p. 17).
- cymose** Of the form or nature of a cyme.
- deciduous** Having leaves that fall off in autumn; not evergreen.
- decompound** More than once compound or divided.
- decumbent** Of a stem, the base lying on the ground, but the tip growing upright.
- decurrent** Of a leaf, the blade extending down the stem (Fig. 5, p. 16).
- deflexed** Turned abruptly downward.
- deltoid** Of a leaf, triangular (Fig. 3, p. 15).
- dentate** Toothed, with teeth pointed and directed outward.
- diadelphous** Of stamens, united in 2, often unequal, bundles.
- dichotomous** Regularly and repeatedly branching in twos.
- dicotyledon** A plant bearing 2 cotyledons, or seed leaves.
- digitate** Of a leaf, with divisions somewhat like fingers (Fig. 4, p. 15).
- dilate** Spreading out in all directions.
- dioecious** Having male and female flowers on separate plants.
- discoid** Having only disk flowers, without ray flowers.
- disk or disc** A more or less fleshy or elevated development of the receptacle about the pistil; the receptacle in the head of Compositae (Fig. 153, p. 524); a flattened extremity, as on tendrils of Virginia creeper.
- dissected** Divided into many segments.
- divaricate** Spreading very far apart; extremely divergent.
- divergent** Spreading away from each other.
- divided** Of a leaf, cleft to the midrib or base (Fig. 4, p. 15).
- dorsal** The backside of an organ, facing away from the axis.
- drupe** A pulpy or fleshy fruit containing a single seed enclosed in a hard shell or stone, as that of the plum.
- drupelet** One part of a fruit composed of aggregate drupes, as in the raspberry.
- edentate** Lacking teeth.
- ellipsoid** Solid but with an elliptical outline.
- elliptical** Of a leaf, oval or oblong with the ends rounded and widest in the middle (Fig. 3, p. 15).

- elongate** Stretched out, lengthened.
- entire** Of a leaf or leaflet, having the margin not toothed or cleft.
- equitant** Of leaves, enfolding each other or borne astride, as in the lily and the iris.
- erose** Appearing as though gnawed at the margin.
- excurrent** Projecting beyond the margin or tip, as an awn.
- exserted** Projecting outward, as stamens from a corolla.
- falcate** Curved and flat; sickle-shaped.
- farinose** Covered with a whitish, mealy powder.
- fascicle** A dense cluster; used of roots, leaves, or flowers.
- fertile** Referring to stamens that bear pollen, and fruits that contain seeds.
- filament** The stalk of a stamen below the anther (Fig. 7, p. 17).
- filiform** Thread-like, long and very slender.
- flaccid** Limp, floppy.
- flexuous** Curved or bending in alternate directions, zigzag.
- floret** A single flower, usually used of a composite head or cluster.
- foliate** Of a leaf, composed of 2 or more parts.
- follicle** A fruit with a single chamber that opens along one side, as in milkweed (Fig. 9, p. 19).
- frond** The expanded leaf-like portion of a fern.
- fruit** The seed-bearing product of a plant.
- funnelform** Of a flower, shaped like a funnel, cone-shaped.
- galea** A hooded or helmet-shaped portion of a perianth, especially the upper part of some 2-lipped corollas, such as in elephant's-head.
- geniculate** Bent, like a knee.
- glabrate** Without pubescence.
- glabrescent** Becoming glabrous with age or at maturity.
- glabrous** Smooth, without hairs.
- gland** An organ that secretes sticky or resinous matter.
- glandular** Bearing glands.
- glaucous** Covered with a bloom; bluish white or bluish gray.
- globose** Spherical or nearly so.
- globular** Globe-like, spherical.
- glomerule** A small compact cluster of flowers, as in Compositae.
- glume** A scaly bract on the floral parts of grasses and sedges.
- glutinous** Sticky.
- grain** A fruit resembling an achene but having the seed coat and thin pericarp fused into one body, particularly the fruit of grasses.
- granular** Covered with very small grains.
- gynaecandrous** Of spikes in *Carex*, having both staminate and pistillate flowers, the latter at the apex.
- hastate** Resembling an arrowhead; of a leaf, with basal lobes protruding sideways.
- head** A dense cluster of flowers or fruits on a very short axis or receptacle.
- herb** A plant without a woody stem above the ground.
- hip** The berry-like enlarged calyx tube containing many achenes, found in roses.

- hirsute** Having coarse spreading hairs.
- hispid** Bearing stiff hairs or bristles.
- hispidulous** Minutely hispid.
- hoary** Grayish white.
- hyaline** Transparent or translucent, glassy.
- hypanthium** The place on a flower head where the sepals and petals are attached.
- imbricate** Overlapping like shingles.
- incised** Having a margin that is cut or slashed irregularly; between toothed and lobed.
- indehiscent** Not opening by valves; remaining closed.
- indurate** Hardened.
- indusium** In ferns, a membranous cover of a spore cluster or sorus; plural, *indusia*.
- inferior** Of an ovary, situated below the calyx and corolla (Fig. 7, p. 17).
- inflorescence** Arrangement of flowers in a cluster (Fig. 6, p. 17).
- internode** The part of an axis between 2 nodes.
- interrupted** Not continuous.
- introduced** Imported from another region for ornamental or cultivation purposes.
- involucre** The whorl of bracts below a flower cluster, or around a flower of Compositae (Fig. 199, p. 684).
- involute** Rolled inward.
- irregular** Of a flower, petals and sepals differing in shape or size (Fig. 8, p. 18).
- keel** The 2 lower united petals of a leguminous flower (Fig. 7, p. 17); a sharp ridge.
- lacerate** Irregularly cleft or cut, as if torn.
- lanate** Covered with soft, intertwined hairs; woolly.
- lanceolate** Of a leaf, much longer than wide, broadest near the base and tapering toward the tip (Fig. 3, p. 15).
- lax** Loose, the opposite of stiff or congested.
- leaflet** A division of a compound leaf (Fig. 4, p. 15).
- legume** A dry pod-like fruit, splitting down one or both sides at maturity (Fig. 9, p. 19).
- lemma** The lower of the 2 bracts enclosing a grass flower (Fig. 16, p. 53).
- lenticular** Lens-shaped, biconvex.
- ligulate** Having a ligule.
- ligule** A strap-shaped organ, as in ray florets of Compositae (Fig. 199, p. 684); also, a collar of a grass blade (Fig. 16, p. 53).
- linear** Of a leaf, long and narrow with parallel margins (Fig. 3, p. 15).
- lip** The main lobe of a 2-lobed corolla or calyx; the odd and peculiar petal of Orchidaceae.
- lobe** A rounded projection of a leaf (Fig. 4, p. 15), or a leaf-like part of a plant.
- locule** The cavity of an ovary or anther.
- loment** A legume or pod, constricted between the seeds, the joints separating at maturity (Fig. 9, p. 19).

lyrate Of a leaf, having a large terminal lobe and smaller lobes toward the base.

malpighian Of hairs, attached in the middle.

membranous Rather soft, thin, and somewhat translucent.

mericarp One of the two parts of the fruit of certain families, especially Umbelliferae.

-merous Having 2 or more parts; -parted.

midrib The central vein of a leaf or other organ.

monocotyledon A plant bearing only 1 cotyledon or seed leaf.

monoecious Having male and female flowers on the same plant.

mucro A stiff, sharp point; plural, mucrones.

mucronate Terminated by a mucro.

nectar A sweet liquid secreted by the nectaries of plants.

nerve A simple or unbranched vein or slender rib.

neutral Of a flower, without stamens or pistils.

node The place on a stem where leaves grow or normally arise; the solid part of a culm.

nut A single-seeded fruit with a woody, hard outer coat (Fig. 9, p. 19).

obcordate Of a leaf, heart-shaped with the attachment at the pointed end.

oblanceolate Of a leaf, much longer than wide, broadest near the tip, and tapering toward the place of attachment (Fig. 3, p. 15).

oblong Longer than broad, having the sides nearly parallel for most of their length (Fig. 3, p. 15).

obovate Of a leaf, egg-shaped having the wide part near the tip (Fig. 3, p. 15).

obovoid Of a plant part, such as a fruit, that is wider near the tip.

obpyramidal Inversely pyramidal.

obtuse Blunt or rounded at the end.

ocrea A loose sheath, composed of 1 or 2 membranous stipules at the base of a leaf stalk.

opposite Borne 2 at a node, on opposing sides of an axis (Fig. 5, p. 16).

orbicular Somewhat circular in outline (Fig. 3, p. 15).

oval Of a leaf, egg-shaped (Fig. 3, p. 15).

ovary The part of a pistil or female organ of a flower containing the cells that become seeds after fertilization.

ovate Of a leaf, egg-shaped with the broad part toward the base (Fig. 3, p. 15).

ovoid Egg-shaped, having the wide part near the point of attachment.

ovule The seed-containing unit of the ovary.

palea The inner of 2 bracts enclosing a grass flower (Fig. 16, p. 53).

palmate Of a leaf, having the shape of a hand with the fingers spread (Fig. 4, p. 15).

paludose Of, or growing in, marshes.

panicle A branched cluster of flowers, each stalked, the lower branches longest and opening first (Fig. 6, p. 17).

paniculate Resembling a panicle.

papilla A minute, nipple-shaped projection.

- papillate** Bearing papillae; also called papillose.
- pappus** The bristly or scale-like appendage on fruits of Compositae (Fig. 199, p. 684).
- paraphysis** A slender sterile filament among the spores of some ferns; plural, paraphyses.
- parasitic** Growing on, and deriving nourishment from, another living plant.
- pectinate** Comb-like.
- pedicel** The stalk of a single flower in a cluster.
- pedicellate** Having or attached by a pedicel.
- peduncle** Stem of a solitary flower or of a flower cluster.
- pedunculate** Having a peduncle.
- peltate** Shield-shaped; of a leaf, attached to its stalk inside the margin (Fig. 3, p. 15).
- pendulous** Hanging down.
- pepo** A berry-like fruit of Cucurbitaceae, with a hard rind and pulpy interior filled with seeds.
- perennial** A plant that persists for more than 2 years.
- perfect** Of a flower, complete, having both stamens and pistil.
- perfoliate** A stalkless leaf with basal portions encircling the stem.
- perianth** Petals and sepals referred to together.
- perigynium** The papery sheath that envelopes the fruit in *Carex*; plural, perigynia.
- persistent** Remaining attached.
- petal** A separate part of a corolla or inner floral ring, usually brightly colored (Fig. 7, p. 17).
- petiole** A stalk of a leaf.
- petiolate** Stalked, having a stalk.
- phyllodium** A leaf-like petiole with no leaf; plural, phyllodia.
- pilose** Sparsely pubescent with long straight hairs.
- pinna** A leaflet or primary division of a pinnate leaf or frond; plural, pinnae.
- pinnate** Of a compound leaf, or frond, with leaflets (pinnae) arranged on each side of a common axis.
- pinnatifid** Cleft or parted in a pinnate way.
- pinnatisect** Cleft pinnately to or almost to the midrib.
- pinnule** A secondary pinna.
- pistil** The female part of a flower, composed of style and stigma.
- pistillate** Having pistils, or female organs; generally used when no male parts are present.
- pith** The spongy center of the stems of most angiosperms.
- plicate** Folded lengthwise, as in a fan.
- plumose** Having fine hairs, resembling a feather.
- pod** A dry fruit, opening when mature.
- pollinium** A mass of waxy pollen, as in Orchidaceae; plural, pollinia.
- polygamous** Having both perfect and unisexual flowers on the same plant.
- pome** A fleshy fruit, as the apple.
- procumbent** Trailing along the ground without rooting at the nodes.
- protuberance** A swelling or bulge.
- puberulent** Minutely pubescent, downy.
- pubescent** Covered with hairs.

- punctate** Dotted, with glandular depressions or colored dots.
- punctulate** Dotted with very small dots.
- pyxis** A capsule, the upper part of which falls off as a lid, as in the fruit of purslane (Fig. 9, p. 19).
- raceme** A flower cluster with each flower borne on a short stalk from a common stem (Fig. 6, p. 17).
- racemose** Composed of or resembling racemes.
- rachilla** Axis of a spikelet (Fig. 17, p. 54).
- rachis** The axis of a spike or compound leaf (Fig. 17, p. 54).
- radiate** Having ray florets (Fig. 199, p. 684).
- ray** In some Compositae, a modified marginal floret with a strap-like extension of the corolla.
- receptacle** The part of a flower stalk bearing the floral organs; the part of a capitulate flower cluster bearing the florets (Fig. 199, p. 684).
- recurved** Curved backward.
- reflexed** Bent sharply backward, or downward.
- regular** Of a flower in which all respective parts are the same size and shape (Fig. 8, p. 18).
- reniform** Of a leaf, somewhat kidney-shaped.
- reticulate** Having the appearance of a net.
- retorse** Bent or curved over, backward or downward.
- retuse** Having a shallow notch at an otherwise rounded apex.
- revolute** Rolled backward or downward.
- rhizome** An underground, root-like stem; rootstock.
- rhombic** Having the outline of an equal-sided oblique diamond shape.
- rootstock** A rhizome.
- rosette** A dense cluster of leaves on a very short stem or axis (Fig. 5, p. 16).
- rotate** Of a flower, wheel-shaped (Fig. 8, p. 18).
- rudimentary** Imperfectly developed and nonfunctional.
- rugose** Wrinkled, corrugated.
- runcinate** Coarsely toothed or cut, the pointed teeth turned toward the base of the leaf.
- sac** A pouch, especially the cavity of an anther.
- sagittate** Resembling the head of an arrow; of a leaf, having the basal lobes pointing toward the place of attachment.
- salverform** Of a flower, having a tube with wheel-shaped expansion on top, as in flax and collomia (Fig. 8, p. 18).
- samara** A winged fruit that does not split open at maturity (Fig. 9, p. 19).
- saprophyte** A plant that lives on dead organic matter.
- scaberulous** Somewhat roughened.
- scabrous** Rough to the touch.
- scale** A dry and appressed modified or reduced leaf or bract.
- scape** A flowering stem growing from the root crown and not bearing proper leaves, as in the tulip.
- scapose** Bearing flowers on a scape.
- scarious** Thin and dry, not green, as in margins of sheaths or bracts.
- scorpioid** Of an inflorescence, uncoiling as the flowers develop.
- secund** One-sided; of flowers that appear to be borne on one side.

- seed** The ripened ovule, consisting of the embryo and its proper coats.
- sepal** One of the separate parts of a calyx, usually green and leaf-like.
- sericeous** Silky hairy.
- serrate** Having sharp teeth pointing forward.
- serrulate** Minutely and finely serrate.
- sessile** Without a stalk (Fig. 5, p. 16).
- sheath** A long tubular structure surrounding some part of a plant.
- shrub** A woody plant that remains low and produces shoots or trunks from the base or caudex.
- silicle** A short silique.
- silique** A capsule with 2 valves separating from a thin longitudinal partition (Fig. 9, p. 19).
- simple** Of a leaf, having a single blade not divided into leaflets.
- sinuate** Wavy-margined.
- sorus** A cluster of spore cases, as in the ferns; plural, sori.
- spadix** A dense or fleshy spike of flowers, as in *Calla*.
- spathe** A large leaf-like bract enclosing a flower cluster.
- spatulate** Spoon-shaped; of a leaf, having a broad rounded tip, gradually narrowing to the point of attachment (Fig. 3, p. 15).
- spike** A flower cluster, the individual flowers of which are stalkless, borne on a common stalk (Fig. 6, p. 17).
- spikelet** A secondary spike, especially in grasses and sedges.
- spinose** Bearing many spines.
- spinulose** Bearing small spines.
- sporangium** A spore case; plural, sporangia.
- spore** A reproductive body, usually of a single detached cell without embryo, as in ferns.
- sporocarp** A receptacle containing spores.
- spur** A hollow projection, usually at the base of a flower, as in the snapdragon.
- squarrose** Sharply recurved at the tips, spreading at right angles.
- stalk** The stem of an organ, such as the petiole, peduncle, filament, or stipe.
- stamen** The male organ of a flower (Fig. 7, p. 17).
- staminodium** A false stamen.
- staminate** Having stamens; the term usually used when no female organs are present.
- standard** The large upper petal (or banner) of a leguminous flower (Fig. 7, p. 17).
- stellate** Star-shaped.
- steppe** Native grassland, prairie.
- stigma** The summit of the style; the part that receives the pollen (Fig. 7, p. 17).
- stipe** Any short stalk, especially that of the pistil; the petiole of a fern leaf.
- stipitate** Borne on a stipe or short stalk.
- stipule** An appendage at the base of a leaf (Fig. 5, p. 16).
- stolon** Basal branch that roots at nodes, often underground.
- striate** Marked with fine longitudinal lines or ridges.
- strigose** Pubescent with appressed straight hairs.
- strobilus** A cone-like aggregation of sporophylls; plural, strobili.

- style** The part of the pistil between the stigma and ovary (Fig. 7, p. 17).
- submersed** Being or growing under water.
- subulate** Awl-shaped, tapering along its entire length.
- succulent** Juicy, fleshy; soft and thickened in texture.
- superior** Of an ovary, above the calyx or hypanthium (Fig. 7, p. 17).
- sympetalous** Having the petals wholly or partly united.
- tendrill** A slender outgrowth by which some plants attach themselves to objects (Fig. 5, p. 16).
- terete** Circular in cross section.
- terminal** At the end of a stem or branch.
- ternate** In threes.
- tomentose** Densely hairy with matted wool.
- translucent** Partly transparent.
- trifid** Cleft part way into 3.
- trifoliate** Having 3 leaves (Fig. 4, p. 15).
- trifoliolate** Having a leaf or leaves of 3 leaflets, as most clovers.
- tuber** A thick underground branch bearing buds.
- tubercle** A small tuber; a rounded protruding body.
- tubular** Hollow and of an elongated or pipe-like form.
- tussock** A dense tuft or bunch, mostly used of grasses and sedges.
- umbel** A flower cluster in which all flower stalks arise from a common point (Fig. 6, p. 17).
- umbellet** A secondary umbel.
- unarmed** Without spines, prickles, or other sharp appendages.
- unisexual** Having only male or female organs.
- utricle** A small bladdery one-seeded fruit, as in amaranth.
- valve** The units or pieces of a capsule or pod; the enlarged inner sepals in *Rumex*.
- venation** The arrangement or disposition of veins.
- ventral** The inside of an organ, facing the axis.
- villous** Bearing long straight hairs.
- verticil** A whorl of similar organs, as flowers or leaves, implanted at a node.
- verticillate** Arranged in verticils.
- viscid** Sticky.
- whorl** A group of 3 or more leaves arising from the same node (Fig. 5, p. 16).
- wing** A thin, dry, or membranous expansion or flat extension or appendage of an organ (see silicle, Fig. 9, p. 19); also the lateral petals of a leguminous flower (Fig. 7, p. 17).
- winter annual** A plant that germinates in the fall and produces seed and dies in the following spring or summer.
- woolly** Covered with entangled soft hairs.
- xerophyte** A plant of very dry habitat.

The spelling of common names

1. One word

- (a) When the modified word is *plant* or a type of plant, as in *leadplant*, *skunkbush*, *peatree*, *peavine*, *jewelweed*, *lungwort*; except when the modifier is more than one word, as in *Kentucky coffee tree*, or a hyphenated word, as in *silk-tassel bush*, or a proper noun, as in *Virginia creeper*.
- (b) When the modified word is a part of a plant, as in *juneberry*, *twayblade*, *buffalobur*, *coneflower*, *leatherleaf*, *bladderpod*, *balsamroot*, *bugseed*, *twistedstalk*, *bluestem*, *buckthorn*, *wormwood*.
- (c) When the modified word is a part of animal anatomy, as in *arrowhead*, *bluelips*, *cattail*, *beardtongue*; except when the modifier is in the possessive case, as in *bird's-eye*, *crane's-bill*.
- (d) When the words are figurative or suggestive, as in *beggarticks*, *fairybells*, *meadowsweet*, *paintbrush*; except when the modifier is in the possessive case, as in *baby's-breath*, or when it is a proper noun or the adjectival form of a proper noun, as in *Venus-slipper*, *Indian-pipe*, or when letters demand separation for ease in reading or pronunciation, as in *morning-glory*.

2. Two words

- (a) When the modified word is taxonomically correct, as in *red clover* (genus *Trifolium*), *alkali grass* (family Gramineae), *fringed milkwort* (genus *Polygala*), *common plantain* (genus *Plantago*).
- (b) When the modifier is the word *common*, *false*, *mock*, *wild*, as in *common camas*, *false flax*, *mock pennyroyal*, *wild chives*.
- (c) When the modifier is a proper noun, as in *Douglas hawthorn*, *Mackenzie's hedysarum*.
- (d) Exceptions in 1(a).

3. Hyphenated

- (a) When the modified word is not taxonomically correct, as in *sweet-clover* (genus *Melilotus*, not genus *Trifolium*), *whitlow-grass* (genus *Draba*, not family Gramineae), *sea-milkwort* (genus *Glaux*, not genus *Polygala*), *water-plantain* (genus *Alisma*, not genus *Plantago*), except in a few instances of spelling of long standing, as *burdock*, *buckwheat*.
- (b) When the modifier is a compound, as in *round-leaved hawthorn*, and whether or not the modified word is taxonomically correct, as in *salt-meadow grass* (family Gramineae), *blue-eyed grass* (genus *Sisyrinchium*, not family Gramineae).
- (c) In certain three- or four-word groups, as *balm-of-Gilead*, *butter-and-eggs*, *grass-of-Parnassus*, *lily-of-the-valley*, *touch-me-not*.
- (d) Exceptions in 1(c), 1(d).

Additions and Corrections to the 1979 Edition

- p. 106 *Bromus tectorum*: add to last sentence “. . . and cultivated fields”
- p. 123 second and third lines, change “floret” to “spikelet”
- p. 256 second line on lead 3 in Liliaceae should read
Leaves variously shaped, not more than 5–6 times as long as wide. 12
- p. 271 add to lead 11 in Orchidaceae as an extra choice
Leaves 2–5, ovate-lanceolate; flowers greenish. *Malaxis*
- p. 304 *Parietaria*: first line, change “opposite” to “alternate”
- p. 313 *Polygonum*: fourth line in paragraph, change “seeds” to “fruits”
- p. 314 *Polygonum*: lead 2, change “fruit” to “outer calyx”
- p. 315 Lead 6 should read
6. Achene 1.5–2.0 mm wide; flowers pale pink to white. *P. lapathifolium*
Achene 2.0–3.0 mm wide; flowers greenish. *P. scabrum*
- p. 324 Chenopodiaceae: in last line of lead paragraph, change “one” to “two” (to read “but two species are poisonous”)
- p. 407 lead 2 should read:
2. Petals longer than 10 mm. *E. asperum*
Petals shorter than 10 mm. 3
- p. 587 bottom full line: change “eastern” to “western”
- p. 612 *Lappula*: lead 2 should read
2. Flowers 1.5–3 mm across. *L. deflexa* var. *americana*
Flowers 5–8 mm across. 3
- p. 702 *Taraxacum*: lead 1 in second half should read
Outer involucre bracts reflexed 2
- p. 734 *Aster junciformis* Rydb.: in fourth line of paragraph, change “cm” to “mm” (to read “15–20 mm”)
- p. 738 *Bidens*: lead 2 should read
2. Ray florets usually conspicuous; leaves stalkless and clasping. *B. cernua*

NOTE: The European–Russian classification of perennial Triticeae (mainly *Agropyron* and *Elymus* species) has come into more general acceptance in North America recently. Because in the European–Russian classification many of the *Agropyron* genus are in the *Elymus* genus, and because the new names are increasingly used in scientific and semi-popular articles, cross-references are given in the following table.

Traditional North American name	European–Russian name
<i>Agropyron albicans</i> (Scribn. & Sm.)	<i>Elymus lanceolatus</i> subsp. <i>albicans</i> (Scribn. & Sm.) D.R. Dewey
<i>Agropyron bakeri</i> E. Nels.	<i>Elymus bakeri</i> (E. Nels.) Löve
<i>Agropyron cristatum</i> (L.) Gaertn.	<i>Agropyron cristatum</i> (L.) Gaertn.
<i>Agropyron desertorum</i> (Fisch. ex Link) Schult.	<i>Agropyron desertorum</i> (Fisch. ex Link) Schult.
<i>Agropyron dasystachyum</i> (Hook.) Scribn.	<i>Elymus lanceolatus</i> (Scribn. & Sm.) Gould
<i>Agropyron elongatum</i> (Host) Beauvois	<i>Elytrigia pontica</i> (Podp.) Holub
<i>Agropyron intermedium</i> (Host) Beauvois	<i>Elytrigia intermedia</i> (Host) Nevskii
<i>Agropyron latiglume</i> (Scribn. & Sm.) Rydb.	<i>Elymus alaskanus</i> (Scribn. & Merrill) Löve subsp. <i>scandicus</i> (Nevskii) Melderis
<i>Agropyron repens</i> (L.) Beauvois	<i>Elytrigia repens</i> (L.) Nevskii
<i>Agropyron riparium</i> Scribn. & Sm.	<i>Elymus lanceolatus</i> (Scribn. & Sm.) Gould
<i>Agropyron scribneri</i> Vasey	<i>Elymus scribneri</i> (Vasey) M.E. Jones
<i>Agropyron smithii</i> Rydb.	<i>Pascopyrum smithii</i> (Rydb.) Löve
<i>Agropyron spicatum</i> (Pursh) Scribn. & Sm.	<i>Pseudoroegneria spicata</i> (Pursh) Löve
<i>Agropyron subsecundum</i> (Link) Hitchc.	<i>Elymus subsecundus</i> (Link) A & D Löve
<i>Agropyron trachycaulum</i> (Link) Malte	<i>Elymus trachycaulus</i> (Link) Gould ex Shinn.
<i>Agropyron trichophorum</i> (Link) Richt.	<i>Elytrigia intermedia</i> subsp. <i>barbulato</i> (Schur) Löve
<i>Agropyron triticeum</i> Gaertn.	<i>Eremopyrum triticeum</i> (Gaertn.) Nevskii
<i>Elymus angustus</i> Trin.	<i>Leymus angustus</i> (Trin.) Pilg.
<i>Elymus arenarius</i> L.	<i>Leymus arenarius</i> (L.) Hochst.
<i>Elymus canadensis</i> L.	<i>Elymus canadensis</i> L.
<i>Elymus cinereus</i> Scribn. & Merrill	<i>Leymus cinereus</i> (Scribn. & Merrill) Löve
<i>Elymus glaucus</i> Buckl.	<i>Elymus glaucus</i> Buckl.
<i>Elymus hirtiflorus</i> Hitchc.	considered a sterile hybrid
<i>Elymus imovatus</i> Beal	<i>Leymus innovatus</i> (Beal) Pilg.
<i>Elymus interruptus</i> Buckl.	<i>Elymus interruptus</i> Buckl.
<i>Elymus junceus</i> Fisch.	<i>Psathyrostachys juncea</i> (Fisch.) Nevskii in Komarov
<i>Elymus virginicus</i> L.	<i>Elymus virginicus</i> L.

Index to common names

(Page numbers of illustrations are in boldface.)

- absinthe 723, **724**
- adder's-mouth 278
 - bog 278
 - green 278
- adder's-tongue 32
- agalinis 639
 - purple 640
 - rough 639
 - slender 640
- agrimony 437
- alder 300
 - black 519
 - green 300
 - speckled 300
- alexanders 563
 - golden 563
 - heart-leaved 563
- alfalfa 485, **486**
- alkali grass 119, **120**
- allocarya 618
 - Scouler's 618
- alumroot 421, 422, **423**
 - alpine 422
 - small-leaved 422
 - sticky 422
- alyssum 393
 - hoary- 396
 - small **390**, 393
- amaranth 342
 - prostrate 342, **343**
- amaryllis 268
- anemone 364
 - Canada 365
 - crocus 366, **367**
 - cut-leaved 365
 - Drummond's 365
 - long-fruited 365
 - narcissus 365
 - small wood 366
 - tall 366
 - wood 365
 - yellow 366
- angelica 550
 - kneeling 550
 - Lyll's 550
- anis 551
- apple, balsam- 677
- arbor-vitae 61
- arethusa 271
- arnica 718
 - alpine 719
 - Cordilleran 721
 - heart-leaved 719, **720**
 - lawless 719
 - leafy 719
 - long-leaved 721
 - mountain 721
 - narrow-leaved 722
 - nodding 721
 - rock 721
 - shining 719, **720**
 - spear-leaved 721
 - twin 722
- arrow-grass 68
 - marsh 68
 - seaside 68, **69**
- arrowhead 18, 70
 - arum-leaved 70, **72**
 - broad-leaved 70
- arrowwood, downy 675
- artichoke, Jerusalem 764
- arum 246
 - water- 246
- ash 19, 587
 - black 587
 - green 587, **588**
 - mountain 461
 - western mountain 460
- ash-spiraea 460
- asparagus 258
- aspen, large-toothed 281
- aspen poplar 281
- asphodel 266
 - bog 266
 - sticky 266
- aster 728
 - alpine 730
 - Arctic 737
 - canescent 731
 - Eaton's 731
 - elegant 734
 - few-flowered 736
 - flat-topped white 737
 - golden- 741
 - hairy golden- 742
 - large northern 736
 - leafy-bracted 737
 - Lindley's 731, **732**

Maccalla's 736
 many-flowered 734
 meadow 731
 New England 736
 purple-stemmed 737
 rayless 731
 rush 734
 showy 731, **733**
 small blue 737
 smooth 734, **735**
 western 730
 western mountain 736
 western silvery 737
 white prairie 734
 white wood 736
 willow 734
 wood 734
 atriplex 326
 garden 327, **328**
 Nuttall's 327
 red-leaved garden 327
 silvery 327
 avens 443
 large-leaved 443
 mountain- 442
 purple 443
 three-flowered 444, **445**
 white mountain- 442
 yellow 443
 yellow mountain- 442
 awl-wort 416
 water 416
 awn, red three- 99
 axyris 329
 azalea, alpine 576

 baby-blue-eyes 606
 small 606
 baby's-breath 353
 annual 353
 sticky 353
 balm 626
 honey 626
 balsam-apple 677
 balsam fir 57
 balsamroot 738
 baneberry 363
 red 363
 white 363
 barberry 382, 384
 barley 137
 little 137
 wild 137, **138**
 bartsia 640

 bassia 329
 five-hook 329
 basswood 520
 bay, Lapland rose- 577
 rose- 577
 white rose- 577
 bayberry 299
 beach-pea 480
 beak-rush 241
 slender 241
 white 241
 bean, buck- 592
 golden- 19, 497, **500**
 bearberry 18, 572, 573
 alpine 573
 beardtongue 652, **655**
 blue 653
 creeping 653
 crested 653
 large-flowered 654
 lilac-flowered 654, **655**
 shrubby 653
 slender 654, **655**
 smooth blue 654, **655**
 white 653
 yellow 653
 bear-grass 267
 bedstraw 667
 ladies' 670
 marsh 668
 northern 668, **669**
 small 668
 sweet-scented 668
 beech 303
 beech-fern 42
 beggar's-lice 476
 beggarticks 738
 common 739
 smooth 739
 tall 739
 bellflower 677
 marsh 678
 bells, velvet 640
 bellwort 267
 small 267
 bent grass 92
 northern 92
 bergamot, western wild 628
 wild 626, **627**
 bilberry 579
 dwarf 579
 bindweed 317
 field 597, **598**
 hedge 599
 inland 599

birch 299, 300
 canoe 301
 river 301
 scrub 300
 swamp 300
 white 301
 bird-rape 397
 bird's-eye 658
 birthwort 308
 biscuit-root 557
 bishop's-cap 424
 bistort 314
 alpine 320
 western 317
 bitter-root, dwarf 346
 bittersweet 18, 517, 635, **636**
 bitterweed 699
 black-eyed Susan 773
 black medick 485
 black torch 656
 bladderpod 19, 412, 414
 northern 412
 sand **390**, 412, **413**
 spatulate 412
 twin 414
 bladderwort 659, 660
 flat-leaved 660
 greater 660, **662**
 horned 660
 lesser 660
 blazingstar 766
 dotted 766, **767**
 meadow 766
 blite, erect sea- 339
 sea- 339
 strawberry 331
 western sea- 339
 bloodroot 385
 bluebell 677
 Alaska 678
 alpine 678
 clustered 678
 creeping 678
 bluebells, desert 606
 blueberry 578
 bluebur 612
 bluebuttons 681
 blue-eyed Mary 644
 blue grass 158
 alkali 163
 alpine 160
 annual 160
 arctic 161
 big 160
 bog 163
 Canada 161
 Canby 161, **162**
 early 161
 fowl 164
 glaucous 163
 Kentucky 164
 pacific 163
 plains 161
 rough-stalked 165
 Sandberg's 165
 skyline 161
 timberline 165
 Wheeler's 164
 wood 164
 bluelips 644
 bluestem 96
 big 96, **97**
 little 96, **98**
 sand 96
 bluets 670
 long-leaved 670
 bog-myrtle 299
 bog orchid 276
 Alaska 277
 green-flowered **272**, 277
 Hooker's 276
 round-leaved 277
 slender 277
 small northern 277
 white 276
 bog-rosemary 572
 bog-sedge 241
 simple 241
 bog violet, northern **528**, 529
 bog willow 296
 boisduvalia 536
 smooth 536
 boltonia 739
 boneset 755
 borage 609, 610
 box elder 517
 boxwood 517
 mountain 517
 brake, cliff 44
 purple cliff 44
 rock 37, 38
 bramble, dwarf 460
 breadroot 495
 Indian 497, **498**
 brickellia 739
 large-flowered 739
 brome 104
 Canada 106
 field 106
 fringed 105

northern awnless 106
smooth 105
bronzebells 266
broom-rape 661
 clustered 661
 Louisiana 663
 small 663
broom, sweet- 19, 476
broomweed 758
 common 758
buck-bean 592
buckbrush 674
buckhorn 664
buckthorn 518
 alder-leaved 518
buckwheat 308, 310
 false 320
 Tartary 310, **312**
 wild 314, 317
buffaloberry 534, **535**
 Canada 534
buffalobur 635
bugloss 615
 purple 611
 small 615
 viper's 611
bugseed 335
 villose 335, **336**
bulrush 242
 Clinton's 243
 dwarf 244
 great 245
 green 243
 Nevada 244
 prairie 244
 red 244
 river 244
 small-fruited 244
 three-square 243
 Torrey's 245
 tufted 243
 viscid great 243
bunchberry 565, **566**
bur-clover 485
burdock 717
 common 717
 cotton 717
 lesser 717
bur oak 303
bur-ragweed 685
bur-reed 63
 broad-fruited 65
 many stalked 65
 narrow-leaved 63

northern 65
small 65
stemless 65
bush-cranberry 675
 high 675
 low 675, **676**
bush-honeysuckle 671
butter-and-eggs 645
buttercup 374
 boreal 379
 bristly 380
 celery-leaved 381
 creeping 380
 dwarf 379
 early 377
 graceful 379
 hairy 381
 heart-leaved 377, **378**
 Lapland 379
 Macoun's 379
 northern 380
 Pallas 380
 prairie 380
 pygmee 380
 seaside 377
 shiny-leaved 379
 smooth-leaved 376
 snow 380
 swamp 381
 tall 376
 western 380
butterflyweed 539
butterweed, Arctic 779
 marsh 777
 mountain 777
butterwort 659, 660
 small 660
 western 660

cactus 531
 ball 531
 purple 18, 531
calamus 246
calla, water 246
camas 260, 268
 common 260
 death 268, **269**
 smooth 268
campion 353
 bladder 357
 Macoun's 358
 moss 357
 Siberian 358
cancerroot 661

caper 386
 caragana, common 475
 caraway 551
 carrionflower 265
 carrot 553
 wild 553
 catchfly 355
 Menzies 358
 night-flowering 358
 sleepy 357
 smooth 357
 catnip 628
 cat's-ear 695
 cattail 61
 common 63, **64**
 narrow-leaved 61
 cedar, western red 61
 white 61
 chaffweed 581
 chaliceflower 365
 chamaerhodos 438
 chamomile 766
 scentless 768, **769**
 wild 768
 checkerberry 575
 cherry, black-fruited choke 455, **456**
 choke 19
 ground- 633
 large white ground- 634
 low sand 455
 pin 455
 prairie ground- 634
 red-fruited choke 455
 small yellow ground- 634
 yellow ground- 634
 chess, downy 106, **107**
 Japanese 105
 chickweed 18, 351
 alpine 352
 common 361
 field 352
 long-stalked 352
 mouse-eared 352
 chicory 691
 chives, wild 258
 choke cherry 19
 black-fruited 455, **456**
 red-fruited 455
 cicely, blunt-fruited sweet 559
 smooth sweet 559, **560**
 sweet 19, 559
 western sweet 559
 cinquefoil 444
 alpine 454
 branched 451
 brook 452
 bushy 452
 Drummond's 449
 early 448
 five-fingered 452
 graceful 449
 low 452
 marsh 451
 mountain 449
 plains 448
 prairie 452, 453
 rough 451
 rough-fruited 452
 shrubby 449, **450**
 silvery 448
 smooth-leaved 448
 snow 451
 sticky 449
 three-toothed 454
 white 448
 woolly 451
 clammyweed 389
 cleavers 668
 clematis 369
 cleome, pink 386
 cliff brake 44
 purple 44
 cliff daisy 750
 cloudberry 458
 clover 497
 alsike 501
 bur- 485
 hairy prairie- 495
 owl's- 648
 prairie- 493
 purple prairie- 495, **496**
 red 501
 Spanish 482
 sweet- 487
 white 501
 white prairie- 495
 white sweet- 487, **488**
 yellow 501
 yellow field 501
 yellow sweet- 487
 club-moss 51
 little 55
 stiff 52
 trailing 52, **53**
 cockle, corn 348
 cow 355, **356**
 Drummond's 354
 mountain 354

nodding 354
 purple 348, **349**
 white 354
 cocklebur 19, 686, **688**
 cohosh, blue 384
 collomia 18, 600
 narrow-leaved 600, **601**
 colt's-foot 768
 arrow-leaved **771**
 palmate-leaved **771**
 vine-leaved **771**
 columbine 366
 blue 368
 crimson 368
 small-flowered 368
 wild 368
 yellow 368
 comandra 305
 northern 305
 pale 308
 Richards 308
 comfrey 618, 619
 rough 618
 wild 611
 composite 683, **684**
 coneflower, brown 770
 long-headed 770, **772**
 purple 746
 tall 773
 convolvulus 597
 coralroot 271
 early 273
 large 273
 striped 273
 cornflower 740
 corydalis 386
 golden 386
 pink 386, **387**
 costmary 741
 cotoneaster 438, 440
 cotton-grass 239
 alpine 244
 beautiful 240
 close-sheathed 239
 one-spike 240
 russet 240
 sheathed 240
 slender 240
 tall 239
 thin-leaved 240
 cottonwood 281
 narrow-leaved 280
 couch grass 87
 cow-parsnip 19, 555, 556
 cow-wheat 645, **646**
 crabgrass 119
 smooth 119
 cranberry 578
 bush- 675
 dry-ground 580
 high bush- 675
 low bush- 675, **676**
 swamp 578
 crane's bill, small-flowered 505
 cress, American winter 396
 Austrian 414
 bitter 398, 400
 alpine 400
 meadow 400
 small 400
 garden 411
 hoary 400
 globe-podded 401
 heart-podded 401
 lens-podded 401
 mountain 400
 mouse-ear 417
 rock 394
 Drummond's 395
 hirsute 395
 lyre-leaved 395
 purple 395
 reflexed 396
 slender 415
 spring 400
 winter 396
 yellow 414
 creeping 415
 marsh 415
 spreading 415
 cross-thistle 553
 crowberry 514
 black 514
 crowfoot 362
 crown-vetch 475
 field 475
 cryptanthe 610
 Fendler's 611
 small 611
 cucumber, wild 677
 cudweed 757
 clammy 758
 common 757
 low 758
 western marsh 757
 Culver's-root 659
 currant 427
 golden 428

mountain 429
 northern black 428
 red 428
 sticky 429
 swamp red 429
 wild black 428
 cymopterus 553
 plains 553, **554**
 cypress 60
 summer- 338

daisy
 Arctic 741
 cliff 750
 dwarf 754
 hairy 754
 large-flowered 753
 ox-eye 741
 pale 753
 purple 754
 wandering 754
 wild 753
 woolly 753
 yellow 751
 yellow alpine 753
 dame's-rocket 409
 dandelion 702
 dwarf 695
 false 690, 691
 large-flowered false 691
 orange false 690
 prairie false 699
 red-seeded 702
 darnel 141
 Persian 141, **142**
 dead-nettle 624
 white 624
 devil's-club 545
 dewberry 460
 diapensia 580
 northern 580
 dill 550
 dillseed 550
 ditch-grass 67
 western 67
 dock 320
 curled 321
 field 323
 golden 323
 great water 323
 long-leaved 323
 narrow-leaved 324
 narrow-leaved field 324
 sand 324, **325**
 sour 321
 toothed 323
 western 323
 dodder 599
 common 600
 dogbane 592
 intermediate 592
 spreading 592, **594**
 dogwood 563
 alternate-leaved 565
 panicled 567
 red-osier 565
 spotted 567
 doorweed 313, 315, **316**
 northern 317
 Douglas-fir 60
 douglasia 583
 downingia 680
 dragonhead 623
 American 623
 false 621, 628
 dragon's-mouth 271
 dropseed 174
 annual 176
 prairie 176
 sand 174, **176**
 duckweed 245
 ivy-leaved 245
 larger 245
 lesser 245
 dwarf-primula, mountain 583

earth-smoke 386
 elder 674
 box 517
 marsh- 686
 elderberry, red 674
 elephant's-head 650, **651**
 ellisia 605
 elm 19, 303
 American 303
 enchanter's-nightshade 536
 large 536
 small 536
 endive 691
 endolepis 335
 eriogonum, branched 310
 eryngo 553
 evening-primrose 534, 540
 gumbo 541, **542**
 shrubby 543
 upland 540
 western yellow 541
 white 543
 yellow 541

eveningstar 531, **532**
 everlasting 709
 alpine 713
 brown-bracted mountain 716
 corymbose 713
 cushion 713
 eastern 716
 Howell's 714, **715**
 low 713
 pearly 709
 plantain-leaved 716
 racemose 716
 rosy 716
 Russell's 716
 showy 716
 silvery 714
 small-leaved 716
 tall 713
 woolly 714
 eyebright 644, 648
 late-flowering 648
 northern 644

 fairybells 260, **261**
 Oregon 260
 false hellebore 267
 green 267
 false spikenard 265
 felwort 591
 marsh 592
 fern 32, 35
 beech- 42
 bladder 38
 bracken 44
 brake 44
 chamomile-leaved grape 33
 crested shield 41
 fragile 38, **40**
 fragrant shield 42
 grape 32, 33
 holly 44
 interrupted 35
 lady 37
 lip 37
 maidenhair 36
 male 41
 marsh 42
 moon 33
 mountain bladder 38
 ostrich 42, **43**
 parsley 37, **39**
 royal 35
 sensitive 42
 shield 38
 spinulose shield 41
 thick-leaved grape 33
 Virginia grape 33, **34**
 fescue 127
 Idaho 129, **130**
 meadow 129
 northern rough 128
 plains rough 129
 red 131
 rough 128
 sheep 131
 six-weeks 131
 tall 129
 fiddle heads 42
 fiddle-neck 610
 field-pea, Austrian 472, **473**
 figwort 638, 654
 hare 654
 fir 57
 balsam 57
 Douglas- 60
 ground- 52
 fireweed 19, 537, **538**
 broad-leaved 539
 flag 269
 blue 269
 sweet 246
 water 269
 flax 19, 507
 false **390**, 398
 flat-seeded false 398
 grooved yellow 508
 large-flowered 508, **509**
 Lewis wild 507
 meadow wild 508
 small-seeded false 398
 toad- 644
 broad-leaved 645
 field 645
 Maroccan 645
 yellow 645
 fleabane 748
 Canada 751
 compound 751
 creeping 751
 dwarf 754
 dwarf daisy 754
 hirsute 753
 northern daisy 750
 Philadelphia 754
 rough 750
 showy 754
 smooth 753
 tall 751
 tufted 751, **752**

flixweed 402
 northern 402
 forget-me-not 616, 618
 field 616
 marsh 616
 small 616
 four-o'clock 344
 foxtail 168
 alpine 94
 green 171, **172**
 meadow 96
 short-awned 94
 water 94, 96
 yellow 168
 fritillary 262
 frog's-bit 73
 frostweed 525
 fumitory 385, 386
 climbing 385

 gaillardia 755
 great-flowered 755, **756**
 gale, sweet 299
 galinsoga 757
 garden-rocket 406
 gaura, scarlet 539
 gentian 18, 589
 alpine 591
 closed 590
 closed marsh 591
 downy 591
 fringed 591
 moss 590
 mountain 590
 northern 590
 northern fringed 591
 oblong-leaved 590
 spurred- 592
 geranium 19, 504
 Bicknell's 504
 Carolina wild 505
 sticky purple 505, **506**
 wild white 505
 germander 631
 hairy 631
 giant-hyssop 621, **622**
 gilia 600
 clustered 602
 scarlet 602
 ginger, wild 308
 ginseng 544
 glasswort 339
 globeflower 382
 American 382
 globe-thistle 746, 748

 goat's-beard 702, 703
 yellow 703
 golden-aster 741
 hairy 742
 golden-bean 19, 497, **500**
 goldenrod 780
 alpine 784
 canescent 782
 flat-topped 782
 graceful 782, **783**
 late 782
 low 784
 marsh 785
 mountain 785, **787**
 pale 782
 Riddell's 785
 sharp-toothed 784
 showy 784
 stiff 785, **786**
 upland white 785
 velvety 784
 goldthread 371
 gooseberry 19
 northern 429, **430**
 swamp 428
 goosefoot 324, 329
 Fremont's 333, **334**
 many-seeded 333
 maple-leaved 333
 narrow-leaved 333
 red 333
 saline 333
 spear-leaved 338
 gourd 677
 goutweed 549
 grama 100
 blue 100, **103**
 side-oats 100
 grape 519
 riverbank 519
 grass 73
 alkali 119, **120**
 arrow- 68
 marsh 68
 seaside 68, **69**
 barnyard 121, **122**
 bear- 267
 beard 165
 bent 92
 northern 92
 blue 158
 alkali 163
 alpine 160
 annual 160
 arctic 161

big 160
 bog 163
 Canada 161
 Canby 161, **162**
 early 161
 fowl 164
 glaucous 163
 Kentucky 164
 pacific 163
 plains 161
 rough-stalked 165
 Sandberg's 165
 skyline 161
 timberline 165
 Wheeler's 164
 wood 164
 blue-eyed 270
 common 270
 brook 112
 buffalo 108
 false 146
 canary 154
 reed 154, **155**
 cord 171
 alkali 171, **173**
 prairie 174
 cotton- 239
 alpine 244
 beautiful 240
 close-sheathed 239
 one-spike 240
 russet 240
 sheathed 240
 slender 240
 tall 239
 thin-leaved 240
 couch 87
 creeping love 127
 cut 140
 rice 140
 ditch- 67
 western 67
 hair 117
 annual 117
 mountain 117
 rough 92, **93**
 slender 119
 tufted 117, **118**
 holy 135
 Indian 171
 June 137, **139**
 manna 132, 133
 fowl 135
 graceful 133
 northern 133
 small-flowered 133
 tall 133, **134**
 melic 141
 millet 143
 mutton 163
 needle 175
 Columbia 177
 green 179, **180**
 Richardson 177
 nut- 234
 awned 235
 Houghton's 235
 sand 235
 straw-colored 235
 oat 115, 135
 California 115
 Hooker's 135
 one-spike 115, 117
 Parry 115
 poverty 115
 purple 168, **169**
 timber 115, **116**
 onion 141
 Alaska 143
 purple 143
 orchard 112, **114**
 panic 151
 long-stalked 152
 pepper- 410
 branched 411
 broad-leaved 411
 common 411
 perfoliate 411
 roadside 411
 porcupine 177
 western 177
 quack 87, **88**
 rabbitfoot 165
 reed 108
 common 158
 giant 158
 marsh 109
 narrow 110
 northern 109
 pine 110
 plains 110, **111**
 purple 110
 rice 146
 Canadian 148
 Indian 148, **149**
 little 148
 little-seed 150
 northern 150
 white-grained mountain 148

rye 140
 Italian 140
 perennial 140
salt 119
salt-meadow 166
 Nuttall's 166, **167**
 slender 166
sand 112, **113**
scratch 144
scurvy- 401
sea lime 124
slough 100, **102**
spear 177, **178**
star- 268
 yellow 268
sweet 135, **136**
switch 152, **153**
tumble 166
wedge 174
 prairie 174
 slender 174
whitlow- 403
 alpine 404
 creeping 406
 few-seeded 406
 golden 404
 hairy 405
 thick-leaved 404
 yellow **390**, 405
witch 151
wood 112
 slender 112
wool- 243
grass-of-Parnassus 425
 fringed 425
 glaucous 425
 northern 425, **426**
 small 425
grass-pink 271
 purple 271
greasewood 339, **340**
gromwell 615
 corn 613
 false 618
 western false 618
 woolly 615
ground-cherry 633
 large white 634
 prairie 634
 small yellow 634
 yellow 634
ground-fir 52
ground-ivy 623
ground-pine 52, **54**
ground-plum 19, **469**
groundsel 774
 balsam 779
 common 780
 compact 780
 entire-leaved 777
 silvery 775, **776**
 stinking 780
gumweed 758
hackberry 303
hair grass 117
 annual 117
 mountain 117
 rough 92, **93**
 slender 119
 tufted 117, **118**
harebell 18, 678, **679**
hare's-ear mustard 19, **390**, 401
hawk's-beard 691
 dwarf 692
 green 692
 narrow-leaved 694
 scapose 692
 small-flowered 692
 smooth 692, **693**
hawkweed 694
 alpine 695
 Canada 695
 orange 694
 white 694
 woolly 694
hawthorn 440
 Douglas 440
 long-spined 440
 round-leaved 440, **441**
hazel, American 19
 beaked 19
hazelnut 301
 American 301, **302**
 beaked 301, **302**
heads, fiddle 42
heath 571
heather, false 526
 mountain- 573, 576
 blue 577
 purple 577
 western 573
 white 573
 yellow 577
 sand- 526
hedge-hyssop 644
 clammy 644
hedge-nettle 629
 marsh 18, 629, **630**

- hedysarum, American 476, **478**
 Mackenzie's 479
 northern 479
 yellow 479
 heliotrope 612
 mountain 681
 spatulate-leaved 612
 hellebore 267
 false 267
 green false 267
 hemlock 553
 bulb-bearing water- 551
 ground 56
 poison 553
 water- 551, **552**
 hemp 304
 Indian- 593
 hemp-nettle 623
 yellow 623
 henbane 631
 black 19, 631, **632**
 henbit 624
 hepatica 374
 hoary-alyssum 396
 hog-peanut 464
 hollyhock, mountain 520
 wild 520
 honewort 553
 honeysuckle 670, 671
 blue fly 673
 bush- 671
 hairy 673
 involucrate 673
 swamp fly 673
 Tartarian 673
 twining **672**, 673
 Utah 673
 hop 304
 common 304
 hopbeam, horn 303
 horehound 625
 common 625
 northern water- 625
 western water- 625
 horned-pondweed 67
 hornwort 361
 horse-radish 396
 horsetail 47
 common 48, **49**
 marsh 50
 meadow 50
 swamp 48
 variegated 50
 woodland 50
 hound's-tongue 611
 huckleberry 577
 hymenopappus 764
 tufted 764
 hyssop 624
 clammy hedge- 644
 giant- 621, **622**
 hedge- 644

 Indian-hemp 593
 Indian-pipe 570, 571
 indigo, dwarf false 464
 false 463
 iris 268
 ironplant 758
 spiny 759, **760**
 toothed 759
 ironweed 790
 western 790
 ironwood 303
 ivy, ground- 623
 poison- 515, **516**

 Jacob's-ladder 603
 showy 603, **604**
 sticky 605
 jewelweed, western 510
 jimsonweed 631
 Joe-pye weed, spotted 755
 juneberry 437
 juniper 60
 creeping 61, **62**
 low 60
 rocky mountain 61

 kittentails 640
 knapweed, diffuse 740
 Russian 740
 spotted 740
 knotweed 313
 bushy 318
 Douglas 317
 dwarf 318
 erect 317
 least 318
 slender 317
 small 317
 striate 315
 kochia 338

 Labrador paintbrush 642
 Labrador-tea 575, 576
 laceflower 435

lady's-slipper 18, 273
 mountain 275
 northern 275
 ram's-head 275
 showy 275
 small white 275
 small yellow **272**, 275
 stemless 273, **274**
 yellow 275
 lady's-thumb 318, **319**
 lady's-tresses 279
 hooded 279
 slender 279
 lamb's-quarters 331, **332**
 lantern plant, Chinese 633
 larch 57
 alpine 57
 larkspur 371
 low 18, 19, 371, **372**
 tall 371, **373**
 laurel, pale 575
 sheep- 575
 sticky 518
 lavas 555
 lavauxia, yellow 541
 leadplant 463
 leatherleaf 573, **574**
 lemonwood 723
 lettuce 695
 annual blue 696
 blue 696
 glaucous white 700
 Indian 346
 lobed prickly 696
 tall blue 696
 tall yellow 696
 western 696
 white 699
 licorice, wild 19, 476, **477**
 lily 256, 262
 bluebead 260
 corn 260
 fawn 262
 fragrant water- 362
 glacier 262, **263**
 Mariposa 258
 sand- 531
 small pond- 362
 small water- 362
 water- 361, 362
 western red 262, **264**
 wood 262
 yellow pond- 361, 362
 lily-of-the-valley, wild 262
 linanthus 602
 linden 520
 liverleaf 374
 loasa 531
 lobelia 680
 Kalm's 680
 spiked 680
 water 680
 locoweed 489
 Bessey's 490, **491**
 early yellow 493, **494**
 late yellow 490
 purple 492
 reflexed 492
 showy 493
 viscid 493
 woolly 493
 loosestrife 534, 583
 fringed 584, **585**
 lance-leaved 584
 purple 534
 tufted 586
 whorl-leaved 584
 lopseed 663
 lousewort 648
 coiled-beak 649
 common 649
 flame-colored 650
 Labrador 650
 large-flowered 649
 leafy 652
 Oeder's 650
 purple 652
 swamp 650
 western 649
 woolly 650
 lovage 555
 luetkea 444
 lungwort 615
 lance-leaved 615
 large-flowered 616
 seaside 616
 tall 616, **617**
 lupine 482
 alpine 483
 flexile 485
 large-leaved 483
 Nootka 483
 silvery 483, **484**
 small 483
 madder 667
 madwort 610
 mahonia 384

maidenhair 36
 mallow 520
 common 521
 crisp 521
 false 522
 musk 521
 pink 521
 purple 522, **523**
 round-leaved 521
 scarlet 522, **524**
 small-flowered 521
 mallows, gay 520
 maple 19, 515
 Manitoba 517
 mountain 517
 silver 517
 white 517
 mare's-tail 543
 marigold, floating marsh- 369
 marsh- 368, 369, **370**
 mountain- 368
 water- 739
 marsh-elder 686
 marsh-marigold 368, 369, **370**
 floating 369
 marsilea 47
 Mary, blue-eyed 644
 mayflower 573
 mayweed 717
 stinking 717
 meadow-rue 381
 few-flowered 382
 tall 382, **383**
 veiny 382
 western 382
 meadowsweet 461
 narrow-leaved 461
 pink 461
 shining-leaved 461
 medick 485
 black 485
 menziesia 576
 mignonette 417
 white cut-leaved 417
 yellow cut-leaved 417
 milfoil 708
 pinnate water- 544
 spiked water- 544
 water- 543, 544
 milk-vetch 19, 464
 alpine 467
 American 470, **471**
 ascending purple 474
 Canadian 469
 cicer 469
 cushion 470
 Drummond's 469
 few-flowered 475
 Indian 467
 loose-flowered 475
 low 472
 Missouri 472
 narrow-leaved **468**, 472
 prickly 472
 purple 469
 Pursh's 474
 racemose 474
 slender 470
 timber 472
 tufted 474
 two-grooved 467, **468**
 milkweed 19, 593
 dwarf 595
 green 597
 hairy 595
 showy 595, **596**
 silky 597
 swamp 595
 whorled 597
 milkwort 510
 fringed 510
 sea- 583
 white 510
 whorled 510
 millet 150
 broomcorn 152
 sand 152
 soft 151
 minniebush, western 576
 mint 620, 626
 field 626
 wild 626
 mistletoe 308
 American 308
 dwarf 308
 miterwort 424
 Brewer's 424
 false 435
 monkeyflower 18, 647
 blue 647
 Lewis 647
 smooth 647
 yellow 647
 monkshood 363
 monk's-wort 618
 red 618
 monolepis 338
 moonseed 384

- morning-glory 18, 597
 wild **598**, 599
 moschatel 677
 moss, club- 51
 little club- 55
 mountain 419
 spike- 55
 stiff club- 52
 trailing club- 52, **53**
 motherwort 624
 Siberian 625
 mountain ash, western 460
 mountain-avens 442
 white 442
 yellow 442
 mountain-heather 573, 576
 blue 577
 purple 577
 western 573
 white 573
 yellow 577
 mountain-marigold 368
 mountain moss 419
 mountain parsnip 550
 mousetail 374
 least 374
 mud-purslane 525
 mudwort 644
 muhly 143
 bog 144
 foxtail 144, **145**
 marsh 146
 mat 146, **147**
 prairie 144
 wood 146
 mullein 656
 common 656
 woolly 656
 musineon 559
 leafy 559
 mustard 18, 389, **390**, 396
 ball **390**, 412
 dog 407
 gray tansy 402
 hare's-ear 19, **390**, 401
 hedge 416
 Indian 397
 narrow-leaved 416
 perennial 414
 short-fruited tansy 402
 tall hedge 416
 tansy 401
 tower **390**, 395
 treacle 407
 tumbling 416
 white 397
 wild **390**, 397
 wormseed 407

 naiad 66
 slender 66
 nannyberry 675
 navarretia 602
 small **601**, 602
 nettle 304
 dead- 624
 English 305
 hedge- 629
 hemp- 623
 marsh hedge- 18, 629, **630**
 stinging 305, **306**
 white dead- 624
 wood 304
 yellow hemp- 623
 nettletree 303
 New Jersey tea 518
 nightshade 634
 black 635
 enchanter's- 536
 large enchanter's- 536
 small enchanter's- 536
 ninebark 444
 mallow-leaved 444
 nipplewort 697
 common 697
 nut-grass 234
 awned 235
 Houghton's 235
 sand 235
 straw-colored 235

 oak 19, 303
 bur 303
 oakfern 41
 oat 99, 100
 wild 100, **101**
 oat grass 135
 California 115
 Hooker's 135
 one-spike 115, 117
 Parry 115
 poverty 115
 purple 168
 timber 115, **116**
 oleaster 533
 olive 587
 Russian 533

- onion 258
 - nodding 258
 - pink-flowered 258
 - prairie 258, **259**
- orache 329
 - halberd-leaved 329
- orange, mock 427
- orchard grass 112, **114**
- orchid **272**, 279
 - bog 276
 - Alaska 277
 - green-flowered **272**, 277
 - Hooker's 276
 - round-leaved 277
 - slender 277
 - small northern 277
 - white 276
 - long-bracted 277
 - round leaved **272**, 279
- orchis 270
- oreocarya, clustered 611
- orpine 418
- owl's-clover 648
- ox-eye 764
- ox-eye daisy 741
- oxtongue 610
 - bristly 699

- paintbrush, downy 642
 - hispid 641
 - Indian 640
 - Labrador 642
 - lance-leaved 642
 - purple 642
 - red Indian 642, **643**
 - scarlet 641
 - stiff yellow 641
 - yellow 641
- pansy 530
 - wild 529
- parilla, yellow 384
- parsley 545
 - hairy-fruited 557, **558**
 - long-fruited 557
 - prairie 555
 - Sandberg's wild 557
 - western wild 557
 - white-flowered 557
- parsnip 561
 - cow- 19, 555, **556**
 - mountain 550
 - mountain wild 557
 - water- 563, **564**
 - wild 561

- pea 462
 - Austrian field- 472, **473**
 - beach- 480
- peanut, hog- 464
- pearlwort 354, 355
 - cushion 355
 - mountain 355
 - spreading 355
- pear, prickly- 531
- peatree, Siberian 475
- peavine, wild 480
- pellitory 304
 - American 304
- pennycress 417
- pennyroyal, mock 624
 - rough 624
- pepper, water- 318
- pepper-grass 410
 - branched 411
 - broad-leaved 411
 - common 411
 - perfoliate 411
 - roadside 411
- pepperwort 47, 410
 - hairy 47
- persicaria, pale 318
 - swamp 315
- phlox 600, **601**, 602
 - blue 602
 - downy 603
 - moss **601**, 603
 - slender 603
 - western 603
- picradeniopsis 738
- pigweed, red-root 342
 - Russian 329, **330**
 - winged 335
- pimpernel 581
 - scarlet 581
- pin cherry 455
- pine 56, 58
 - eastern white 60
 - ground- 52, **54**
 - jack 59
 - limber 59
 - lodgepole 59
 - Norway 60
 - running- 52
 - western white 59
 - western yellow 59
 - whitebark 59
- pineappleweed 768
- pinedrops 571
- piresap 571

- pink 347, 352
 - grass- 271
 - purple grass- 271
 - rush- 702
 - steppen 352
 - wood 353
- pinweed 526
 - hairy 526
- pipsissewa 567
- pitcherplant 418
- plantain 663
 - buckhorn 664
 - common 665, **666**
 - common water- 70, **71**
 - hoary 665
 - lesser rattlesnake- 276
 - linear-leaved 665
 - narrow-leaved water- 70
 - Pursh's 667
 - rattlesnake- 275, 276
 - saline 665
 - sand 667
 - seaside 665
 - water- 69
- plum 454
 - American 19, 455
 - Canada 455
 - ground- 19, 469
- poison-ivy 515, **516**
- poison suckleya 339, **341**
- pond-lily, small 362
 - yellow 361, 362
- pondweed 65, 66
 - fennel-leaved 67
 - horned- 67
 - inland 67
 - leafy 67
 - Richardson's 67
 - sago 67
 - sheathed 67
 - various-leaved 67
- poplar 280
 - aspen 281
 - balsam 281
 - white 280
- poppy 384
 - alpine 385
 - common 385
 - corn 385
 - Iceland 385
- potato 631
- povertyweed 686
- prairie-clover 493
 - hairy 495
- purple 495, **496**
 - white 495
- prairie-rocket, small-flowered 409
- prickly-pear 531, 533
 - brittle 533
- primrose 580, **582**, 586
 - dwarf 586
 - erect 586
 - evening- 534, 540
 - gumbo 541, **542**
 - shrubby 543
 - upland 540
 - western yellow 541
 - white 543
 - yellow 541
 - mealy **582**, 586
 - mountain dwarf- 583
- prince's-pine 567
- psoralea
 - lance-leaved 497, **499**
 - silverleaf 495
- puccoon 613
 - hoary 613
 - narrow-leaved 613, **614**
 - red 385
- puffs, sand 344
- purslane 19, 345, 346
 - mud- 525
 - sea- 351
- pussy-toes 713
 - common 714
- pussy willow 294
- pygmyflower 581, **582**
 - western 581
- pyrrocoma, lance-leaved 759

- quack grass 87, **88**
- quillwort 50
 - flowering 66

- rabbitbrush 742
- radish 414
 - horse- 396
 - wild 19, **390**, 414
- ragweed 685
 - bur- 685
 - common 685
 - false 686, **687**
 - great 685
 - perennial 685
- ragwort, brook 780
 - cut-leaved 777, **778**
 - few-flowered 779
 - golden 775

- large-flowered 779
- marsh 775
- northern 779
- rayless 777
- rape, bird- 397
 - broom- 661
 - clustered 661
 - Louisiana 663
 - small 663
- raspberry 19, 458
 - stemless 458
 - wild red 458, **459**
- rattle, Arctic 650
 - Lapland 650
 - purple 652
- rattlesnake-plantain 275, 276
 - lesser 276
- rattlesnakeroot 699
 - purple 700
- redtop 94, **95**
 - alpine 94
 - spike 92
- reed, bur- 64
 - broad-fruited bur- 65
 - many stalked bur- 65
 - narrow-leaved bur- 65
 - northern bur- 65
 - small bur- 65
 - stemless bur- 65
- ribgrass 665
 - western 664
- rice, annual wild- 181
 - wild- 181
- rock brake 37, 38
- rock cress 394
 - Drummond's 395
 - hirsute 395
 - lyre-leaved 395
 - purple 395
 - reflexed 396
- rockcress 416
 - silver 416
- rocket 406, 409
 - dame's- 409
 - garden- 406
 - gray 409
 - purple 409
 - sand- 402
 - small-flowered prairie- 409
- rock-rose 525
- rockstar 422
 - small-flowered 424
- rock tripe 44
 - common 44, **45**
- romanzoffia 607
 - Sitka 607
- rose 436, 455
 - low prairie 457
 - prickly 457
 - rock- 525
 - smooth 457
 - wood's 457
- rose-bay 577
 - Lapland 577
 - white 577
- rose mandarin 266
- rose-root 420
- rowan 460
- rowan tree 461
- rubberweed 764
 - Colorado 764, **765**
 - stemless 764
- rudbeckia 773
- running-pine 52
- rush 246, 247
 - alpine 249
 - arctic 249
 - Baltic 250, **251**
 - beak- 241
 - big-head 254
 - Canada 250
 - chestnut 252
 - common scouring- 48
 - Drummond's 252
 - Dudley's 252
 - dwarf scouring- 50
 - equitant-leaved 252
 - few-flowered 252
 - flattened 252
 - jointed 250
 - knotted 253
 - long-styled 253
 - mud 254
 - Parry's 253
 - Rocky Mountain 253
 - short-tail 250
 - slender 253
 - slender beak- 241
 - slender-stemmed 253
 - smooth scouring- 50
 - spike- 235
 - bald 236
 - beaked 237
 - creeping 237, **238**
 - Englemann's 237
 - few-flowered 237
 - flattened 236
 - one-glumed 237

- slender 237
- Small's 237
- thread 252
- toad 250
- Torrey's 254
- two-glumed 250
- white 249
- white beak- 241
- wood- 254
 - field 255
 - Greenland 255
 - hairy 255
 - mountain 256
 - northern 255
 - small-flowered 255
 - smooth 255
 - spiked 256
- rush-pink 702
- Russian-thistle 339
- rye 168
 - Altai wild 124
 - blue wild 124
 - Canada wild 124, **125**
 - giant wild 124
 - hairy wild 126
 - Macoun's wild 84
 - Russian wild 126
 - smooth wild 124
 - variable-glumed wild 126
 - Virginia wild 126
 - wild 84, 123
- sage 628
 - Herriot's 727
 - lance-leaved 628
 - long-leaved 725
 - Michaux's 727
 - mountain 727
 - pasture 725, **726**
 - prairie 727
 - slender 727
 - wood 628
- sagebrush 728
 - hoary 725
- salsify 703
- samphire 338
 - red 338
- sandalwood 305
- sand-heather 526
- sand-lily 531
- sand-rocket 402
- sand puffs 344
- sandwort 348
 - blunt-leaved 350
 - Dawson 351
 - early 351
 - large-leaved 350
 - rocky-ground 350
 - thyme-leaved 351
- sarsaparilla, bristly 545, **546**
 - wild 544, 545, **547**
- saskatoon 438, **439**
- sawwort 773
 - dwarf 773
 - tall 773
- saxifrage 420, 429
 - alpine brook 433
 - early 435
 - golden 421
 - leather-leaved 422
 - marsh 433, **434**
 - red-flowered 433
 - rhomboid-leaved 433
 - spotted 433
 - three-toothed 435
 - wedge-leaved 431
 - yellow mountain 431
- scabious, field 681
- scorpionweed 606
 - Franklin's 606
 - glandular 607
 - linear-leaved 607
 - Lyall's 607
 - silky 607, **608**
 - silver-leaved 607
 - tansy 607
- scorzonella 697
 - nodding 699
- scouring-rush, common 48
 - dwarf 50
 - smooth 50
- scurvy-grass 401
- sea-blite 339
 - erect 339
 - western 339
- sea-milkwort 583
- sea-purslane 351
- sedge 181, 182, **183, 184**
 - alpine 223
 - Assiniboia 202
 - awl-fruited 229
 - awned 202, **203**
 - Back's 204
 - bald 232
 - beaked 227
 - beautiful 206
 - bent 207
 - blackening 220

black-margined 220
blister 233
blunt 220
bog 217, **218**
bog- 241
bristle-leaved 208
bristle-stalked 215
broad-fruited 208
broom 229
brown 205
browned 201
brownish 205
capitate 205
chaffy 221
chestnut 205
Crawe's 206
Crawford's 206
cyperus-like 223
Dewey's 207
Douglas 208
erect 231
few-flowered 221
few-fruited 220
fox 234
foxtail 201
fragrant 219
Geyer's 210
glacier 210
golden 204
graceful 223
granular 210
green 234
hair-like 205
hairy-fruited 215
hay 229, **230**
head-like 222
heavy 212
Hood's 212
Hooker's 212
Hudson Bay 212
inland 219
Kellogg's 214
lakeshore 214
lens-fruited 215
livid 216
long-beaked 231
long-bracted 202
long-fruited 219
long-haired 206
low 229
low northern 206
Mackenzie 216
mud 216
narrow 201

Nebraska 220
nodding 219
northern bog 212, **213**
Norway 220
open 201
Parry's 221
pasture 222
pleasing 215
porcupine 214
prairie 223
Presl 223
prickly 233
prostrate 206
purple 217
Raynold's 224, **225**
reddish 227
Richardson's 224
rigid 232
rock 227
rocky-ground 228
Ross' 227
round 227
rush-like 228
rye-grass 216
salt 227
sand 212
Sartwell's 228
scant 224
seaside 217
sheathed 233
short 207
short-awned 219
silvery-flowered 201
simple bog- 241
slender 210, **211**
smooth-fruited 214
spiked 224
Sprengel's 229
stalked 221
stiff 204
stone 222
straw-colored 229
sun-loving 221
swollen 214
thick-spike 216
thin-flowered 232
thread-leaved 208, **209**
three-seeded 233
tinged 232
Torrey's 232
turned 224, **226**
two-colored 204
two-seeded 207
two-stamened 207

umbellate 233
 water 201
 weak 231
 white-scaled 234
 wire-grass 208
 woolly 214
 yellow 210
 selaginella, prairie 55
 selfheal 628
 serviceberry 438
 eastern 438
 sheep-laurel 575
 shepherd's-needle 561
 shepherd's-purse 19, **390**, 398, **399**
 shieldwort, pygmy 581
 shinleaf, common 570
 white-veined 570
 shootingstar 18, 19, 583
 mountain **582**, 583
 saline 583
 sibbaldia 454
 Siberian peatree 475
 silverberry 533
 silverplant 310
 silverweed 448
 sisymbrium 415
 skeletonweed 697, **698**, 702
 annual 697
 skullcap 629
 blue 629
 marsh 629
 small 629
 skunkberry 428
 skunkbush 515
 small-snapdragon 642
 smartweed 314
 green 320
 water 318
 snakeroot 561, **562**
 seneca 510
 snapdragon, small- 642
 sneezeweed 709, **710**, 761
 mountain 761
 snowberry 674
 creeping 575
 few-flowered 674
 western 674
 snow-in-summer 352
 snow-on-the-mountain 512
 soapweed 267
 soapwort 355
 solomon's-seal 262
 common 262
 star-flowered 265
 three-leaved 265
 two-leaved **261**, 262
 sorrel, alpine sheep 323
 Bush's yellow wood- 507
 creeping wood- 507
 mountain 310
 sheep 321, **322**
 wood- 505
 yellow wood- 507
 sow-thistle 700
 annual 700
 perennial 700, **701**
 prickly 700
 smooth perennial 700
 spangletop 168, **170**
 spearmint 626
 spearwort, creeping 377
 speedwell 656
 alpine 657
 American 658
 broad-leaved 659
 germander 658
 hairy 658
 marsh 658
 prostrate 657
 smooth 658
 spiked 658
 thyme-leaved 659
 water 658
 spiderflower 386, **388**
 spiderwort 246
 western 246
 spike-moss 55
 spikenard 545
 false 265
 western wild 265
 wild 265
 spike-rush 235
 bald 236
 beaked 237
 creeping 237, **238**
 Engelmann's 237
 few-flowered 237
 flattened 236
 needle 236
 one-glumed 237
 slender 237
 Small's 237
 spiraea, ash- 460
 false 460
 spleenwort 36
 alpine 37
 green 36

springbeauty 345
 alpine 346
 lance-leaved 345
 linear-leaved 346
 small-leaved 346
 spruce 57
 black 58
 western white 58
 white 58
 spurge 511
 cypress 511
 leafy 18, 512, **513**
 petty 512
 prostrate 512
 shining 512
 sun 512
 thyme-leaved 512
 spurred-gentian 592
 spurry 358
 corn 358
 salt-marsh sand 358
 sand 358
 squawroot 561, 661
 squirreltail 171
 staff-tree 517
 starflower 422, 586, 587
 northern **582**, 587
 star-grass 268
 yellow 268
 star-thistle, yellow 741
 starwort, northern water- 514
 vernal water- 514
 water- 514
 stenanthium 266
 stenotus, narrow-leaved 759
 sticky 759
 stickseed 612
 Jessica's 613
 large-flowered 613
 nodding 612
 stinkgrass 127
 stinkweed 19, **390**, 417
 stitchwort 358
 fleshy 360
 long-leaved 360
 long-stalked 360
 northern 360
 St. John's-wort 522
 Canada 522
 large Canada 525
 marsh 525
 western 525
 stonecrop 418, 420, 427
 ditch 427
 lance-leaved 419
 narrow-petaled 420
 stork's-bill 504
 stramonium 631
 strawberry 19, 442
 American wild 442
 smooth wild 442
 suckleya 339
 poison 339, **341**
 sugarscoop 435
 suksdorfia 435
 blue 435
 sumach 514
 smooth 514
 summer-cypress 338
 sundew 418
 oblong-leaved 418
 round-leaved 418
 slender-leaved 418
 sundrops 543
 large 543
 sunflower 761
 beautiful 762
 narrow-leaved 762, **763**
 prairie 762
 rough false 764
 showy 761
 tuberous-rooted 762
 Susan, black-eyed 773
 swamp-candles 584
 sweet-broom 19, 476
 sweet cicely 19, 559
 blunt-fruited 559
 smooth 559, **560**
 western 561
 sweet-clover 487
 white 487, **488**
 yellow 487
 sweet flag 246
 sweet gale 299
 sweet grass 135, **136**
 sweet William 352
 tamarack 57
 western 57
 tansy 788, **789**
 Indian 788
 taraxia 541
 tarweed 766
 tea, Labrador- 575, 576
 New Jersey 518
 trapper's- 576
 tear-thumb, arrow-leaved 318
 teasel 681
 thimbleberry 460

thistle 742
 bull 746
 Canada 743, **744**
 cross- 553
 dwarf 745
 entire-leaved Canada 745
 field 743
 Flodman's 745
 globe- 746, 748
 Hooker's 745
 nodding 739
 plumeless 739
 Russian- 339
 short-stemmed 745
 sow- 700
 annual 700
 perennial 700, **701**
 prickly 700
 smooth perennial 700
 swamp 745
 wavy-leaved 746, **747**
 yellow star- 741
 thorough-wax 550
 thoroughwort 755
 three-awn, red 99
 tickseed 746, 788
 common 746
 tick-trefoil 476
 timothy 154, **157**
 alpine 154, **156**
 toadflax 18
 bastard 305, **307**
 toad-flax 638, 644
 broad-leaved 645
 field 645
 Maroccan 645
 yellow 645
 tomatillo 634
 tomato, wild 18, 635, **637**
 torch, black 656
 torchflower 444
 touch-me-not 508
 spotted 508
 townsendia 788
 low 788
 Parry's 790, **791**
 trailplant 709
 trapper's-tea 576
 tree, rowan 461
 trefoil 480, 482
 bird's-foot 482
 tick- 476
 tripe, common rock 44, **45**
 rock 44

trisetum 179
 awnless 181
 nodding 179
 spike 181
 tall 179
 trumpetweed, Bruner's 755
 tumbleweed 342
 turkey-beard 267
 turtlehead 644
 twayblade 277
 broad-lipped 278
 heart-leaved 278
 northern **272**, 278
 western 278
 twinflower 671, **672**
 twistedstalk 266
 clasping-leaved **261**, 266

 umbrellaplant 309, 310
 cushion 309
 nodding 309
 yellow 310, **311**
 umbrellawort 344
 hairy 344
 heart-leaved 345
 linear-leaved 344

 valerian 681
 northern 681, **682**
 velvetleaf 520
 Venus'-comb 561
 Venus-slipper 271, **272**
 verbena, sand 344
 vervain 619
 blue 619
 bracted 619
 nettle-leaved 619
 vetch 19, 501, 502
 American 502, **503**
 chicken 480
 crown- 475
 field crown- 475
 hairy 502
 milk- 19, 464
 alpine 467
 American 470
 ascending purple 474
 Canadian 469
 cicer 469
 cushion 470
 Drummond's 469
 few-flowered 475
 Indian 467
 loose-flowered 475

low 472
 Missouri 472
 narrow-leaved 472
 prickly 472
 purple 469
 Pursh's 474
 racemose 474
 slender 470
 timber 472
 tufted 474
 two-grooved 467
 narrow-leaved 502
 Oregon 502
 tufted 502
 vetchling 18, 479
 cream-colored 480, **481**
 marsh 480
 tuberous 480
 vine, matrimony 633
 violet 18, 19, 526, **528**
 crowfoot 530
 downy yellow 530
 early blue 527, **528**
 kidney-shaped 530
 long-spurred 530
 marsh 529
 northern bog **528**, 529
 Nuttall's yellow **528**, 529
 round-leaved wood 529
 sweet white 529
 Western Canada **528**, 530
 yellow wood 529
 Virginia creeper 519
 large-toothed 519
 virgin's-bower 369, 371
 purple 369
 western 369

 wakerobin 267
 nodding 267
 western 267
 wallflower, western 407, **408**
 water-arum 246
 water calla 246
 watercrowfoot, large-leaved 377
 small yellow 379
 water-hemlock 551, **552**
 bulb-bearing 551
 water-horehound 625
 northern 625
 western 625
 waterleaf 605
 water-lily 361, 362
 fragrant 362
 small 362

 water-marigold 739
 water-milfoil 543, 544
 pinnate 544
 spiked 544
 water-parsnip 563, **564**
 water-pepper 318
 water-plantain 69
 common 70, **71**
 narrow-leaved 70
 waterpod 605
 water-starwort 514
 northern 514
 vernal 514
 waterweed 73
 Canada 73
 waterwort 525
 weed, spotted Joe-pye 755
 wheat 181
 cow- 645, **646**
 wheatgrass 84
 awned 89, **90**
 awned northern 85
 Baker's 85
 bluebunch 89
 broad-glumed 87
 crested 85, **86**
 northern 87
 slender 89, **91**
 streambank 87
 western 89
 white geranium, wild 505
 white lettuce 699
 glaucous 700
 whitetop 750
 whitlow-grass 403
 alpine 404
 creeping 406
 few-seeded 406
 golden 404
 hairy 405
 thick-leaved 404
 yellow **390**, 405
 whitlowwort 354
 low 354
 whortleberry 579
 bog 579
 red 579
 wild barley 137, **138**
 wild bergamot 626, **627**
 western 628
 wild black currant 428
 wild buckwheat 314, 317
 wild chamomile 768
 wild chives 258
 wild columbine 368

wild cucumber 677
 wild flax, Lewis 507
 meadow 508
 wild geranium, Carolina 505
 wild ginger 308
 wild licorice 19, 476, **477**
 wild lily of-the-valley 262
 wild mint 626
 wild morning-glory **598**, 599
 wild mustard **390**, 397
 wild oat 100, **101**
 wild pansy 529
 wild parsley, Sandberg's 557
 western 557
 wild parsnip 561
 mountain 557
 wild peavine 480
 wild radish 19, **390**, 414
 wild red raspberry 458, **459**
 wild-rice 181
 annual 181
 wild rye 84, 123
 Altai 124
 blue 124
 Canada 124, **125**
 giant 124
 hairy 126
 Macoun's 84
 Russian 126
 smooth 124
 variable-glumed 126
 Virginia 126
 wild sarsaparilla 544, 545, **547**
 wild spikenard 265
 western 265
 wild strawberry, American 442
 smooth 442
 wild tomato 18, 635, **637**
 wild white geranium 505
 willow 279, 281, **283**
 Alaska 290
 arctic 291
 autumn 298
 balsam 298
 Barclay's 291
 Barratt's 292
 basket 297
 bay-leaved 297
 beaked 292
 bog 296
 brittle 294
 changeable 294
 diamond 294
 flat-leaved 297
 gray 295
 hoary 292
 lime 292
 Mackenzie 296
 mountain 296
 peach-leaved 290
 pussy 294
 rock 299
 sandbar 295
 satin 297
 shining 295
 short-capsuled 292, **293**
 shrubby 291
 Sitka 298
 smooth 294
 snow 298
 trailing 291
 velvet-fruited 296
 white 290
 wolf- 533
 yellow 295
 willowherb 537
 alpine 537
 annual 539
 low 540
 marsh 539
 northern 539
 winterfat **337**, 338
 wintergreen 567, **568**, 569, 575
 alpine 575
 common 570
 greenish-flowered 570
 large 569
 lesser 570
 one-flowered 567, **568**
 one-sided **568**, 570
 pink **568**, 569
 wolf-willow 533
 wonderberry 635
 wood nettle 304
 woodruff 667
 field 667
 wood-rush 254
 field 255
 Greenland 255
 hairy 255
 mountain 256
 northern 255
 small-flowered 255
 smooth 255
 spiked 256
 wood-sorrel 505
 Bush's yellow 507
 creeping 507
 yellow 507
 wool-grass 243

woollen-breeches 605
wormwood 722
 biennial 723
 common 728
 linear-leaved 725
 plains 725
 Roman 727
yarrow 708

 many-flowered 708
 woolly 709
yellowbell 262
yellowrattle 654
yew 56
 western 56
youngia 692
yucca 267

Index to scientific names

(Page numbers of illustrations are in boldface; synonyms are in italic type.)

- Abies* 56, 57
 balsamea 57
 var. *fallax* 57
 lasiocarpa 57
Abronia 344
 micrantha 344
Abutilon theophrasti 520
Acer 515
 glabrum var. *douglasii* 515, 517
 negundo var. *interius* 515, 517
 saccharinum 515, 517
 spicatum 515, 517
Aceraceae 25, 28, 515
Acerates viridiflora 597
Achillea 708
 borealis 709
 lanulosa 709
 millefolium 708
 var. *nigrescens* 709
 var. *occidentalis* 708
 f. *purpurea* 708
 multiflora 709
 ptarmica 708, **710**
 sibirica 708
Aconitum 362, 363
 delphinifolium 363
Acorus 246
 calamus 246
Actaea 363
 rubra 363
 f. *neglecta* 363
Actinea richardsonii 764
Acutae 188, 198
Adenocaulon 704, 709
 bicolor 709
Adiantum 35, 36
 pedatum 36
Adlumia 385
 fungosum 385
Adoxa 677
 moschatellina 677
Adoxaceae 30, 677
Aegopodium 548, 549
 podagraria 549
Agalinis 639
 aspera 639
 purpurea 639
 var. *parviflora* 640
 tenuifolia 639
 var. *parviflora* 640
Agastache 620, 621
 anethiodora 621
 foeniculum 621, **622**
Agoseris 689, 690
 aurantiaca 690
 cuspidata 699
 glauca 690
 var. *dasycephala* 691
 grandiflora 690, 691
Agrimonia 437
 striata 437
Agrohordeum 81, 84
 macounii 84
Agropyron 80, 84, 804
 albicans 84, 85
 var. *griffithsii* 85
 bakeri 85
 crisatiforme 87
 crisatum 84, 85, **86**
 dasystachyum 84, 87
 desertorum 85
 latiglume 85, 87
 pectiniforme 87
 repens 84, 87, **88**
 riparium 84, 87
 scribneri 85, 87
 smithii 84, 89
 var. *molle* 89
 spicatum 85, 89
 var. *inermis* 89
 subsecundum 85, 89, **90**
 trachycaulum 85, 89, **91**
 trachycaulum 89
 var. *unilaterale* 89
 triticeum 84, 89
Agrostemma 347, 348
 githago 348, **349**
Agrostideae 77, 78, 81
Agrostis 82, 92
 alba 94
 borealis 92
 exarata 92
 scabra 92, **93**
 stolonifera 92, 94
 f. *palustris* 94
 var. *genuina* 94, **95**

- var. *major* 92, 94
thurberiana 92, 94
variabilis 92, 94
Albae 188, 190, 196
Alisma 69, 70
 geyeri 70
 gramineum 70
 plantago-aquatica 70, **71**
 Alismaceae 22, 69
Allium 256, 258
 cernuum 258
 schoenoprasum var. *sibiricum* 258
 stellatum 258
 textile 258, **259**
Allocarya californica 618
Alnus 299, 300
 crispa 300
 incana 300
 rugosa var. *americana* 300
 tenuifolia 300
 viridis var. *sinuata* 300
Alopecurus 82, 94
 aequalis 94
 alpinus 94
 geniculatus 94, 96
 pratensis 94, 96
Althea rosea 520
Alyssum 391, 393
 alyssoides **390**, 393
 Amaranthaceae 24, 342
Amaranthus 342
 albus 342
 blitoides 342
 graecizans 342, **343**
 retroflexus 342
 Amaryllidaceae 22, 268
Ambrosia 685
 artemisiifolia var. *elatior* 685
 coronopifolia 685
 elatior 685
 psilostachya var. *coronopifolia* 685
 trifida 685
Amelanchier 436, 437
 alnifolia 438, **439**
 florida 438
 sanguinea 438
Amorpha 28, 462, 463
 canescens 463
 fruticosa 463
 nana 463, 464
Amphicarpa 462, 464
 bracteata 464
Amsinckia 610
 menziesii 610
 Anacardiaceae 28, 514
Anagallis 581
 arvensis 581
Anaphalis 704, 709
 margaritacea 709
 var. *subalpina* 709
Androcera rostrata 635
Andromeda 571, 572
 polifolia 572
Andropogon 84, 96
 gerardi 96, **97**
 hallii 96
 scoparius 96, **98**
 Andropogoneae 77, 79, 84
Androsace 580, 581
 chamaejasme 581
 occidentalis 581
 septentrionalis 581, **582**
 var. *diffusa* 581
 var. *puberulenta* 581
Anemone 24, 363, 364
 canadensis 364, 365
 cylindrica 364, 365
 drummondii 364, 365
 multifida 365
 var. *richardsiana* 364, 365
 narcissiflora 364, 365
 nemorosa var. *bifolia* 364, 365
 occidentalis 364, 365
 parviflora 364, 366
 patens var. *wolfgangiana* 364, 366,
 367
 quinquefolia var. *interior* 365
 richardsonii 364, 366
 riparia 366
 virginiana 364, 366
Anethum 549, 550
 graveolens 550
Angelica 548, 550
 arguta 550
 dawsonii 550
 genuflexa 550
 lyallii 550
Angiospermae 20, 21, 62, 790
Anogra nuttallii 543
Anoplanthus fasciculatus 661
Antennaria 704, 709
 aizoides 716
 alpina 712, 713
 var. *canescens* 713
 anaphaloides 711, 713
 angustata 712, 713
 aprica 712, 713
 canadensis 714

- corymbosa 712, 713
 dimorpha 711, 713
 glabrata 712, 713
 howellii 712, 714, **715**
 var. athabascensis 714
 var. campestris 714
 var. howellii 714
 lanata 711, 714
 luzuloides 711, 714
 monocephala 712, 714
 neodioica 712, 714, 716
 var. randii 714
 obovata 714
 parlinii 712, 716
 parvifolia 712, 716
 plantaginifolia 712, 716
 pulcherrima 711, 716
 racemosa 711, 716
 rosea 712, 716
 russellii 712, 716
 umbrinella 711, 716
 Anthemis 708, 717
 cotula 717
 tinctoria 717
Anthopogon crinitus 591
Anticlea elegans 268
 Apocynaceae 31, 592
 Apocynum 592
 androsaemifolium 592
 var. incanum 593, **594**
 cannabinum 592
 var. hypericifolium 593
 medium 592, 593
 sibiricum 593
 Aquilegia 363, 366
 brevistyla 366, 368
 canadensis 368
 flavescens 366, 368
 formosa 368
 jonesii 366, 368
 Arabis 392–394, 409
 arenicola 394
 var. arenicola 394
 var. pubescens 394
 divaricarpa 394, 395
 var. dacotica 395
 drummondii 394, 395
 glabra **390**, 394, 395
 hirsuta 394, 395
 var. glabrata 395
 lemmonii 394, 395
 var. drepanoloba 395
 lyallii 394, 395
 lyrata var. kamchatica 394, 395
 nuttallii 394, 396
 retrofracta 394, 396
 var. collinsii 396
 var. multicaulis 396
 Araceae 22, 246
 Aralia 544
 hispidula 544, 545, **546**
 nudicaulis 544, 545, **547**
 racemosa 544, 545
 Araliaceae 27, 544
 Arceuthobium 308
 americanum 308
 pusillum 308
 Arctagrostis 82, 99
 latifolia 99
 Arctium 704, 717
 lappa 717
 minus 717
 tomentosum 717
 Arctophila 79, 99
 fulva 99
 Arctostaphylos 572
 alpina 572, 573
 var. rubra 573
 uva-ursi 573
 Arenaria 348
 capillaris var. americana 348, 350
 congesta var. lithophila 348, 350
 humifusa 348, 350
 laricifolia var. occulta 348, 350
 lithophila 350
 lateriflora 348, 350
 macrophylla 348, 350
 nuttallii 348, 350
 obtusiloba 350
 peploides var. diffusa 348, 351
 rossii var. columbiana 350, 351
 serpyllifolia 348, 351
 stricta ssp. dawsonensis 350, 351
 verna 350, 351
 Arenariae 186
 Arethusa 270, 271
 bulbosa 271
 Aristida 82, 99
 longiseta 99
 Aristolochiaceae 24, 308
 Armoracia 391, 396
 rusticana 396
 Arnica 705, 706, 718
 alpina var. ungavensis 719
 var. vestita 719
 chamissonis 718, 719
 cordifolia 718, 719, **720**
 diversifolia 718, 719

fulgens 719, **720**
 latifolia 718, 721
 lonchophylla 718, 721
 longifolia 721
 louiseana 719, 721
 mollis 718, 721
 var. aspera 721
 parryi 718, 721
 rydbergii 719, 722
 sororia 719, 722
Artemisia 661, 704, 705, 722
 abrotanum 723
 absinthium 722, 723, **724**
 biennis 722, 723
 borealis 725
 bourgeauiana 725
 campestris 722, 725
 var. douglasiana 725
 var. scouleriana 725
 var. wormskjoldii 725
 camporum 725
 cana 722, 725
 canadensis 725
 caudata 725
 dracunculus 722, 725
 frigida 722, 725, **726**
 herriotii 727
 longifolia 723, 725
 ludoviciana 723
 var. gnaphalodes 727
 var. ludoviciana 727
 michauxiana 723, 727
 norvegica var. saxatilis 723, 727
 pontica 723, 727
 tilesii ssp. unalascensis 723, 727
 tridentata 722, 728
 vulgaris 723, 728
Asarum 308
 canadense 308
Asclepiadaceae 31, 593
Asclepias 593
 incarnata 595
 lanuginosa 595
 ovalifolia 595
 speciosa 595, **596**, 597
 syriaca 595, 597
 verticillata 595, 597
 viridiflora 595, 597
 var. linearis 597
Asparagus 256, 258
 officinalis 258
Asperugo 609, 610
 procumbens 610
Asperula 667
 arvensis 667

Asplenium 36
 viride 36
Aster 705, 707, 708, 728
 adscendens 729, 730
 adsurgens 734
 alpinus 729, 730
 angustus 731
 brachyactis 728, 731
 campestris 728, 731
 canescens 728, 731
 ciliolatus 729, 731, **732**
 commutatus 734
 conspicuus 728, 731, **733**
 eatoni 730, 731
 engelmannii 730, 734
 ericoides 730, 734
 falcatus 730, 734
 foliaceus var. apricus 737
 frondeus 737
 hesperius 730, 734
 junciformis 730, 734, 804
 laevis 729, 734, **735**
 lateriflorus 730, 734
 lindleyanus 731
 maccallae 729, 736
 macrophyllus 728, 736
 modestus 729, 736
 multiflorus 734
 novae-angliae 728, 736
 occidentalis 729, 736
 osterhoutii 734
 pansus 734
 pauciflorus 728, 736
 ptarmicoides 785
 var. lutescens 785
 puniceus 730, 737
 sericeus 729, 737
 sibiricus 729, 737
 simplex 730, 737
 subspicatus 729, 737
 var. apricus 737
 umbellatus 729, 737
 var. pubens 737
Astragalus 463, 464, 489
 aboriginorum 467
 aboriginum 466, 467
 var. major 467
 adsurgens 475
 alpinus 465–467
 var. brunetianus 467
 americanus 470
 bisulcatus 464, 467, **468**
 bourgovii 467, 469
 caespitosus 474
 canadensis 466, 469

- caryocarpus* 469
cicer 466, 469
crassicarpus 465, 469
danicus 465, 466, 469
 var. *dasyglottis* 469
 f. *virgultulus* 469
drummondii 465, 469
eucosmus 467, 470
 var. *eucosmus* 470
 f. *leucocarpus* 470
flexuosus 465, 470
frigidus 466, 470
 var. *americanus* 470, **471**
 var. *frigidus* 470
gilviflorus 462, 465, 470
goniatus 469
hypoglottis 469
iochrous 466, 472, **473**
kentrophyta 464, 472
lotiflorus 465, 472
miser 464, 466, 472
 var. *serotinus* 472
missouriensis 465, 472
pectinatus 464, 465, **468**, 472
purshii 465, 474
racemosus 466, 474
robbinsii 467, 474
serotinus 472
spatulatus 462, 465, 474
striatus 466, 474
tenellus 466, 475
triphyllus 470
vexilliflexus 465, 475
yukonis 466, 475
Atelophragma aboriginum 467
 alpinum 467
Atenia gairdneri 561
Athyrium 36
 alpestre 36, 37
 filix-femina 37
 var. *cyclosorum* 37
 var. *filix-femina* 37
Atratae 191, 198
Atriplex 23, 326
 argentea 327
 dioica 329
 hortensis 327, **328**
 var. *atrosanguinea* 327
 nuttallii 326, 327
 patula 327, 329
 var. *hastata* 329
 powellii 327, 329
Avena 81, 99
 fatua 99, 100, **101**
 sativa 99, 100
Aveneae 77, 78, 81
Avicularia 313
Axyris 326, 329
 amaranthoides 329, **330**

Bahia 706, 738
 oppositifolia 738
Balsaminaceae 27, 508
Balsamorhiza 707, 738
 sagittata 738
Barbarea 392, 396
 orthoceras 396
 vulgaris 396
Bartsia 639, 640
 alpina 640
Bassia 326, 329
 hyssopifolia 329
Batidophaca lotiflora 472
Beckmannia 83, 100
 syzigachne 100, **102**
Berberidaceae 26, 29, 382
Berberis 382, 384
 aquifolium 384
 f. *repens* 384
Berteroa 391, 396
 incana 396
Besseyia 638, 640
 cinerea 640
 wyomingensis 640
Betula 299, 300
 fontinalis 301
 glandulifera 300
 glandulosa 300
 nana var. *sibirica* 300
 occidentalis 300, 301
 papyrifera 300, 301
 var. *cordifolia* 301
 var. *neoalaskana* 301
 var. *papyrifera* 301
 pumila var. *glandulifera* 300
Betulaceae 25, 299
Bicolores 188, 196
Bidens **684**, 705, 707, 738, 804
 beckii 738
 cernua 738, 739, 804
 frondosa 738, 739
 var. *puberula* 739
 glaucescens 739
 tripartita 738, 739
Bilderdykia 313, 314
Bistorta 313, 314
Blysmus rufus 244
Boisduvalia 536
 glabella 536

- Bolelia laeta* 680
 Boltonia 708, 739
 asteroides 739
 Boraginaceae 31, 609
 Borago 609, 610
 officinalis 610
 Botrychium 32
 boreale 32
 var. obtusilobum 32
 lanceolatum 32
 lunaria 32, 33
 matricariifolium 32, 33
 multifidum 32, 33
 simplex 32, 33
 virginianum 32, 33, **34**
 Bouteloua 83, 100
 curtipendula 100
 gracilis 100, **103**
Brachyactis angusta 731
 Bracteosae 187, 191
 Brassica 392, 396
 campestris 397
 hirta 397
 juncea 397
 kaber **390**, 397
 napus 397
Braxilia minor 570
 Braya 393, 397
 humilis 397
 Brickellia 705, 739
 grandiflora 739
 Bromus 80, 104
 ciliatus 104, 105
 inermis 104, 105
 japonicus 104, 105
 kalmii 104, 105
 marginatus 104, 105
 porteri 104, 106
 pumpellianus 104, 106
 purgans 104, 106
 squamosus 104, 106
 tectorum 104, 106, **107**, 804
 vulgaris 104, 106
 Buchloe 83, 108
 dactyloides 108
 Bupleurum 548, 550
 americanum 550
 Cactaceae 25, 531
 Calamagrostis 82, 108
 canadensis 109
 var. robusta 109
 var. scabra 109
 deschampsiioides 108, 109
 inexpansa 109
 lapponica 109, 110
 var. nearctica 110
 montanensis 108, 110, **111**
 neglecta 109, 110
 purpurascens 108, 110
 rubescens 108, 110
 Calamovilfa 82, 112
 longifolia 112, **113**
 Calla 246
 palustris 246
 Callistachys 185, 191
 Callitrichaceae 23, 514
 Callitriche 514
 autumnalis 514
 hermaphroditica 514
 palustris 514
 Calochortus 256, 258
 apiculatus 258
 Calopogon 271
 pulchellus 271
 Caltha 363, 368
 leptosepala 368
 natans 368, 369
 palustris 368, 369, **370**
 Calypso 270, 271
 bulbosa 271, **272**
 Camassia 256, 260
 quamash 260
 Camelina 391, 398
 dentata 398
 microcarpa 398
 sativa **390**, 398
 Campanula 677
 aparinoides 678
 glomerata 677, 678
 lasiocarpa 677, 678
 rapunculoides 677, 678
 rotundifolia 678, **679**
 uliginosa 678
 uniflora 677, 678
 Campanulaceae 31, 677
 Cannabinaceae 23, 304
 Capillares 190
 Capitatae 185
 Capparidaceae 27, 28, 386
 Caprifoliaceae 29, 31, 670
 Capsella 389, 398
 bursa-pastoris **390**, 398, **399**
 Caragana 462, 475
 arborescens 475
 Cardamine 393, 398
 bellidifolia 398, 400
 bulbosa 398, 400

parviflora 400
 pennsylvanica 400
 pratensis 400
 var. angustifolia 400
 umbellata 400
 Cardaria 389, 400
 draba 401
 var. repens 401
 pubescens var. elongata 401
 Carduus 705, 739
 nutans 739
 Carex 182, 185
 adusta 194, 201
 aenea 202
 albo-nigra 204
 alopecoidea 192, 201
 angustior 219
 aperta 199, 201
 aquatilis 199, 201
 var. altior 201
 var. stans 201
 arcta 193, 201
 argyrantha 194
 assiniboinensis 189, 197, 202
 atherodes 200, 202, **203**
 athrostachya 193, 202
 atrata 197, 202
 atrofusca 197, 204
 atrosquama 204
 aurea 196, 204
 backii 186, 189, 204
 bebbii 208
 bicknellii 231
 bicolor 196, 204
 bigelowii 199, 204
 brevior 231
 brunnescens 193, 205
 buxbaumii 198, 205
 canescens 207
 capillaris 190, 205
 var. capillaris 205
 var. elongata 205
 capitata 185, 205
 castanea 190, 197, 205
 chordorrhiza 186, 206
 concinna 196, 206
 concinoides 196, 206
 crawei 197, 206
 crawfordii 193, 206
 crinita 199, 206
 cristatella 233
 cumulata 231
 curta 193, 207
 deflexa 195, 207

deweyana 187, 207
 diandra 192, 207
 disperma 187, 192, 207
 douglasii 191, 208
 eburnea 190, 196, 208
 eleusinoides 198, 208
 epapillosa 204
 festivella 217
 festucacea 194, 208
 filifolia 186, 208, **209**
 flava 197, 210
 foenea 229
 garberi var. bifaria 204
 geyeri 186, 210
 glacialis 190, 196, 210
 gracillima 190, 210, **211**
 granularis 197, 210
 gravida 192, 212
 gynocrates 185, 212, **213**
 haydeniana 217
 heleonastes 193, 212
 heliophila 222
 hoodii 192, 212
 hookerana 191, 212
 houghtonii 197, 212
 hystricina 200, 214
 incondita 217
 interior 219
 intumescens 189, 214
 kelloggii 199, 214
 lacustris 200, 214
 laeviconica 200, 214
 lanuginosa 197, 215
 lasiocarpa 197, 215
 laxiflora 190
 var. varians 215
 lenticularis 199, 215
 leptalea 186, 215
 limnophila 217
 limosa 198, 216
 livida 196, 216
 loliacea 193, 216
 mackenziei 193, 216
 macloviana 194, 216, 217
 magellanica 198, 217, **218**
 maritima 186, 217
 merritt-fernaldii 231
 mertensii 198, 217
 michauxiana 188, 219
 microglochin 199, 219
 microptera 217
 misandra 197, 219
 molesta 231
 muricata 188, 219

- nardina 186, 219
 var. hepburnii 219
 nebraskensis 199, 220
 nigricans 191, 220
 nigromarginata 195
 var. elliptica 220
normalis 208
 norvegica 198, 220
 obtusata 186, 195, 220
 oligosperma 191, 200, 220
pachystachya 217
 paleacea 199, 221
 parryana 198, 221
 pauciflora 199, 221
paupercula 217
 pedunculata 195, 221
 pennsylvanica 195, 221
 var. digyna 222
 var. distans 222
 var. pennsylvanica 222
 petasata 194, 222
 petricosa 197, 222
 var. franklinii 222
 phaeocephala 194, 222
platylepis 222
 podocarpa 198, 223
 praegracilis 191, 223
 prairea 192, 223
praticola 222
 preslii 194, 223
projecta 233
 pseudo-cyperus 200, 223
 pyrenaica 191, 224
 rariflora 198, 224
raymondii 202, 204
 raynoldsii 198, 224, 225
 retrorsa 200, 224, 226
 richardsonii 196, 224
 rossii 195, 227
 rostrata 200, 227
 rotundata 200, 227
 rufina 188, 196, 227
 rupestris 196, 227
 salina 199, 227
 var. kattegatensis 228
 var. salina 228
 var. subspathacea 228
 sartwellii 187, 228
 saxatilis 188, 200, 228
 var. major 228
 var. saxatilis 228
 scirpoidea 185, 228
 scoparia 194, 229
 siccata 186, 229, 230
 sprengelii 189, 229
 stenophylla 191
 ssp. eleocharis 229
 stipata 192, 229
 straminea 194, 229
 stricta 199, 231
 supina 191, 195
 ssp. spaniocarpa 231
 sychnocephala 193, 231
tenera 208
 tenuiflora 192, 232
 tetanica 196, 232
 var. meadii 232
 var. woodii 232
 tinctoria 194, 232
 tonsa 195, 232
 torreyi 189, 232
 tribuloides 194, 233
 trisperma 192, 233
 umbellata 195, 233
 vaginata 196, 233
 vesicaria 200, 233
 viridula 197, 234
 vulpinoidea 187, 234
 xerantica 194, 234
 Carum 549, 551
 carvi 551
 Caryophyllaceae 24, 26, 347
 Cassiope 572, 573
 mertensiana 573
 tetragona 573
 var. saximontana 573
 var. tetragona 573
 Castilleja 638, 640
 acuminata 642
 coccinea 640, 641
 cusickii 641
 hispida 641
 lutea 641
 lutescens 641
 miniata 641, 643
 occidentalis 641
 pallida 641
 var. septentrionalis 642
 raupii 640
 sessiliflora 640
 sulphurea 642
 Catabrosa 79, 112
 aquatica 112
Cathartolinum rigidum 508
 sulcatum 508
 Caulophyllum 26, 382, 384
 thalictroides 384

- Ceanothus 29, 518
 ovatus 518
 velutinus 518
 Celastraceae 28, 517
 Celastrus 517
 scandens 517
 Celtis 303
 occidentalis 303
 Centaurea 704, 705, 740
 diffusa 740
 maculosa 740
 picris 741
 repens 740
 solstitialis 740, 741
 Centunculus 580, 581
 minimus 581
 Cerastium 347, 351
 alpinum 352
 arvense 351, 352
 nutans 351, 352
 tomentosum 351, 352
 vulgatum 351, 352
 Ceratophyllaceae 23, 361
 Ceratophyllum 361
 demersum 361
 Chaenorrhinum 638, 642
 minus 642
 Chamaedaphne 571, 573
 calyculata 573, **574**
Chamaenerion angustifolium 537
 latifolium 537
 spicatum 537
Chamaepericlymenum canadense 565
 Chamaerhodos 437, 438
 erecta var. parviflora 438
 nuttallii 438
Chamaesaracha grandiflora 634
Chamomilla suaveolens 768
 Cheilanthes 36, 37
 feei 37
Cheirinia aspera 407
 cheiranthoides 407
 Chelone 639, 644
 glabra var. linifolia 644
 Chenopodiaceae 23–25, 324, 804
 Chenopodium 326, 329
 album 331, **332**
 capitatum 329, 331
 dacoticum 331
 fremontii 331, 333, **334**
 glaucum var. pulchrum 331, 333
 hybridum var. gigantospermum 331,
 333
 lanceolatum 331
 leptophyllum 331, 333
 polyspermum 331, 333
 rubrum 331, 333
 salinum 333
 subglabrum 331, 333
 Chimaphila 29, 567
 umbellata var. occidentalis 567
Chiogenes hispidula 575
 Chlorideae 77, 78, 83
Chondrophylla fremontii 590
 Chordorrhizae 186
 Chrysanthemum 704, 708, 741
 arcticum 741
 var. polaris 741
 balsamita 741
 f. tanacetoides 741
 leucanthemum 741
 var. pinnatifidum 741
Chrysobotrya aurea 428
 Chrysopsis 706, 741
 hispidula 742
 villosa 742
 Chrysosplenium 24, 420, 421
 iowense 421
 tetrandrum 421
 Chrysothamnus 705, 742
 frigidus 742
 nauseosus 742
 Cichorium 689, 691
 endivia 691
 intybus 691
 Cicutula 548, 551, 555
 bulbifera 551
 douglasii 553
 mackenzieana 551
 maculata var. angustifolia 551, **552**
 var. maculata 553
 occidentalis 553
 Cinna 82, 112
 latifolia 112
 Circaea 536
 alpina 536
 quadrisulcata var. canadensis 536
 Cirsium **684**, 704, 740, 742
 altissimum var. discolor 743
 arvense 742, 743, **744**
 var. integrifolium 745
 drummondii 743, 745
 flodmanii 743, 745
 foliosum 743, 745
 hookerianum 743, 745
 lanceolatum 746
 muticum 743, 745

- undulatum 743, 745, 746, **747**
 - var. megacephalum 746
 - vulgare 742, 746
- Cistaceae 27, 28, 525
- Claytonia 345
 - caroliniana var. lanceolata 345
 - fontana 345, 346
 - linearis 345, 346
 - megarrhiza 345, 346
 - parvifolia 345, 346
- Clematis 24, 25, 28, 362, 369
 - columbiana 369
 - ligusticifolia 369
 - tangutica 369
 - verticellaris var. columbiana 369
 - virginiana 369, 371
- Cleome 386
 - serrulata 386, **388**
- Clintonia 257, 260
 - borealis 260
 - uniflora 260
- Cnemidophacos pectinatus* 474
- Cochlearia 389, 401
 - officinalis 401
- Cogswellia macrocarpa* 557
 - orientalis 557
- Collinsia 639, 644
 - parviflora 644
- Collomia 600
 - linearis 600, **601**
- Comandra 305
 - livida 305
 - pallida 308
 - richardsiana 308
 - umbellata 305, **307**
 - var. angustifolia 305, 308
 - var. pallida 308
 - var. umbellata 305, 308
- Comarum palustre* 451
- Commelinaceae 22, 246
- Compositae 29, 30, 683
- Conimitella 420, 421
 - williamsii 421
- Conium 549, 553
 - maculatum 553
- Conopholis 661
 - americana 661
- Conringia 391–393, 401
 - orientalis **390**, 401
- Convolvulaceae 30, 31, 597
- Convolvulus 597
 - americanus 599
 - arvensis 597, **598**
 - interior 599
 - sepium 597, 599
 - var. americanus **598**, 599
 - var. pubescens 599
- Coptis 363, 371
 - trifolia 371
- Corallorhiza 270, 271
 - maculata 271, 273
 - striata 271, 273
 - trifida 271, 273
- Coreopsis 707, 746
 - atkinsoniana 746
 - tinctoria 746
- Corispermum 326, 335
 - hyssopifolium 335
 - marginale 335
 - orientale var. emarginatum 335, **336**
 - villosum 335
- Cornaceae 26, 28, 563
- Cornus 563
 - alba 565
 - var. baileyi 565
 - var. interior 565
 - alternifolia 565
 - canadensis 565, **566**
 - racemosa 565, 567
 - rugosa 565, 567
 - stolonifera 565
- Coronilla 463, 475
 - varia 475
- Corydalis 385, 386
 - aurea 386
 - sempervirens 386, **387**
- Corylus 25, 299, 301
 - americana 301, **302**
 - cornuta 301, **302**
- Cotoneaster 436, 438
 - acutifolia 438, 440
 - melanocarpa 440
- Crassulaceae 28, 418
- Crataegus 436, 440
 - chrysoarpa 440
 - columbiana 440
 - douglasii 440
 - rotundifolia 440, **441**
 - succulenta 440
- Crepis 689–691
 - capillaris 691, 692
 - elegans 692
 - intermedia 692
 - nana 691, 692
 - occidentalis 692
 - runcinata 692
 - var. glauca 692, **693**
 - var. hispida 692

- tectorum 691, 694
 Cruciferae 28, 389, **390**
 Cryptantha 609, 610
 bradburiana 611
 fendleri 610, 611
 minima 610, 611
 nubigena 610, 611
 Cryptocarpaceae 188, 199
 Cryptogramma 35, 37, 38
 crispa 37
 var. *acrostichoides* 38, **39**
 stelleri 37, 38
 Cryptotaenia 548, 553
 canadensis 553
 Cucurbitaceae 30, 677
 Cupressaceae 21, 60
 Cuscuta 597, 599
 curta 600
 gronovii 600
 pentagona 600
 planiflora 600
Cyclachaena xanthifolia 686
 Cycloloma 326, 335
 atriplicifolium 335
 Cymopterus 549, 553
 acaulis 553, **554**
 Cynoglossum 609, 611
 boreale 611
 officinale 611
 Cyperaceae 22, 74, 181
 Cyperus 182, 234
 houghtonii 235
 inflexus 234, 235
 schweinitzii 234, 235
 strigosus 234, 235
 Cypridium 270, 273
 acaule 273, **274**
 arietinum 275
 calceolus 275
 var. *parviflorum* **272**, 275
 var. *pubescens* 275
 candidum 275
 montanum 275
 passerinum 275
 reginae 275
 Cystopteris 36, 38
 fragilis 38, **40**
 montana 38
Cytherea bulbosa 271
 Dactylis 79, 112
 glomerata 112, **114**
 Danthonia 81, 112
 californica 112
 var. *americana* 112
 intermedia 112, **116**
 parryi 112
 spicata 112
 unispicata 112
Dasiphora fruticosa 449
Dasystephana affinis 590
 andrewsii 590
 flavida 590
 puberula 591
 Datura 631
 stramonium 631
 Daucus 549, 553
 carota 553
 Delphinium 363, 371
 bicolor 371, **372**
 brownii 371
 canmorense 371
 glaucum 371, **373**
 Deschampsia 81, 117
 atropurpurea 117
 caespitosa 117, **118**
 danthonioides 117
 elongata 117, 119
 Descurainia 391, 392, 401
 pinnata var. *brachycarpa* 402
 richardsonii 402
 sophia 402
 sophioides 402
 Desmodium 463, 476
 canadense 476
 Deweyanae 187
 Dianthus 347, 352
 barbatus 352
 deltoides 352
 sylvestris 352, 353
 Diapensia 580
 lapponica 580
 Diapensiaceae 29, 580
 Dicotyledoneae 12, 20, 23, 279, 790
 Diervilla 670, 671
 lonicera 671
 Digitaria 84, 119
 ischaemum 119
 sanguinalis 119
 Digitatae 189, 195
Diholcos bisulcatus 467
 Dioicae 185
 Diplotaxis 392, 402
 muralis 402
 Dipsacaceae 30, 681
 Disporum 257, 260
 hookeri var. *oreganum* 260
 trachycarpum 260, **261**
Distegia involucrata 673

- Distichlis 79, 119
 stricta 119, **120**
 Divisae 187, 191
 Dodecatheon 580, 583
 conjugens 583
 var. beamishii **582**, 583
 cylindrocarpum 583
 pulchellum 583
Doellingeria pubens 738
 umbellatus 737
 Douglasia 580, 583
 montana 583
 Downingia 680
 laeta 680
 Draba 391
 alpina 404
 aurea 403, 404
 var. leiocarpa 404
 cinerea 403, 404
 crassifolia 404
 fladnizensis 404, 405
 hirta 403, 405
 incana 403, 405
 incerta 404, 405
 lanceolata 403, 405
 lutea 405
 mccallae 403, 405
 nemorosa 403, 405
 var. leiocarpa **390**, 405
 nivalis 404, 406
 var. elongata 406
 oligosperma 404, 406
 paysonii var. *treleasii* 406
 praealta 403, 406
 reptans var. micrantha 403, 406
 stenoloba 403, 406
 var. nana 406
 stenopetala 404, 406
 Dracocephalum 620, 621
 nuttallii 628
 parviflorum 621
 thymiflorum 621
 Drosera 418
 anglica 418
 linearis 418
 rotundifolia 418
 Droseraceae 26, 418
 Dryas 436, 442
 drummondii 442
 integrifolia 442
 octopetala var. hookeriana 442
Drymocallis agrimonioides 448
 Dryopteris 36, 38
 austriaca 41
 var. spinulosa 41
 cristata 41
 disjuncta 38, 41
 filix-mas 41
 fragrans 41, 42
 phegopteris 38, 42
 spinulosa 41
 thelypteris 38, 42
 Dupontia 79, 121
 fisheri 121
 Echinacea 708, 746
 angustifolia 746
 Echinochloa 83, 121
 crusgalli 121, **122**
 f. breviseta 121
 f. longiseta 121
 var. depressa 121
 var. erecta 121
 var. frumentacea 121
 pungens 121
 Echinocystis 677
 lobata 677
 Echinops 705, 746
 exaltatus 746, 748
 sphaerocephalus 746, 748
 Echium 609, 611
 plantagineum 611
 vulgare 611
 Elaeagnaceae 25, 533
 Elaeagnus 533
 angustifolia 533
 commutata 533
 Elatinaceae 26, 525
 Elatine 525
 triandra var. americana 525
 Eleocharis 182, 235
 acicularis 236
 calva 236
 compressa 236
 elliptica 237
 engelmannii 235, 237
 palustris 237, **238**
 pauciflora var. fernaldii 237
 rostellata 237
 smallii 237
 uniglumis 237
 Ellisia 605
 nyctelea 605
 Elodea 73
 canadensis 73
 Elsholtzia 620, 623
 ciliata 623
 Elymus 81, 123, 804
 angustus 124

- arenarius 123, 124
 ssp. mollis 124
 canadensis 123, 124, **125**
 cinereus 123, 124
 glaucus 123, 124
 hirtiflorus 123, 124
 innovatus 123, 126
 interruptus 123, 126
 junceus 123, 126
 virginicus 123, 126
 Empetraceae 25, 514
 Empetrum 514
 nigrum 514
 Endolepis 326, 335
 suckleyi 335
 Epigaea 572, 573
 repens 573
 Epilobium 536, 537
 adenocaulon 539
 adenocladon 539
 alpinum 537
 angustifolium 537, **538**
 ciliatum 537
 glandulosum var. *adenocaulon* 539
 latifolium 537
 leptophyllum 539
 lineare 539
 palustre 537
 paniculatum 537
 Equisetaceae 20, 47
 Equisetum 47
 arvense 48, **49**
 fluviatile 48
 hyemale 48
 var. affine 48
 kansanum 48
 laevigatum 48
 limosum 48
 palustre 48
 pratense 48
 prealtum 48
 scirpoides 48
 sylvaticum 47
 variegatum 48
 Eragrostis 79, 126
 cilianensis 127
 hypnoides 127
 poaeoides 127
 Ericaceae 29, 30, 571
 Erigeron 705, 707, 708, 748
 acris var. *asteroides* 748, 750
 annuus 749, 750
 arthurii 749, 750
 asper 750
 aureus 749, 751
 caespitosus 749, 751, **752**
 canadensis 748, 751
 compositus 748, 749, 751
 elatus 748, 751
 flagellaris 749, 751
 glabellus 750, 753
 grandiflorus 749, 750, 753
 hyssopifolius 749, 750, 753
 lanatus 749, 753
 lonchophyllus 748, 753
 ochroleucus var. *scribneri* 749, 753
 pallens 749, 753
 peregrinus ssp. *callianthemus* var.
 scaposus 750, 754
 philadelphicus 750, 754
 pumilus 749, 754
 radicatus 749, 754
 scotteri 748, 749, 754
 speciosus 750, 754
 uniflorus var. *unalaschkensis* 748, 754
 Eriogonum 24, 308, 309
 androsaceum 309
 cernuum 309
 flavum 309, 310, **311**
 multiceps 309, 310
 ovalifolium 309, 310
 subalpinum 310
 umbellatum 309, 310
 Eriophorum 182, 239
 angustifolium 239
 brachyantherum 239
 callitrix 239, 240
 chamissonis 239, 240
 f. *albidum* 240
 gracile 239, 240
 scheuchzeri 239, 240
 vaginatum 239, 240
 ssp. *spissum* 240
 viridi-carinatum 239, 240
 Erodium 504
 cicutarium 504
 Eruca 391, 406
 sativa 406
 Erucastrum 392, 407
 gallicum 407
 Eryngium 548, 553
 planum 553
 Erysimum 392, 393, 407
 asperum 407, **408**, 804
 cheiranthoides 407
 hieracifolium 407, 409
 inconspicuum 407, 409
 pallassii 407, 409
 parviflorum 409

- Erythronium 257, 262
 grandiflorum 262, **263**
 Eupatorium 705, 755
 bruneri
 maculatum 755
 var. *bruneri* 755
 perfoliatum 755
 Euphorbia 511
 cyparissias 511
 esula 511, 512, **513**
 geyeri 511, 512
 glyptosperma 512
 helioscopia 511, 512
 lucida 511, 512
 marginata 511, 512
 peplus 511, 512
 serpyllifolia 511, 512
 var. *hirtella* 512
 Euphorbiaceae 23, 511
 Euphrasia 639, 644
 arctica 644
 var. *dolosa* 644
 Eurotia 326, 338
 lanata **337**, 338
 Eutrema 393, 409
 edwardsii 409
 Extensae 189, 191, 197

 Fagaceae 25, 303
 Fagopyrum 309, 310
 esculentum 310
 tataricum 310, **312**
 Ferrugineae 189, 197
 Festuca 80, 127
 altaica 128
 f. *scabrella* 128
 doreana 128
 elatior 127, 129
 var. *arundinacea* 129
 hallii 128, 129
 idahoensis 128, 129, **130**
 obtusata 127, 129
 occidentalis 128, 131
 octoflora 127, 131
 ovina 128, 131
 ssp. *eu-ovina* 131
 var. *saximontana* 131
 var. *vulgaris* 131
 rubra 128, 131
 ssp. *eu-rubra* 131
 subvar. *arenaria* 131
 var. *genuina* 131
 saximontana 131
 scabrella 128
 subulata 127, 131
 viridula 127, 132
 Festuceae 77–79
 Festucoideae 78, 79
 Filifoliae 186
 Firmiculmes 186
 Foetidae 186
 Folliculatae 188
 Fragaria 437, 442
 americana 442
 glauca 443
 vesca var. *americana* 442
 virginiana 442
 virginiana var. *glauca* 443
 Franseria 685
 acanthicarpa 685
 Fraxinus 587
 campestris 587
 nigra 587
 pennsylvanica var. *austini* 587, **588**
 var. *subintegerrima* 587
 Fritillaria 257, 262
 pudica 262
 Fumaria 385, 386
 officinalis 386
 Fumariaceae 28, 385

 Gaillardia **684**, 706, 755
 aristata 755, **756**
 Galeopsis 621, 623
 speciosa 623
 tetrahit 623
 Galinsoga 708, 757
 ciliata 757
 parviflora 757
 Galium 667
 aparine 667, 668
 aparine var. *vaiillantii* 668
 boreale 667, 668, **669**
 var. *intermedia* 668
 palustre 668
 septentrionale 668
 trifidum 668
 triflorum 667, 668
 vaiillantii 668
 verum 667, 670
 Gaultheria 572, 575
 hispidula 575
 humifusa 575
 procumbens 575
 Gaura 536, 539
 coccinea 539
 var. *glabra* 539

- Gayophytum 536, 540
 humile 540
 Gentiana 589
 acuta 590
 affinis 589, 590
 amarella 590
 andrewsii 589, 590
 f. *albiflora* 590
 aquatica 589, 590
 calycosa 589, 590
 crinita 589, 591
 var. *browniana* 591
 var. *tonsa* 591
 detonsa 589
 var. *raupii* 591
 flavida 590
 fremontii 590
 glauca 589, 591
 interrupta 590
 linearis 589, 591
 macounii 591
 plebeia 590
 procera 591
 propinqua 590, 591
 prostrata 590
 puberula 591
 puberulenta 589, 591
 scopulorum 590
 strictiflora 590
 Gentianaceae 31, 589
Geocaulon lividum 305
Geoprimum crassicarpum 469
 Geraniaceae 27, 504
 Geranium 504
 bicknellii 504, 505
 carolinianum 504, 505
 pusillum 504, 505
 richardsonii 504, 505
 strigosum 505
 viscosissimum 504, 505, **506**
 Geum 437, 443
 aleppicum 443
 perincisum 443
 rivale 443
 strictum 443
 triflorum 443, 444, **445**
 Gilia 600
 aggregata 600, 602
 congesta 600, 602
 Glaux 24, 580, 583
 maritima 583
 Glechoma 621, 623
 hederacea 623
 Glyceria 80, 132
 borealis 132, 133
 elata 133
 fluitans 132, 133
 grandis 132, 133, **134**
 maxima 133
 ssp. *grandis* 133
 var. *americana* 133
 pauciflora 132, 133
 pulchella 132, 133
 striata 132, 135
 Glycyrrhiza 463, 476
 lepidota 476, **477**
 Gnaphalium 704, 757
 macounii 758
 microcephalum 757
 palustre 757, 758
 uliginosum 757, 758
 viscosum 757, 758
 Goodyera 271, 275
 oblongifolia 275, 276
 repens 275, 276
 var. *ophioides* 276
 Gracillimae 190
 Gramineae 22, 73, 74
 Granulares 190, 197
 Gratiola 638, 644
 neglecta 644
 Grindelia 706, 758
 perennis 758
 squarrosa 758
 Gutierrezia 706, 758
 diversifolia 758
 sarothrae 758
 Gymnospermae 20, 21, 56, 790
 Gypsophila 347, 353
 acutifolia 353
 elegans 353
 paniculata 353
 Habenaria 270, 276
 bracteata 277
 dilatata 276
 hookeri 276
 hyperborea **272**, 276, 277
 obtusata 276, 277
 orbiculata 276, 277
 saccata 276, 277
 unalascensis 276, 277
 viridis var. *bracteata* 276, 277
Hackelia americana 612
 floribunda 613
 Halenia 589, 592
 deflexa 592

- Halimolobos 393, 409
 virgata 409
- Haloragaceae 23, 26, 543
- Haplopappus 705, 706, 758
 armerioides 759
 lanceolatus 759
 lyallii 759
 nuttallii 758, 759
 spinulosus 758, 759, **760**
- Hedeoma 621, 624
 hispida 624
- Hedysarum 463, 476
 alpinum 479
 var. americanum 476, **478**
americanum 476
 boreale 476, 479
 var. boreale 479
 var. cinerascens 479
 var. mackenzii 479
 sulphurescens 476, 479
- Helenium 706, 761
 autumnale var. montanum 761
montanum 761
- Heleonastes 187, 192
- Helianthemum 525
 bicknellii 525
- Helianthus **684**, 707, 761
 annuus 761
 f. lenticularis 761
 var. giganteus 761
annuus 762
 ssp. *lenticularis* 762
aridus 762
 couplandii 761, 762
 laetiflorus var. subrhomboideus 761,
 762
 maximiliani 761, 762, **763**
 nuttallii 761, 762
 var. subtuberosus 762
petiolaris 762
 tuberosus 761, 764
- Helictotrichon 81, 135
 hookeri 135
- Heliopsis 707, 764
 helianthoides var. scabra 764
- Heliotropium 609, 612
 curassavicum var. obovatum 612
spathulatum 612
- Hepatica 363, 374
 nobilis 374
- Heracleum 548, 555
 lanatum 555, **556**
maximum 555
- Hesperis 393, 409
 matronalis 409
- Heuchera 421
 cylindrica 421, 422
 var. glabella 422
 glabra 421, 422
 parvifolia var. dissecta 421, 422
 richardsonii 421, 422, **423**
- Hieracium 689, 690, 694
albertinum 694
 albiflorum 694
 aurantiacum 694
canadense 695
 cynoglossoides 694, 695
 triste var. gracile 694, 695
 umbellatum 694, 695
- Hierochloa 83, 135
 alpina 135
 odorata 135, **136**
- Hippuris 23, 543
 vulgaris 543
- Hirtae 188, 197
- Homalobus caespitosus* 474
serotinus 472
tenellus 475
vexilliflexus 475
- Hordeae 77, 78, 80
- Hordeum 81, 137
brachyantherum 137
 distichon 137
 hexastichon 137
 jubatum 137, **138**
 var. boreale 137
 var. caespitosum 137
 pusillum 137
 vulgare 137
- Houstonia 667, 670
 longifolia 670
- Hudsonia 525, 526
 tomentosa 526
- Humulus 304
 lupulus 304
- Hutchinsia procumbens* 410
- Hydrocharitaceae 22, 73
- Hydrophyllaceae 31, 605
- Hydrophyllum 605
 capitatum 605
- Hymenolobus 389, 409
 procumbens 409
- Hymenopappus 705, 764
 filifolius 764
 var. polycephalus 764
- Hymenoxys 706
 acaulis 764
 richardsonii 764, **765**
- Hyoscyamus 631
 niger 631, **632**

- Hypericaceae 26, 522
 Hypericum 522
 canadense 522
 formosum var. *nortoniae* 522, 525
 majus 522, 525
 virginicum var. *fraseri* 522, 525
 Hypochoeris 689, 695
 radicata 695
Hypopithys latisquama 571
 Hypoxis 268
 hirsuta 268
 Hyssopus 620, 624
 officinalis 624

 Iliamna 520
 rivularis 520
 Impatiens 508
 biflora 508, 510
 capensis 508
 noli-tangere 508, 510
 occidentalis 510
 Intermediae 187
 Iridaceae 22, 268
 Iris 268, 269
 pseudacorus 269
 versicolor 269
 Isoetaceae 20, 50
 Isoetes 50
 bolanderi 50, 51
 echinospora 51
 var. *braunii* 51
 Iva 683, 686
 axillaris 686
 xanthifolia 686, **687**

 Juncaceae 22, 74, 246
 Juncaginaceae 22, 68
 Juncus 247
 albescens 248, 249
 alpinus 249
 arcticus 247, 249
 articulatus 249, 250
 balticus 247, 250
 var. *littoralis* 250
 var. *montanus* 250, **251**
 biglumis 248, 250
 brevicaudatus 249, 250
 bufonius 247, 250
 canadensis 249, 250
 castaneus 248, 252
 compressus 248, 252
 confusus 248, 252
 drummondii 247, 252
 dudleyi 248, 252
 ensifolius 248, 252
 filiformis 247, 252
 longistylis 248, 253
 mertensianus 248, 253
 nodosus 248, 253
 parryi 247, 253
 saximontanus 248, 253
 tenuis 248, 253
 var. *dudleyi* 252
 torreyi 249, 254
 tracyi 248, 254
 vaseyi 248, 254
 Juniperus 60
 communis 60
 var. *depressa* 60
 horizontalis 60, 61, **62**
 scopulorum 60, 61
 sibirica 60

 Kalmia 28, 571, 575
 polifolia 575
Kentrophyta montana 472
 Knautia 681
 arvensis 681
 Kobresia 182, 241
 myosuroides 241
 simpliciuscula 241
 Kochia 326, 338
 scoparia 338
 Koeleria 81, 137
 gracilis 137, **139**
 ssp. *eugracilis* 140
 ssp. *nitida* 140
 var. *glabra* 140
 var. *typica* 140
 Krigia 690, 695
 biflora 695

 Labiatae 30, 31, 620
 Lactuca **684**, 690, 695
 biennis 695, 696
 canadensis 696
 floridana 696
 ludoviciana 695, 696
 pulchella 695, 696
 serriola 695, 696
 var. *integrata* 697
 spicata 696
 virosa 697
 Lamium 621, 624
 album 624
 amplexicaule 624
 Laportea 304
 canadensis 304
 Lappula 609, 612, 804
 americana 612

- deflexa var. americana 612, 804
 echinata 612
 var. occidentalis 612
 floribunda 612, 613
 jessicae 612, 613
 occidentalis 612
 Lapsana 689, 697
 communis 697
 Larix 56, 57
 laricina 57
 lyallii 57
 occidentalis 57
 Lathyrus 462, 479
 japonicus 479, 480
 ssp. maritimus 480
 ochroleucus 479, 480, **481**
 odoratus 479
 palustris 479, 480
 sativus 479, 480
 tuberosus 479, 480
 venosus 479, 480
 var. intonsus 480
 Lavatera thuringiaca 520
Lavauxia flava 541
 Laxiflorae 190
 Lechea 525, 526
 intermedia 526
 stricta 526
 Ledum 571, 575
 glandulosum 575, 576
 groenlandicum 576
 palustre 575, 576
 var. decumbens 576
 var. latifolium 576
 Leersia 83, 140
 oryzoides 140
 Leguminosae 27, 28, 462
 Lemna 245
 minor 245
 trisulca 245
 Lemnaceae 21, 245
 Lentibulariaceae 31, 659
 Leonurus 621, 624
 cardiaca 624
 sibiricus 624, 625
Lepachys columnifera 770
 Lepidium 389, 410
 apetalum 411
 campestre 410
 densiflorum 410, 411
 var. bourgeauanum 411
 latifolium 410, 411
 perfoliatum 410, 411
 ramosissimum 410, 411
 ruderales 410, 411
 sativum 410, 411
 Leptarrhena 421, 422
 pyrifolia 422
Leptilon canadense 751
Leptotaenia multifida 557
 Lesquerella 391, 412
 alpina var. spathulata 412
 arctica 412
 var. purshii 412
 ludoviciana var. arenosa **390, 412, 413**
Leucanthemum vulgare 741
Leucophysalis grandiflora 634
 Levisticum 548, 555
 officinale 555
 Lewisia 345, 346
 pygmaea 346
 Liatris 705, 766
 ligulistylis 766
 punctata 766, **767**
 Lilaea 65, 66
 scillioides 66
 subulata 66
 Liliaceae 22, 256, 804
 Lilium 256, 262
 philadelphicum 262
 var. andinum 262, **264**
Limnobotrya lacustris 428
Limnorchis dilatata 276
 viridiflora 277
 Limosae 191, 198
 Limosella 638, 644
 aquatica 644
 Linaceae 28, 507
 Linanthus 600, 602
 harknessii var. septentrionalis 602
 septentrionalis 602
 Linaria 638, 644
 canadensis 645
 var. texana 645
 dalmatica 645
 maroccana 645
 vulgaris 645
 Linnaea 670, 671
 americana 671
 borealis var. americana 671, **672**
 Linum 507
 compactum 508
 lewisii 507
 pratense 507, 508
 rigidum 507, 508, **509**
 sulcatum 507, 508
 Liparis 271, 277
 loeselii 277
 Listera 270, 277
 borealis **272, 278**

- caurina 278
 convallarioides 278
 cordata 277, 278
 Lithophragma 420, 422
 bulbifera 422
 parviflora 422, 424
 Lithospermum 609, 613
 angustifolium 615
 arvense 613
 canescens 613
 incisum 613, **614**
 officinale 613, 615
 ruderales 613, 615
 Loasaceae 27, 531
 Lobelia 680
 dortmanna 680
 kalmii 680
 spicata var. *hirtella* 680
 strictiflora 680
 Lobeliaceae 31, 680
 Loiseleuria 572, 576
 procumbens 576
 Lolium 80, 140
 multiflorum 140
 perenne 140
 persicum 141, **142**
 temulentum 141
 Lomatium 548, 549, 555, 559
 cous 555, 557
 dissectum var. *multifidum* 555, 557
 foeniculaceum 557
 macrocarpum 555, 557
 orientale 555, 557
 sandbergii 555, 557
 simplex 557
 triternatum 555, 557
 villosum 555, 557, **558**
 Lomatogonium 589, 592
 rotatum 592
 Longirostres 189
 Lonicera 670, 671
 caerulea var. *villosa* 671, 673
 dioica 673
 var. *glaucescens* 671, **672**, 673
 glaucescens 673
 hirsuta var. *schindleri* 671, 673
 involucrata 671, 673
 oblongifolia 671, 673
 tatarica 671, 673
 utahensis 671, 673
 villosa var. *solonis* 673
 Loranthaceae 23, 308
 Lotus 463, 480
 americanus 482
 corniculatus 482
 pedunculatus 482
 purshianus 482
 Luetkea 437, 444
 pectinata 444
 Lupinus 463, 483
 argenteus 483, **484**
 flexuosus 485
 minimus 482, 483
 nootkatensis 483
 polyphyllus 483
 pusillus 482, 483
 sericeus 482, 485
 Lupulinae 189
 Luzula 254
 campestris 254, 255
 confusa 254, 255
 glabrata 254, 255
 groenlandica 255
 parviflora 254, 255
 pilosa 254
 var. *saltuensis* 255
 spicata 255
 wahlenbergii 254
 Lychnis 347, 353
 affinis 353, 354
 alba 353, 354
 apetala 353, 354
 ssp. *attenuata* 354
 var. *arctica* 354
 drummondii 353, 354
 Lycium 631, 633
 halimifolium 633
 Lycopodiaceae 20, 51
 Lycopodium 51
 alpinum 51, 52
 annotinum 51, 52
 clavatum 51, 52
 complanatum 51, 52, **53**
 inundatum 51, 52
 obscurum 51, 52, **54**
 sabinifolium 52
 var. *sitchense* 55
 selago 51, 55
 Lycopsis 610, 615
 arvensis 615
 Lycopus 621, 625
 americanus 625
 asper 625
 uniflorus 625
 virginicus 625
 var. *pauciflorus* 625
 Lygodesmia 690, 697, 702
 juncea 697, **698**
 rostrata 697
 Lysiella *obtusata* 277

- Lysimachia* 14, 581, 583
 ciliata 13, 14, 584, **585**
 hybrida 584
 quadrifolia 584
 terrestris 583, 584
 thyrsiflora 584, 586
Lythraceae 26, 534
Lythrum 534
 salicaria 534
- Machaeranthera canescens* 731
 pulverulenta 731
Madia 704, 707, 766
 glomerata 766
Maianthemum 256, 257, 262
 canadense var. *interius* **261**, 262
Malaxis 271, 278, 804
 monophylla var. *brachypoda* 278
 paludosa 278
 unifolia 278
Malva 520
 alcea 520, 521
 crispa 521
 moschata 520, 521
 neglecta 521
 parviflora 521
 pusilla 521
 sylvestris 520, 522, **523**
Malvaceae 27, 520
Malvastrum 520, 522
 coccineum 522, **524**
Mamillaria 531
 vivipara 531
Marrubium 621, 625
 vulgare 625
Marsilea 47
 mucronata 47
 vestita 47
Marsileaceae 21, 47
Maruta cotula 717
Matricaria 705, 708, 766
 chamomilla 768
 inodora 717, 768
 maritima 768, **769**
 var. *maritima* 717
 matricarioides 766, 768
Matteuccia 35, 42
 struthiopteris 42
 var. *pennsylvanica* 42, **43**
Medicago 462, 485
 lupulina 485
 polymorpha 485
 sativa 485
 ssp. *falcata* 485, 487
 ssp. *sativa* 485, **486**, 487
- Megalodonta* 707
 beckii 739
Melampyrum 639, 645
 lineare 645, **646**
Melica 80, 141
 bulbosa 141
 smithii 141
 spectabilis 141, 143
 subulata 141, 143
Melilotus 462, 487
 alba 487, **488**
 indica 487
 officinalis 487
 wolgica 487
Melissa 621, 626
 officinalis 626
Menispermaceae 28, 384
Menispermum 384
 canadense 384
Mentha 620, 621, 626
 arvensis 626
 var. *villosa* 626
 spicata 626
Mentzelia 531
 decapetala 531, **532**
Menyanthes 589, 592
 trifoliata 592
Menziesia 572, 576
 ferruginea var. *glabella* 576
Meriolix serrulata 543
Mertensia 610, 615
 lanceolata 615
 linearis 615
 longiflora 615, 616
 maritima 615, 616
 paniculata 615, 616, **617**
Micrampelis lobata 677
Micranthes rhomboidea 433
Microseris 689, 690, 697
 cuspidata 697, 699
 nutans 697, 699
Milium 82, 143
 effusum 143
Mimulus 639, 647
 floribundus 647
 glabratus 647
 guttatus 647
 langsдорffii 647
 lewisii 647
 ringens 647
Minuartia laricifolia 350
Mirabilis 344
 hirsuta 344
 linearis 344
 nyctaginea 344, 345

- Mitella 420, 424
 breweri 424
 nuda 424
 pentandra 424
 trifida 424
Moehringia lateriflora 350
Moldavica parviflora 623
 thymiflora 623
 Monarda 620, 626
 fistulosa 626, **627**
 var. *menthaefolia* 628
 menthaefolia 628
 Moneses 567
 uniflora 567, **568**
 Monocotyledoneae 20, 21, 62, 790
 Monolepis 326, 338
 nuttalliana 338
 Monotropa 570, 571
 hypopithys var. *latisquama* 571
 uniflora 571
 Monotropaceae 30, 570
 Montanae 188, 189, 195
Montia linearis 346
 Muhlenbergia 82, 143
 andina 144, **145**
 asperifolia 143
 cuspidata 143
 glomerata 144
 var. *cinnoides* 144
 mexicana 144, 146
 racemosa 144, 146
 richardsonis 143, 146, **147**
 Multiflorae 187
 Munroa 79, 83, 146
 squarrosa 146
 Musineon 549, 559
 divaricatum 559
 trachyspermum 559
 Myosotis 610, 616
 arvensis 616
 laxa 616
 scorpioides 616
 sylvatica var. *alpestris* 616, 618
 Myosurus 363, 374
 minimus 374
 Myrica 299
 gale 299
 Myricaceae 25, 299
 Myriophyllum 23, 543, 544
 alterniflorum 544
 exalbescens 544
 pinnatum 544
 scabratum 544
 spicatum 544
 spicatum var. *exalbescens* 544
Myzorrhiza ludoviciana 663

Nabalus albus 699
 racemosus 700
 Najas 66
 flexilis 66
 Nardinae 185, 186
Naumburgia thyrsoflora 586
 Navarretia 600, 602
 minima **601**, 602
Negundo interius 517
 Nemophila 605, 606
 breviflora 606
Neomamillaria vivipara 531
 Nepeta 620, 628
 cataria 628
 Neslia 391, 412
 paniculata **390**, 412
 Nonea 610, 618
 vesicaria 618
 Nuphar 361
 microphyllum 361, 362
 variegatum 361, 362
Nuttallia decapetala 531
 Nyctaginaceae 24, 344
 Nymphaea 361, 362
 odorata 362
 tetragona ssp. *leibergii* 362
 Nymphaeaceae 26, 361

 Obtusatae 186, 191, 195
Ochrocodon pudicus 262
 Odontites 639, 648
 serotina 648
 Oenothera 536, 540
 andina 540
 biennis 540, 541
 var. *canescens* 541
 var. *hirsutissima* 541
 breviflora 540, 541
 caespitosa 540, 541, **542**
 var. *montana* 541
 flava 540, 541
 fruticosa var. *linearis* 540, 543
 nuttallii 540, 543
 perennis 540, 543
 serrulata 540, 543
 Oleaceae 25, 29, 587
 Onagraceae 26, 27, 534
 Onoclea 35, 42
 sensibilis 42
 struthiopteris 42

- Onosmodium 609, 618
 molle var. hispidissimum 618
 var. occidentale 618
 Ophioglossaceae 21, 32
 Oplopanax 544, 545
 horridum 545
 Opuntia 531
 fragilis 533
 polyacantha 533
 Orchidaceae 22, 270, 804
 Orchis 270, 279
 rotundifolia 272, 279
Oreocarya glomerata 611
 Orobanchaceae 30, 661
 Orobanche 661
 fasciculata 661
 ludoviciana 661, 663
 uniflora 661, 663
Orophaca caespitosa 470
Orthilia secunda 570
 Orthocarpus 638, 648
 luteus 648
 Orthocerates 185, 199
 Oryzae 78, 83
 Oryzopsis 82, 146
 asperifolia 148
 canadensis 148
 exigua 148
 hymenoides 146, 149
 micrantha 148, 150
 pungens 148, 150
 Osmorhiza 548, 559
 aristata 559
 var. brevistylis 559
 var. longistylis 559, 560
 chilensis 559
 var. chilensis 559, 561
 var. cupressimontana 559, 561
 var. purpurea 559, 561
 divaricata 559
 obtusa 561
 occidentalis 559
 purpurea 561
 Osmunda 35
 claytoniana 35
 Osmundaceae 21, 35
 Ostrya 299, 303
 virginiana 303
 Ovales 187, 193
 Oxalidaceae 27, 505
 Oxalis 505
 corniculata 505, 507
 europaea 505, 507
 stricta 505, 507
Oxybaphus hirsutus 344
 linearis 345
 nyctagineus 345
 pilosus 344
 Oxycoccus 577, 578
 microcarpus 578
 palustris 578
 quadripetalus 578
 Oxyria 309, 310
 digyna 310
 Oxytropis 463, 489
 bellii 489, 490
 besseyi 489, 490, 491
 campestris 489, 490
 ssp. campestris 490
 ssp. gracilis 490
 var. cusickii 490
 var. dispar 490
 var. gracilis 490
 var. johannensis 490
 var. varians 490
 deflexa 489, 492
 var. deflexa 492
 var. foliolosa 492
 var. sericea 492
 lagopus 489, 492
 lambertii 489, 492
 podocarpa 489, 493
 sericea 489, 493
 var. sericea 493
 var. spicata 493, 494
 splendens 489, 493
 var. richardsonii 493
 viscida 463, 489, 493

Pachylophus caespitosus 541
 montanus 541
 Pachystima 517
 myrsinites 517
 Paludosae 189, 200
 Paniceae 77, 79, 83, 190, 196
 Panicoideae 78, 83
 Paniculatae 187, 192
 Panicum 84, 150
 capillare 150, 151
 var. occidentale 151
 depauperatum 151
 lanuginosum 150, 151
 leibergii 150, 152
 linearifolium 151, 152
 miliaceum 150, 152
 oligosanthes 151
 var. scribnerianum 152
 perlongum 151, 152

- virgatum 150, 152, **153**
 wilcoxianum 151, 152
 xanthophysum 151, 154
Papaver 384
 nudicaule 384, 385
 pygmaeum 384, 385
 rheas 384, 385
 somniferum 384, 385
Papaveraceae 26, 384
Parietaria 304, 804
 pennsylvanica 304
Parnassia 420, 425
 americana 425
 fimbriata 425
 glauca 425
 kotzebuei 425
 palustris var. *montanensis* 425
 var. *parviflora* 425
 var. *tenuis* 425, **426**
Paronychia 24, 347, 354
 depressa 354
 sessiliflora 354
Parthenocissus 519
 inserta 519
 quinquefolia 519
Pastinaca 548, 561
 sativa 561
Pedicularis 638, 648
 bracteosa 649
 canadensis 649
 capitata 648, 649
 contorta 648, 649
 flammea 648, 650
 groenlandica 648, 650, **651**
 labradorica 649, 650
 lanata 649, 650
 lanceolata 648, 650
 langsдорffii 649, 650
 lapponica 648–650
 montanensis 649
 oederi var. *albertae* 649, 650
 parviflora 648, 652
 racemosa 648, 652
 sudetica 649, 652
Pedimelum esculentum 497
Pellaea 35, 44
 glabella 44
 var. *nana* 44
 var. *simplex* 44
Penstemon 638, 652
 albertinus 653
 albidus 652, 653
 confertus 652, 653
 ellipticus 652, 653
 eriantherus 652, 653
 fruticosus 652, 653
 gracilis 653, 654, **655**
 lyallii 652, 654
 nitidus 652–654, **655**
 procerus 652, 654, **655**
Pentastemon 652
Penthorum 420, 427
 sedoides 427
Pentstemon 652
Perideridia 549, 561
 gairdneri 561
Peritoma serrulatum 386
Persicaria 313, 314
Petalostemon 463, 493
 candidum 495
 purpureus 495, **496**
 var. *pubescens* 495
 villosum 493, 495
Petasites 705, 707, 768
 palmatum 768, 770, **771**
 sagittatum 768, 770, **771**
 vitifolium 768, 770, **771**
Phaca americana 470
Phacelia 605, 606
 campanularia 606
 franklinii 606
 hastata 606, 607
 linearis 606, 607
 lyallii 606, 607
 sericea 606, 607, **608**
 tanacetifolia 606, 607
 thermalis 606, 607
Phalarideae 77, 78, 83
Phalaris 83, 154
 arundinacea 154, **155**
 canariensis 154
Philadelphus 420, 427
 lewisii 427
Phleum 82, 154
 alpinum 154, **156**
 pratense 154, **157**
Phlox 600, 602
 alyssifolia 602
 gracilis 602, 603
 hoodii **601**, 602, 603
 pilosa 602, 603
Phragmites 79, 158
 communis 158
Phryma 663
 leptostachya 663
Phrymaceae 30, 663
Phyllodoce 572, 576
 caerulea 576, 577

- empetriformis 576, 577
 glanduliflora 576, 577
 Phyllostachyae 186, 189
 Physalis 631, 633
 alkekengi 633
 grandiflora 633, 634
 heterophylla 633, 634
 ixocarpa 633, 634
 pubescens 633, 634
 virginiana 633, 634
 Physaria 389, 414
 didymocarpa 414
 Physocarpus 436, 444
 malvaceus 444
 Physostegia 620, 628
 parviflora 628
 virginiana var. *formosior* 628
 var. *ledinghamii* 628
 var. *parviflora* 628
 Picea 56, 57
 glauca 58
 var. *albertiana* 58
 mariana 58
Picradeniopsis oppositifolia 738
 Picris 689, 699
 echioides 699
 Pinaceae 21, 56
 Pinguicula 659
 macroceras 659, 660
 villosa 659, 660
 vulgaris 659, 660
 Pinus 56, 58
 albicaulis 59
 banksiana 58, 59
 contorta 58
 var. *latifolia* 59
 divaricata 59
 var. *latifolia* 59
 flexilis 59
 monticola 59
 murrayana 59
 ponderosa 58, 59
 resinosa 58, 60
 strobus 59, 60
Pisophaca flexuosa 470
 Plagiobothrys 609, 618
 scouleri var. *penicillatus* 618
 Plantaginaceae 30, 663
 Plantago 663
 aristata 664
 canescens var. *cylindrica* 664
 coronopus 663, 664
 elongata 664, 665
 eriopoda 664, 665
 lanceolata 664, 665
 major 664, 665, **666**
 var. *asiatica* 665
 maritima 664, 665
 media 664, 665
 patagonica 664, 667
 psyllium 663, 667
 purshii 667
 spinulosa 667
Pleurogyne rotata 592
 Poa 80, 158
 agassizensis 165
 alpina 160
 ampla 158, 160
 annua 159, 160
 arctica 158, 159, 161
 arida 158, 161
 canbyi 159, 161, **162**
 compressa 158, 161
 cusickii 159, 161
 epilis 159, 161
 fendleriana 158, 163
 glauca 159, 163
 glaucifolia 159, 163
 gracillima 160, 163
 interior 164
 juncifolia 160, 163
 leptocoma 159, 163
 nemoralis 159, 164
 var. *firmula* 164
 var. *interior* 164
 nervosa 158, 164
 palustris 160, 164
 pattersonii 159, 164
 pratensis 158, 164
 var. *arenaria* 165
 var. *humilis* 165
 rupicola 159, 165
 secunda 159, 165
 stenantha 160, 165
 trivialis 160, 165
 Polanisia 386, 389
 dodecandra 389
 var. *trachysperma* 389
 graveolens 389
 Polemoniaceae 30, 31, 600
 Polemonium 600, 603
 occidentale 603
 pulcherrimum 603, **604**
 viscosum 603, 605
 Politrichoideae 186
 Polygala 510
 alba 510
 paucifolia 510

- senega 510
 verticillata 510
 Polygalaceae 28, 510
 Polygonaceae 24, 308
 Polygonatum 257, 262
 canaliculatum 262
 commutatum 262
 Polygonum 309, 313, 804
 achoreum 313, 315
 amphibium 314, 315
 var. *emersum* 315
 amphibium var. *stipulaceum* 315
 austinae 313, 315
 aviculare 314, 315, **316**
 bistortoides 314, 317
 boreale 314, 317
 cilinode 314, 317
 coccineum f. *natans* 315
 f. *terrestre* 315
 confertiflorum 313, 317
 convolvulus 314, 317
 douglasii 313, 317
 engelmannii 313, 317
 erectum 314, 317
 hydropiper 315, 318
 kelloggii 313, 318
 lapathifolium 315, 318, 804
 minimum 313, 318
 natans f. *genuinum* 315
 f. *hartwrightii* 315
 persicaria 315, 318, **319**
 punctatum 315
 var. *confertiflorum* 318
 ramosissimum 314, 318
 sagittatum 315, 318
 scabrum 315, 320, 804
 scandens 314, 320
 tomentosum 318
 viviparum 314, 320
 Polypodiaceae 21, 35
 Polypodium 36, 44
 vulgare 44
 var. *columbianum* 44
 var. *virginianum* 44, **45**
 Polypogon 82, 165
 monspeliensis 165
 Polystichum 36, 44
 lonchitis 44
 Populus 279, 280
 alba 280
 angustifolia 280
 balsamifera 280, 281
 candicans 281
 deltoides 280, 281
 var. *occidentalis* 281
 gileadensis 281
 grandidentata 280, 281
 tacamahacca 281
 tremuloides 280, 281
 trichocarpa 281
 Portulaca 345, 346
 oleracea 346
 Portulacaceae 26, 28, 345
 Potamogeton 65, 66
 foliosus 66, 67
 gramineus 66, 67
 interior 66, 67
 pectinatus 66, 67
 richardsonii 66, 67
 vaginatus 66, 67
 Potentilla 436, 437, 444
 anserina 446, 448
 f. *sericea* 448
 argentea 448
 arguta 446, 448
 bipinnatifida 447, 448
 concinna 447, 448
 diversifolia var. *glaucophylla* 448
 drummondii 446, 449
 flabellifolia 447, 449
 var. *emarginata* 449
 floribunda 449
 fruticosa 449
 fruticosa ssp. *floribunda* 444, 449, **450**
 glandulosa var. *intermedia* 446, 449
 gracilis 447-449
 var. *flabelliformis* 449
 var. *gracilis* 449
 hippiana 446, 451
 var. *argyrea* 451
 millegrana 452
 monspeliensis 451
 multifida 447, 451
 nivea 447, 451
 var. *pulchella* 451
 var. *villosa* 451
 norvegica 446, 451
 palustris 444, 451
 paradoxa 446, 452
 pennsylvanica 446, 452, **453**
 var. *atrovirens* 452
 var. *pennsylvanica* 452
 pennsylvanica
 var. *bipinnatifida* 448
 plattensis 446, 452
 quinquefolia 447, 452
 recta 447, 452
 rivalis 446, 447, 452

- saximontana 447, 454
 sibbaldii 447, 454
 tridentata 444, 454
 Prenanthes 690, 699
 alba 699
 racemosa 699, 700
 sagittata 699, 700
 Primula 580, 586
 egaliksensis 586
 incana **582**, 583, 586
 mistassinica 586
 stricta 586
 Primulaceae 14, 24, 31, 580
 Prunella 620, 628
 vulgaris 628
 Prunus 436, 454
 americana 454, 455
 nigra 454, 455
 pennsylvanica 454, 455
 pumila 454, 455
 virginiana 454, 455
 var. melanocarpa 454, 455, **456**
 Psedera *quinquefolia* 519
vitacea 519
 Pseudo-Cyperae 189, 200
 Pseudotsuga 56, 60
 menziesii 60
 Psilocarphus 704, 770
 elatior 770
 Psoralea 462, 495
 argophylla 495
 esculenta 495, 497, **498**
 lanceolata 495, 497, **499**
 Psoralidium *argophyllum* 495
lanceolatum 497
 Pteretis *nodulosa* 42
pennsylvanica 42
 Pteridium 35, 44
 aquilinum 44
 var. pubescens 46
 Pteridophyta 20, 32, 790
 Pterospora 570, 571
 andromedeae 571
 Puccinellia 80, 166
 distans 166
 nuttalliana 166, **167**
 phryganodes 166
 Pulsatilla *ludoviciana* 366
occidentalis 365
patens ssp. *teklae* 366
 Pyrola 567, 569
 asarifolia 569
 var. incarnata **568**, 569
 bracteata 569
chlorantha 570
 elliptica 569, 570
 minor 569, 570
 picta 569, 570
 rotundifolia 569, 570
 secunda **568**, 569, 570
uniflora 567
 virens 569, 570
 Pyrolaceae 26, 29, 31, 567
 Pyrrocoma *lanceolata* 759
 Quercus 303
 macrocarpa 303
 Ranunculaceae 24–28, 362
 Ranunculus 363, 374
 abortivus 375, 376
 acris 375, 376
affinis 380
 aquatilis 374, 377
 var. capillaceus 377
 var. eradicatus 377
 var. longirostris 377
 var. subrigidus 377
buddii 379
 cardiophyllus 375, 377, **378**
 cymbalaria 375, 377
 fascicularis 376, 377
 flammula 374
 var. filiformis 377
 var. ovalis 377
 gelidus 376, 379
 glaberrimus 374, 379
 var. ellipticus 379
 gmelinii 374, 376, 379
 hyperboreus 375, 379
 var. intertextus 379
 inamoenus 375, 379
 lapponicus 375, 379
 macounii 376, 379
 nivalis 375, 379, 380
 var. eschscholtzii 380
 var. nivalis 380
 occidentalis 375, 380
 var. brevistylis 380
ovalis 381
 pallassii 374, 380
 pedatifidus var. leiocarpus 375, 380
 pennsylvanicus 376, 380
 pygmaeus 375, 380
 repens 376, 380
 rhomboideus 375, 380
 sceleratus 376, 380, 381
 septentrionalis 376, 381

- uncinatus 375, 381
verecundus 380
 Raphanus 391, 414
 raphanistrum **390**, 414
 Rapistrum 391, 414
 perenne 414
 Ratibida **684**, 707, 770
 columnifera 770, **772**
 f. *pulcherrima* 770
 Reseda 417
 alba 417
 lutea 417
 Resedaceae 27, 417
 Rhamnaceae 29, 518
 Rhamnus 518
 alnifolius 518
 catharticus 518
 frangula 518, 519
 Rhinanthus 639, 654
 crista-galli 654
 Rhododendron 572, 577
 albiflorum 577
 lapponicum 577
 Rhus 514
 aromatica var. *trilobata* 515
 glabra 514, 515
 radicans var. *rydbergii* 515, **516**
 Rhynchospora 182, 241
 alba 241
 capillacea 241
 Ribes 29, 420, 427
 americanum 428
 aureum 427, 428
 diacanthum 428
 floridum 428
 glandulosum 427, 428
 hirtellum 429
 hudsonianum 428
 lacustre 427, 428
 laxiflorum 427, 429
 oxyacanthoides var. *oxyacanthoides*
 427, 429, **430**
 var. *saxosum* 429
 rubrum var. *propinquum* 429
 setosum 429
 triste 429
 viscosissimum 427, 429
 Romanzoffia 605, 607
 sitchensis 607
 Rorippa 391–393, 414
 austriaca 414
 hispida 415
 islandica 415
 var. *fernaldiana* 415
 var. *hispida* 415
 palustris 415
 sinuata 415
 sylvestris 415
 tenerrima 415
 Rosa 437, 455
 acicularis 457
 arkansana 457
 blanda 457
 woodsii 457
 Rosaceae 27, 29, 436
 Rubiaceae 31, 667
 Rubus 436, 437, 458
 acaulis 458
 chamaemorus 458
 idaeus var. *aculeatissimus* 458, **459**
 melanolasius 458
 parviflorus 458, 460
 pedatus 458, 460
 pubescens 458, 460
 strigosus 458
 Rudbeckia 707, 773
 hirta 773
 laciniata 773
 serotina 773
 Rumex 309, 320
 acetosa 320, 321
 acetosella 320, 321, **322**
 confertus 321
 crispus 321
 dentatus 320, 323
 domesticus var. *pseudonatronatus* 323
 fennicus 321, 323
 longifolius 321, 323
 maritimus var. *fueginus* 320, 323
 mexicanus 324
 occidentalis 321, 323
 orbiculatus 321, 323
 paucifolius 320, 323
 salicifolius 321, 324
 stenophyllus 321, 324
 triangulivalvis 324
 venosus 320, 324, **325**
 Rupestres 186, 190, 196
 Ruppia 65, 67
 maritima 67
 occidentalis 67

Sabina horizontalis 61
 Sagina 348, 354
 caespitosa 355
 decumbens 354, 355
 nodosa 355
 saginoides 355

- Sagittaria 69, 70
 cuneata 70, **72**
 latifolia 70
 Salicaceae 25, 279
 Salicornia 24, 324, 338
 europaea 338
 rubra 338
 Salix 280–282, 286
 alaxensis 284, 287, 290
 alba 286, 287, 290
 amygdaloides 286, 288, 290
 arbusculoides 284, 288, 291
 arctica 284, 289, 291
 arctophila 284, 289, 291
 barclayi 284, 285, 288, 291
 barrattiana 284, 292
 bebbiana 282, 287, 290, 292
 brachycarpa 284, 287, 292, **293**
 calicicola 285, 288, 289, 292
 candida 284, 287, 292
 commutata 285, 287, 294
 discolor 285, 287, 290, 294
 drummondiana 297
 exigua 295
 fluviatilis 295
 f. *hindsiana* 295
 var. *sericans* 295
 fragilis 286, 288, 294
 glauca 284, 287, 289, 290, 294
 herbacea 286, 289, 294
 humilis 285, 287, 294
 interior 282, 286–288, 295
 var. *pedicellata* 295
 var. *wheeleri* 295
 lasiandra 295
 lucida 286, 288, 295
 lutea 286, 289, 290, 295
 maccalliana 284, 289, 296
 mackenzieana 285, 289, 296
 monticola 285, 289, 290, 296
 myrtillofolia 286, 288, 296
 nivalis 298
 pedicellaris 286, 290, 296
 var. *athabascensis* 297
 pellita 285, 287, 290, 297
 var. *angustifolia* 297
 var. *psila* 285, 297
 pentandra 286, 288, 297
 petiolaris 284, 288, 297
 phylicifolia 298
 ssp. *planifolia* 298
 planifolia 285, 290, 297
 pyrifolia 285, 286, 288, 298
 reticulata 288, 289, 298
 var. *nivalis* 298
 serissima 286, 288, 298
 sitchensis 284, 287, 298
 vestita 284, 287, 289, 298
 Salsola 324, 339
 kali var. *tenuifolia* 339
 pestifer 339
 Salvia 620, 628
 nemorosa 628
 reflexa 628, 629
 Sambucus 670, 674
 racemosa 674
 var. *arborescens* 674
 var. *pubens* 674
 Sanguinaria 384, 385
 canadensis 385
 Sanicula 548, 561
 marilandica 561, **562**
 Santalaceae 24, 305
 Saponaria 347, 355
 vaccaria 355, **356**
 Sarcobatus 25, 324, 339
 vermiculatus 339, **340**
 Sarracenia 418
 purpurea 418
 Sarraceniaceae 26, 418
 Saussurea 705, 773
 glomerata 773
 nuda var. *densa* 773
 Saxifraga 421, 429
 adscendens ssp. *oregonensis* 431
 aizoides 431
 bronchialis var. *austromontana* 431, 432
 cernua 431, 432
 cespitosa 431, 432
 ferruginea 431, 432
 f. *vreelandii* 432
 flagellaris 431, 432
 hirculus 431, 432
 lyallii 431, 432
 var. *hultenii* 432
 var. *laxa* 433
 mertensiana 429, 433
 occidentalis 431, 433
 oppositifolia 429, 433
 pensylvanica 429, 433, **434**
 punctata var. *porsildiana* 431, 433
 rhomboidea 433
 rivularis 431, 433
 tricuspidata 431, 435
 virginiensis 431, 435
 Saxifragaceae 24, 28, 29, 420
 Scabiosa *arvensis* 681
 Scandix 549, 561
 pecten-veneris 561

- Schedonnardus 83, 166
 paniculatus 166
 Scheuchzeria 68
 palustris 68
 Schizachne 79, 168
 purpurascens 168, **169**
 Scirpinae 185
 Scirpus 182, 242
 acutus 242, 243
 americanus 242, 243
 atrovirens 243
 caespitosus 242
 var. *callosus* 243
 clintonii 242, 243
 cyperinus 243
 fluviatilis 242, 244
 hudsonianus 242, 244
 microcarpus 243, 244
 nevadensis 242, 244
 paludosus 243, 244
 pumilus 242, 244
 rufus 242, 244
 torreyi 242, 244
 validus 242, 244
 Scolochloa 79, 168
 festucea 168, **170**
 Scrophularia 639, 654
 lanceolata 654
 Scrophulariaceae 30, 31, 638
 Scutellaria 621, 629
 epilobiifolia 629
 galericulata 629
 lateriflora 629
 parvula var. *leonardii* 629
 Secale 80, 168
 cereale 168
 Sedum 418
 acre 419
 aizoon 419
 hybridum 419
 lanceolatum 419
 rosea 419
 var. *integrifolium* 420
 stenopetalum 419, 420
 telephium 419, 420
 Selaginella 55
 densa 55
 rupestris 55
 selaginoides 55
 wallacei 55
 Selaginellaceae 20, 55
 Senecio 705, 706, 774
 aureus 775
 canus 775, **776**
 congestus 774, 775
 eremophilus 774, 777, **778**
 foetidus 774, 777
 fremontii 774, 777
 indecorus 775, 777
 integerrimus 774, 777
 var. *exaltatus* 777
 var. *integerrimus* 777
 var. *lugens* 777
 megacephalus 774, 779
 palustris 775
 pauciflorus 775, 779
 pauperculus 775, 779
 var. *firmifolius* 779
 var. *pauperculus* 779
 var. *thomsoniensis* 779
 resedifolius 774, 779
 streptanthifolius 774, 775, 779
 triangularis 774, 780
 tridenticulatus 774, 775, 780
 viscosus 774, 780
 vulgaris 774, 780
 Setaria 83, 168
 glauca 168
 viridis 168, 171, **172**
 Shepherdia 534
 argentea 534, **535**
 canadensis 534
Sibbaldiopsis tridentata 454
Sideranthus grindelioides 759
 spinulosus 759
Sieversia triflora 444
 Silene 347, 355
 acaulis var. *exscapa* 357
 anthirrina 357
 cserei 357
 cucubalus 357
 menziesii 357, 358
 noctiflora 357, 358
 parryi 357, 358
 sibirica 357, 358
 vulgaris 357
 Sisymbrium 392, 415
 altissimum 415, 416
 linifolium 415, 416
 loeselii 416
 officinale 415, 416
 var. *leiocarpum* 416
 Sisyrinchium 268, 270
 angustifolium 270
 montanum 270
 Sitanion 81, 171
 hystrix 171
 Sium 548, 555, 563
 cicutae-folium 563
 suave 563, **564**

- Smelowskia 392, 416
 calycina var. *americana* 416
 Smilacina 257, 265
 racemosa 265
 var. *amplexicaulis* 265
 stellata 265
 trifolia 265
 Smilax 256, 265
 herbacea var. *lasioneura* 265
 Solanaceae 31, 631
 Solanum 631
 dulcamara 634, 635, **636**
 interius 635
 nigrum 634, 635
 rostratum 634, 635
 sarachoides 635
 triflorum 634, 635, **637**
 Solidago 706, 780
 bicolor 781, 782
 var. *concolor* 781, 782
 canadensis 782
 var. *canadensis* 782, **783**
 var. *gilvocanescens* 782
 decumbens 785
 gigantea var. *serotina* 782
 glaberrima 784
 graminifolia 781
 var. *graminifolia* 782
 var. *major* 784
 hispida var. *lanata* 782
 juncea 782, 784
 lutescens 785
 missouriensis 782, 784, 785
 var. *fasciculata* 785
 mollis 781, 782, 784
 multiradiata 781, 784
 nemoralis 781, 784
 oreophila 785
 pruinosa 782
 ptarmicoides 780, 785
 riddellii 781, 785
 rigida var. *humilis* 781, 785, **786**
 serotina 782
 spathulata var. *spathulata* 781, 785, **787**
 uliginosa 781, 785
 Sonchus 690, 700
 arvensis 700, **701**
 var. *glabrescens* 700
 asper 700
 oleraceus 700
 f. *integrifolius* 702
 uliginosus 700
 Sophia *brachycarpa* 402
 multifida 402
 richardsonii 402
 Sorbaria 437, 460
 sorbifolia 460
 Sorbus 437, 460
 americana 460
 aucuparia 460, 461
 scopulina 461
 sitchensis 460, 461
 Sorghastrum 84, 171
 nutans 171
 Sparganiaceae 22, 63
 Sparganium 63
 angustifolium 63
 chlorocarpum var. *acaule* 63
 eurycarpum 63, 65
 fluctuans 63, 65
 hyperboreum 63, 65
 minimum 63, 65
 multipedunculatum 63, 65
 Spartina 83, 171
 gracilis 171, **173**
 pectinata 171, 174
 Spargula 347, 358
 arvensis 358
 Spargularia 347, 358
 marina var. *leiosperma* 358
 salina 358
 Spermatophyta 12, 20, 21, 56, 790
 Sphenopholis 81, 174
 intermedia 174
 obtusata 174
 Spiraea 436, 461
 alba 461
 densiflora 461
 lucida 461
 Spiranthes 271, 279
 gracilis 279
 romanzoffiana 279
 Spirodela 245
 polyrhiza 245
 Sporobolus 82, 174
 cryptandrus 174, 175, **176**
 heterolepis 174, 175
 neglectus 174, 175
 Stachys 620, 629
 palustris var. *pilosa* 629, **630**
 Steironema 14
 ciliatum 584
 hybridum 584
 Stellaria 347, 358
 americana 359, 360
 borealis 360
 calycantha 359, 360

ciliatisepala 359, 360
 crassifolia 359, 360
 crispa 359, 360
 humifusa 359, 360
 laeta 359, 360
 longifolia 359, 360
 longipes 359, 360
 media 359, 361
 obtusa 359, 361
 umbellata 359, 361
 Stellulatae 188
 Stenanthium 257, 266
 occidentale 266
Stenotus armerioides 759
 Stephanomeria 689, 702
 runcinata 702
 Stipa 82, 175
 columbiana 177
 comata 175, 177, **178**
 richardsonii 177
 spartea 175, 177
 var. *curtiseta* 175, 177
 viridula 177, 179, **180**
 Streptopus 257, 266
 amplexifolius **261**, 266
 var. *americanus* 266
 var. *denticulatus* 266
 roseus var. *perspectus* 266
 Suaeda 326, 339
 depressa 339
 var. *erecta* 339
 maritima var. *americana* 339
 Subularia 389, 416
 aquatica 416
 Suckleya 326, 339
 suckleyana 339, **341**
 Suksdorfia 421, 435
 violacea 435
Svida stolonifera 565
Swainsona salsula 472
 Sylvaticae 189, 190, 197
 Symphoricarpos 670, 674
 albus 674
 var. *pauciflorus* 674
 occidentalis 674
 pauciflorus 674
 Symphytum 610, 618
 asperum 618
 officinale 618, 619
 Tanacetum 704, 788
 huronense 788
 vulgare 788, **789**
 Taraxacum 689, 702, 804
 ceratophorum 702
 erythrospermum 702
 laevigatum 702
 officinale 702
Taraxia breviflora 541
 Taxaceae 21, 56
 Taxus 56
 brevifolia 56
 canadensis 56
 Telesonix 421, 435
 jamesii var. *heucheriformis* 435
Tetraneuris acaulis 764
 Teucrium 620, 631
 canadense var. *occidentale* 631
 Thalictrum 363, 381
 dasycarpum 381, 382, **383**
 occidentale var. *palousense* 381, 382
 sparsiflorum var. *richardsonii* 381,
 382
 venulosum 381, 382
 var. *lunellii* 382
 var. *turneri* 382
 var. *venulosum* 382
 Thelesperma 704, 788
 marginatum 788
 Thellungiella 393, 417
 salsuginea 417
 Thermopsis 463, 497
 rhombifolia 497, **500**
 Thlaspi 389, 417
 arvense **390**, 417
 Thuja 61
 occidentalis 61
 plicata 61
 Tiarella 421, 435
 trifoliata 435
 unifoliata 435
 Tilia 520
 americana 520
 Tiliaceae 29, 520
Tium drummondii 470
 racemosum 474
 Tofieldia 257, 266
 glutinosa 266
 pusilla 266
 Townsendia 707, 788
 exscapa 788
 hookeri 788
 parryi 788, 790, **791**
 sericea 788
Toxicodendron rydbergii 515
Toxicoscordion gramineum 268
 Tradescantia 246
 occidentalis 246

- Tragopogon **684**, 689, 702
 dubius 703
 porrifolius 703
 pratensis 703
 Trientalis 580, 586
 americana 587
 borealis **582**, 586, 587
 europaea 587
 Trifolium 462, 497
 aureum 501
 campestre 501
 hybridum 501
 pratense 501
 repens 497, 501
 Triglochin 68
 maritima 68, **69**
 palustris 68
Trigonella americana 482
 Trillium 257, 267
 cernuum var. *macranthum* 267
 ovatum 267
 Trisetum 81, 179
 canescens 179
 cernuum 179
 spicatum 179, 181
 wolfii 179, 181
 Triticeae 804
 Triticum 80, 181
 aestivum 181
 Trollius 363, 382
 laxus 382
 Typha 61
 angustifolia 61
 latifolia 61, 63, **64**
 Typhaceae 22, 61
- Ulmaceae 25, 303
 Ulmus 303
 americana 303
 Umbelliferae 27, 545
Unamia alba 785
 Urtica 304
 dioica var. *procera* 304, 305, **306**
 urens 304, 305
 Urticaceae 24, 304
 Utricularia 659, 660
 cornuta 660
 intermedia 660
 macrorhiza 661
 minor 660
 vulgaris var. *americana* 660, 661, **662**
 Uvularia 257, 267
 sessilifolia 267
- Vaccaria vulgaris* 355
 Vacciniaceae 29, 577
 Vaccinium 577, 578
 angustifolium 578, 579
 var. *angustifolium* 579
 var. *myrtilloides* 579
 caespitosum 578, 579
 membranaceum 578, 579
 myrtilloides 579
 myrtilus 578, 579
 oxycoccus var. *microphyllum* 578
 var. *oxycoccus* 578
 scoparium 578, 579
 uliginosum 578, 579
 vitis-idaea 578
 var. *minus* 580
 Valeriana 681
 dioica var. *sylvatica* 681, **682**
 septentrionalis 681
 sitchensis 681
 var. *scouleri* 681
 Valerianaceae 31, 681
 Veratrum 257, 267
 eschscholtzii 267
 Verbascum 638, 656
 nigrum 656
 phlomoides 656
 thapsus 656
 Verbena 619
 bracteata 619
 bracteosa 619
 hastata 619
 urticifolia 619
 Verbenaceae 30, 619
 Vernonia 705, 790
 fasciculata var. *corymbosa* 790
 Veronica 638, 656
 agrestis 657
 alpina var. *unalaschensis* 657
 americana 657, 658
 catenata 658
 chamaedrys 657, 658
 comosa var. *glaberrima* 657, 658
 connata ssp. *glaberrima* 658
 longifolia 657, 658
 peregrina var. *xalapensis* 657, 658
 persica 657, 658
 polita 657, 658
 scutellata 657, 658
 serpyllifolia var. *humifusa* 657, 659
 teucrium 656, 659
 xalapensis 658
 Veronicastrum 638, 659
 virginicum 659

- Vesicariae 188, 189, 191, 200
 Viburnum 670, 675
 affine var. *hypomalacum* 675
 edule 675, **676**
 eradiatum 675
 lentago 675
 opulus var. *americanum* 675
 rafinesquianum 675
 trilobum 675
 Vicia 462, 479, 501
 americana 502, **503**
 var. *minor* 502
 var. *truncata* 502
 americana var. *angustifolia* 502
 cracca 502
 oregana 502
 sativa ssp. *nigra* 502
 villosus 502
 Viola 526
 adunca 527, **528**
 arvensis 526
 blanda 527
 cucullata 527, **528**
 glabella 527
 nuttallii 527, **528**
 orbiculata 527
 palustris 527
 pedatifida 527, 530
 pubescens 527, 530
 renifolia 527, 530
 rugulosa 527, **528**, 530
 selkirkii 527, 530
 tricolor 527, 530
 Violaceae 27, 526
 Virescentes 189
 Vitaceae 28, 519
 Vitis 519
 riparia 519
 Vitis-idaea punctata 580
 Vulpinae 187, 192

Wahlbergella drummondii 354
 Woodsia 36, 46
 alpina 46
 glabella 46
 ilvensis 46
 oregana 47
 var. *cathcartiana* 47
 var. *oregana* 47
 scopolina 47

 Xanthium 685, 686
 commune 686
 echinatum 686
 glanduliferum 686
 italicum 686
 macounii 686
 strumarium 686, **688**
Xanthoxalis corniculata 507
 stricta 507
 Xerophyllum 257, 267
 tenax 267
Xylophacos missouriensis 472
 purshii 474
Xylosteon oblongifolium 673
 tataricum 673

 Yucca 257, 267
 glauca 267

 Zannichellia 66, 67
 palustris 67
 Zizania 83, 181
 aquatica 181
 Zizanieae 78, 83
 Zizia 548, 563
 aptera 563
 aurea 563
 cordata 563
 Zosteraceae 21, 65
 Zygodenus 257, 268
 elegans 268
 gramineus 268, **269**

